

This is a repository copy of *Economic and socio-cultural consequences of changing political rule on human and faunal diets in medieval Valencia (c. fifth–fifteenth century AD) as evidenced by stable isotopes*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/143483/>

Version: Accepted Version

Article:

Alexander, Michelle Marie orcid.org/0000-0001-8000-3639, Gutiérrez, Alejandra, Millard, Andrew et al. (2 more authors) (2019) Economic and socio-cultural consequences of changing political rule on human and faunal diets in medieval Valencia (c. fifth–fifteenth century AD) as evidenced by stable isotopes. *Archaeological and Anthropological Sciences*. pp. 3875-3893. ISSN 1866-9557

<https://doi.org/10.1007/s12520-019-00810-x>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Economic and socio-cultural consequences of changing political rule on human and faunal diets in medieval Valencia (c. 5th-15th century AD) as evidenced by stable isotopes

Authors

Michelle M. Alexander^{1*}, Alejandra Gutiérrez², Andrew R. Millard², Michael P. Richards³, Christopher M. Gerrard²

Affiliation

1 BioArCh, Department of Archaeology, University of York, York, UK

2 Department of Archaeology, Durham University, Durham, UK

3 Department of Archaeology, Simon Fraser University, Burnaby, BC, Canada

*Correspondence to:

Michelle M Alexander, University of York, BioArCh, Environment Building 2nd Floor, Wentworth Way, Heslington, York, YO10 5DD, Email: michelle.alexander@york.ac.uk

ORCID

Keywords (3 to 5): Muslim, Christian, Visigothic, Spain, collagen, diet

Acknowledgements

The authors gratefully acknowledge Pepa Pascual (*Servicio de Investigación Arqueológica Municipal de Valencia*) and Javier Martí (*Museo de Historia de Valencia*) for all their help and enabling access to the skeletal collections in Valencia. Thanks to Olaf Nehlich (MPI-EVA) for consultation and collagen preparation and analysis. Thanks go to Helen Goodchild, (University of York), for aid in ARCGIS. We are also grateful to the reviewers for their constructive comments on the manuscript. This research was funded by an AHRC PhD studentship awarded to Alexander (Ref: 2006/125150/University of Durham) with contributions from a Durham Doctoral Fellowship and a Society for Medieval Archaeology

ABSTRACT

This paper explores the impact of changing religious political rule on subsistence within a single city through time using stable isotope analysis of human and animal bone collagen. The diet and economy of the medieval city of Valencia (Spain) is examined over a 1,000 year period during successive periods of Visigothic, Muslim and Christian rule. Bulk stable isotope analysis of carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) was carried out on 67 humans and 47 animals sampled from several archaeological sites dating between the 5th and 15th centuries AD. Terrestrial C_3 resources dominated the diet in all periods. However, an increase in consumption of C_4 plants (e.g. millet, sorghum) and/or marine resources is detected amongst individuals dating to the Islamic period. Differences in the isotopic values of humans and animals between the three periods indicates a significant dietary diversity during the Islamic phases (11th to 13th century), compared with the earlier Visigothic phase (5th-7th century) and the later Christian phase (14th-15th century). Observed diachronic changes in isotopic results provide evidence for a shift in diet and subsistence, particularly during the Muslim and later Christian periods. This is linked with change in population and economic focus. Dietary diversity amongst Muslim individuals is hypothesised to indicate the polyculture that was reflected in varied Islamic agricultural practices and the presence of potential migrants from the rest of the Islamic world.

The influence of socio-political and economic change on diet over the long term is just beginning to be directly investigated using isotope analysis. Investigations have centred on a single location, such as the city of York (Müldner and Richard 2007), or multiple sites across a large area, e.g. in Croatia (Lightfoot et al., 2012) and Mallorca (Fuller et al., 2010). Such studies have highlighted human dietary responses to these shifts. Recent work has, however, emphasized resilience and continuity in dietary practice despite dramatic shifts in socio-political circumstances, e.g. Christianisation in Poland (Reitsema et al., 2017). The present research explores the dietary impact of the major religious, socio-political, economic and population transformations that occurred over 1,000 years in the medieval city of Valencia to the level of an individual. Both human and animal remains were analysed to examine aspects of day-to-day subsistence, agriculture and animal husbandry under Visigothic, Islamic and subsequent Christian control in eastern Spain.

During the medieval period, the Iberian Peninsula harboured a multi-faith society that was subject to dramatic socio-cultural, economic and demographic shifts associated with changes in religious political rule. The long and eventful settlement history of Valencia, a major city in Spain during the medieval period, provides the ideal opportunity to investigate the diet of its urban population over the long term in the context of these changes. In this article we explore a period of roughly 1,000 years from the 5th to the 15th centuries AD through the analysis of carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) isotopes in collagen from human and animal bone: these samples are taken from multiple sites across the medieval city. The application of isotopic methodologies to medieval populations is growing in Spain, with many concentrating on Islamic populations (e.g. Salazar-García et al., 2014, 2016; Pickard et al., 2017; Guede et al., 2017), status (Jiménez-Brobeil et al., 2016), the Christian North (e.g. Sirignano et al., 2014) or the interplay of faith and diet between contemporaneous populations of Christians and Muslims under later Christian rule (Alexander et al., 2015). Although others have explored diachronic changes in diet in Spain, for example following the Islamic conquest in the Balearic Islands (Fuller et al. 2010), this is the first time any bioarchaeological analysis of historic diet has investigated such an extended chronology for any major Spanish city and in particular the transition from Islamic to Christian rule.

[Figure 1 here]

Development of the Medieval City

The city of Valencia lies on the banks of the River Turia on Spain's Mediterranean coastline (Fig. 1). Founded in 138 BC, the Roman city of *Valentia* declined in importance during the Visigothic period (5th-8th century) but maintained its strategic value as the see of a Bishopric and remained the administrative nucleus of a wide territory (Ribera, 2006; 2008; Jiménez Salvador, 2009). In AD 711 the city was taken by Muslims and became fully established with an Islamic ruler in AD 800. After Rodrigo Díaz de Vivar ('El Cid') briefly interrupted Islamic rule, Valencia was re-taken by the Almoravids in AD 1109 and remained under Islamic control (Kennedy, 1996). In 1238 Valencia fell to the Aragonese King James I and was annexed to the Crown of Aragón. The city now became capital of the Kingdom of Valencia between Murcia and the Tortosa region of the Ebro valley. Throughout the 13th and 14th centuries, the city was populated by Christian immigrants from the northern territories of

Aragón (Guinot 2009, p. 176). Muslims who stayed in the area moved to a *morería* or Muslim quarter established outside the walls of Valencia (Baydal and Esquilache, 2014).

Economy and Agriculture

Throughout the medieval period, Valencia was an important port and commercial centre. By the 12th century, Genoese merchants used the city to export goods from al-Andalus (the area of the Iberian Peninsula under Islamic control) to Christian ports throughout the Mediterranean (Constable, 1994, p. 20). Agriculture, however, was at the heart of the region's economy and the changes to the ethnic and religious make-up of the population and shifting political power outlined above would have had a dramatic effect on agriculture, economy and diet. Agrarian transformation occurred across the Iberian Peninsula under Islamic rule. This 'agricultural revolution' (first coined as such by Watson in 1974, but see Decker, 2009 and Squatriti, 2014 for criticism), has been attributed to societal change and formation of a tributary economy, innovations in agricultural and hydraulic technology and the introduction of new crops, such as pomegranate, rice, cotton, sorghum, and citrus fruits (Ruas et al., 2017). There is little evidence for a revolution in the archaeobotanical record so far, but the study of medieval plant remains has lagged behind that of the prehistoric periods in Spain (Peña-Chocarro et al., 2017).

The extensive irrigated area that surrounds Valencia today (known as the *Horta*) has its origins in the Islamic conquest of the 8th century (Guinot and Esquilache 2017). The development of intensive irrigation enabled the cultivation of a variety of crops at any given time and agricultural diversity in particular was a defining feature of the Islamic agrarian economy (Glick 1995). The introduction of sorghum (a C₄ crop) in particular might be visible through isotopic data if it not yet archaeologically. The later Christian conquest saw the establishment of a feudal society and the reorganisation of cultivated land (Torró 2012a; b). Irrigation was maintained, however new systems emerged, and the *Horta* was extended, while vines and cereal crops, particularly wheat, received new emphasis. There was also an increased emphasis on stock-keeping and this period in particular saw the rise of long-range transhumance (Guinot and Esquilache 2017; Guinot 2009; Torró 2009a).

Zooarchaeological data from sites across the Iberian Peninsula indicate that sheep/goat were the most common species at both Islamic and Christian period sites. Variation in kill-off patterns suggests that younger meat was preferred in the Islamic period, but these animals were increasingly used for secondary resources in the Christian period (e.g. wool). Conversely, cattle seem to have undergone the reverse trend (Morales et al., 2011). Pigs have been reported at Islamic period sites, although in low frequencies compared to Christian sites (Morales et al. 2011; Grau-Sologestoa, 2017). These generalisations however belie the diversity of animal management practiced in the medieval period; there is no clear 'Muslim' signal for a zooarchaeological assemblage, although an urban/rural divide does appear to exist in the data (Morales et al., 2011; Grau-Sologestoa, 2017; Estaca-Gómez et al., 2018). Large scale shifts in agriculture and stock keeping have the potential to be read from the isotopic data, providing direct evidence for discernible shifts in diet and economy that would be applicable at the individual and population level.

Medieval diet

A brief historical and archaeological overview of medieval diet in Spain is provided by

Alexander et al. (2015; but see also, for example, de Castro, 1993; García Sánchez, 1996; 2002; Adamson, 2004). Wheat, a C₃ crop, was the most desirable grain, although mixed-grain bread would have been the most commonly eaten; C₄ crops such as millet (broomcorn and/or foxtail) or, from the Islamic period, sorghum, were considered secondary. These crops tended to be relied upon particularly in times of scarcity or by those of lower social status (de Castro, 1993 p. 286; García Sánchez, 2002, p. 279). A wide range of vegetables and fruit were also available from markets and kitchen gardens, but these contribute little in terms of the isotopic evidence considered here. Meat and dairy products made up a smaller, but important, part of the diet according to social status, geographical location/environment and faith. Lamb and mutton were most commonly consumed meats throughout the medieval period, and beef, chicken, pork (in the case of Christians) and game could also be found on the table (Waines, 1992; de Castro, 1993; García Sánchez, 1996).

Freshwater and/or marine resources provided a valuable supplement to terrestrial resources and fish was often dried, salted, smoked or pickled (Gallart et al., 2005). Fish also served a liturgical requirement for Christians, who were obliged to avoid meat when fasting for a third of the year (Tomás, 2009, p. 465). It is possible that a discernible increase in fish consumption might be detected in the later Christian population in line with religious doctrine, although alternatives, such as eggs, could have also been used instead (Gerrard, 2003, p. 301). Consumption of terrestrial animal protein, marine resources and C₃ and C₄ plants, particularly if the latter were fed to animals whose products were consumed by humans, will all have an impact on the carbon and nitrogen isotope signature of consumer bone collagen.

Stable Isotope Analysis

In-depth reviews of stable isotopes and dietary reconstruction of archaeological populations are available elsewhere (e.g. Katzenberg, 2000; Lee Thorpe, 2008; Schwarcz and Schoeninger, 2011). In general terms, the composition of carbon and nitrogen isotopes in bone collagen is representative of a long-term average of the main protein sources of the diet in the decades before death, where protein intake is adequate (Hedges, 2004; Hedges et al., 2007). The use of carbon isotopes to distinguish between the consumption of common, temperate C₃ crops and arid-adapted C₄ plants (O'Leary, 1981) is notable in the context of medieval Spain where C₄ crops such as millet, sorghum and sugarcane were present. $\delta^{13}\text{C}$ values also differ between marine and terrestrial environments (Schoeninger and DeNiro, 1984).

Nitrogen isotope ratios depend on the position of an organism within the food chain. It is widely accepted that $\delta^{15}\text{N}$ values increase 3-5‰ between trophic levels (Bocherens and Drucker, 2003), although spacing of up to 6‰ has been suggested for humans (O'Connell, et al., 2012). There is also a slight increase in $\delta^{13}\text{C}$ values with trophic level of 0-2‰ (Bocherens and Drucker, 2003). Values for $\delta^{15}\text{N}$ tend to be higher in aquatic ecosystems due to the longer length of food chains (Schoeninger et al., 1983). Therefore, although $\delta^{13}\text{C}$ values for marine and C₄ consumers can overlap, $\delta^{15}\text{N}$ values are usually able to distinguish predominantly marine from terrestrial C₄ feeders. The Mediterranean is particularly problematic in this regard because $\delta^{15}\text{N}$ values for marine and terrestrial resources are often

similar (Craig et al., 2013). As a result, the identification of consumption of marine foods, as opposed to indirect consumption of C₄ crops via C₄-fed terrestrial animal protein (with corresponding higher $\delta^{15}\text{N}$ values), is not always straightforward using bulk isotope analysis due to the problem of equifinality when analysing consumer isotope values (Alexander et al., 2015).

MATERIALS

Human populations

Human bone from 67 individuals was sampled from five sites representing six phases within and outside the medieval city walls (Table 1, Fig. 2); 10 individuals from the Visigothic period, 38 from the Islamic period and 19 from the later Christian period (see Table 1 for dates). Osteological analysis was undertaken by Alexander following established methods to ascertain adult age (ca. 18 years of age and above) based on the development and fusion of bones (Scheuer and Black 2000) and dental eruption and wear (Ubelaker 1989; Brothwell 1981). Sex was determined using standard osteological techniques outlined in Mays and Cox (2000) assessing the morphology of the pelvis and skull and the measurements of bones to support morphological examination.

Individuals were identified as Christian or Muslim from their distinctive burial rites. In general, Christians were buried supine, orientated E-W with their heads to the west, while the Muslims were buried on their right-side orientated SW-NE with their face towards Mecca (Lozoya et al., 2015). Unfortunately, no *mozarab* or later *mudéjar* (Muslims under Christian rule) burials were available for analysis. It should also be noted that a Jewish community was present in Valencia throughout the periods examined here, but as no Jewish burials have been analysed, there will be no discussion of this population in this paper.

[Figure 2 here]

[Table 1 here]

Both Visigothic and selected Muslim burials were sampled from excavations in the Plaza de la Almoina, the political and religious nucleus of Valencia from the Roman period onwards. Given that this was the area of the Cathedral and Bishop's residence in the 6th-7th centuries, the Visigothic burials sampled here have been interpreted as members of the social elite (Martin and Ribera, 2006). Burials were sampled from two phases of interments representing both individual and multiple graves (Table 1). The main mosque, originally located here next to the Islamic palace (*alcázar*), was later converted into the present cathedral. The palace has an adjoining Islamic cemetery (known now as La Rauda) that was situated within the city walls, unlike other cemeteries of the period (Pascual, 1989, p. 406; Ribera, 2006, p. 82). Historical sources identify this cemetery as the high-status burial ground for the Taifa kingdom (Pascual and Serrano, 1996). Samples taken here allow any associations between diet and status within the Islamic population of Valencia to be explored.

Further 11th-13th century Muslim burials derive from the large cemetery of Bab al-

Hanax, and a smaller cemetery uncovered in Carrer Sagunt. The latter is likely to be related to a small extramural suburb and allows comparison between populations of different quarters of the city. Three Muslim burials were also sampled from a group of interments in Carrer Pintor Sorolla. These burials lay within the Islamic city walls and given that this is a practice that was prohibited under normal circumstances, it seems likely that these particular people were interred during the Christian siege of 1238.

Christian burials were sampled from two early parishes. San Andrés was one of the first parishes created after the Christians took control of the city and was one of the wealthiest during the 13th century (Burns, 1967, p. 97; García and Ruiz, 1997 p. 5). The burial ground excavated near Carrer Sant Vicent was part of a complex of a church, monastery and hospital that was established in the 13th century by royal patronage near the shrine dedicated to St Vincent. The monastic group that attended and administered the complex was chosen by the king and underwent several changes including monks from the monastery of San Victorián, Mercedarians, followed by Cistercians (Burns, 1967, pp. 283 and 294). The monastery served as a retirement home for civil servants, families or old soldiers, while the hospital included a hostel for the poor and infirmaries for the sick (Burns, 1967 p. 285-286).

Faunal remains

The animal remains analysed in this study come from well-stratified refuse deposits in the Plaza de la Almoina (n=42). Samples were taken from a selection of contexts dating to the Visigothic (6th-8th century), Islamic (11th-13th century) and Christian (13th-16th century) periods excavated between 1997 and 2001. Previously published marine fish $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values from other medieval sites in eastern Spain, Albarracín (Teruel) and Gandía (Valencia) (Alexander et al. 2015) were also included.

METHODS

All human samples derived from adult (>18yrs) individuals (based on osteological analysis). Bone (200-500 mg) was preferentially sampled from ribs or cortical bones when ribs were unavailable. Cortical bone was chosen from animal specimens (300-500 mg), ensuring that only one sample was taken from one individual in each context. Collagen was isolated from bone following a modified Longin (1971) method (Brown et al., 1988) as described in Alexander et al. (2015).

Collagen (0.5mg) was analysed in duplicate EA/IRMS with a Delta XP (Thermo-Finnigan®, Bremen, Germany) mass spectrometer interfaced with a Costech ESC 4010 elemental analyser at the Alaska Stable Isotope facility (University of Alaska Fairbanks's Water & Environmental Research Centre; WERC) for humans or coupled with a Flash EA 2112 at the Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology (MPI-EVA, Leipzig, Germany) for animals. Stable isotope ratios were reported as δ values expressed as per mille (‰) deviation from the international standards VPDB (carbon) and AIR (nitrogen) following the equation [$\delta = (R_{\text{sample}} - R_{\text{standard}}) / R_{\text{standard}}$] (Coplen, 2011). For both laboratories, analytical error (calculated from repeated measurements of each sample and an in-house laboratory gelatine control from multiple extracts) was <0.2 ‰ (1 σ) for both $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$. Collagen samples were exchanged between laboratories to ensure instrument consistency and as outlined in Alexander *et al.* (2015), a slight difference (0.6‰ \pm 0.2‰, 1 σ) was detected between $\delta^{13}\text{C}$ values at each laboratory and though is at

the threshold of the Minimum Meaningful Difference (MMD, 0.6‰) between laboratories suggested by Pestle et al., (2014), it is not at the magnitude of dietary interpretation (typically 1‰; Sealy et al., 2014).

Statistical tests were performed using PAleontological STatistics software (PAST 3.08; Hammer et al., 2001). The non-parametric Mann-Whitney U and Kolmogorov-Smirnov tests were employed, with Kruskal-Wallis tests used to assess overall variability between the three groups.

RESULTS AND DATA INTERPRETATION

The results are presented here alongside an initial interpretation of human diet and animal management within and between periods. The later discussion and synthesis is focused on a diachronic comparison and the wider implications of the results in terms of economy and diet under changing socio-political rule.

Sample preservation

Sufficient collagen (>1% yield) was found to be preserved in all but two samples (11025 and VCP2, Table 2 and 3 respectively), however all samples met published collagen quality criteria (DeNiro, 1985; van Klinken, 1999) and, as ultrafiltration can reduce collagen yields (by up to 50%; Jørkov et al., 2007), compared to unfiltered samples, the data from the two samples with low yield were retained.

All periods

Human values for all periods (n=67) range from -19.8‰ to -14.2‰ (range of 5.6‰) in $\delta^{13}\text{C}$ with a mean of $-18.0\text{‰} \pm 1.1\text{‰}$ (1σ) and from 8.0‰ to 17.1‰ (range of 9.1‰) in $\delta^{15}\text{N}$ with a mean of $11.3\text{‰} \pm 1.5\text{‰}$ (1σ). Contemporaneous animals (n=42) range from -22.3‰ to -10‰ (range of 12.3‰) in $\delta^{13}\text{C}$ with a mean of $-19.1\text{‰} \pm 1.9\text{‰}$ (1σ) and from 2.5‰ to 11.5‰ (range of 9‰) in $\delta^{15}\text{N}$ with a mean of $6.6\text{‰} \pm 2.3\text{‰}$ (1σ). The human and animal stable isotope data for all periods are presented in Table 2 and 3 respectively and plotted together in Figure 3; summary statistics for humans and animals in each period are listed in Table 4 and 5 respectively. Within the Christian period dataset there is an extreme human outlier, VSVC3622, with a $\delta^{15}\text{N}$ value (17.1‰) that is more than three standard deviations (3.3) from the population mean. VSVC3622 was therefore omitted from any statistical comparisons as this individual is considered to be unrepresentative of the local population.

[Table 2 here]

[Table 3 here]

[Table 4 here]

[Table 5 here]

[Figure 3 here]

In general, herbivores (cattle, sheep/goat) in all periods tend to have lower $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values than omnivores (chicken, dog, pig) (Table 5). This breadth in values for all species is indicative of a variety of geographic and environmental origins and long-term feeding strategies and must reflect the wide catchment of resources the city was drawing upon, including the possibility of the presence of non-local animals imported from further afield. No one region supplied all the city's needs.

Low $\delta^{13}\text{C}$ values for the majority of domestic animals indicate that C_3 plants were the main resource throughout the medieval period and the diets of sheep/goat, the most popular source of meat throughout this time period are consistently C_3 -based. Higher $\delta^{13}\text{C}$ values indicative of C_4 consumption, however, are seen among cattle, chickens and pigs and a dog (Table 5), indicating some presence of C_4 resources in animal diets. In the Islamic and Christian periods, the highest $\delta^{15}\text{N}$ values for herbivores overlap with humans, a relatively uncommon observation amongst published medieval faunal data. Given that an effort was made to sample animals older than weaning age, an enrichment of the plant baseline is the most probable explanation here. There is a complex combination of environmental (aridity, salinity) and anthropogenic influences (e.g. manuring) on $\delta^{15}\text{N}$ in food webs (Heaton, 1987; Bogaard et al., 2007). Manuring was widely practiced and, even in the recent past, *corrales* or *bordas* (dedicated enclosures for the stabling and foddering of animals) were important for the production of manure to fertilise meadows and hillside terraces (Christie et al., 2004, p. 109; Montserrat and Fillat, 1990, p. 58). $\delta^{15}\text{N}$ increases may also be attributed to herbivores grazing close to urban settlements in the vicinity of accumulations of human and animal nitrogenous waste (Vassberg, 1974; Butzer, 1988; Hedges et al., 2005; Triantaphyllou et al., 2008). Grazing of coastal areas and salt marshes is an additional factor to be considered (Britton et al., 2008). To the south of Valencia, the Albufera is an extensive wetland with salt marshes around a lagoon (Fig. 1) used as pasture for sheep and cattle (Mombloch, 2003; Torr , 2012a; b). Aridity may also play a part; some parts of the Valencian region today are dry and semi-arid bioclimatic contexts, with a mean annual rainfall ranging from 500mm/yr to less than 300mm/yr (Hidalgo et al., 2003), particularly areas to the south. Environmental evidence indicates that the Mediterranean region of the Iberian Peninsula experienced particularly dry conditions during the Medieval Warm Period (900-1300 AD) (Moreno et al., 2012).

Values for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ for humans indicate that a C_3 terrestrial diet predominated in all periods under study. The Islamic data in particular, however, stand apart in demonstrating a variable pattern of $\delta^{13}\text{C}$ values, and a tendency towards elevated $\delta^{15}\text{N}$ values in comparison to the other periods. A Kruskal-Wallis test comparing sites medians by period for humans indicates there is no difference for $\delta^{13}\text{C}$, (H(2):3.373, p=0.184) or $\delta^{15}\text{N}$ (H(2):5.922, p=0.051). Kolmogorov-Smirnov tests for the distribution of isotopic values between sites, however, indicate that there is a significant difference in the distributions of values between $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values for the Islamic period compared to both the Visigothic and Christian periods (Table S1). Examination of patterning in human data between periods needs to take into account potential shifts in the animal baseline, indicative of the environment and husbandry in each period. To that end, herbivores and humans from all periods are plotted together for comparison in Figure 4. Offsets between humans and herbivores remain similar through all three periods (Table 6) indicating that the shift to C_4 plants in particular is broadly reflected in both animals and humans. The dietary shift

exhibited in the Islamic period is therefore indicative of an economic shift towards the use of C₄ crops in both in human and animal diets rather than solely a change in human food choice. There was no difference between the sexes identified in the isotopic data from any period.

[Figure 4 here]

[Table 6 here]

Visigothic Period

All animals from this period possess a C₃-based diet bar one dog (VVD1, Fig 3., Tables 3 and 5). This dog possesses a $\delta^{13}\text{C}$ value of -16.4‰, which, combined with a relatively low $\delta^{15}\text{N}$ value (9‰) most probably reflects an input of C₄-based terrestrial protein in the diet. This is puzzling because dogs are often used as a proxy for human diet (see Guiry 2012), but none of the humans sampled from this period have isotopic values indicative of C₄ consumption. Only a small number of humans have been analysed from this period, however, and given that these individuals have been identified as the social elite, they may not reflect the popular diet of the time. Dogs may also have not been fed the 'best' foods from the table. Analysis of further humans and animals from this period are needed to investigate this more thoroughly.

While the results for humans in the Visigothic period are more tightly clustered than later periods (Fig. 3), the two phases of burial fall into two distinct groups (Fig. 6). Burials of the later 6th-7th century are slightly enriched in ¹³C and ¹⁵N and this is statistically significant for $\delta^{13}\text{C}$ values (Mann Whitney U=2.5, p=0.032), although not for $\delta^{15}\text{N}$ values (Mann Whitney U=22, p=0.056). While the sample is small, the significant difference in $\delta^{13}\text{C}$ values is of interest when related to burial practice among these individuals. The earlier burials on site follow the preceding Hispano-Roman burial tradition with single interments, an absence of grave goods and tiles covering the grave. The later burials, on the other hand, tend to be multiple interments with artefacts (amber pendants, coins and silver objects) and walled tombs, possibly representing family mausolea (Ribera, 2008). These changes also correlate with morphological differences observed between the skeletons buried in these two phases: individuals of the later phase are of higher stature and greater robusticity (Martin and Ribera, 2006). These findings may reflect an increasing Germanic influence in this quarter of the city and it is recorded, for example, that there was an Arian bishop in the city in AD 589 (Martin and Ribera, 2006; Ribera, 2008). The isotopic patterning seen here could therefore be a reflection of shifting dietary practice in line with a change in population, although the limited sample size prevents further interpretation.

[Figure 6 here]

More broadly, the Valencian Visigothic data are compared to 6th-7th century 'Late Roman' individuals from Tossal de les Basses in Alicante (Salazar-García et al., 2016) and 4th-7th century Late Antique/Early Byzantine individuals from Ibiza (Fuller et al. 2010) in Figure 5. Of

these two, the values for Valencia are remarkably similar to Tossal de les Basses, indicating that diets in these two places were comparable, with C₃-based diets for humans at both sites.

Islamic Period

Among the animal data from the Islamic period, both cattle and chickens stand out for their wide range in $\delta^{13}\text{C}$ values (4.3‰ and 7.9‰ respectively, Table 5) indicative of diverse feeding strategies in both species. Some individuals possess low $\delta^{13}\text{C}$ values indicative of C₃ consumption whereas others have values reflecting a significant proportion of C₄ plants in their diets, including one chicken (VMGa2) with an extreme $\delta^{13}\text{C}$ value of -10‰. The same C₃/C₄ patterning among cattle was found in the rural Muslim settlement of Gandía in a later period (Alexander et al., 2015). In line with the evidence presented here, species of millet are mentioned among the grains used to fatten both cattle and chickens by the agronomist Ibn al-Awwam in his 12th century *Treatise on Agriculture* (Ibn al-Awwam, 1802). In north-west Spain, ethnographic evidence demonstrates that millet is still traditionally used for foddering animals, particularly stabled cattle and chickens (Moreno-Larrazabal et al., 2015).

Human data from Islamic Valencia exhibit a strikingly wide range in both $\delta^{13}\text{C}$ (5.5‰) and $\delta^{15}\text{N}$ (7.0‰, Table 4). When the isotopic data is plotted (Fig. 3) a core group of individuals show $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values (-19.0‰ to -18.0‰ and 10‰ to 12‰ respectively) indicative of a predominantly terrestrial, C₃ based diet. Isotopic values for the remainder of the sampled individuals suggest the consumption of marine fish and/or C₄ protein (directly and/or indirectly). As previously noted, it is difficult to disentangle the consumption of marine fish rather than C₄ resources in the Mediterranean environment, although trends are discernible that could indicate the consumption of one over the other. One group of individuals possess increasingly higher $\delta^{13}\text{C}$ values >-18‰ but relatively constant $\delta^{15}\text{N}$ values of around 12‰, indicating input of C₄ resources, whereas other individuals exhibit considerable enrichment in ¹⁵N ($\delta^{15}\text{N}$ values of up to ~17‰) coupled with corresponding higher $\delta^{13}\text{C}$ values (-18‰ to -16‰) suggesting marine protein consumption.

This variability is also visible by site (Fig. 7), suggesting that a range of dietary resources were available across the city and between communities. Only the individuals from the Bab al-Hanax cemetery plot in a relatively cohesive group with some individuals tailing out towards higher $\delta^{13}\text{C}$ values. For the most part, this population may represent locals of middling status who subsisted mainly on C₃ terrestrial foods.

[Figure 7 here]

Burials from Carrer Sagunt, associated with the artisan suburb of la Alcudia outside the city, are especially notable for their extremely variable $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values; these individuals have some of the highest $\delta^{15}\text{N}$ values exhibited by the population. The three individuals from Carrer Pintor Sorolla, believed to date to the late 13th century siege, all show higher $\delta^{13}\text{C}$ values indicative of C₄ protein consumption. High-status burials from the Rauda are again highly variable, with some plotting with the majority in Carrer de Dalt, whereas others possess higher $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values. The latter group may have consumed a

greater proportion of marine resources (>50% for individual VRM1356 following Richards et al., 2006), alternatively these individuals may have originated elsewhere.

The presence of non-locals may explain the variation in the population overall. As a successful port, Valencia was at the centre of political, familial, trade and travel links between al-Andalus and the Maghreb during the 11th to 13th centuries (Glick 2005, p. 357). The people buried at the Rauda are believed to have been those related either politically or by kinship to the Almoravid and Almohad rulers of Valencia and it is very probable that some of these elites may have originated from areas such as the Almohad capital of Marrakesh on the edge of the Sahara in North Africa (Burns, 1973; Boone and Benco, 1999; Torr , 2009b). Written records indicate that both members of the nobility and inhabitants of other cities in the Valencian territory were buried at Bab al-Hanax (Pascual and Serrano, 1996, p. 232). There were continuous waves of Berber immigrants from North Africa into al-Andalus throughout the Islamic period and their frequency increased from the 10th century onwards (Guichard 2000). Slaves would have also been a major presence in Valencia during this period. The presence of, and access to, black slaves in al-Andalus purportedly goes back to the 11th century and is associated with Islamic penetration of Sub-Saharan areas and the arrival of the Almoravid and Almohad dynasties from North Africa (Constable, 1996, p. 269). It may be that some of the individuals identified with relatively high $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values originated from the Maghreb. Substantial enrichment in ^{15}N of consumer bone collagen has been reported from arid areas such as Dakheleh Oasis in the Sahara ($\delta^{15}\text{N}$ values from around 15‰ to 18‰, Schwarcz et al., 1999; Dupras and Schwarcz, 2001) and $\delta^{13}\text{C}$ values, similar to those exhibited here, have been reported in human collagen samples from archaeological populations from North Africa, for example from Sudanese Nubia (Turner et al., 2007).

In the wider context, this variability in isotopic values, particularly for $\delta^{13}\text{C}$, is also a feature of published isotopic values for Islamic populations in other parts of Spain (Fig. 5). This is often due to the presence of specific outliers increasing the standard deviation of these populations, particularly in those cases with smaller samples, e.g. La Torrecilla, Granada (Jim nez-Brobeil et al., 2016). In the 10th-13th century Islamic population from Can Fonoll, Ibiza, there was a wide range of $\delta^{13}\text{C}$ values, but only four individuals had values above -16‰ indicative of significant input of C_4 resources and these were therefore interpreted as migrants (Pickard et al., 2017). In Valencia, by contrast, many individuals exhibit increased $\delta^{13}\text{C}$ values and, although the presence of migrants is highly likely, individual choice or restriction in diet cannot be ruled out.

Local socio-economic factors must therefore also be considered, where low status individuals and/or slaves may have relied on C_4 grain in particular. On the other hand, individuals with higher $\delta^{13}\text{C}$ values may be local migrants from rural areas where C_4 grain was utilized more widely. Isotopic evidence points to a rural-urban divide where later medieval Islamic communities from rural contexts exhibit higher $\delta^{13}\text{C}$ values indicative of C_4 signatures (Fig. 5) e.g. La Torrecilla (Jim nez-Brobeil et al., 2016), Benipeixcar (Alexander et al., 2015) and El Raval (Salazar-Garc a et al., 2014).

Christian Period

Animal data for the Christian period indicates a return to a predominantly C₃-based diet, although no chickens and only two cows were sampled from contexts dating to this period (Fig. 3, Table 5). Perhaps more samples of these species in particular would reveal further diversity. Data for pigs are most notable among the animal data for this period, demonstrating more enrichment in ¹⁵N in comparison to earlier periods (minimum δ¹⁵N value 8‰, maximum δ¹⁵N value 9‰, n=4, Table 5). This is perhaps indicative of more intensive husbandry and inclusion of animal protein in the diet of pigs that may signal a renewed importance for animal for the Christian populace. One pig (VCP4, Table 3) is also enriched in ¹³C, with a δ¹³C value of -17.0‰. This suggests the input of a source of ¹³C-enriched carbon in the diet, probably from C₄-derived protein rather than marine sources as the pigs have lower δ¹⁵N values than the majority of humans (Fig. 3).

Human individuals from the later medieval Christian Valencia (Table 4) display a relatively narrow range (2.5‰) in δ¹³C values but a wide range (9.1‰) in δ¹⁵N, due to the presence of an extreme outlier from Carrer Sant Vicent, VSVC3622 (Table 2). The δ¹³C and δ¹⁵N values for the majority indicate a C₃ terrestrial-based diet with animal protein deriving from herbivores and little contribution from pigs, which have values similar to the humans themselves (Fig. 3, human δ¹⁵N values are on average only 1.8‰ higher in than the mean for pigs). However, there may be an additional source of ¹³C-enriched carbon and ¹⁵N-enriched nitrogen in the diet of some individuals, perhaps from C₄ fed pork or marine sources.

Inter-site comparison in Valencia is limited by San Andrés being represented by just four individuals (Fig. 8). One group of four individuals from the Sant Vicent cemetery are identifiable as possessing δ¹⁵N values that are on par with some of the higher herbivore values. Their consumption of animal protein was therefore limited, and they were probably poor people housed by the hospital at the Sant Vicent complex.

[Figure 8 here]

The outlier from Carrer Sant Vicent, VSVC3622, with a remarkably high δ¹⁵N value of 17‰ lacks a corresponding enrichment in ¹³C, which rules out the consumption of marine fish. This extreme δ¹⁵N value is unlikely to represent a local who consumed freshwater fish, given that the value is higher than even Mesolithic individuals who are hypothesised to have primarily subsisted on freshwater fish from the Danube (Bonsall et al., 2004). A recent study of an Islamic population from Aragón (8th-10th century), however, showed similar δ¹⁵N values (mean of 15‰) correlated with a C₃ carbon value (mean -19.1‰) so that such signatures are possible in some areas of Spain (Guede et al., 2017). Although data from later Christian periods are lacking from that area, northern Aragón is a possible place of origin that ties with population movement during the initial phase of Christian control and repopulation of Valencia. However, any other area with high aridity and C₃ crops remains a possibility.

Across the Valencian region (Fig. 5), there is a greater reliance of on C₄ plants exhibited by the broadly contemporaneous Christian population from Gandía and the *rudéjar* populations of Benipeixcar and El Raval (Alexander et al., 2015; Salazar-García et al., 2014) in comparison to the inhabitants of Christian Valencia. This may again reflect a difference between urban and rural subsistence in this later phase. During the 13th and 14th centuries, the city of Valencia itself had authority over cereal production and distribution

(Barrio, 2009, p. 63) but in times of shortage of wheat grain was imported from outside the kingdom (García Marsilla 1993, p. 7). Because of this, it is highly probable that the availability of wheat in the more rural areas was limited to those who could afford it. The bulk of the populace would have had to (or chosen to) rely on grains produced locally, including millet or sorghum.

DISCUSSION AND SYNTHESIS

Although sample sizes are relatively small for the Visigothic and, to some extent, later Christian periods, the consistency of the core isotopic data for all periods ($\sim -19\text{‰}$ to -17‰ $\delta^{13}\text{C}$, $\sim 10\text{‰}$ to 12‰ $\delta^{15}\text{N}$, Fig. 3) and the stark contrast we see in the Islamic period in particular lends support to our interpretations.

Diachronic trends

The diet for the majority of the sampled population was characterised by a reliance on terrestrial C_3 resources which is similar throughout all three periods under discussion. Despite this, significant shifts in subsistence occurred between each phase, which can be linked to cultural and socioeconomic changes in the historic record. The Islamic period is noteworthy for a wide variation in isotopic values and the greater reliance on C_4 crops compared to the periods both before and after.

The variability in subsistence exhibited by the Islamic population may reflect an increased non-local presence in the population, made visible by migrants from more arid areas where C_4 crops were cultivated, such as the Maghreb. When Christians took control of much of the Iberian Peninsula from the 13th century onwards there was some re-orientation of trade towards northern Christian Europe although some trade links between the Sultanate of Granada and the Islamic Maghreb did continue, if diminished (Constable, 1994; Ferrer, 2012). This change in trade may have had an impact on the origin of travellers, merchants and traders present in the city. The presence of African slaves, however, continued into the Christian period and Valencia became the principal port for the import and export of slaves in the western Mediterranean (Montalvo 2000; Mummey and Reyerson, 2011; Ferrer 2012).

The shift towards the greater use of C_4 crops among local humans and animals in the Islamic period is another factor behind diversity in isotopic values that may be linked to the use of crops such as sorghum (Watson, 1974; Squatriti, 2014). In a recent survey of available archaeobotanical data for the Iberian Peninsula, it was reported that evidence for millets has been recorded on many medieval sites particularly in the west/north-west and the north-east although the archaeobotanical evidence from Valencia itself is extremely patchy and does not extend into the Christian period (Peña-Chocarro et al. 2017). Sorghum is yet to be identified at any site, although evidence has been found in medieval France and North Africa (Peña-Chocarro et al. 2017). It is not until the Christian period (14th and 15th centuries) that we have historical records confirming the presence of C_4 crops such as sorghum in the *Horta* of Valencia, where a white form was used primarily for human consumption, and a red

variety was used as a fodder for cattle and poultry (Rubio, 1995, p. 169). Focus on wheat as the most profitable commercial crop, however, meant that summer grains such as millet and sorghum may have been more limited to the tenants of the irrigated *Horta* (Toró 2009). The lower $\delta^{13}\text{C}$ values exhibited by humans and animals in the Christian population confirm a decrease in C_4 plant consumption, perhaps reflecting a decline in millet and sorghum cultivation during this period.

Socio-cultural factors may have also combined with these economic circumstances. Following the 13th century Christian conquest, the newly arrived population would have comprised of settlers from northern Christian kingdoms, principally Aragón, who were encouraged to migrate by the offer of reduced taxes and enhanced privileges (Burns, 1975; Ruiz, 2014). Written sources suggest that C_4 plants were regarded as minority crops in these regions (Aguilera, 1991, p. 9; Glick, 1982, p. 82), although isotopic data does reveal a reliance on millet in other areas of northern Spain such as Galicia (Kaal et al., 2016; López-Costas and Müldner, 2016) and the Basque Country (Quirós Castillo., 2013). As a result, the consumption of C_4 crops such as sorghum may have decreased after the Christians took control, at least among the urban population (Rubio, 1995, p. 162).

The growing picture of diet in southern and eastern regions of Spain (Fig. 5) suggests that, despite millet being present in the Iberian Peninsula since the Middle Bronze Age (Buxó and Piqué, 2008), C_4 crops only rose in importance as a food source in these areas during the period of Islamic rule. This is evidenced in the comparatively low $\delta^{13}\text{C}$ values seen so far in Late Antique/Visigothic populations and may not have occurred immediately after the Islamic conquest as the largely C_3 diet of the earlier 8th-9th century site in Alicante site attests (Salazar-García et al., 2016). Although C_4 crops continued to be grown under later Christian rule, their importance was diminished and seem firmly associated with the lower social orders and rural populations.

Despite faunal data exhibiting high variation in all periods, shifts in isotopic values were detected, indicating a baseline economic shift and husbandry between all three periods. While there may have been much continuity in husbandry between periods, the *scale* of agriculture and animal husbandry differed with changing production strategies and reorganization of agricultural landscapes, particularly during the later Christian period (Torró, 2012b; López and Retamero, 2017). As mentioned above, there was a shift in economic strategy towards a more intensive focus on economic crops and an emphasis on animal livestock and long-range transhumance (Montalvo, 1992; Trillo San José, 1999; Malpica, 2012). It is in the later Christian period that large sheep flocks were driven through the peninsula from Aragón and Castile to the lowland pastures of southern Valencia (Klein, 1920; Butzer, 1988; Montalvo, 1992). Similarly, cattle ranching had developed earlier in the northern Christian kingdoms and was introduced to Valencia in the 13th century. Herds from Valencia were driven to grazing pastures some 170km to the south at Elche (O'Callaghan, 1975, p. 478; Montalvo, 1992, p. 169). This change towards large-scale animal husbandry in the Christian period may serve to lessen the variability in livestock practices and homogenize dietary signals among livestock when compared to the Islamic period. For example, transhumant cattle and sheep flocks driven between summer and winter grazing lands would have effectively fed on C_3 pastures all year round which may have led to a decline in C_4 fodder fed to local sedentary animals. This would particularly have been the case at times of the year when there was only poor-quality local pasture available. To determine whether

the differences in livestock practices suggested here are truly significant, further samples would need to be analysed from different quarters of the city and be combined with zooarchaeological study.

Although the isotopic data indicate the potential for certain individuals to have consumed marine resources in the Islamic period, the lack of evidence for significant marine fish consumption for the later Christian period is somewhat surprising, particularly given that fish would have fulfilled a liturgical requirement. Port archives of the 15th century indicate the importance of fishing to the city, mentioning the availability of 27 different types of marine and freshwater fish (Guiral, 1989, 381). Fresh fish either originated from the Mediterranean or from brackish lagoons such as the Albufera; sardines in particular were a cheap fish that could be afforded by those of little wealth in Valencia during this period (Granero and Amparo, 1962; Gallart et al., 2005; García Marsilla, 1993). The dietary contribution of low trophic level marine fish and minor quantities of marine and especially brackish resources is, however, difficult to detect using isotopic analysis, particularly for the Mediterranean (Garvie-Lok, 2001; Hedges, 2004; Triantaphyllou *et al.*, 2008). There is only one published instance of increased fish consumption in the later medieval period in the Mediterranean, from 15th century Rome, where the consumption of traded Atlantic fish has been hypothesised (Salamon et al., 2008). Valencia and Alicante served as redistribution centres for fish caught in Galicia and being transported inland to Castile and Aragón (Ferreira, 1988), although high trophic level Atlantic fish with corresponding higher $\delta^{15}\text{N}$ values (e.g. López-Costas and Müldner, 2016) do not appear to have been a major component in the diet. The isotopic results particularly for the later Christian period in Valencia might be sufficient to question how far marine resources contributed to the local diet in the city.

It is notable that no differentiation in diet was detected between sexes at any period, something which is not detected universally in medieval populations. This may be due to a lack of dietary differentiation despite the differing roles of males and females purported for medieval Muslim and Christian societies, where women were typically associated with food preparation and domestic tasks (Shatzmiller, 1994; Ward, 2002). On the other hand, this may simply reflect the broad nature of isotopic evidence for diet and/or the small sample sizes often involved in such comparisons.

CONCLUSION

Diet in Valencia over 1,000 years is characterised by significant isotopic variability between Visigothic, Muslim and later medieval Christian individuals, revealing distinct dietary regimes between these culturally contrasting phases of medieval Valencia. Diachronic trends in the data support historical evidence for a significant change in diet and subsistence, particularly during the Muslim and later Christian periods. Differences in both human and animal data between periods are the result of a complex interplay of factors including socioeconomic shifts and developing travel and trade routes.

Although many individuals in the Islamic period have isotopic values which are similar to those exhibited by individuals in the earlier Visigothic period, there is a far greater range in the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of the Muslim population. This may be due to an increased

number of migrants from areas with distinctive enrichment of foods in ^{13}C and ^{15}N such as the Maghreb, to which Valencia was politically and culturally aligned in this period. This diversity in values also almost certainly reflects the variety in choice or access to isotopically distinct foods at this time, as well as the introduction of new crops, an emphasis on polyculture and the balance of local herding and cereal farming in the region. The growth of urban Valencia under Islamic rule may also have encouraged diversification, particularly the introduction of C_4 crops to the diets of both humans and animals. These foodstuffs are traditionally associated with lower status social groups and have the effect of increasing the isotopic 'visibility' of socioeconomic differences among the populations sampled here, although issues around the detection of marine protein consumption obscure a clear interpretation.

A reduction in observed dietary diversity in the Christian period may be linked with a change in economic focus, both in terms of the reorganisation of trade routes but also agriculture and animal husbandry. The transition to Christian feudal rule emphasised commercial crops and flocks, potentially homogenising isotopic values, although rural and urban differences in the use of C_4 crops remained. Given the reorientation of trade routes, people from the medieval Islamic world with dietary isotopic signatures indicative of C_4 consumption and/or living in arid environments were less likely to be present in the city.

The potential for differential consumption of marine fish complicating the subsistence patterns interpreted here cannot be disregarded although identifying this with bulk isotope analysis is confounded by the C_4 animal protein/Mediterranean fish phenomenon. In the future, techniques utilising compound-specific analysis of bone collagen amino acid $\delta^{13}\text{C}$ could be implemented to identify diets with high marine protein input (Corr et al., 2005). These developing techniques, combined with a thorough regional review of existing medieval archaeozoological and archaeobotanical datasets, offer the best prospect for the future.

LITERATURE CITED

- Adamson M. W. (2004). *Food in medieval times*. Connecticut & London: Greenwood Press.
- Aguilera, J. (1991). *La cocina hispano-árabe y las cocinas españolas y del norte de África*. Madrid: Darek-Nyumba, D.L.
- Albiach, R., Badia, Á., Calvo, M., Marín, C., & Ribera, A. (1998). *Memòria de l'excavació arqueològica al solar de L'Almoina (Valencia)*. 8ª Campaña. Valencia: Ayuntamiento de Valencia.
- Alexander, M. M., Gerrard, C. M., Gutiérrez, A., & Millard, A. R. (2015). Diet, society, and economy in late medieval Spain: stable isotope evidence from Muslims and Christians from Gandía, Valencia. *American Journal of Physical Anthropology*, 156(2), 263-273.
- Álvarez, N., & Ribera, A. (2002). *Informe arqueológico de la excavación de L'Almoina (Valencia)*. 11ª Campaña. Valencia: Ayuntamiento de Valencia.
- Barrio, J. A. (2009). La producción, el consumo y la especulación de los cereales en una ciudad de frontera: Orihuela, siglos XII-XIV. In B. Arízaga & J. A. Solárzano (eds) *Alimentar la ciudad en la Edad Media* (59-86). Logroño: Instituto de Estudios Riojanos.
- Baydal, V., & Esquilache, F. (2014). Exploitation and differentiation: Economic and social stratification in the rural Muslim communities of the Kingdom of Valencia, 13th–16th centuries. In F. Aparisi, & V. Royo (eds) *Beyond lords and peasants: rural elites and economic differentiation in pre-modern Europe* (37-67). Valencia: Publicacions de la Universitat de València.
- Bisson, T. N. (1986). *The medieval Crown of Aragon: a short history*. Oxford University Press.
- Blumenthal, D. (2009). *Enemies and familiars: slavery and mastery in fifteenth-century Valencia*. Ithaca and London: Cornell University Press.
- Bocherens, H., & Drucker, D. (2003). Trophic level isotopic enrichment of carbon and nitrogen in bone collagen: case studies from recent and ancient terrestrial ecosystems. *International Journal of Osteoarchaeology*, 13(1-2), 46-53.
- Bogaard, A., Heaton, T. H., Poulton, P., & Merbach, I. (2007). The impact of manuring on nitrogen isotope ratios in cereals: archaeological implications for reconstruction of diet and crop management practices. *Journal of Archaeological Science*, 34(3), 335-343.
- Britton, K., Müldner, G., & Bell, M. (2008). Stable isotope evidence for salt-marsh grazing in the Bronze Age Severn Estuary, UK: implications for palaeodietary analysis at coastal sites. *Journal of Archaeological Science*, 35(8), 2111-2118.
- Brothwell, D. R. (1981). *Digging up bones*. New York: Cornell University Press.
- Brown, T. A., Nelson, D. E., Vogel, J. S., & Southon, J. R. (1988). Improved collagen extraction by modified Longin method. *Radiocarbon* 30, 171–177.
- Burns, R. I. (1967). *The Crusader Kingdom of Valencia: reconstruction of a thirteenth-century frontier, Volume 1*. Cambridge, Mass.: Harvard University Press.

- Burns, R. I. (1981). Los mudéjares de Valencia: temas y metodología. *Actas del I Simposio Internacional de Mudejarismo* (pp. 453–497). Teruel: Instituto de Estudios Turolenses.
- Burns, R. I. (2003). Valencia, Kingdom and city of. In E. M. Gerli (ed.) *Medieval Iberia: an Encyclopaedia (829-831)*. New York, London: Routledge.
- Butzer, K. W. (1988). Cattle and sheep from Old to New Spain: historical antecedents. *Annals of the Association of American Geographers*, 78(1), 29-56.
- Buxó, R., & Piqué, R. 2008. *Arqueobotánica: los usos de las plantas en la Península Ibérica*. Barcelona: Ariel prehistoria.
- de Castro, T. (1993). *La alimentación en las crónicas castellanas bajomedievales*. Granada: Universidad de Granada.
- Christie, N., Beavitt, P., Gisbert, J. A., Seguí, J. & Gil, M. V. (2004). *Ethnography and archaeology in upland Mediterranean Spain*. Leicester: School of Archaeology and Ancient History, University of Leicester.
- Constable, O. R. (1994). *Trade and traders in Muslim Spain*. Cambridge: Cambridge University Press.
- Coplen, T. B. (2011). Guidelines and recommended terms for expression of stable-isotope-ratio and gas-ratio measurement results. *Rapid Communications in Mass Spectrometry*, 25(17), 2538-2560.
- Corr, L. T., Sealy, J. C., Horton, M. C., & Evershed, R. P. (2005). A novel marine dietary indicator utilising compound-specific bone collagen amino acid $\delta^{13}\text{C}$ values of ancient humans. *Journal of Archaeological Science*, 32(3), 321-330.
- Craig, O. E., Bondioli, L., Fattore, L., Higham, T., & Hedges, R. (2013). Evaluating marine diets through radiocarbon dating and stable isotope analysis of victims of the AD79 eruption of Vesuvius. *American Journal of Physical Anthropology*, 152(3), 345-352.
- Decker, M. (2009). Plants and progress: rethinking the Islamic agricultural revolution. *Journal of World History*, 20(2), 187-206.
- DeNiro, M. J. (1985). Postmortem preservation and alteration of in vivo bone collagen isotope ratios in relation to palaeodietary reconstruction. *Nature* 317, 806–809.
- Dupras, T. L., & Schwarcz, H. P. (2001). Strangers in a strange land: stable isotope evidence for human migration in the Dakhleh Oasis, Egypt. *Journal of Archaeological Science*, 28(11), 1199-1208.
- Estaca-Gómez, V., Malalana-Ureña, A., Yravedra, J., Matás, G.J.L. & de Pablos, J.M. (2018). Economic implications of livestock management strategies in the center of the Iberian Peninsula, Tagus Basin, and Mancha Alta region between the VIII and XI centuries AD. *Archaeological and Anthropological Sciences*, 1-17. doi:10.1007/s12520-018-0607-9
- Ferreira, E. (1988). *Galicia en el comercio marítimo medieval*. Colección de documentos históricos. La Coruña.
- Ferrer, M. T. 2012. Catalan commerce in the later Middle Ages. *Catalan Historical Review*, 29-65.

- Fuller, B. T., Márquez-Grant, N., & Richards, M. P. (2010). Investigation of diachronic dietary patterns on the islands of Ibiza and Formentera, Spain: evidence from carbon and nitrogen stable isotope ratio analysis. *American Journal of Physical Anthropology*, 143(4), 512-522.
- Gallart, L., Roberto, I. E. & Maupoey, P. F. (2005). *La salazón de pescado, una tradición en la dieta mediterránea*. Valencia: Universidad Politécnica de Valencia.
- García, M. I. & Ruiz, E. (1997). *Informe arqueológico preliminar Calle Pintor Sorolla, Nº 5 y 7 de Valencia*. Valencia: Ayuntamiento de Valencia.
- García Marsilla, J. V. (1993). *La jerarquía de la mesa*. Valencia: Centre d'Estudis d'Història Local.
- García Sánchez, E. (1996). La alimentación popular urbana en al-Andalus. *Arqueología Medieval*, 4, 219 - 235.
- García Sánchez, E. (2002). Dietic aspects of food in al-Andalus. In D. Waines (ed.) *Patterns of everyday life: the formation of the Classical Islamic world, vol. 10*, (pp. 275-288). Aldershot: Ashgate.
- Gerrard, C. M. (2003) *Paisaje y señorío. La casa conventual de Ambel (Zaragoza): arqueología, arquitectura e historia de las Órdenes militares del Temple y del Hospital*. Zaragoza: Institución Fernando el Católico.
- Glick, T. F. (1970). *Irrigation and society in medieval Valencia*. Cambridge, Mass.: Belknap Press of Harvard University Press.
- Glick, T. F. (1982). Agriculture and nutrition: the Mediterranean region. In R. Strauyer (ed.) *Dictionary of the Middle Ages* (pp. 79-88). New York: Charles Scribner's Sons for the American Council of Learned Societies.
- Glick, T. F. (1995). *From Muslim fortress to Christian castle: social and cultural change in medieval Spain*. Manchester: Manchester University Press.
- Glick, T. F. (2005). *Islamic and Christian Spain in the Early Middle Ages*. Leiden: Brill.
- Granero, C., & Amparo, M. (1962). Abastecimientos de la ciudad de Valencia durante la Edad Media. *Saitabi* 12, 1-27.
- Grau-Sologestoa, I. (2017). Socio-economic status and religious identity in medieval Iberia: the zooarchaeological evidence. *Environmental Archaeology*, 22(2), 189-199.
- Guede, I, Ortega, L. A., Zuluaga, M. C., Alonso-Olazabal, A., Murelaga, X., Pina, M., Gutiérrez, F. J., & Iacumin, P. (2017). Isotope analyses to explore diet and mobility in a medieval Muslim population at Tauste (NE Spain). *PLoS One* 12(5), e0176572.
- Guinot, E. (2009). La construcción de una ciudad feudal: Valencia (1238-1300). In J. Hermosilla Pla (coord.) *La ciudad de Valencia, historia, geografía y arte de la ciudad de Valencia. Vol. 1: Historia* (pp. 169-179). Valencia: Universitat de Valencia.
- Guinot, E. and Esquilache, F., (2017). Not only peasants: the myth of continuity in the irrigation communities of Valencia, Spain, in the medieval and early modern periods. *Continuity and Change*, 32(2), 129-156.

- Guichard, P. (2000). *Al-Andalus (711-1492)*. Paris: Hachette Littératures.
- Guiral, J. (1989). *Valencia puerto mediterráneo en el siglo XV (1410-1525)*. Valencia: Edicions Alfons el Magnànim.
- Guiry, E. J. (2012). Dogs as analogs in stable isotope-based human paleodietary reconstructions: a review and considerations for future use. *Journal of Archaeological Method and Theory*, 19(3), 351-376.
- Hammer, Ø., Harper, D. & Ryan P. D. (2001). PAST: Paleontological statistics software package for education and data analysis. *Palaeontology Electronic* 4, 9.
- Heaton, T., H. (1987). The $^{15}\text{N}/^{14}\text{N}$ ratios of plants in South Africa and Namibia: relationship to climate and coastal/saline environments. *Oecologia*, 74, 236-246.
- Hedges, R. E. M. (2004). Isotopes and red herrings: comments on Milner et al. & Lidén et al. *Antiquity*, 78(299), 34-37.
- Hedges, R. E. M., Thompson, J., & Hull, B. D. (2005). Stable isotope variation in wool as a means to establish Turkish carpet provenance. *Rapid Communications in Mass Spectrometry*, 19(22), 3187-3191.
- Hedges, R. E. M., Clement, J. G., Thomas, C. D. L., & O'Connell, T. C. (2007). Collagen turnover in the adult femoral mid-shaft: modeled from anthropogenic radiocarbon tracer measurements. *American Journal of Physical Anthropology*, 133(2), 808-816.
- Herreros, A. (1999). *Excavación realizada en la Calle Sagunto no. 165 y 167 de Valencia*. Valencia: Ayuntamiento de Valencia.
- Hidalgo, J. G., De Luis, M., Raventós, J., & Sánchez, J. R. (2003). Daily rainfall trend in the Valencia Region of Spain. *Theoretical and Applied Climatology*, 75(1-2), 117-130
- Houston, J. M., 1949. Urban geography of Valencia: the regional development of a Huerta City. *Transactions and Papers (Institute of British Geographers)* 15, 19–35.
- Jiménez-Brobeil, S. A., Laffranchi, Z., Maroto, R. M., Sánchez, F. L., & Huertas, A. D. (2016). How royals feasted in the court of Pedro I of Castile: A contribution of stable isotope study to medieval history. *Journal of Archaeological Science: Reports*, 10, 424-430.
- Jiménez Salvador, J. L. 2009. La ciudad visigoda: un nuevo paisaje urbano. In J. Hermsilla Pla (coord.) *La ciudad de Valencia, historia, geografía y arte de la ciudad de Valencia. Vol. 1: Historia* (pp. 139-147). Valencia: Universitat de Valencia.
- Jørkov, M. L. S., Heinemeier, J., & Lynnerup, N., 2007. Evaluating bone collagen extraction methods for stable isotope analysis in dietary studies. *Journal of Archaeological Science*, 34(11), 1824-1829.
- Kaal, J., López-Costas, O., & Martínez Cortizas, A. (2016). Diagenetic effects on pyrolysis fingerprints of extracted collagen in archaeological human bones from NW Spain, as determined by pyrolysis-GC-MS. *Journal of Archaeological Science*, 65, 1–10.
- Katzenberg, M. A. (2000). Stable isotope analysis: a tool for studying past diet, demography and history. In: M. A. Katzenberg & S. R. Saunders (eds) *Biological anthropology of the human skeleton* (pp. 305–328). New York: Wiley Periodicals.

- Klein, J. (1920). *The Mesta: a study in Spanish economic history 1273-1836*. Cambridge: Harvard University Press.
- Kennedy, H. (1996). *Muslim Spain and Portugal: a political history of al-Andalus*. London: Longman.
- van Klinken, G. (1999). Bone Collagen Quality Indicators for Palaeodietary and Radiocarbon Measurements. *Journal of Archaeological Science* 26, 687–695.
- Lee-Thope, J. A. (2008). On isotopes and old bones. *Archaeometry*, 50(6), 925-950.
- Lightfoot, E., Slaus, M., & O'Connell, T. C. 2012. Changing cultures, changing cuisines: cultural transitions and dietary change in Iron Age, Roman, and Early Medieval Croatia. *American Journal of Physical Anthropology* 148, 543-556.
- Longin, R. (1971). New method of collagen extraction for radiocarbon dating. *Nature*, 230, 241–242.
- Lopez, E., & Retamero, F. (2017). Segregated fields. Castillian and Morisco peasants in Moclón (Málaga, Spain, sixteenth century). *International Journal of Historical Archaeology*, pp.1-18. doi: 10.1007/s10761-016-0390-1
- López-Costas, O., & Müldner, G. (2016). Fringes of the empire: diet and cultural change at the Roman to post-Roman transition in NW Iberia. *American Journal of Physical Anthropology*, 161(1), 141-154.
- Lozoya, M. C., Gallego, R. S. & Pérez, J. P. (2006). Ensayo de rituales de enterramiento islámicos en Al-Andalus. *Anales de prehistoria y arqueología* 22, 153-164.
- Malpica, A. (2012). La vida agrícola y la ganadería en al-Andalus y en el reino nazarí de Granada. In R. Marín López (coord.) *Homenaje al Profesor José Ignacio Fernández de Viana* (213-228). Granada: University of Granada.
- Marsilla, J. V. G. (1993). *La jerarquía de la mesa: los sistemas alimentarios en la Valencia bajomedieval*. Valencia: Center d'Estudis d'Història Local.
- Martin, L. & Ribera, A. V. (2006). Cementerios tardoantiguos de Valencia: arqueología y antropología. *Anales de Arqueología Cordobesa* 17(2), 161-194.
- Mays, S. and Cox, M. (2000). Sex determination in skeletal remains. In M. Cox and S. Mays (eds) *Human osteology in Archaeology and Forensic Science* (pp. 117-130). London: Cambridge University Press.
- Momblanch, F, P. (2003). *Historia de la Albufera de Valencia*. Valencia: Ayuntamiento de Valencia.
- Montalvo, J. H. (1992). Aproximación a la ganadería alicantina en la Edad Media. *Anales de la Universidad de Alicante, Historia medieval* 9, 161-178.
- Montalvo, J. H. (2000). Esclavos, nobles y corsarios en el Alicante medieval. In *De la esclavitud a la libertad en el reino de Valencia durante los siglos medievales*, Barcelona: CSIC, 431–71.
- Montserrat, P. & Fillat, F. (1990). The systems of grassland management in Spain. In A. I.

- Breymeyer (ed.) *Managed grasslands: regional studies, ecosystems of the World* v.17A (37-70). Amsterdam: Elsevier Publications.
- Morales, A., Moreno-García, M., Roselló, E., Llorente, L., & Morales, D. C. (2011). 711 AC: ¿el origen de una disyunción alimentaria?. *Zona Arqueológica* 15(2) 303-322.
- Moreno, A., Pérez, A., Frigola, J., Nieto-Moreno, V., Rodrigo-Gámiz, M., Martrat, B., González-Sampériz, P., Morellón, M., Martín-Puertas, C., Corella, J.P. & Belmonte, Á. (2012). The medieval climate anomaly in the Iberian Peninsula reconstructed from marine and lake records. *Quaternary Science Reviews*, 43, 16-32.
- Moreno-Larrazabal, A., Teira-Brión, A., Sopelana-Salcedo, I., Arranz-Otaegui, A., & Zapata, L. (2015). Ethnobotany of millet cultivation in the north of the Iberian Peninsula. *Vegetation History and Archaeobotany*, 24(4), 541-554.
- Mummey, K. & Reyerson, K. (2011). Whose city is this? Hucksters, domestic servants, wet-nurses, prostitutes, and slaves in late medieval western Mediterranean urban society. *History Compass*, 9(12), 910-922.
- O'Callaghan, J. F. (1975). *A history of medieval Spain*. Ithaca and London: Cornell University Press.
- O'Connell, T. C., Kneale, C. J., Tasevska, N., & Kuhnle, G. G. (2012). The diet-body offset in human nitrogen isotopic values: A controlled dietary study. *American Journal of Physical Anthropology*, 149(3), 426-434.
- O'Leary, M. H. (1981). Carbon isotope fractionation in plants. *Phytochemistry*, 20(4), 553-567.
- Ortega, D. & Gálego, A. (2004). *Informe preliminar de la intervención arqueológica en la C/ Sant Vicent 134, Valencia*. Valencia: Ayuntamiento de Valencia.
- Pascual, J. (1989). La necrópolis islámica de L'Almoina (Valencia). Primeros resultados arqueológicos. *III Congreso de Arqueología Medieval Española II* (pp. 406-412). Oviedo: Universidad de Oviedo.
- Pascual, J. & Ribera, A. (1990). Plaça de L'Almoina in *Excavaciones arqueológicas de selvament a la Comunitat Valenciana 1984-1988, 1, intervencions urbanes*. Valencia: Generalitat de Valencia.
- Pascual, J. & Serrano, M. L. (1996). Necrópolis islámicas en la ciudad de Valencia. *Saitabi*, 46, 231-252.
- Peña-Chocarro, L., Pérez-Jordà, G., Alonso, N., Antolín, F., Teira-Brión, A., Tereso, J. P., Moya, E. M. M., & Reyes, D. L. (2017). Roman and medieval crops in the Iberian Peninsula: A first overview of seeds and fruits from archaeological sites. *Quaternary International*, In Press, doi: 10.1016/j.quaint.2017.09.037
- Pestle, W. J., Crowley, B. E. & Weirauch, M. T. (2014). Quantifying inter-laboratory variability in stable isotope analysis of ancient skeletal remains. *PLOS One*, 9(7), p.e102844.
- Pickard, C., Girdwood, L. K., Kranjoti, E., Márquez-Grant, N., Richards, M. P., & Fuller, B. T. (2017). Isotopic evidence for dietary diversity at the mediaeval Islamic necropolis of Can

- Fonoll (10th to 13th centuries CE), Ibiza, Spain. *Journal of Archaeological Science: Reports*, 13, 1-10.
- Quirós Castillo, J. A. (2013). Los comportamientos alimentarios del campesinado medieval en el País Vasco y su entorno (siglos VIII-XIV). *Historia Agraria*, 59, 13-41.
- Reitz, E. J., Speller, C., McGrath, K., & Alexander, M. (2016). A sixteenth-century turkey (Meleagris gallopavo) from Puerto Real, Hispaniola. *Journal of Archaeological Science: Reports*, 10, 640-646.
- Ribera, A. (2005). El contexto histórico y arqueológico de las emisiones monetarias en el País Valenciano. *Gaceta numismática*, 157, 45-62.
- Ribera, A. (2006). The Roman foundation of Valencia and the town in the 2nd-1st c. B.C. In L. Abad, S. Keay & S. Ramallo (eds.) *Early Roman Towns in Hispania Tarraconensis* (pp.75-90). Journal of Roman Archaeology Supplementary Series Number 62, Portsmouth, Rhode Island.
- Ribera, A. (2008). La ciudad de Valencia durante el periodo visigodo. *Zona arqueológica*, 9, 303-320.
- Richards, M. P., Fuller, B. T., & Molleson, T. I. (2006). Stable isotope palaeodietary study of humans and fauna from the multi-period (Iron Age, Viking and Late Medieval) site of Newark Bay, Orkney. *Journal of Archaeological Science*, 33(1), 122-131.
- Ruas, M.-P., Mane, P., Hallavant, C., & Lemoine, M. (2017). Citrus fruit in historical France. Written sources, iconographic and archaeological remains. In V. Zech-Matterne & G. Fiorentino (eds.) *AGRUMED: Archaeology and history of citrus fruit in the Mediterranean* (pp. 157-184). Naples: Publications du Centre Jean Bérard.
- Rubio, A. (1995). El consumo de pan en la Valencia bajomedieval. *Col.loqui d'història de l'alimentació a la Corona d'Aragó. Actes*, Vol. 1, 153-184.
- Ruiz, T. F. (2014). *Spanish Society, 1400-1600*. London and New York: Routledge.
- Ruiz-Taboada, A., (2015). Comments on Muslim, Jewish and Christian Burial Practices in medieval Toledo (Spain). *Journal of Islamic Archaeology*, 2(1), 51-71.
- Salazar-García, D. C., Richards, M. P., Nehlich, O., & Henry, A. G. (2014). Dental calculus is not equivalent to bone collagen for isotope analysis: a comparison between carbon and nitrogen stable isotope analysis of bulk dental calculus, bone and dentine collagen from same individuals from the Medieval site of El Raval (Alicante, Spain). *Journal of Archaeological Science*, 47,70–77.
- Salazar-García, D. C., Romero, A., García-Borja, P., Subirà, M. E., & Richards, M. P. (2016). A combined dietary approach using isotope and dental buccal-microwear analysis of human remains from the Neolithic, Roman and Medieval periods from the archaeological site of Tossal de les Basses (Alicante, Spain). *Journal of Archaeological Science: Reports*, 6, 610-619.
- Sealy, J., Johnson, M., Richards, M., & Nehlich, O. (2014). Comparison of two methods of extracting bone collagen for stable carbon and nitrogen isotope analysis: comparing whole bone demineralization with gelatinization and ultrafiltration. *Journal of*

- Archaeological Science*, 47, 64-69.
- Serrano, L. (1997). *Memoria preliminar de la intervención arqueológica situada en los solares nºs 48 al 58 de la Calle Alta Valencia*. Valencia: Ayuntamiento de Valencia
- Schoeninger, M. J., DeNiro, M. J., & Tauber, H. (1983). Stable nitrogen isotope ratios of bone collagen reflect marine and terrestrial components of prehistoric human diet. *Science*, 220(4604), 1381-1383.
- Schoeninger, M. J., & DeNiro, M. J. (1984). Nitrogen and carbon isotopic composition of bone collagen from marine and terrestrial animals. *Geochimica et Cosmochimica Acta*, 48(4), 625-639.
- Scheuer, L., & Black, S. (2000). *Developmental juvenile osteology*. San Diego: Academic Press.
- Schwarcz H. P. & Schoeninger, M. J. (2011). Stable isotopes of carbon and nitrogen as tracers for palaeo-diet reconstruction. In: M. Baskaran (ed.) *Handbook of environmental isotope geochemistry* (pp. 725–742). Berlin: Springer-Verlag.
- Schwarcz, H. P., Dupras, T. L., & Fairgrieve, S. I. (1999). ¹⁵N enrichment in the Sahara: in search of a global relationship. *Journal of Archaeological Science*, 26(6), 629-636.
- Shatzmiller, M. (1994). *Labour in the medieval Islamic world*. Leiden: Brill.
- Sirignano, C., Grau Sologestoa, I., Ricci, P., García-Collado, M. I., Altieri, S., Quirós Castillo, J. A., Lubritto, C. (2014). Animal husbandry during Early and High Middle Ages in the Basque Country (Spain). *Quaternary International* 346,138–148.
- Squatriti, P. (2014). Of seeds, seasons, and seas: Andrew Watson’s medieval agrarian revolution forty years later. *The Journal of Economic History*, 74(4), 1205-1220.
- Szpak, P. (2014). Complexities of nitrogen isotope biogeochemistry in plant-soil systems: implications for the study of ancient agricultural and animal management practices. *Frontiers in Plant Science*, 5, 288.
- Tomás, M. S. (2009). El uso terapéutico de la alimentación en la Baja Edad Media. In B. Arízaga & J. A. Solórzano (eds) *Alimentar la ciudad en la Edad Media* (pp. 459-490). Logroño: Instituto de Estudios Riojanos.
- Torró, J. (2009a). Field and canal building after the conquest: modifications to the cultivated ecosystem in the Kingdom of Valencia, ca. 1250-ca. 1350. In: B. A. Catalos (ed.) *Worlds of History and Economics* (pp. 77-108). Valencia: Universitat de Valencia.
- Torró, J. (2009b). *Madíña Balansiya: la Valencia andalusí. Siglos VIII-XIII*. In J. Hermosilla Pla (coord.) *La ciudad de Valencia, historia, geografía y arte de la ciudad de Valencia. Vol. 1: Historia* (pp. 159-169). Valencia: Universitat de Valencia.
- Torró, J. (2012a). One aspect of the Christian settlement of the Kingdom of Valencia: the drainage and placing under cultivation of coastal wetlands (c. 1270-1320). In P. Galetti (ed.) *Paesaggi, Comunità, Villaggi Medievali* (pp.225-237). Fondazione Centro Italiano di Studi Sull’Alto Medioevo, Spoleto.
- Torró, J. (2012b). La conquista del Reino de Valencia. Un proceso de colonización medieval desde la Arqueología del territorio. In J. A. Eiroa (ed.) *La conquista de al-Andalus en el*

siglo XIII (pp. 9-40). Murcia: Editum-Centro de Estudios Medievales de la Universidad de Murcia.

Triantaphyllou, S., Richards, M. P., Zerner, C., & Voutsaki, S. (2008). Isotopic dietary reconstruction of humans from Middle Bronze Age Lerna, Argolid, Greece. *Journal of Archaeological Science*, 35(11), 3028-3034.

Trillo San José, C. (1999). El paisaje vegetal en la Granada islámica y sus transformaciones tras la conquista castellana. *Historia Agraria*, 17, 131-152.

Turner, B. L., Edwards, J. L., Quinn, E. A., Kingston, J. D., & Van Gerven, D. P. (2007). Age-related variation in isotopic indicators of diet at medieval Kulubnarti, Sudanese Nubia. *International Journal of Osteoarchaeology*, 17(1), 1-25.

Ubelaker, D. H. (1989). *Human skeletal remains: excavation, analysis, interpretation*. Washington: Taraxacum.

Vassberg, D. E. (1974). The 'Tierras Baldías': Community property and public lands in 16th century Castile. *Agricultural History*, 48(3), 383-401.

Waines, D. (1992). The culinary culture of al-Andalus. In S. K. Jayyusi (ed.) *The Legacy of Muslim Spain* (pp. 725-740). Lieden: Brill.

Ward, J. (2002). *Women in medieval Europe 1200-1500*. Harlow: Longman.

Watson, A. M. (1974). The Arab agricultural revolution and its diffusion, 700-1100. *The Journal of Economic History*, 34(1), 8-35.

Watson, A. M. (1983). *Agricultural innovation in the early Islamic world: the diffusion of crops and farming techniques 700-1100*. London: Cambridge University Press.

FIGURE CAPTIONS

Figure 1: Location map of Valencia. Map created with data from Natural Earth and the Shuttle Radar Topography Mission (SRTM).

Figure 2: Map of the modern centre of Valencia with the location of sites sampled. Drawing by A. Gutiérrez after CartoCiudad 2006-2017 CC-BY-4.0 scne.es.

Figure 3: Stable carbon and nitrogen isotope data for animal and human bone collagen for each period, Visigothic (A), Islamic (B) and Christian (C).

Figure 4: Stable carbon and nitrogen isotope data for herbivore (cattle, sheep/goat, white boxes) and human (grey boxes) bone collagen plotted by period as box and jitter plots. The boxes indicate the inter-quartile range (IQR), the whiskers comprise the values that are with 1.5x the IQR and the dots beyond these are outliers. Sample numbers: herbivores 7, 13, 9 and humans 10, 38, 18, for Visigothic, Islamic and Christian respectively.

Figure 5: Mean values for stable carbon and nitrogen isotope data for human bone collagen for the later medieval period published from southern and eastern Spain and Ibiza. Error bars

represent $\pm 1\sigma$. References: Gandía and Benipeixcar (Alexander et al. 2015), El Raval (Salazar-García et al. 2014), Tossal de les Basses (Salazar-García et al. 2016), La Torrecilla and Seville Royalty (Jiménez-Brobeil et al. 2016), Can Fonoll (Pickard et al. 2016), Es Soto and S'Hort des Llimoners (Fuller et al. 2010).

Figure 6: Stable carbon and nitrogen isotope data for human bone collagen from the Visigothic period (5th-7th century) separated by phase.

Figure 7: Stable carbon and nitrogen isotope data for human bone collagen from Almoina, z and y sites in the Muslim period (11th- 13th century)

Figure 8: Stable carbon and nitrogen isotope data for human bone collagen from San Andrés and San Vicent sites in the Christian period (14th-15th century).