



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/143617/>

Version: Accepted Version

Article:

Tierney, GJ and Simms, CK (2018) Can tackle height influence head injury assessment risk in elite rugby union? *Journal of Science and Medicine in Sport*, 21 (12). pp. 1210-1214. ISSN: 1440-2440

<https://doi.org/10.1016/j.jsams.2018.05.010>

© 2018 Sports Medicine Australia. Published by Elsevier Ltd. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Can tackle height influence head injury assessment risk in elite rugby union?

Gregory J. Tierney^[1]

gtierne@tcd.ie

Ciaran K. Simms^[1]

csimms@tcd.ie

^[1]Trinity Centre for Bioengineering, Trinity College Dublin, Ireland

Word Count: 2569

Abstract Word Count: 250

Key Words: Concussion, Head Impact, Tackling, Injury Prevention

Number of Figure: 2

Number of Tables: 1

Corresponding Author:

Gregory Tierney

Trinity Centre for Bioengineering

Trinity College Dublin

Dublin 2

Ireland

gtierne@tcd.ie

+353 89 2381688

1 **Can tackle height influence head injury assessment risk in elite rugby union?**

2 **Abstract**

3 **Objectives:** Tackle height laws are an area of controversy in rugby union. It is reported that the tackler is at
4 most risk of a Head Injury Assessment (HIA). Therefore, the aim of this study was to use match video
5 evidence of tackles in elite level rugby union to examine the effect of tackle heights on HIA risk for the
6 tackler.

7 **Design:** Qualitative observational case-control study

8 **Methods:** Each HIA (n=74) and control tackle (n=965) was categorised based on tackle direction (front- or
9 side-on), tackle type (arm, shoulder or smother) and tackle height (upper trunk, mid-trunk, lower trunk,
10 upper leg or lower leg). The Relative Risk (RR), 95% Confidence Interval (CI) and probability (*p*) values
11 were calculated for each tackle height.

12 **Results:** Intended primary contact at the upper trunk of the ball carrier had a greater propensity to result in
13 a HIA for the tackler for front-on upper body shoulder tackles (RR=1.48; 95%CI=1.16-1.90; $p<0.01$) and
14 side-on upper body smother tackles (RR=2.30; 95%CI=1.82-2.92; $p<0.01$). Intended primary contact at the
15 upper leg of the ball carrier had a greater propensity to result in a HIA for the tackler for front-on (RR=2.60;
16 95%CI=1.70-3.97; $p<0.01$) and side-on (RR=3.34; 95%CI=1.65-6.79; $p<0.01$) lower body shoulder
17 tackles.

18 **Conclusions:** To reduce tackler HIA risk, the results suggest tackling below the upper trunk for upper body
19 tackles. The results also suggest tackling at the lower trunk for lower body tackles and avoiding the upper
20 legs. Prevention strategies should place emphasis on tackling lower risk body regions such as the mid- and
21 lower trunk.

22 **Word Count:** 3214

23 **Key Words:** Concussion, Head Impact, Tackling, Injury Prevention

24 1. Introduction

25 Rugby union is a territorial sport characterised by frequent contact between opposing players and/or the
26 ground^{1,2}. For the sixth consecutive season, concussion was the most commonly reported match injury for
27 English Premiership rugby (incidence rate of 20.9/1000 player hours, contributing to 22% of all match
28 injuries during the 2016-17 season)³. Tackling is regarded as the most common cause of injury and
29 concussion in rugby union^{4,5}. Correct technique in the tackle is essential for safe and successful tackle
30 outcomes⁶⁻⁸, whilst poor tackle technique is a reported risk factor for injury^{9,10} and direct head impact
31 causation^{2,11}. This indicates the importance for concussion injury prevention strategies.

32 In rugby union, retrospective analyses of match video evidence have previously been used to identify
33 injury^{1,9} and head impact/concussion risk factors^{2,11,12} as well as guide prevention techniques^{2,9,11}. Recent
34 tackle technique studies^{2,13} have categorised legal tackles as either upper body or lower body tackles. An
35 upper body tackle is defined by the tackler's intended initial contact being above the ball carrier's hip while
36 a lower body tackle is defined as the tackler's intended initial contact being at or below the ball carrier's hip
37². One study² found that tacklers were at most risk of sustaining a direct head impact and that upper body
38 tackles were the greatest cause of this. However, the upper body and lower body tackle definitions utilised
39 in this study covered a wide range of body regions. Therefore, the study reports limited information on
40 specific tackle height analysis for direct head impact aetiology. In particular, this study did not conduct a
41 direct head impact risk analysis for specific tackle heights. The tackle height law in rugby union is set at the
42 line of the ball carrier's shoulder and any contact above this line is regarded as foul play¹⁴. It has been an
43 area of concern with respect to injury for many years¹⁵. Lowering the maximum legal tackle height has
44 been recommended since the 1970s¹⁵, but the evidence base for this is limited. To guide concussion
45 prevention strategies and before tackle laws can be changed, it is essential to examine the effect of tackle
46 heights on head impact and concussion risk.

47 The Head Injury Assessment (HIA) was introduced in 2012 by World Rugby as the pitch side assessment
48 process for concussion injuries and has previously been described in detail¹⁶. Briefly, the HIA provides a

49 standardised tool for the medical assessment of concussion injuries in rugby ¹⁶. A player enters the HIA
50 protocol by exhibiting on-field signs and symptoms of concussion ¹⁶. The HIA examines a range of
51 concussive symptoms ¹⁷ including both immediate and delayed memory difficulties, cognitive ability,
52 balance and player discomfort. If a player does not pass the HIA, he or she is removed from play and must
53 follow the return-to-play protocol. Players diagnosed with a concussion are managed through the the
54 graduated return to play protocol set out by World Rugby ³. The protocol encompasses six stages which
55 each player must complete sequentially: (1) physical and cognitive rest until asymptomatic; (2) light
56 aerobic exercise; (3) sport specific exercise; (4) non-contact training drills; (5) full contact practice; (6)
57 return to play. Players can only progress to the next stage if they remained asymptomatic for an unbroken
58 period of 24 hours ³. The minimum time to return to play is 6 days from the day of injury ³. It is clear that a
59 reduction in tackle-related HIAs would be evidence of concussion injury reduction within rugby union.
60 Accordingly, the aim of this study was to use match video evidence of tackles in elite level rugby union to
61 examine the effect of tackle heights on HIA risk for the tackler (based on intended primary contact location
62 on the ball carrier ²), for both upper body and lower body tackles.

63 2. Methods

64 A qualitative observational case-control study design was used to identify the risks associated with specific
65 tackle heights on HIA aetiology in men's professional rugby union using match video evidence. A tackle
66 was defined as "when the ball-carrier was contacted (hit and/or held) by an opponent without reference to
67 whether the ball-carrier went to ground" ¹. A HIA tackle was defined as "when a tackler received a
68 direct/indirect head impact in the tackle and was subsequently removed from play for a HIA and did not
69 return to play for the remainder of the game." ¹¹. All data was freely available online and no medical data on
70 individuals is reported for this study. As a result of this, no ethical permission was required similar to other
71 rugby union video analysis studies on head impacts ^{2,18} and knee injuries ¹⁹. A non-HIA tackle was defined
72 as "when a player did not receive an injury/head impact/HIA (including HIAs that resulted in temporary
73 and permanent removal from the game) as a result of the tackle."

74 The HIA video data from Tierney et al. ¹¹ was utilised for this study. In brief this data consists of 74 tackles
75 that resulted in a HIA for the tackler (19 upper body and 19 lower body for front-on tackles and 23 upper
76 body and 13 lower body for side-on tackles) from elite level competitions including Pro 12 (2014-2017),
77 European Rugby Champions Cup games (2014-2017), RBS 6 Nations (2014-2017), Guinness Autumn Test
78 Series (2013-2016), Rugby World Cup warm-up games (2015), the Rugby World Cup (2015) and the
79 British and Irish Lions Tour (2017). Although a HIA can occur from an impact to the body ¹⁷, a direct head
80 impact was identified in every video. A non-HIA tackle was defined as “when a player did not receive an
81 injury/head impact in the tackle and was not removed from play for the remainder of the game ¹¹.” To
82 provide non-HIA cases as a control cohort, the dataset from Tierney et al. ²⁰ was utilised. In brief, this
83 dataset consisted of five randomly selected games from the abovementioned competitions. Every non-HIA
84 tackle (n=965) from these five games was analysed for the control cohort dataset. **No tackles that resulted in**
85 **a temporary HIA (player returned to play during the game) occurred in the five games of the control cohort.**

86 Each tackle analysed was categorised based on tackle direction (front- or side-on), tackle type (arm,
87 collision, jersey, lift, shoulder, smother or tap ¹⁴) and tackle height (upper trunk, mid-trunk, lower trunk,
88 upper leg or lower leg, see Figure 1). Fuller et al. ¹⁴ defined the following for arm, shoulder and smother
89 tackles; Arm Tackle - “Tackler impedes/stops ball carrier with upper limb(s)”; Shoulder tackle - “Tackler
90 impedes/stops ball carrier with shoulder as the first point of contact followed by use of arm(s)”; Smother
91 tackle - “Tackler uses chest and wraps both arms around ball carrier”. The following definitions were
92 utilised for the tackle height analysis; Upper trunk - line of the shoulders to base of the chest/pectorals;
93 Mid-trunk - base of chest/pectorals to top of pelvis; Lower trunk - top of pelvis to base of pelvis; Upper leg
94 - base of pelvis to base of knees; Lower leg - below base of knees, see Figure 1. One reviewer
95 (Biomechanist) analysed each video. The videos were analysed using Sports Code (Version 8) enabling a
96 frame-by-frame viewing of the tackle. The video had a minimum frame rate of 25 fps and could be watched
97 as many times as necessary.

98

Insert Figure 1 near here

99 Sixty tackles (including HIA and non-HIA cases) were randomly selected using a random number
100 generator (<http://www.random.org/>). The reviewer then conducted the analysis on these 60 cases, for each
101 tackle variable (tackle height, direction and type), at least one week after conducting the initial set of cases.
102 Intra-rater reliability was then assessed using Cohen's Kappa (K). To assess inter-rater reliability, an
103 external reviewer (ex-player) conducted the analysis on the same 60 cases using the same protocol as the
104 main reviewer. Similarly, inter-rater reliability was then assessed using Cohen's Kappa (K). A Cohen's
105 Kappa value greater than 0.8 indicates almost perfect agreement ²¹. For intra-rater reliability, Cohen's
106 kappa values of 0.93, 0.97 and 0.92 were achieved for tackle height, direction and type, respectively. For
107 inter-rater reliability, Cohen's kappa values of 0.83, 0.83, and 0.82 were achieved for tackle height,
108 direction and type, respectively.

109 For upper and lower body front- and side-on tackles, only the main tackle type that resulted in a HIA for
110 each of these categories (see Figure 2) and the tackle type matched control cases were utilised for the
111 statistical analysis. The Relative Risk (RR), 95% Confidence Interval (CI) and probability (*p*) values were
112 calculated for each tackle height ^{2 14}. The RR for each height was calculated by comparing the frequency of
113 occurrence for HIA cases with the frequency of occurrence in non-HIA cases. An RR=1 indicates that the
114 tackle height has no greater propensity to cause a HIA than that anticipated by chance; an RR>1 and RR<1
115 indicates that the tackle height has a greater and lesser propensity to cause a HIA than expected by chance,
116 respectively ^{2 14}. A variable was considered to have statistical significance if the 95% CI for the RR value
117 did not include 1 and the *p*-value was <0.05. The 95% CI was also reported for the frequency of occurrence
118 results ²². RR values >1 and <1.11, ≥1.11 and <1.43, ≥1.43 and <2.00, ≥2.00 and <3.33, ≥3.33 and <10 were
119 indicative of trivial, small, medium, large and very large, respectively ²³. Similarly, RR values <1 and
120 >0.90, ≤0.90 and >0.70, ≤0.70 and >0.50, ≤0.50 and >0.30, ≤0.30 and >0.10 were indicative of trivial,
121 small, medium, large and very large, respectively ²³.

122

123 3. Results

124 Figure 2 shows that shoulder (79%; n=15) and smother tackles (65%; n=15) account for the majority of
125 upper body front- and side-on tackles, respectively. Also, shoulder tackles account for the majority of lower
126 body front- and side-on tackles (95%; n=18 and 71%; n=10, respectively).

127 **Insert Figure 2 near here**

128 Table 1 shows that intended primary contact at the upper trunk of the ball carrier had a greater propensity to
129 result in a HIA for the tackler for front-on upper body shoulder tackles (RR=1.48; 95% CI=1.16-1.90;
130 $p<0.01$) and side-on upper body smother tackles (RR=2.30; 95% CI=1.82-2.92; $p<0.01$). However,
131 intended primary contact at the mid-trunk of the ball carrier had a lower propensity to result in a HIA for the
132 tackler for side-on smother tackles (RR=0.11; 95% CI=0.02-0.75; $p=0.02$).

133 Table 1 illustrates that for front-on lower body shoulder tackles, intended primary contact at the lower trunk
134 of the ball carrier had a lower propensity to result in a HIA for the tackler (RR=0.45; 95% CI=0.23-0.88;
135 $p<0.02$). Intended primary contact at the upper leg of the ball carrier had a much greater propensity to result
136 in a HIA for the tackler for front-on (RR=2.60; 95% CI=1.70-3.97; $p<0.01$) and side-on (RR=3.34; 95%
137 CI=1.65-6.79; $p<0.01$) lower body shoulder tackles.

138 **Insert Table 1 near here**

139 4. Discussion

140 The aim of this study was to use match video evidence of tackles in elite level rugby union to examine the
141 effect of tackle heights on HIA risk for the tackler for both upper and lower body tackles. The results
142 suggest **avoiding** the upper trunk for upper body tackles. Tackles to the upper trunk had a greater propensity
143 to result in a HIA for the tackler in both front- and side-on upper body tackles. For lower body tackles, the
144 results suggest tackling at the lower trunk for lower body tackles and avoiding the upper legs. Tackles to the
145 lower trunk had a lower propensity to result in a HIA for the tackler in front-on shoulder tackles. However,

146 tackles to the upper legs had a higher propensity to result in a HIA for the tackler in both front- and side-on
147 shoulder tackles. These findings can be utilised by coaches to develop tackle height specific prevention
148 strategies and training drills.

149 Tierney et al.²⁴ found that tackles to the upper trunk accounted for almost half (46%) of all tackle related
150 direct head impacts to the tackler, however, that study did not assess tackle height risk. It has been
151 previously reported that the head and shoulders of the ball carrier were the main body parts to strike the
152 tackler's head in direct head impact² and HIA-related²⁵ upper body tackles. This is consistent with the
153 findings of this study as tackling the upper trunk puts the tackler's head closer to these impacting body
154 parts. Furthermore, Tucker et al.²⁵ found that an upright tackler, which corresponds to a higher tackle
155 height, was the most common body position to result in a HIA for the tackler. In this study, it was found that
156 in certain upper trunk tackles, the ball carrier entered the tackle in a crouched position meaning that an
157 upper trunk tackle was almost unavoidable, particularly for front-on tackles. Therefore, if a change to the
158 tackle height law was to be made, it would have to ensure that this is mitigated against.

159 **The results of this study indicate that tackling at the upper legs has a higher propensity to result in a HIA**
160 **than tackling at the upper trunk. Lowering the maximum legal tackle height to below the upper trunk of the**
161 **ball carrier could reduce HIA risk during upper body tackles. However, it may increase the likelihood of**
162 **upper leg related HIAs as lowering the tackle height law could increase the number of tackles to the upper**
163 **leg region. This could have an adverse effect on HIA reduction. Therefore, effective coaching strategies**
164 **that place emphasis on tackling lower risk body regions such as the mid- and lower trunk are paramount.**

165 The intended primary contact at the upper legs for both front- and side-on lower body tackles showed a
166 greater propensity to result in a HIA for the tackler. The judgement made by the tackler arises in a dynamic
167 situation in which the ball carrier can adjust both running speed and direction. Part of the skill of the ball
168 carrier is to be unpredictable ensuring that the tackler does not make an effective tackle. Therefore, the
169 upper legs of the ball carrier can be moving rapidly which could increase the risk of a head impact for the
170 tackler in comparison to the lower trunk, for example, which reflects the bulk movement of the player. The

171 results also demonstrate that tackling the mid/lower trunk of the ball carrier has a lower propensity to result
172 in a HIA. This supports the recommendation of contacting the ball carrier in the centre of gravity proposed
173 in previous contact technique based studies^{9,26}. Quarrie and Hopkins¹ found that tackling high (roughly at
174 the upper trunk) was the main legal tackle to cause general injury for the ball carrier in terms of injury rate
175 (3.4 injuries per 1000 tackle events). However, they also found that tackling low (roughly at the upper and
176 lower legs) was the main cause of general injury for the tackler in terms of injury rate (2.2 injuries per 1000
177 tackle events). These findings are consistent with the results from the current paper.

178 The tackle is a dynamic and open phase of play by nature and this must be considered when analysing
179 tackling⁹. This is a limitation of technical video analysis. In some tackle scenarios, tackles were executed
180 because of a defensive system error. The tackler may have been forced to execute a tackle as a result of a
181 teammate's missed tackle or poor positioning in the defensive line. In these circumstances, the tackler may
182 not have optimally identified the ball carrier in a timely fashion as their attention was focused on another
183 opposing player. Therefore, this may have prevented them from reacting to the ball carrier's motion and
184 executing a tackle at the intended tackle height. This highlights the importance of on-field communication
185 and clearly defined defensive systems where players have defined roles and responsibilities. Future work
186 should analyse this aspect of the game.

187 This study utilised an inclusion criterion based on a player being removed for a HIA and not returning to
188 play for the remainder of the game. This could be considered a strong indication of concussion, however, it
189 cannot be fully regarded as a concussion diagnosis. Access to player medical data would have clarified this.
190 This study utilised elite level rugby union games however further research is required to apply these results
191 towards HIA prevention in both youth and amateur level rugby union. To fully understand the concussion
192 injury mechanism further work must study the biomechanics of rugby union head impacts and tackles. This
193 can be achieved by utilising wearable head sensors or model-based approaches^{27,28}. The approach
194 undertaken in this study focused on the tackler as previous literature has reported that the tackler is at most
195 risk of a head impact², HIA^{25,29} and concussion³⁰. However, Cross et al.³⁰ reported that in 30% of cases, it

196 is the ball carrier who sustains a concussion. Furthermore, Tucker et al.²⁹ reported 0.54 HIAs per 1000
197 tackles for the ball carrier in a study with a large HIA sample size (464 tackle-related HIAs). Therefore,
198 further work should assess the effect of tackle height on ball carrier HIA risk. Also, it is possible that
199 lowering the tackle height law could increase the risk of general injury to players.

200 Only tackle height was analysed in this study. Modelling the combination and interaction of other technical
201 characteristics and match situation characteristics, such as tackle speed³⁰, could allow for an even greater
202 understanding of HIA risk. Although the HIA sample size was larger than the injury sample size utilised by
203 Burger et al⁹, the study would have benefited from a larger HIA sample size. The study could be considered
204 underpowered due to the small sample size and this should be considered when interpreting the results. This
205 is a limitation to the data collection approach utilised in this study. Access to open source video data of head
206 impact/HIA/concussion events would have greatly benefited this study.

207 **5. Conclusion**

208 Analysis of match video evidence from elite level rugby union games shows that tackle heights can
209 influence HIA aetiology for the tackler. For front-on upper body shoulder tackles and side-on upper body
210 smother tackles, intended primary contact at the upper trunk of the ball carrier had a greater propensity to
211 result in a HIA for the tackler. However, for side-on upper body smother tackles, intended primary contact
212 at the mid-trunk of the ball carrier had a lower propensity to result in a HIA for the tackler. For front- and
213 side-on lower body shoulder tackles, intended primary contact at the upper leg of the ball carrier had a
214 greater propensity to result in a HIA for the tackler. However, for side-on tackles, intended primary contact
215 at the lower trunk had a lower propensity to result in a HIA for the tackler. To reduce tackler HIA risk, the
216 results suggest tackling below the upper trunk for upper body tackles. The results also support tackling at
217 the lower trunk for lower body tackles and avoiding the upper legs. These findings can be utilised to
218 develop tackle height specific coaching strategies and training drills that place emphasis on tackling lower
219 HIA risk body regions such as the mid- and lower trunk.

220 6. Practical Implications

- 221 • Lowering the tackle height law to below the upper trunk of the ball carrier could reduce the HIA
222 risk during upper body tackles. However, lowering the tackle height law may increase the
223 likelihood of upper leg related HIAs.
- 224 • The results of this study suggest that tackling at the upper legs has a higher propensity to result in a
225 HIA than tackling at the upper trunk. Lowering the tackle height law could subsequently increase
226 the number of tackles to the upper leg region. This could have an adverse effect on HIA reduction.
- 227 • The results suggest tackling below the upper trunk for upper body tackles and tackling at the lower
228 trunk for lower body tackles to reduce HIA risk.
- 229 • These findings can be utilised to develop tackle height specific coaching strategies and training
230 drills that place emphasis on tackling lower HIA risk body regions such as the mid- and lower
231 trunk.

232 7. Acknowledgements

233 8. References

- 234 1. Quarrie KL, Hopkins WG. Tackle Injuries in Professional Rugby Union. *Am J Sports Med*
235 2008;36(9):1705-16. doi: 10.1177/0363546508316768
- 236 2. Tierney GJ, Lawler J, Denvir K, et al. Risks associated with significant head impact events in elite rugby
237 union. *Brain Inj* 2016;30(11):1350-61. doi: 10.1080/02699052.2016.1193630
- 238 3. Rugby Football Union. England Professional Rugby Injury Surveillance Project 2016-17. Available at:
239 [http://www.englandrugby.com/mm/Document/General/General/01/32/91/95/InjurySurveillanceR
240 eport2016-17_English.pdf](http://www.englandrugby.com/mm/Document/General/General/01/32/91/95/InjurySurveillanceReport2016-17_English.pdf). Accessed 17 April 2018.
- 241 4. Brooks JHM, Fuller CW, Kemp SPT, et al. Epidemiology of injuries in English professional rugby
242 union: part 1 match injuries. *Br J Sports Med* 2005;39(10):757-66. doi: 10.1136/bjsm.2005.018135

- 243 5. Fuller CW, Taylor A, Raftery M. Epidemiology of concussion in men's elite Rugby-7s (Sevens World
244 Series) and Rugby-15s (Rugby World Cup, Junior World Championship and Rugby Trophy,
245 Pacific Nations Cup and English Premiership). *Br J Sports Med* 2015;49(7):478-83.
- 246 6. Hendricks S, Lambert M. Theoretical model describing the relationship between the number of tackles in
247 which a player engages, tackle injury risk and tackle performance. *J Sports Sci Med*
248 2014;13(3):715.
- 249 7. Tierney GJ, Denvir K, Farrell G, et al. Does player time-in-game affect tackle technique in elite level
250 rugby union? *J Sci Med Sport* 2017: 1-5. doi: 10.1016/j.jsams.2017.06.023
- 251 8. Tierney GJ, Denvir K, Farrell G, et al. The effect of technique on tackle gainline success outcomes in
252 elite level rugby union. *Int J Sports Sci Coach* 2017:1-10. doi:10.1177/1747954117711866
- 253 9. Burger N, Lambert MI, Viljoen W, et al. Tackle technique and tackle-related injuries in high-level South
254 African Rugby Union under-18 players: real-match video analysis. *Br J Sports Med*
255 2016;50(15):932-38.
- 256 10. Hendricks S, O'Connor S, Lambert M, et al. Contact technique and concussions in the South African
257 under-18 Coca-Cola Craven Week Rugby tournament. *Eur J Sport Sci* 2015;15(6):557-64.
- 258 11. Tierney GJ, Denvir K, Farrell G, et al. The Effect of Tackler Technique on Head Injury Assessment
259 Risk in Elite Rugby Union. *Med Sci Sports Exerc* 2017. doi: 10.1249/MSS.0000000000001461.
- 260 12. Hendricks S, O'Connor S, Lambert M, et al. Video analysis of concussion injury mechanism in
261 under-18 rugby. *BMJ Open Sport Exerc Med* 2016;2(1):e000053.
- 262 13. Tierney GJ, Simms CK. The effects of tackle height on inertial loading of the head and neck in Rugby
263 Union: A multibody model analysis. *Brain Inj* 2017:1-7. doi: 10.1080/02699052.2017.1385853
- 264 14. Fuller CW, Ashton T, Brooks JH, et al. Injury risks associated with tackling in rugby union. *Br J Sports*
265 *Med* 2010;44(3):159-67.
- 266 15. Scher A. The High Rugby Tackle-an Avoidable Cause of Cervical Spinal Injury? *Drug Res* 1978;5:197.
- 267 16. Fuller GW, Kemp SPT, Raftery M. The accuracy and reproducibility of video assessment in the
268 pitch-side management of concussion in elite rugby. *J Sci Med Sport* 2017;20(3):246-49.

- 269 17. McCrory P, Johnston K, Meeuwisse W, et al. Summary and agreement statement of the 2nd
270 International Conference on Concussion in Sport, Prague 2004. *Br J Sports Med*
271 2005;39(4):196-204.
- 272 18. Tierney GJ, Krosshaug T, Wilson F et al. An Assessment of a Novel Approach for Determining the
273 Player Kinematics in Elite Rugby Union Players. IRCOBI Conference proceedings; 2015; Lyon,
274 France.
- 275 19. Montgomery C, Blackburn J, Withers D, et al. Mechanisms of ACL injury in professional rugby union:
276 a systematic video analysis of 36 cases. *Br J Sports Med* 2016;bjsports-2016-096425.
- 277 20. Tierney GJ, Simms CK. Can tackle height influence tackle gainline success outcomes in elite level
278 rugby union? *Int J Sports Sci Coach* 2017: 1-6. doi: 10.1177/1747954117751345
- 279 21. Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics*
280 1977:159-74.
- 281 22. Wilson EB. Probable inference, the law of succession, and statistical inference. *J Am Stat Assoc*
282 1927;22(158):209-12.
- 283 23. Hopkins WG. Linear models and effect magnitudes for research, clinical and practical applications.
284 *Sportsmedicine* 2010;14(1):49-57.
- 285 24. Tierney GJ, Simms CK. The effect of intended primary contact location on tackler head impact risk.
286 IRCOBI Conference proceedings; 2017; Antwerp, Belgium.
- 287 25. Tucker R, Raftery M, Kemp S, et al. Risk factors for head injury events in professional rugby union: a
288 video analysis of 464 head injury events to inform proposed injury prevention strategies. *Br J*
289 *Sports Med* 2017;51(15):1152-57.
- 290 26. Gabbett T, Kelly J. Does fast defensive line speed influence tackling proficiency in collision sport
291 athletes? *Int J Sports Sci Coach* 2007;2(4):467-72.
- 292 27. Tierney GJ, Joodaki H, Krosshaug T, et al. The kinematics of head impacts in contact sport: an initial
293 assessment of the potential of model based image matching. ISBS-Conference Proceedings
294 Archive; 2016; Tsukuba, Japan.

- 295 28. Tierney GJ, Joodaki H, Krosshaug T, et al. Assessment of model-based image-matching for future
296 reconstruction of unhelmeted sport head impact kinematics. *Sports Biomech* 2017:1-15. doi:
297 10.1080/14763141.2016.1271905
- 298 29. Tucker R, Raftery M, Fuller GW, et al. A video analysis of head injuries satisfying the criteria for a head
299 injury assessment in professional Rugby Union: a prospective cohort study. *Br J Sports Med*
300 2017;51(15):1147-51.
- 301 30. Cross MJ, Tucker R, Raftery M, et al. Tackling concussion in professional rugby union: a case-control
302 study of tackle-based risk factors and recommendations for primary prevention. *Br J Sports Med*
303 2017:1-6. doi: 10.1136/bjsports-2017-097912.
- 304
- 305
- 306
- 307

308 **Table 1: The Relative Risk (RR) of tackle heights on HIA aetiology with 95% Confidence Interval**
 309 **(CI) and p-value for front- and side-on upper- and lower body tackles. Percentages reported include**
 310 **95% CI.**

	HIA	Non-HIA	RR (95% CI)	p-value
Upper Body				
Tackles				
	(n=15)	(n=130)		
Front-On				
(Shoulder Tackle)				
Upper Trunk	13 (87%; 62%-96%)	76 (58%; 50%-67%)	1.48 (1.16-1.90)	<0.01
Mid-Trunk	2 (13%; 4%-38%)	54 (42%; 33%-50%)	0.32 (0.09-1.19)	0.09
Side-On (Smother Tackle)				
	(n=15)	(n=148)		
Upper Trunk	14 (93%; 70%-99%)	60 (41%; 33%-49%)	2.30 (1.82-2.92)	<0.01
Mid-Trunk	1 (7%; 1%-30%)	88 (59%; 51%-67%)	0.11 (0.02-0.75)	0.02
Lower Body				
Tackles				
	(n=18)	(n=152)		
Front-On				
(Shoulder Tackle)				
Lower Trunk	6 (33%; 16%-56%)	112 (74%; 66%-80%)	0.45 (0.23-0.88)	0.02
Upper Leg	12 (67%; 44%-84%)	39 (25%; 19%-33%)	2.60 (1.70-3.97)	<0.01
Lower Leg	0 (0%; 0%-18%)	1 (1%; 0%-4%)	2.68 (0.11-63.6)	0.54
Side-On				
(Shoulder Tackle)				
	(n=10)	(n=43)		
Lower Trunk	3 (30%; 11%-60%)	33 (77%; 62%-87%)	0.39 (0.15-1.02)	0.06
Upper Leg	7 (70%; 40%-89%)	9 (21%; 11%-35%)	3.34 (1.65-6.79)	<0.01
Lower Leg	0 (0%; 0%-28%)	1 (2%; 0%-12%)	1.33 (0.06-30.6)	0.86

311

312

313 **Figure Captions**

314 Figure 1: The ball carrier's body split into (a) upper body and (b) lower body regions.

315 Figure 2: The distribution of HIA tackles for front- and side-on Upper Body Tackles (UBT) and Lower
316 Body Tackles (LBT) based on tackle type.

317

318

319

Figure 1
[Click here to download high resolution image](#)









Figure 2
[Click here to download high resolution image](#)



Acknowledgements

The authors would like to thank the Irish Research Council for funding this study.