



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/143775/>

Version: Accepted Version

Article:

McNeill, A. (2019) Movement disorders in adults with 22q11 deletion syndrome. *Movement Disorders Clinical Practice*, 6 (4). p. 339. ISSN: 2330-1619

<https://doi.org/10.1002/mdc3.12752>

This is the peer reviewed version of the following article: McNeill, A. (2019), Movement disorders in adults with 22q11 deletion syndrome. *Mov Disord Clin Pract.*, which has been published in final form at <https://doi.org/10.1002/mdc3.12752>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Movement disorders in adults with 22q11 deletion syndrome

Alisdair McNeill FRCP Edin PhD DCH,

Department of Neuroscience,

The University of Sheffield.

385a Glossop Road.

Sheffield.

S10 2HQ.

a.mcneill@sheffield.ac.uk

word count: 303 words

Key words:

22q11 deletion syndrome, Parkinson's disease

Accepted Article

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/mdc3.12752

Dear Sir,

I read with great interest the recent review article by Boot et al on Parkinson's disease in 22q11 deletion syndrome¹. There is evidence to suggest an increased risk of Parkinson's disease among people with 22q11 deletion syndrome. In our study of 50 adults with 22q11 deletion syndrome, we identified an increased prevalence of potential clinical markers of prodromal neurodegeneration (e.g. hyposmia)². We fully agree with Boot's argument that the core clinical features of 22q11 deletion syndrome may account for the presence of some of these apparent prodromal markers. We also identified a significantly increased prevalence of symptoms of REM sleep behaviour disorder in our cohort. Formal sleep studies in people with 22q11 deletion should be performed to try and identify the prevalence and significance of REM sleep disorder in 22q11. The 22q11 deletion syndrome offers a genetically defined group in which to study prodromal PD and development of the motor features of PD. However, we need to consider which clinical features to use as prodromal markers to make sure we are not simply detecting core clinical features of 22q11 (e.g. hyposmia associated with upper airways malformation).

Chromosome microdeletions frequently cause disease through haploinsufficiency for genes in the deletion region (or deletion of genes at the breakpoint)^{3,4}. There are several interesting candidates within the 22q11 deletion; involved in dopamine metabolism, mitochondrial function or microRNA processing. No single nucleotide variants in any of these candidate genes have been reported in PD cases to my knowledge. The deletion is due to non-allelic homologous recombination (NAHR) and therefore is of a consistent size, precluding identification of a critical region. The molecular mechanisms explaining increased PD risk in 22q11 remain open for exploration. Once again it will be crucial to disentangle mechanisms which are core to the 22q11 deletion phenotype from those which might explain increased PD risk.

References

1. Boot E, Bassett AS, Marras C. 22q11.2 deletion syndrome-associated Parkinson's disease. *Mov Disord Clin Pract*; <https://doi.org/10.1002/mdc3.12687>
2. Buckley E, Siddique A, McNeill A. Hyposmia, symptoms of rapid eye movement sleep behaviour disorder, and parkinsonian motor signs suggest prodromal neurodegeneration in 22q11 deletion syndrome. *Neuroreport* 2017; 28: 677-681.
3. Hempel A, Pagnamenta AT, Blyth M, et al. Deletions and de novo mutations of SOX11 are associated with a neurodevelopmental disorder with features of Coffin-Siris syndrome. *J Med Genet* 2016; 53: 152-162.
4. Blanchet P, Bebin M, Bruet S, et al. MYT1L mutations cause intellectual disability and variable obesity by dysregulating gene expression and development of the neuroendocrine hypothalamus. *PLoS Genet* 2017; 13: e1006957.

Disclosures:

Funding sources and conflicts of interest: No specific funding for this work.

Financial disclosures for previous 12 months: Salary support from University of Sheffield and NHS in the last 12 months.

Author Roles

Alisdair McNeill 1) Research project: A. Conception, B. Organization, C. Execution;

3) Manuscript: A. Writing of the first draft, B. Review and Critique.

Ethical Compliance Statement:

The authors confirm that the approval of an institutional review board was not required for this work. Informed consent was not obtained as no patients are described in this manuscript. We confirm that we have read the Journal's position on issues involved in ethical publication and affirm that this work is consistent with those guidelines.