



UNIVERSITY OF LEEDS

This is a repository copy of *Effects of Uterine Cells-Conditioned Media on Expression of DNMT3B and DNMT3C in Mouse Embryos Cultured in a Microfluidic Device.*

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/143902/>

Version: Accepted Version

Proceedings Paper:

Mancini, V, Lu, J, Colucci, F et al. (4 more authors) (2019) Effects of Uterine Cells-Conditioned Media on Expression of DNMT3B and DNMT3C in Mouse Embryos Cultured in a Microfluidic Device. In: Reproductive Sciences. 66th Annual Meeting of the Society for Reproductive Investigation, 12-16 Mar 2019, Paris, France. SAGE Publications , 377A-377A.

<https://doi.org/10.1177/1933719119834079>

© 2019, Author(s). This is an author produced version of a paper published in Reproductive Sciences. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Effects of uterine cells-conditioned media on expression of DNMT3 β and DNMT3C in mouse embryos cultured in a microfluidic device

V. Mancini¹, J. Lu², F. Colucci¹, P.J McKeegan², J.D. Huntriss², H.M Picton², V. Pensabene^{1,2}

¹School of Electronic and Electrical Engineering, University of Leeds, Leeds, UK, LS2 9JT;

²Reproduction and Early Development Research Group, Discovery and Translational Science Department, Leeds Institute of Cardiovascular and Metabolic Medicine, School of Medicine, University of Leeds, UK, LS2 9JT;

KEYWORDS: microfluidics, in vitro culture, real-time PCR

INTRODUCTION:

DNA methylation is an epigenetic mechanism that plays an important role in early embryo development in mammals. We hypothesize that the de novo DNA methyltransferase genes, DNMT3 β and DNMT3C, can be affected by medium composition during in vitro mouse embryo development. Herein, we describe the culture of murine embryos in a microfabricated polydimethylsiloxane (PDMS) device and we investigated the effect of cells-conditioned media on gene expression of individual blastocysts by qPCR.

METHODS: The microfluidic device was fabricated in PDMS using traditional soft-lithographic technique. Groups of 12, 1-cell murine embryos (B6C3F1xB6D2F1 strain, EmbryoTech, USA) were cultured in mouse uterine epithelial cells-conditioned media into a microfluidic device (n=15 devices, 180 embryos). Control embryos were cultured in the device using control media (n=15 devices, 180 embryos). We analysed expression of two genes involved in the de novo DNA methylation process, DNA methyltransferase 3 β (DNMT3 β) and DNA (cytosine-5-)-methyltransferase 3c (DNMT3C), in blastocysts cultured in cells-conditioned media compared with those cultured in control media. To determine transcript abundance of the selected genes we performed qPCR of cDNAs generated from individual, stage matched late-blastocysts cultured in the microfluidic device.

RESULTS: From the analysis of qPCR results we observed that both DNMT3 β (n=12, p=0.006) and DNMT3C (n=12, p=0.048) were highly expressed in blastocysts cultured in cell-conditioned media compared with those cultured in control media. The upregulation of DNMT3 β and DNMT3C suggests that the use of cells-conditioned media might be an efficient approach to improve quality of embryos produced in vitro. However, the expression pattern of those genes may vary during embryonic development and our findings need to be confirmed by investigating transcript abundance in blastocyst from different stages.

CONCLUSIONS: We conclude that mRNA relative expression levels of DNMT3 β and DNMT3C were significantly affected by the use of cells-conditioned media during preimplantation in vitro development of mouse embryo in a microfluidic device. Further studies will be required to elucidate the effect of this variation in mouse embryo development.