

# **Dietary protein and bone health across the life-course: an updated systematic review and meta-analysis over 40 years**

**Andrea L. Darling<sup>1</sup>, Ralph J. F. Manders<sup>1</sup>, Shivani Sahni<sup>2-3</sup>, Kun Zhu<sup>4-5</sup>, Catherine E. Hewitt<sup>6</sup>, Richard L. Prince<sup>4-5</sup>, D. Joe Millward<sup>1</sup> and Susan A. Lanham-New<sup>1</sup>**

## **Author affiliations and addresses:**

<sup>1</sup> Department of Nutritional Sciences, School of Biosciences and Medicine, University of Surrey, Guildford, United Kingdom

<sup>2</sup> Hinda and Arthur Marcus Institute for Aging Research, Hebrew SeniorLife, 1200 Centre St, Boston MA 02131, USA

<sup>3</sup> Department of Medicine, Beth Israel Deaconess Medical Center, Harvard Medical School, 330 Brookline Ave, Boston, MA 02215, USA

<sup>4</sup>Department of Endocrinology and Diabetes, Sir Charles Gairdner Hospital, Nedlands, Western Australia

<sup>5</sup>Medical School , University of Western Australia, Crawley, Western Australia

<sup>6</sup> York Trials Unit, Department of Health Sciences, University of York, York, United Kingdom

## **ORCID ID:**

ALD 0000-0002-7038-3161

RJFM 0000-0003-0199-2120

SS 0000-0002-9827-0986

KZ 0000-0002-8723-7574

CEH 0000-0002-0415-3536

RLP None

DJM 0000-0003-0736-8396

SLN 0000-0003-3153-6345

**Word Count:** 4131 (excluding title page, abstract, tables, references), 4 tables and 2 figures

**Corresponding author:** Dr Andrea Darling, Department of Nutritional Sciences, School of Biosciences and Medicine, Faculty of Health and Medical Sciences, University of Surrey, Guildford, GU2 7XH +44 (0)1483 689222 email: [a.l.darling@surrey.ac.uk](mailto:a.l.darling@surrey.ac.uk)

**Acknowledgements:** This work received no funding. The authors are grateful to Dr Helen Lambert for her help with translation of papers.

**Conflict of Interest Statement:** SS has received institutional grants from Dairy Management Inc. and she serves as a member of the National Dairy Council's Nutrition Research Scientific Advisory Committee. SLN has received consultancy fees from the following companies/organisations: National Dairy Council UK, Yoplait, Kelloggs, Danone, Nestle. All other authors have nothing to declare.

**Authors' contribution:** ALD, RM and S-LN designed the research: ALD, RM and S-LN conducted the research. ALD analysed data. ALD wrote the paper. ALD and S-LN have primary responsibility for final content. ALD, CEH, DJM, KZ, RM, RP, SS, S-LN interpreted data and aided with statistical analyses. ALD, CEH, DJM, KZ, RM, RP, SS and S-LN revised manuscript content for intellectual content. ALD, CEH, DJM, KZ, RM, RP, SS and S-LN approved the final version of manuscript.

**Awards:** ALD received a Young Investigator Travel Award from the 10th International Symposium on the Nutritional Aspects of Osteoporosis Conference in Hong Kong, 28th November to 1st December 2017 for this work.

**Please cite this article in press as:** Darling AL et al. (2019) Dietary protein and bone health across the life-course: an updated systematic review and meta-analysis over 40 years. *Osteoporosis International* (in press).

**Link to publisher version:** <http://link.springer.com/article/10.1007/s00198-019-04933-8>

**DOI :** 10.1007/s00198-019-04933-8

## 1 **Abstract**

2 **Purpose:** This systematic review and meta-analysis analysed the relationship between dietary  
3 protein and bone health across the life-course.

4 **Methods:** The PubMed database was searched for all relevant human studies from the 1st  
5 January 1976 to 22nd January 2016, including all bone outcomes except calcium metabolism.

6 **Results:** The searches identified 127 papers for inclusion, including 74 correlational studies,  
7 23 fracture or osteoporosis risk studies and 30 supplementation trials. Protein intake  
8 accounted for 0 - 4% of areal BMC and areal BMD variance in adults and 0-14% of areal  
9 BMC variance in children and adolescents. However, when confounder adjusted (5 studies)  
10 adult lumbar spine and femoral neck BMD associations were not statistically significant.

11 There was no association between protein intake and relative risk (RR) of osteoporotic  
12 fractures for total ( $RR_{(random)} = 0.94$ ; 0.72 to 1.23,  $I^2=32\%$ ), animal ( $RR_{(random)} = 0.98$ ; 0.76 to  
13 1.27,  $I^2 = 46\%$ ) or vegetable protein ( $RR_{(fixed)} = 0.97$  (0.89 to 1.09,  $I^2 = 15\%$ ). In total protein  
14 supplementation studies, pooled effect sizes were not statistically significant for LSBMD  
15 (total n=255,  $MD_{(fixed)}=0.04$  g/cm<sup>2</sup> (0.00 to 0.08, P=0.07),  $I^2=0\%$ ) or FNBMD (total n=435,  
16  $MD_{(random)}=0.01$  g/cm<sup>2</sup> (-0.03 to 0.05, P=0.59),  $I^2=68\%$ ).

17 **Conclusions:** There appears to be little benefit of increasing protein intake for bone health in  
18 healthy adults but there is also clearly no indication of any detrimental effect, at least within  
19 the protein intakes of the populations studied (around 0.8-1.3 g/Kg/day). More studies are  
20 urgently required on the association between protein intake and bone health in children and  
21 adolescents.

22

23 **Key Words:** Aging, Epidemiology, IGF-1, Nutrition, Osteoporosis, Diet

24

25 **Mini Abstract:** We undertook a systematic review and meta-analysis of published papers  
26 assessing dietary protein and bone health. We found little benefit of increasing protein intake  
27 for bone health in healthy adults but no indication of any detrimental effect, at least within the  
28 protein intakes of the populations studied.

29

30

## 31 **Introduction**

32

33 The association between dietary protein intake and bone health has been debated worldwide  
34 for decades. It is unclear as to whether dietary protein exerts a positive or detrimental  
35 influence on bone and there are competing theories as to the effects of dietary protein on bone  
36 health. Proponents of a positive link between protein intake and bone health cite the known  
37 anabolic effects of dietary protein on bone, via dietary protein's known ability to increase  
38 secretion of insulin-like growth factor 1 (IGF-1). Dietary protein may also increase calcium  
39 absorption from the gut [1], which is likely to be beneficial for bone mineralisation.

40 On the other hand, proponents of a detrimental association between protein intake and bone  
41 argue that high protein intakes may be bad for bone health. This is on the basis of known in-  
42 vitro increased osteoclast activity with increasing body acidity[2] and predicted subsequent  
43 bone demineralisation. However, it is important to consider dietary protein type as only  
44 proteins that are rich in sulphur amino acids are likely to increase net physiological acid  
45 production. Also, protein is consumed in the diet with other components which may modify  
46 the overall net acid concentration (e.g. fruits and vegetables which have an alkalizing effect,  
47 soy isoflavones which have estrogenic effects, and dairy products which contain calcium).  
48 These nutrients may modify the association between protein intake and bone health [3, 4].

49 A previous systematic review and meta-analysis published in 2009, covering published papers  
50 from January 1966 to July 2008 [5], showed a small beneficial effect of dietary protein  
51 supplementation on areal bone mineral density (aBMD) in human adults, as well as showing  
52 that nearly all published cross-sectional studies demonstrate a positive association between  
53 dietary protein intake and bone health. However, there was no association between dietary  
54 protein intake and fracture risk in cohort studies, and much of the findings of the cross-  
55 sectional studies may be due to inadequate controlling for confounding factors. Also, the

56 previous systematic review and meta-analysis did not assess dietary protein and bone health  
57 in children and adolescents, which is an important area of research due to the known  
58 importance of obtaining peak bone mass at this time of life.

59 Since the publication of the dietary protein and bone health systematic review and meta-  
60 analysis in 2009 [5] the number of relevant articles on the topic of dietary protein and bone  
61 health has doubled. This highlights the need for an updated systematic review of this topic,  
62 including consideration of peak bone mass development in children and adolescents, as well  
63 as bone health in adults. Further systematic reviews have been conducted on the topic since  
64 2009 [6-10], but none to date have included all study designs (cross-sectional, longitudinal  
65 and randomized control trials) and included both adults and children. Our objective was to  
66 perform a systematic review and meta-analysis of the evidence published over the last 40  
67 years assessing dietary protein intake and all indices of bone health across the human life-  
68 course, including all relevant observational and intervention studies examining intakes of all  
69 types of protein.

70

## 71 **Materials and Methods**

### 72 Search Strategy

73 The PUBMED databases, as well as reference lists of relevant journal papers were searched to  
74 ensure broad coverage. The search phrase used for the electronic search was “(protein intake  
75 OR dietary protein OR protein supplement OR protein consumption) AND (bone OR fracture  
76 OR BMD OR bone turnover)” limited to human studies, in any language, from 1<sup>st</sup> January  
77 1976 to 22<sup>nd</sup> January 2016. Two of the authors (ALD, SLN) identified potential studies for  
78 inclusion by first screening titles and abstracts, and then looking at the full version of papers  
79 if necessary. Any disagreements were resolved by consensus and deferred to a third party if  
80 required.

81

82 Study Design and Criteria

83 Papers assessing relevant bone outcomes (specifically BMD or BMC, bone turnover markers  
84 and/or fracture risk) in healthy human adults, adolescents or children were considered for  
85 inclusion. Studies investigating subjects with a pre-existing medical condition affecting bone  
86 or calcium metabolism, or studying animals, infants, and pregnant or lactating women were  
87 excluded as were studies involving only calcium balance or calcium metabolism.

88 All cross-sectional, longitudinal and intervention studies were eligible for inclusion, including  
89 intervention studies of any design. However, due to problems of statistically combining  
90 crossover and parallel intervention designs, crossover studies were not included in the meta-  
91 analysis and were discussed in the systematic review only. Studies were excluded if they were  
92 published before 1975, were weight loss studies, did not have usable protein or bone data,  
93 were not on the relevant topic, were review articles or correspondence, or only had data on  
94 dietary patterns. They were also excluded if they had confounders present in the central study  
95 design (e.g. differing exercise levels in the intervention and control groups as part of the  
96 intervention, for example protein and exercise as intervention vs. no protein and exercise). All  
97 bone sites in the body were eligible for inclusion in the review, as were all markers of bone  
98 metabolism.

99 Soy protein studies were excluded if they did not have data on isoflavone devoid soy protein  
100 isolate. Studies on whole dietary measures (e.g. meat intake, frequency of high protein meals)  
101 were also excluded due to the confounding potential of other dietary constituents (such as  
102 isoflavones in soy protein; fat or iron in meat). However, whole diet intervention studies were  
103 included as long as the nutritional contents of the diets were shown in the paper and the diets  
104 had been designed to minimise the impact of confounders (e.g. ensuring high and low protein



105 diets were isocaloric, similar in dietary calcium etc.). No ethical approval was required as the  
106 analysis was conducted only on already published data.

107

### 108 Data synthesis

109 See Online Resource for full details of the data extraction process. In the meta-analyses, two  
110 sided tests with an alpha value of 5% were used, and Bonferroni adjustment for multiple  
111 testing was applied when necessary.

112 The systematic review included all eligible studies, but only studies with data suitable for  
113 meta-analysis were analysed quantitatively. The R software (version 3.1.2) [11] add-on  
114 packages ‘metacor’[12], ‘meta’[13], ‘metafor’[14] and ‘metagen’[15] were used to pool  
115 correlation coefficients, intervention and fracture risk data for meta-analysis, as well as  
116 creating all plots. For cross-sectional and longitudinal data, multivariate adjusted analyses  
117 were used wherever possible, rather than unadjusted or age adjusted measures. Linear and  
118 non-linear modelling was undertaken to assess whether there was an association between  
119 correlation coefficients or actual BMD values with the following: calcium: protein ratio  
120 (mg/g/d); protein dose (g/kg/d) and calcium dose (mg/kg/d).

121 It was planned *a priori* to run a combined analysis of hip and non-hip fracture data as all  
122 fracture outcome first, then a subsequent analysis of hip fracture data only. Unfortunately the  
123 calcium: protein ratios (calcium mg/protein g) in the fracture risk studies were too constant  
124 (10-12 mg/g/d) to be able to perform an assessment of calcium: protein ratio vs. fracture risk.

125 Also, there were too few studies within the same protein type to be able to plot protein or  
126 calcium dose against fracture risk.

127 The intervention trials meta-analysis examined the main effects of protein supplementation  
128 (all protein types but excluding studies specifically looking at soy protein isolate and milk  
129 basic protein (MBP)) on BMD, BMC and bone turnover markers, based on mean differences.

130 The decision was made *a priori* to also perform a separate analysis for MBP trials as this type  
131 of protein is highly concentrated and is supplemented in very high doses. Similarly,  
132 experience from the previous meta-analysis suggested that soy protein studies need to be  
133 analysed separately than non-soy studies, as soy studies usually assess the effect of soy  
134 protein vs. another protein type (hence the actual protein intake is the same, just different  
135 protein type). Conversely, studies of mixed protein supplementation usually compare higher  
136 with lower doses of protein (or have another type of control, e.g. maltodextrin), hence the  
137 protein intake varies in each arm of the study.

138 Finally, due to the differences in skeletal biology between early life and in older age, as well  
139 as differences in nutritional requirements, children and adults were analysed separately in all  
140 meta-analyses. Also, all adult groups were broken down into gender and life stage specific  
141 analyses wherever there was enough data to do so. Only for the cross-sectional studies were  
142 there enough data to perform a meta-analysis in children.

143

#### 144 Heterogeneity, Sensitivity and Publication Bias- meta-analysis

145 It was pre-specified that the  $I^2$  statistic was to be used to assess heterogeneity between studies  
146 as this is recommended for analyses with smaller numbers of studies[16].  $I^2$  values of 25%,  
147 50% and 75% were considered low, moderate and high, respectively[16]. Random-effects  
148 (heterogeneous comparisons) and fixed effects (homogenous comparisons) models were used  
149 accordingly, with the decision based on size of  $I^2$  (%), rather than the  $P_{(\text{heterogeneity})}$  value from  
150 the Q test, as number of studies in the meta-analyses were predicted to be small. It was also  
151 pre-specified that funnel plots were used to assess possible biases in meta-analyses containing  
152 10 or more studies[17]. Sensitivity analyses were also planned to assess the impact of removal  
153 of each study in turn on the pooled effect size.

154

155 Quality Analysis

156 Intervention study quality analysis was undertaken at the study level. This was conducted  
157 independently by 2 authors (AD, SLN) using the Jadad Scale[18]. Due to the extensive  
158 number of observational studies, only one author (AD) assessed quality for these papers,  
159 using the Newcastle-Ottawa adapted scales [19] for cohort and case-control study designs as  
160 appropriate. Cross-sectional studies were not assessed for study quality due to the very large  
161 number of papers included in the review. For all study types, due to the small predicted  
162 numbers of potential studies for the fracture and intervention study meta-analyses, we decided  
163 *a priori* that the quality analysis information was to be used to guide result interpretation  
164 rather than to exclude studies *per se*.

165

166

## 167 **Results**

168

169 **Figure 1** shows the Quality of Reporting of Meta-analyses (QUOROM) flow diagram[20]  
170 illustrating the search and selection process. The systematic review included 127 studies,  
171 including 74 studies reporting correlation or regression coefficients (**Table 1**). These included  
172 either just cross-sectional data (55 studies), just longitudinal change data (4 studies) or both (9  
173 studies). These also included 2 studies reporting both longitudinal change in BMD data and  
174 fracture data and 4 studies reporting cross-sectional data and odds of low BMD or  
175 osteoporosis diagnosis (**Table 2**). In addition there were 23 studies reporting only fracture or  
176 osteoporosis risk (**Table 3**), and 30 intervention studies (**Table 4**). Due to the extensive  
177 number of studies to be discussed here, see the Online Resource for the systematic review and  
178 the study quality assessments.

179

### 180 Pooling of studies reporting correlation or regression coefficients

#### 181 *By population subgroup*

182 See Table 1 (and **Online Resource Table S1**) for details of the studies reporting correlation  
183 or regression coefficients. Of these 74 studies, 30 studies in adults [21-50] and 5 studies in  
184 children [51-55] gave cross-sectional R correlation coefficients suitable for pooling. Of note,  
185 only 5 of these studies [22, 24, 25, 36, 56] reported adjusted data, so the majority (n=30) were  
186 non-adjusted for confounders. When pooling by population subgroup the R<sup>2</sup> values suggested  
187 that <1-8% of adult bone BMD or BMC (depending on age and biological sex) as well as  
188 10% of child BMC was explained by protein intake (**Online Resource Tables S2-S3**, Online  
189 Resource Material Text). Heterogeneity was highly variable with values ranging from 0-74%  
190 depending on population subgroup.

191

192 *By bone parameter*

193 For adults, **Online Resource Figures 1-2** illustrate the pooled effect size for FNBMD and  
 194 LSBMD. FNBMD (total n=4786) showed  $r_{(fixed)}=0.07$  (0.04 to 0.09)  $R^2=0.005$  (0.5%)  
 195  $P<0.0001$ ,  $I^2=26\%$   $P_{(heterogeneity)}=0.15$ . LSBMD (total n=4257) showed  $r_{(random)}=0.09$  (0.04 to  
 196 0.14)  $R^2=0.008$  (0.8%)  $P<0.001$ ,  $I^2=58\%$   $P_{(heterogeneity)}=0.001$ . Of note, when only studies  
 197 reporting multivariate adjusted data were shown, the pooled effect sizes for FNBMD (total  
 198 n=1169,  $r_{(random)}=0.04$  (-0.05 to 0.12)  $R^2=0.0016$  (0.2%)  $P=0.37$ ,  $I^2=45\%$   $P_{(heterogeneity)}=0.14$ ),  
 199 and for LSBMD (total n=728,  $r_{(fixed)}=0.0$  (-0.07 to 0.07),  $R^2=0$  (0%)  $P=0.97$ ,  $I^2=0\%$   
 200  $P_{(heterogeneity)}=0.61$ ) were no longer statistically significant. Other bone outcomes ranged in  $R^2$   
 201 value from 0-10% (including unadjusted data). See Online Resource Material for full results  
 202 of pooling for other bone sites in adults. Heterogeneity was highly variable with values  
 203 ranging from 0-94% depending on adult bone site.

204 In children and adolescents, the  $R^2$  values suggested that 0-14% of child/adolescent BMC and  
 205 21% child/adolescent total body bone area (TBBA) were explained by protein intake (all  
 206 unadjusted data). Heterogeneity was 0% (All BMC and total body bone mineral content  
 207 (TBBMC)), 79% (TBBA) and 87% total body bone mineral density (TBBMD)). See Online  
 208 Resource Material for further details of all the above correlational analyses including  
 209 sensitivity analyses and funnel plots, as well as associations with protein and calcium dose,  
 210 and calcium: protein ratio (**Online Resource Table S4**).

211

212 Fracture risk meta-analysis

213 See Table 3 and **Online Resource Table S5** for details of all studies presenting fracture or  
 214 osteoporosis risk data. Of 29 studies (6 of which have already appeared in correlational  
 215 section of review), 5 studies were pooled in the meta-analysis for cohort studies assessing  
 216 relative risk of fracture [57-61] (**Figure 2**), all of which provided multivariate adjusted risk

217 estimates. No statistically significant association was found between total protein intake and  
218 relative risk (RR) of all fractures ( $RR_{(random)}=0.94$  (0.72 to 1.23,  $P=0.55$   $n=4$  studies,  $I^2=32\%$   
219  $P_{(heterogeneity)}=0.30$ ). Similar results were seen for animal protein intake,  $RR_{(random)}=0.98$  (0.76 to  
220 1.27,  $P=0.87$ ,  $n=4$  studies,  $I^2=46\%$   $P_{(heterogeneity)}=0.13$ ) and vegetable protein intake:  
221  $RR_{(fixed)}=0.97$  (0.89 to 1.09,  $P=0.61$ ,  $n=3$  studies,  $I^2=15\%$   $P_{(heterogeneity)}=0.31$ ). For three cohort  
222 studies reporting hazard ratios (HR) (**Online Resource Figure S3**)[62-64], no significant  
223 association was found between total protein intake and HR for all fractures ( $HR_{(random)}=0.82$   
224 (0.59 to 1.14,  $P=0.24$ ),  $n=4$  studies (5 data points as 2 studies had independent subgroups  
225 which can both be entered),  $I^2=35\%$   $P_{(heterogeneity)}=0.19$ ). Of note, removal of the Sahni (Low  
226 calcium) data[64] led to a reduced risk of fracture with increased protein intake (HR  
227 (fixed)=0.79 (0.64 to 0.97,  $P=0.02$ ),  $n=4$  studies,  $I^2=0\%$   $P_{(heterogeneity)}=0.66$ ), which would  
228 be of borderline statistical significance after adjustment for multiple testing (using a cut-off of  
229 0.02 for 3 fracture meta-analyses). For three case-control studies reporting odds ratio (OR) for  
230 fracture[65-67] (**Online Resource Figure S4**) there was no significant association between  
231 total protein intake and odds of fracture: ( $OR_{(random)}=0.69$  (0.30 to 1.58,  $P=0.38$ ),  $n=4$  studies  
232 (4 data points as 1 study had independent subgroups which can both be entered)  $I^2=65\%$   
233  $P_{(heterogeneity)}=0.03$ ). See Online Resource Material for details of relevant sensitivity analyses.

234

### 235 Intervention Studies Meta-analysis

236 Thirty papers were intervention studies (Table 4, **Online Resource Table S6**). See Online  
237 Resource Material for details of study quality assessment for these studies as well as  
238 systematic review of intervention studies.

239

240

241

242 *Meta-analysis- summary of included studies*

243 Nine of the 30 intervention studies [68-76] were suitable for meta-analysis. The other 20  
 244 studies were excluded due to missing data [77], cross over design study with no report of data  
 245 prior to crossover [1, 78-85], and not having data that is compatible with other studies to  
 246 allow meta-analysis [86-93], being a food-based intervention [94] or presenting data for  
 247 change in bone indices only[95, 96].

248

249 *Total protein intake*

250 The pooled effect size for protein supplementation on LSBMD[68, 72] showed no statistically  
 251 significant effect ( $P=0.07$ ) (total  $n=255$ ,  $MD_{(fixed)}=0.04$  (0.00 to 0.08,  $P=0.07$ ),  $I^2=0\%$   
 252  $P_{(heterogeneity)}=0.47$ ; **Online Resource Figure S5**) Equivalent data for FNBMD (3 studies)[68,  
 253 72, 76], were also not statistically significant (total  $n=435$ ,  $MD_{(random)}=0.01$  (-0.03 to 0.05,  
 254  $P=0.59$ )  $I^2=68\%$   $P_{(heterogeneity)}=0.04$ )(**Online Resource Figure S6**).

255

256 *MBP and Soy Protein*

257 For MBP studies, comparing MBP supplementation with a nutrient matched drink [69, 70] or  
 258 no control[71], the pooled effects sizes were as follows: LSBMD Mean Difference  
 259  $(MD)_{(fixed)}= 0.02$  (0.00-0.04,  $P=0.08$ ),  $I^2=0\%$   $P_{(heterogeneity)}=0.87$ , 3 studies) (**Online Resource**  
 260 **Figure S7**). For soy protein studies no statistically significant effect was found for LSBMD  
 261  $(MD_{(random)} = -0.01$  (-0.07 to 0.06,  $P=0.82$ ),  $I^2=51\%$   $P_{(heterogeneity)}=0.13$ ) , FNBMD  $(MD_{(random)}$   
 262  $=0.01$  (-0.06 to 0.07,  $P=0.87$ ),  $I^2=74\%$   $P_{(heterogeneity)}=0.05$ ) or BAP  $(MD_{(random)} = -1.75$  (-10.50  
 263 to 7.01,  $P=0.70$ ),  $I^2=91\%$   $P_{(heterogeneity)}=0.0009$ ). There were not enough studies with  
 264 compatible data to assess any other bone sites or bone markers for soy protein or MBP  
 265 supplementation.

266 Overall heterogeneity was low for the influence of MBP on LSBMD at  $I^2=0\%$  but high for  
267 soy protein LSBMD, FNBMD and BAP ( $I^2=51\%$ ,  $71\%$  and  $91\%$  respectively). For both  
268 MBP and soy protein sensitivity analyses showed that removal of each study in turn had no  
269 effect on the pooled effect sizes (see Online Resource Material). The results for the soy and  
270 MBP subgroup analyses are unlikely to remain statistically significant after Bonferroni  
271 adjustment for multiple testing.

272

273



**274 Discussion**

275 This systematic review and meta-analysis reviewed evidence, published over the last 40 years,  
276 assessing the association between dietary protein and bone health across the life-course.

277 Dietary protein intake explained 0 to 4% of adult BMD, BMC and bone markers, as well as 0-  
278 14% of child/adolescent BMC and 21% child/adolescent TBBA. However, when only studies  
279 adjusted for confounders (5 studies) were included in the adult analyses for FNBMD and  
280 LSBMD these effects did not remain statistically significant, suggesting the effect sizes seen  
281 could be caused by potential confounding. The larger pooled effect sizes for the children and  
282 adolescents suggest a stronger relationship between protein and bone health at this life stage  
283 than in later life. Indeed, nutritional requirements for skeletal growth in childhood and  
284 adolescence are different to those for reducing senescence related bone loss, as the cellular  
285 mechanisms involved differ at these two life stages. Protein intake may have a strong  
286 relationship with bone health in childhood due to the involvement of amino acids in  
287 endochondral ossification during bone growth. However, this finding of a stronger association  
288 between protein intake and bone health in children may also be an artefact of the smaller  
289 number of participants included in the pooled effect sizes in the child and adolescent analyses  
290 compared with adults.

291 In adults, neither calcium: protein ratio [calcium (mg)/protein (g /d)] nor protein (g/kg/d) or  
292 calcium (mg/kg/d) dose were associated with strength of effect for the association between  
293 protein intake and bone health. As seen in the previous 2009 systematic review and meta-  
294 analysis, there was no association between dietary protein intake and fracture risk in cohort  
295 and case-control studies, for total protein, animal or vegetable protein. In intervention  
296 studies, no statistically significant effects of total protein supplementation (excluding MBP  
297 and soy protein) were seen for FNBMD and LSBMD, or for higher vs. lower soy protein  
298 supplementation.

299 The lack of an effect of dietary protein on bone seen in the above fracture risk and  
300 supplementation trials may be for a variety of reasons. It could be that the opposing anabolic  
301 and catabolic actions of dietary protein cancel each other out to some extent. Alternatively  
302 dietary protein, at the protein intakes seen in the populations in this review, may not affect  
303 bone health and it may be that there is a small window whereby very low protein intakes are  
304 detrimental to bone, due to low IGF-1 production and dietary calcium absorption from the  
305 gut. Few studies in the review contained populations with low population intakes, with most  
306 studies reporting intakes of over 0.75g/Kg/d-0.84g/Kg/d which is considered adequate by  
307 western countries such as Australia [97], Europe [98] and the US [99]. Some studies that  
308 contained older people or Buddhist Nuns had intakes nearer to, or slightly lower than,  
309 0.75g/Kg/d [66, 95, 100-102]. Twelve of the trials had no baseline protein intake data which  
310 makes it difficult to fully assess the degree of sufficiency. However, observational studies  
311 included this information with only one observational study not reporting overall protein  
312 intake [42]. In only 2 of the childhood studies was there a deficient protein intake [53, 92],  
313 and these were studies of malnourished children so this is as would be expected. However,  
314 none of the other child or adolescent studies showed deficient protein intakes.

315 Equally, other aspects of the diet that the authors of the original studies did not control for in  
316 the analysis (e.g. calcium intake) may be affecting the results. Of note, the study by Sahni et  
317 al. (2010) [64] showed a higher risk of hip fracture with higher protein intake when calcium  
318 intake was low (<800 mg/d) but conversely a lower risk of fracture with higher protein intake  
319 when calcium was higher ( $\geq$ 800 mg/d). Failure to control for calcium intake in the fracture  
320 analyses could therefore obscure any relationship between protein intake and fracture risk. Of  
321 importance, the meta-analysis for cohort studies reporting hazard ratios was borderline  
322 statistically significant for a reduced risk of fracture with increasing protein intake, when the  
323 low calcium arm of the study by Sahni et al. 2010 [64] was removed, suggesting this study

324 arm may be masking an association between protein intake and fracture which was evident  
325 when only the high calcium arm was included. Moreover, the meta-analysis result for the  
326 case-control studies (hazard ratios, see Online Resource Figure 4) would have been  
327 statistically significant if using fixed models, which were not used due to heterogeneity  
328 present in our analysis. This may explain some of the differing results of our study from other  
329 meta-analyses, in that other analyses have combined data from cohort and case-control  
330 studies, which we did not, and the results of other meta-analyses may only be significant  
331 because of the inclusion of case-control data.

332 Since the publication of our original meta-analysis in 2009 [5] there have been a series of  
333 meta-analyses published on the subject of dietary protein and bone [6-9]. However, these  
334 differ from our meta-analysis in significant ways. Wu et al. (2015) only assessed fracture  
335 risk, and Wallace and Frankenfeld (2017) did not assess cross-sectional studies. Santesso et  
336 al. (2012) only included bone data from weight loss trials and Shams-White et al. (2017) [6]  
337 included a mixture of weight loss and non-weight loss trials, which may partially explain their  
338 finding of a protective effect of higher protein intake against LSBMD loss. Moreover, only  
339 one study in the latter review's change in LSBMD meta-analysis showed a positive effect, so  
340 the analysis was reliant on this one study for the significant result. The most recent meta-  
341 analysis, Shams-White (2018)[10] included only studies on animal protein vs. soy protein  
342 with isoflavones, so differs substantially from our current review. Finally, none of these meta-  
343 analyses assessed data from children and adolescents.

344 There have also been 13 relevant protein and bone health papers [103-115] published since  
345 our electronic searches were completed. All of these studies have found either a positive  
346 association, or no association of protein take with bone health, except for one study which  
347 found poorer bone health with increased protein intake [107]. All these studies were of  
348 observational study design except for two randomised control trials; one of soy protein in

349 children which found no effect on bone marker concentration [110] and one of collagen  
350 supplementation in postmenopausal women which found a positive effect of supplementation  
351 on aBMD [108]. Overall, our current meta-analysis reported here is the most comprehensive  
352 to date on the subject and the inclusion of studies published since 2016 are unlikely to have  
353 significantly changed our results. As stated above, the separation of cohort and case-control  
354 studies may explain differences from our study to that of other meta-analyses.

355 The strengths of this analysis include the fact that we included papers published in both  
356 English and non-English languages, and from both observational and intervention studies.  
357 Moreover, inclusive coverage both in terms of timescale (i.e. 1975 to the present day; 40  
358 years), topic area (all types of protein, BMD, BMC, BA and bone turnover markers) and  
359 population age subgroups (children, adolescents, adults, older adults) was achieved. This  
360 makes this analysis a comprehensive overview of dietary protein and bone health across the  
361 life-course, using data from the last 40 years. This makes the work novel and of importance  
362 to scientists, public health providers and clinicians alike.

363 However, the review was limited by the following factors. First, studies did not vary enough  
364 in calcium intake, as well as too few studies, to be able to assess whether the association  
365 between dietary protein and fracture risk varies by calcium intake. Second, there were  
366 concerns due to the low methodological quality of some intervention and cross-sectional  
367 studies. Many cross-sectional studies did not report any multivariate adjusted correlation  
368 coefficients in the paper (only unadjusted r values). Third, there is a clear lack of intervention  
369 studies, particularly parallel trials and cross-over studies reporting data from before the cross-  
370 over point, so effect sizes in the intervention meta-analysis are likely to have been  
371 underpowered due to a small number of studies with a small number of participants.  
372 Fourth, the diets of the populations represented in this analysis generally showed protein  
373 intakes that meet (and usually slightly exceed) protein intake recommendations. Results of

374 the analysis could differ if persons with a low protein intake (or very high protein intake)  
375 were included. Indeed, as discussed above, few studies had populations with inadequate  
376 intake and more research is required into bone health in these populations. Moreover, it is also  
377 difficult to answer the question as to whether there is a threshold whereby protein intake  
378 becomes so high that is detrimental to bone due to the lack of data in persons with high  
379 intakes. Fifth, the Newcastle Ottawa tool was not always a good fit to the studies, especially  
380 for nested case-control studies where it is not clear whether they should be scored on the case-  
381 control or cohort scales. Therefore, the results of the quality analysis should be seen as a  
382 general guide to study quality, rather than a definitive and accurate score. Finally, the funnel  
383 plots for the correlation studies showed some evidence of publication bias in terms of a lack  
384 of small to medium studies showing negative associations between protein intake and bone  
385 health.

386 Future research needs to include larger-scale intervention studies, particularly in understudied  
387 population sub-groups such as children and adolescents and persons with very low protein  
388 and calcium intakes (e.g. the frail elderly) or very high protein and calcium intakes. Fracture  
389 risk studies need to be undertaken in the elderly, followed up to increased age (e.g. 80s) to  
390 truly see the effect of increased protein intake on long term fracture risk.

391 Overall, this systematic review and meta-analysis of the association between dietary protein  
392 and bone health across the life-course assessed evidence from the last 40 years. A positive  
393 cross-sectional association was shown between protein intake and bone health in most pooled  
394 analyses of  $r$  values in adults. However, these associations disappeared when only covariate  
395 adjusted data were used and there was no association between fracture risk and intake of any  
396 protein type in the fracture risk meta-analysis. There was no effect of any form of protein  
397 supplementation on any indices of bone health. The public health and clinical implications of  
398 this work are that there may be little benefit of increasing protein intake for bone health in

399 healthy adults with an adequate protein intake but there is clearly no indication of any  
400 detrimental effect.

401

402

403 **References**

404

- 405 1. Cao JJ, Johnson LK, Hunt JR (2011) A diet high in meat protein and potential renal  
 406 acid load increases fractional calcium absorption and urinary calcium excretion without  
 407 affecting markers of bone resorption or formation in postmenopausal women. *J Nutr* 141:391-  
 408 397
- 409 2. Arnett TR, Dempster DW (1986) Effect of pH on bone resorption by rat osteoclasts in  
 410 vitro. *Endocrinology* 119:119-124
- 411 3. Mangano KM, Walsh SJ, Kenny AM, Insogna KL, Kerstetter JE (2014) Dietary acid  
 412 load is associated with lower bone mineral density in men with low intake of dietary calcium.  
 413 *J Bone Miner Res* 29:500-506
- 414 4. Massey LK (2003) Dietary animal and plant protein and human bone health: a whole  
 415 foods approach. *J Nutr* 133:862S-865S
- 416 5. Darling AL, Millward DJ, Torgerson DJ, Hewitt CE, Lanham-New SA (2009) Dietary  
 417 protein and bone health: a systematic review and meta-analysis. *Am J Clin Nutr* 90:1674-  
 418 1692
- 419 6. Shams-White MM, Chung M, Du M, et al. (2017) Dietary protein and bone health: a  
 420 systematic review and meta-analysis from the National Osteoporosis Foundation. *Am J Clin*  
 421 *Nutr* 105:1528-1543
- 422 7. Wu AM, Sun XL, Lv QB, Zhou Y, Xia DD, Xu HZ, Huang QS, Chi YL (2015) The  
 423 relationship between dietary protein consumption and risk of fracture: a subgroup and dose-  
 424 response meta-analysis of prospective cohort studies. *Sci Rep* 5:9151
- 425 8. Wallace TC, Frankenfeld CL (2017) Dietary Protein Intake above the Current RDA  
 426 and Bone Health: A Systematic Review and Meta-Analysis. *J Am Coll Nutr* 1-16
- 427 9. Santesso N, Akl EA, Bianchi M, Mente A, Mustafa R, Heels-Ansdell D, Schunemann  
 428 HJ (2012) Effects of higher- versus lower-protein diets on health outcomes: a systematic  
 429 review and meta-analysis. *Eur J Clin Nutr* 66:780-788
- 430 10. Shams-White MM, Chung M, Fu Z, et al. (2018) Animal versus plant protein and  
 431 adult bone health: A systematic review and meta-analysis from the National Osteoporosis  
 432 Foundation. *PLoS One* 13:e0192459
- 433 11. Team RC (2014) R: A language and environment for statistical computing. . 3.1.2 edn
- 434 12. Laliberté E (2011) metacor: Meta-analysis of correlation coefficients, R package  
 435 version 1.0-2. <http://CRAN.R-project.org/package=metacor>. Accessed 10 April 2017
- 436 13. Schwarzer G (2007) meta: An R package for meta-analysis. *R News* 7:40-45
- 437 14. Viechtbauer W (2010) Conducting Meta-Analyses in R with the metafor Package.  
 438 *Journal of Statistical Software* 36:1-48
- 439 15. Möbius TWD (2014) metagen: Inference in Meta Analysis and Meta Regression. R  
 440 package version 1.0. <http://CRAN.R-project.org/package=metagen>. Accessed 10 April 2017
- 441 16. Higgins JP, Thompson SG, Deeks JJ, Altman DG (2003) Measuring inconsistency in  
 442 meta-analyses. *BMJ* 327:557-560
- 443 17. Egger M, Davey Smith G, Schneider M, Minder C (1997) Bias in meta-analysis  
 444 detected by a simple, graphical test. *BMJ* 315:629-634
- 445 18. Jadad AR, Moore RA, Carroll D, Jenkinson C, Reynolds DJ, Gavaghan DJ, McQuay  
 446 HJ (1996) Assessing the quality of reports of randomized clinical trials: is blinding necessary?  
 447 *Control Clin Trials* 17:1-12
- 448 19. Wells GA, Shea B, O'Connell D, Peterson J, Welch V, Losos M, Tugwell P (2014)  
 449 The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in  
 450 meta-analyses. [http://www.ohri.ca/programs/clinical\\_epidemiology/oxford.asp](http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp). Accessed 11  
 451 April 2017

- 452 20. Moher D, Cook DJ, Eastwood S, Olkin I, Rennie D, Stroup DF (1999) Improving the  
453 quality of reports of meta-analyses of randomised controlled trials: the QUOROM statement.  
454 Quality of Reporting of Meta-analyses. *Lancet* 354:1896-1900
- 455 21. Alissa EM, Alnahdi WA, Alama N, Ferns GA (2014) Relationship Between  
456 Nutritional Profile, Measures of Adiposity, and Bone Mineral Density in Postmenopausal  
457 Saudi Women. *Journal of the American College of Nutrition* 33:206-214
- 458 22. Chan R, Woo J, Lau W, Leung J, Xu L, Zhao XH, Yu W, Lau E, Pocock N (2009)  
459 Effects of lifestyle and diet on bone health in young adult Chinese women living in Hong  
460 Kong and Beijing. *Food Nutr Bull* 30:370-378
- 461 23. Chiu JF, Lan SJ, Yang CY, Wang PW, Yao WJ, Su LH, Hsieh CC (1997) Long-term  
462 vegetarian diet and bone mineral density in postmenopausal Taiwanese women. *Calcif Tissue*  
463 *Int* 60:245-249
- 464 24. Cooper C, Atkinson EJ, Hensrud DD, Wahner HW, O'Fallon WM, Riggs BL, Melton  
465 LJ, 3rd (1996) Dietary protein intake and bone mass in women. *Calcif Tissue Int* 58:320-325
- 466 25. Coin A, Perissinotto E, Enzi G, Zamboni M, Inelmen EM, Frigo AC, Manzato E,  
467 Busetto L, Buja A, Sergi G (2008) Predictors of low bone mineral density in the elderly: the  
468 role of dietary intake, nutritional status and sarcopenia. *Eur J Clin Nutr* 62:802-809
- 469 26. Devine A, Dick IM, Islam AFM, Dhaliwal SS, Prince RL (2005) Protein consumption  
470 is an important predictor of lower limb bone mass in elderly women. *American Journal of*  
471 *Clinical Nutrition* 81:1423-1428
- 472 27. Freudenheim JL, Johnson NE, Smith EL (1986) Relationships between Usual Nutrient  
473 Intake and Bone-Mineral Content of Women 35-65 Years of Age - Longitudinal and Cross-  
474 Sectional Analysis. *American Journal of Clinical Nutrition* 44:863-876
- 475 28. Gunn CA, Weber JL, Kruger MC (2014) Diet, weight, cytokines and bone health in  
476 postmenopausal women. *J Nutr Health Aging* 18:479-486
- 477 29. Gregg EW, Kriska AM, Salamone LM, Wolf RL, Roberts MM, Ferrell RE, Anderson  
478 SJ, Kuller LH, Cauley JA (1999) Correlates of quantitative ultrasound in the women's healthy  
479 lifestyle project. *Osteoporosis Int* 10:416-424
- 480 30. Henderson NK, Price RI, Cole JH, Gutteridge DH, Bhagat CI (1995) Bone-Density in  
481 Young-Women Is Associated with Body-Weight and Muscle Strength but Not Dietary  
482 Intakes. *Journal of Bone and Mineral Research* 10:384-393
- 483 31. Hirota T, Nara M, Ohguri M, Manago E, Hirota K (1992) Effect of diet and lifestyle  
484 on bone mass in Asian young women. *Am J Clin Nutr* 55:1168-1173
- 485 32. Horiuchi T, Onouchi T, Takahashi M, Ito H, Orimo H (2000) Effect of soy protein on  
486 bone metabolism in postmenopausal Japanese women. *Osteoporosis Int* 11:721-724
- 487 33. Jaime PC, Latorre Mdo R, Florindo AA, Tanaka T, Zerbini CA (2006) Dietary intake  
488 of Brazilian black and white men and its relationship to the bone mineral density of the  
489 femoral neck. *Sao Paulo Med J* 124:267-270
- 490 34. Knurick JR, Johnston CS, Wherry SJ, Aguayo I (2015) Comparison of Correlates of  
491 Bone Mineral Density in Individuals Adhering to Lacto-Ovo, Vegan, or Omnivore Diets: A  
492 Cross-Sectional Investigation. *Nutrients* 7:3416-3426
- 493 35. Kumar A, Mittal S, Orito S, Ishitani K, Ohta H (2010) Impact of dietary intake,  
494 education, and physical activity on bone mineral density among North Indian women. *J Bone*  
495 *Miner Metab* 28:192-201
- 496 36. Lacey JM, Anderson JJB, Fujita T, Yoshimoto Y, Fukase M, Tsuchie S, Koch GG  
497 (1991) Correlates of Cortical Bone Mass among Premenopausal and Postmenopausal  
498 Japanese Women. *Journal of Bone and Mineral Research* 6:651-659
- 499 37. Lau EMC, Kwok T, Woo J, Ho SC (1998) Bone mineral density in Chinese elderly  
500 female vegetarians, vegans, lacto-vegetarians and omnivores. *Eur J Clin Nutr* 52:60-64



- 501 38. Michaelsson K, Holmberg L, Mallmin H, Wolk A, Bergstrom R, Ljunghall S (1995)  
502 Diet, bone mass, and osteocalcin: a cross-sectional study. *Calcif Tissue Int* 57:86-93
- 503 39. Nakamura K, Hori Y, Nashimoto M, Okuda Y, Miyazaki H, Kasai Y, Yamamoto M  
504 (2004) Dietary calcium, sodium, phosphorus, and protein and bone metabolism in elderly  
505 Japanese women: a pilot study using the duplicate portion sampling method. *Nutrition*  
506 20:340-345
- 507 40. New SA, BoltonSmith C, Grubb DA, Reid DM (1997) Nutritional influences on bone  
508 mineral density: A cross-sectional study in premenopausal women. *American Journal of*  
509 *Clinical Nutrition* 65:1831-1839
- 510 41. Orozco Lopez P, Ruiz Gil E, Nolla Sole JM (1998) [Are food intake and life styles  
511 related with bone mass in fertile women?]. *An Med Interna* 15:63-69
- 512 42. Orwoll ES, Weigel RM, Oviatt SK, Meier DE, McClung MR (1987) Serum protein  
513 concentrations and bone mineral content in aging normal men. *Am J Clin Nutr* 46:614-621
- 514 43. Quintas ME, Ortega RM, Lopez-Sobaler AM, Garrido G, Requejo AM (2003)  
515 Influence of dietetic and anthropometric factors and of the type of sport practised on bone  
516 density in different groups of women. *Eur J Clin Nutr* 57 Suppl 1:S58-62
- 517 44. Rapuri PB, Gallagher JC, Haynatzka V (2003) Protein intake: effects on bone mineral  
518 density and the rate of bone loss in elderly women. *American Journal of Clinical Nutrition*  
519 77:1517-1525
- 520 45. Rubinacci A, Porrini M, Sirtori P, Galli L, Tessari L (1992) [Nutrients, anthropometric  
521 characteristics and osteoporosis in women in the recent and late postmenopausal period].  
522 *Minerva Med* 83:497-506
- 523 46. Teegarden D, Lyle RM, McCabe GP, McCabe LD, Proulx WR, Michon K, Knight  
524 AP, Johnston CC, Weaver CM (1998) Dietary calcium, protein, and phosphorus are related to  
525 bone mineral density and content in young women. *Am J Clin Nutr* 68:749-754
- 526 47. Thorpe M, Mojtahedi MC, Chapman-Novakofski K, McAuley E, Evans EM (2008) A  
527 positive association of lumbar spine bone mineral density with dietary protein is suppressed  
528 by a negative association with protein sulfur. *J Nutr* 138:80-85
- 529 48. Wang MC, Villa ML, Marcus R, Kelsey JL (1997) Associations of vitamin C, calcium  
530 and protein with bone mass in postmenopausal Mexican American women. *Osteoporosis Int*  
531 7:533-538
- 532 49. Ho SC, Chan SG, Yip YB, Chan CSY, Woo JLF, Sham A (2008) Change in bone  
533 mineral density and its determinants in pre- and perimenopausal Chinese women: the Hong  
534 Kong perimenopausal women osteoporosis study. *Osteoporosis Int* 19:1785-1796
- 535 50. Loenneke JP, Balapur A, Thrower AD, Syler G, Timlin M, Pujol TJ (2010) Short  
536 report: Relationship between quality protein, lean mass and bone health. *Ann Nutr Metab*  
537 57:219-220
- 538 51. Bounds W, Skinner J, Carruth BR, Ziegler P (2005) Current research - The  
539 relationship of dietary and lifestyle factors to bone mineral indexes in children. *J Am Diet*  
540 *Assoc* 105:735-741
- 541 52. Chevalley T, Bonjour JP, Ferrari S, Rizzoli R (2008) High-protein intake enhances the  
542 positive impact of physical activity on BMC in prepubertal boys. *Journal of Bone and Mineral*  
543 *Research* 23:131-142
- 544 53. Ekbote VH, Khadilkar AV, Chiplonkar SA, Khadilkar VV (2011) Determinants of  
545 bone mineral content and bone area in Indian preschool children. *J Bone Miner Metab*  
546 29:334-341
- 547 54. Hoppe C, Molgaard C, Michaelsen KF (2000) Bone size and bone mass in 10-year-old  
548 Danish children: effect of current diet. *Osteoporosis Int* 11:1024-1030

- 549 55. Jones G, Riley MD, Whiting S (2001) Association between urinary potassium, urinary  
550 sodium, current diet, and bone density in prepubertal children. *American Journal of Clinical*  
551 *Nutrition* 73:839-844
- 552 56. Ho SC, Woo J, Lam S, Chen Y, Sham A, Lau J (2003) Soy protein consumption and  
553 bone mass in early postmenopausal Chinese women. *Osteoporosis Int* 14:835-842
- 554 57. Dargent-Molina P, Sabia S, Touvier M, Kesse E, Breart G, Clavel-Chapelon F,  
555 Boutron-Ruault MC (2008) Proteins, dietary acid load, and calcium and risk of  
556 postmenopausal fractures in the E3N French women prospective study. *J Bone Miner Res*  
557 23:1915-1922
- 558 58. Munger RG, Cerhan JR, Chiu BC (1999) Prospective study of dietary protein intake  
559 and risk of hip fracture in postmenopausal women. *Am J Clin Nutr* 69:147-152
- 560 59. Feskanich D, Willett WC, Stampfer MJ, Colditz GA (1996) Protein consumption and  
561 bone fractures in women. *Am J Epidemiol* 143:472-479
- 562 60. Mussolino ME, Looker AC, Madans JH, Langlois JA, Orwoll ES (1998) Risk factors  
563 for hip fracture in white men: the NHANES I Epidemiologic Follow-up Study. *J Bone Miner*  
564 *Res* 13:918-924
- 565 61. Meyer HE, Pedersen JI, Loken EB, Tverdal A (1997) Dietary factors and the  
566 incidence of hip fracture in middle-aged Norwegians. A prospective study. *Am J Epidemiol*  
567 145:117-123
- 568 62. Langsetmo L, Barr SI, Berger C, et al. (2015) Associations of Protein Intake and  
569 Protein Source with Bone Mineral Density and Fracture Risk: A Population-Based Cohort  
570 Study. *J Nutr Health Aging* 19:861-868
- 571 63. Misra D, Berry SD, Broe KE, McLean RR, Cupples LA, Tucker KL, Kiel DP, Hannan  
572 MT (2011) Does dietary protein reduce hip fracture risk in elders? The Framingham  
573 osteoporosis study. *Osteoporosis Int* 22:345-349
- 574 64. Sahni S, Cupples LA, McLean RR, Tucker KL, Broe KE, Kiel DP, Hannan MT  
575 (2010) Protective effect of high protein and calcium intake on the risk of hip fracture in the  
576 Framingham offspring cohort. *J Bone Miner Res* 25:2770-2776
- 577 65. Martinez-Ramirez MJ, Delgado-Martinez AD, Ruiz-Bailen M, de la Fuente C,  
578 Martinez-Gonzalez MA, Delgado-Rodriguez M (2012) Protein intake and fracture risk in  
579 elderly people: a case-control study. *Clin Nutr* 31:391-395
- 580 66. Nieves JW, Grisso JA, Kelsey JL (1992) A Case-Control Study of Hip Fracture -  
581 Evaluation of Selected Dietary Variables and Teenage Physical-Activity. *Osteoporosis Int*  
582 2:122-127
- 583 67. Wengreen HJ, Munger RG, West NA, Cutler DR, Corcoran CD, Zhang J, Sassano NE  
584 (2004) Dietary protein intake and risk of osteoporotic hip fracture in elderly residents of Utah.  
585 *J Bone Miner Res* 19:537-545
- 586 68. Kerstetter JE, Bihuniak JD, Brindisi J, et al. (2015) The Effect of a Whey Protein  
587 Supplement on Bone Mass in Older Caucasian Adults. *J Clin Endocrinol Metab* 100:2214-  
588 2222
- 589 69. Aoe S, Koyama T, Toba Y, Itabashi A, Takada Y (2005) A controlled trial of the  
590 effect of milk basic protein (MBP) supplementation on bone metabolism in healthy  
591 menopausal women. *Osteoporosis Int* 16:2123-2128
- 592 70. Unishi K, Ishida H, Toba Y, Aoe S, Itabashi A, Takada Y (2007) Milk basic protein  
593 increases bone mineral density and improves bone metabolism in healthy young women.  
594 *Osteoporosis Int* 18:385-390
- 595 71. Zou ZY, Lin XM, Xu XR, Xu R, Ma L, Li Y, Wang MF (2009) Evaluation of milk  
596 basic protein supplementation on bone density and bone metabolism in Chinese young  
597 women. *Eur J Nutr* 48:301-306

- 598 72. Tkatch L, Rapin CH, Rizzoli R, Slosman D, Nydegger V, Vasey H, Bonjour JP (1992)  
599 Benefits of oral protein supplementation in elderly patients with fracture of the proximal  
600 femur. *J Am Coll Nutr* 11:519-525
- 601 73. Alekel DL, St Germain A, Pererson CT, Hanson KB, Stewart JW, Toda T (2000)  
602 Isoflavone-rich soy protein isolate attenuates bone loss in the lumbar spine of perimenopausal  
603 women. *American Journal of Clinical Nutrition* 72:844-852
- 604 74. Kenny AM, Mangano KM, Abourizk RH, Bruno RS, Anamani DE, Kleppinger A,  
605 Walsh SJ, Prestwood KM, Kerstetter JE (2009) Soy proteins and isoflavones affect bone  
606 mineral density in older women: a randomized controlled trial. *Am J Clin Nutr* 90:234-242
- 607 75. Vupadhyayula PM, Gallagher JC, Templin T, Logsdon SM, Smith LM (2009) Effects  
608 of soy protein isolate on bone mineral density and physical performance indices in  
609 postmenopausal women--a 2-year randomized, double-blind, placebo-controlled trial.  
610 *Menopause* 16:320-328
- 611 76. Zhu K, Meng X, Kerr DA, Devine A, Solah V, Binns CW, Prince RL (2011) The  
612 effects of a two-year randomized, controlled trial of whey protein supplementation on bone  
613 structure, IGF-1, and urinary calcium excretion in older postmenopausal women. *J Bone*  
614 *Miner Res* 26:2298-2306
- 615 77. Yamamura J, Aoe S, Toba Y, Motouri M, Kawakami H, Kumegawa M, Itabashi A,  
616 Takada Y (2002) Milk basic protein (MBP) increases radial bone mineral density in healthy  
617 adult women. *Biosci Biotechnol Biochem* 66:702-704
- 618 78. Evans EM, Racette SB, Van Pelt RE, Peterson LR, Villareal DT (2007) Effects of soy  
619 protein isolate and moderate exercise on bone turnover and bone mineral density in  
620 postmenopausal women. *Menopause* 14:481-488
- 621 79. Spence LA, Lipscomb ER, Cadogan J, Martin B, Wastney ME, Peacock M, Weaver  
622 CM (2005) The effect of soy protein and soy isoflavones on calcium metabolism in  
623 postmenopausal women: a randomized crossover study. *Am J Clin Nutr* 81:916-922
- 624 80. Kerstetter JE, Mitnick ME, Gundberg CM, Caseria DM, Ellison AF, Carpenter TO,  
625 Insogna KL (1999) Changes in bone turnover in young women consuming different levels of  
626 dietary protein. *J Clin Endocr Metab* 84:1052-1055
- 627 81. Roughead ZK, Johnson LK, Lykken GI, Hunt JR (2003) Controlled high meat diets do  
628 not affect calcium retention or indices of bone status in healthy postmenopausal women. *J*  
629 *Nutr* 133:1020-1026
- 630 82. Ceglia L, Harris SS, Abrams SA, Rasmussen HM, Dallal GE, Dawson-Hughes B  
631 (2009) Potassium bicarbonate attenuates the urinary nitrogen excretion that accompanies an  
632 increase in dietary protein and may promote calcium absorption. *J Clin Endocrinol Metab*  
633 94:645-653
- 634 83. Hunt JR, Johnson LK, Fariba Roughead ZK (2009) Dietary protein and calcium  
635 interact to influence calcium retention: a controlled feeding study. *Am J Clin Nutr* 89:1357-  
636 1365
- 637 84. Shapses SA, Robins SP, Schwartz EI, Chowdhury H (1995) Short-term changes in  
638 calcium but not protein intake alter the rate of bone resorption in healthy subjects as assessed  
639 by urinary pyridinium cross-link excretion. *J Nutr* 125:2814-2821
- 640 85. Jenkins DJ, Kendall CW, Vidgen E, Augustin LS, Parker T, Faulkner D, Vieth R,  
641 Vandenbroucke AC, Josse RG (2003) Effect of high vegetable protein diets on urinary  
642 calcium loss in middle-aged men and women. *Eur J Clin Nutr* 57:376-382
- 643 86. Khalil DA, Lucas EA, Juma S, Smith BJ, Payton ME, Arjmandi BH (2002) Soy  
644 protein supplementation increases serum insulin-like growth factor-I in young and old men  
645 but does not affect markers of bone metabolism. *J Nutr* 132:2605-2608
- 646 87. Arjmandi BH, Khalil DA, Smith BJ, Lucas EA, Juma S, Payton ME, Wild RA (2003)  
647 Soy protein has a greater effect on bone in postmenopausal women not on hormone

- 648 replacement therapy, as evidenced by reducing bone resorption and urinary calcium excretion.  
649 J Clin Endocr Metab 88:1048-1054
- 650 88. Cuneo F, Costa-Paiva L, Pinto-Neto AM, Morais SS, Amaya-Farfan J (2010) Effect of  
651 dietary supplementation with collagen hydrolysates on bone metabolism of postmenopausal  
652 women with low mineral density. *Maturitas* 65:253-257
- 653 89. Dalais FS, Ebeling PR, Kotsopoulos D, McGrath BP, Teede HJ (2003) The effects of  
654 soy protein containing isoflavones on lipids and indices of bone resorption in postmenopausal  
655 women. *Clin Endocrinol (Oxf)* 58:704-709
- 656 90. Toba Y, Takada Y, Matsuoka Y, et al. (2001) Milk basic protein promotes bone  
657 formation and suppresses bone resorption in healthy adult men. *Biosci Biotechnol Biochem*  
658 65:1353-1357
- 659 91. Dawson-Hughes B, Harris SS, Rasmussen H, Song LY, Dallal GE (2004) Effect of  
660 dietary protein supplements on calcium excretion in healthy older men and women. *J Clin*  
661 *Endocr Metab* 89:1169-1173
- 662 92. Lampl M, Johnston FE (1978) The effects of protein supplementation on the growth  
663 and skeletal maturation of New Guinean school children. *Ann Hum Biol* 5:219-227
- 664 93. Martin-Bautista E, Martin-Matillas M, Martin-Lagos JA, Miranda-Leon MT, Munoz-  
665 Torres M, Ruiz-Requena E, Rivero M, Quer J, Puigdueta I, Campoy C (2011) A nutritional  
666 intervention study with hydrolyzed collagen in pre-pubertal spanish children: influence on  
667 bone modeling biomarkers. *J Pediatr Endocrinol Metab* 24:147-153
- 668 94. Ince BA, Anderson EJ, Neer RM (2004) Lowering dietary protein to U.S.  
669 Recommended dietary allowance levels reduces urinary calcium excretion and bone  
670 resorption in young women. *J Clin Endocrinol Metab* 89:3801-3807
- 671 95. Schurch MA, Rizzoli R, Slosman D, Vadas L, Vergnaud P, Bonjour JP (1998) Protein  
672 supplements increase serum insulin-like growth factor-I levels and attenuate proximal femur  
673 bone loss in patients with recent hip fracture. A randomized, double-blind, placebo-controlled  
674 trial. *Ann Intern Med* 128:801-809
- 675 96. Aoe S, Toba Y, Yamamura J, Kawakami H, Yahiro M, Kumegawa M, Itabashi A,  
676 Takada Y (2001) Controlled trial of the effects of milk basic protein (MBP) supplementation  
677 on bone metabolism in healthy adult women. *Biosci Biotech Bioch* 65:913-918
- 678 97. Ministry of Health (2014) Nutrient Reference Values for Australia and New Zealand:  
679 Nutrients, Protein. <https://www.nrv.gov.au/nutrients/protein>. Accessed 22nd February 2019
- 680 98. EFSA (2012) Scientific Opinion on Dietary Reference Values for protein. *EFSA*  
681 *Journal* 10:2557
- 682 99. IOM (2005) Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty  
683 Acids, Cholesterol, Protein and Amino Acids. The National Academies Press, Washington  
684 D.C.
- 685 100. Geinoz G, Rapin CH, Rizzoli R, Kraemer R, Buchs B, Slosman D, Michel JP, Bonjour  
686 JP (1993) Relationship between Bone-Mineral Density and Dietary Intakes in the Elderly.  
687 *Osteoporosis Int* 3:242-248
- 688 101. Ho-Pham LT, Nguyen PLT, Le TTT, Doan TAT, Tran NT, Le TA, Nguyen TV (2009)  
689 Veganism, bone mineral density, and body composition: a study in Buddhist nuns.  
690 *Osteoporosis Int* 20:2087-2093
- 691 102. Oh SM, Kim HC, Rhee Y, Park SJ, Lee HJ, Suh I, Feskanich D (2013) Dietary protein  
692 in relation to bone stiffness index and fat-free mass in a population consuming relatively low  
693 protein diets. *J Bone Miner Metab* 31:433-441
- 694 103. Fung TT, Meyer HE, Willett WC, Feskanich D (2017) Protein intake and risk of hip  
695 fractures in postmenopausal women and men age 50 and older. *Osteoporos Int* 28:1401-1411

- 696 104. Langsetmo L, Shikany JM, Cawthon PM, et al. (2017) The Association Between  
697 Protein Intake by Source and Osteoporotic Fracture in Older Men: A Prospective Cohort  
698 Study. *J Bone Miner Res* 32:592-600
- 699 105. Durosier-Izart C, Biver E, Merminod F, van Rietbergen B, Chevalley T, Herrmann  
700 FR, Ferrari SL, Rizzoli R (2017) Peripheral skeleton bone strength is positively correlated  
701 with total and dairy protein intakes in healthy postmenopausal women. *Am J Clin Nutr*  
702 105:513-525
- 703 106. Chevalley T, Bonjour JP, Audet MC, Merminod F, van Rietbergen B, Rizzoli R,  
704 Ferrari S (2017) Prepubertal Impact of Protein Intake and Physical Activity on Weight-  
705 Bearing Peak Bone Mass and Strength in Males. *J Clin Endocrinol Metab* 102:157-166
- 706 107. Isanejad M, Sirola J, Mursu J, Kroger H, Tuppurainen M, Erkkila AT (2017)  
707 Association of Protein Intake with Bone Mineral Density and Bone Mineral Content among  
708 Elderly Women: The OSTPRE Fracture Prevention Study. *J Nutr Health Aging* 21:622-630
- 709 108. Konig D, Oesser S, Scharla S, Zdzieblik D, Gollhofer A (2018) Specific Collagen  
710 Peptides Improve Bone Mineral Density and Bone Markers in Postmenopausal Women-A  
711 Randomized Controlled Study. *Nutrients* 10:
- 712 109. Steell L, Sillars A, Welsh P, Iliodromiti S, Wong SC, Pell JP, Sattar N, Gill JMR,  
713 Celis-Morales CA, Gray SR (2018) Associations of dietary protein intake with bone mineral  
714 density: An observational study in 70,215 UK Biobank participants. *Bone* 120:38-43
- 715 110. Mejia W, Cordoba D, Duran P, Chacon Y, Rosselli D (2018) Effect of Daily Exposure  
716 to an Isolated Soy Protein Supplement on Body Composition, Energy and Macronutrient  
717 Intake, Bone Formation Markers, and Lipid Profile in Children in Colombia. *J Diet Suppl* 1-  
718 13
- 719 111. Langsetmo L, Shikany JM, Burghardt AJ, et al. (2018) High dairy protein intake is  
720 associated with greater bone strength parameters at the distal radius and tibia in older men: a  
721 cross-sectional study. *Osteoporos Int* 29:69-77
- 722 112. Johansson J, Hult A, Morseth B, Nordstrom A, Nordstrom P (2018) Self-reported  
723 protein intake and properties of bone in community-dwelling older individuals. *Arch*  
724 *Osteoporos* 13:10
- 725 113. Kim BJ, Lee SH, Isales CM, Koh JM, Hamrick MW (2018) The positive association  
726 of total protein intake with femoral neck strength (KNHANES IV). *Osteoporos Int* 29:1397-  
727 1405
- 728 114. Cauley JA, Cawthon PM, Peters KE, et al. (2016) Risk Factors for Hip Fracture in  
729 Older Men: The Osteoporotic Fractures in Men Study (MrOS). *J Bone Miner Res* 31:1810-  
730 1819
- 731 115. Mangano KM, Sahni S, Kiel DP, Tucker KL, Dufour AB, Hannan MT (2017) Dietary  
732 protein is associated with musculoskeletal health independently of dietary pattern: the  
733 Framingham Third Generation Study. *Am J Clin Nutr* 105:714-722
- 734 116. Alexy U, Remer T, Manz F, Neu CM, Schoenau E (2005) Long-term protein intake  
735 and dietary potential renal acid load are associated with bone modeling and remodeling at the  
736 proximal radius in healthy children. *American Journal of Clinical Nutrition* 82:1107-1114
- 737 117. Alissa EM, Qadi SG, Alhujaili NA, Alshehri AM, Ferns GA (2011) Effect of diet and  
738 lifestyle factors on bone health in postmenopausal women. *J Bone Miner Metab* 29:725-735
- 739 118. Beasley JM, Ichikawa LE, Ange BA, Spangler L, LaCroix AZ, Ott SM, Scholes D  
740 (2010) Is protein intake associated with bone mineral density in young women? *American*  
741 *Journal of Clinical Nutrition* 91:1311-1316
- 742 119. Beasley JM, LaCroix AZ, Larson JC, et al. (2014) Biomarker-calibrated protein intake  
743 and bone health in the Women's Health Initiative clinical trials and observational study. *Am J*  
744 *Clin Nutr* 99:934-940

- 745 120. Budek AZ, Hoppe C, Michaelsen KE, Bugel S, Molgaard C (2007) Associations of  
746 total, dairy, and meat protein with markers for bone turnover in healthy, prepubertal boys. *J*  
747 *Nutr* 137:930-934
- 748 121. Budek AZ, Hoppe C, Ingstrup H, Michaelsen KF, Bugel S, Molgaard C (2007)  
749 Dietary protein intake and bone mineral content in adolescents-The Copenhagen Cohort  
750 Study. *Osteoporos Int* 18:1661-1667
- 751 122. Chan R, Woo J, Leung J (2011) Effects of Food Groups and Dietary Nutrients on  
752 Bone Loss in Elderly Chinese Population. *Journal of Nutrition Health & Aging* 15:287-294
- 753 123. Chevalley T, Bonjour JP, van Rietbergen B, Ferrari S, Rizzoli R (2014) Tracking of  
754 Environmental Determinants of Bone Structure and Strength Development in Healthy Boys:  
755 An Eight-Year Follow Up Study on the Positive Interaction Between Physical Activity and  
756 Protein Intake From Prepuberty to Mid-Late Adolescence. *Journal of Bone and Mineral*  
757 *Research* 29:2182-2192
- 758 124. Dawson-Hughes B, Harris SS (2002) Calcium intake influences the association of  
759 protein intake with rates of bone loss in elderly men and women. *Am J Clin Nutr* 75:773-779
- 760 125. Fairweather-Tait SJ, Skinner J, Guile GR, Cassidy A, Spector TD, MacGregor AJ  
761 (2011) Diet and bone mineral density study in postmenopausal women from the TwinsUK  
762 registry shows a negative association with a traditional English dietary pattern and a positive  
763 association with wine. *American Journal of Clinical Nutrition* 94:1371-1375
- 764 126. Genaro PD, Pinheiro MD, Szejnfeld VL, Martini LA (2015) Dietary Protein Intake in  
765 Elderly Women: Association With Muscle and Bone Mass. *Nutr Clin Pract* 30:283-289
- 766 127. Hannan MT, Tucker KL, Dawson-Hughes B, Cupples LA, Felson DT, Kiel DP (2000)  
767 Effect of dietary protein on bone loss in elderly men and women: the Framingham  
768 Osteoporosis Study. *J Bone Miner Res* 15:2504-2512
- 769 128. Hernandezavila M, Stampfer MJ, Ravnkar VA, Willett WC, Schiff I, Francis M,  
770 Longscope C, Mckinlay SM (1993) Caffeine and Other Predictors of Bone-Density among  
771 Premenopausal and Perimenopausal Women. *Epidemiology* 4:128-134
- 772 129. Ho-Pham LT, Vu BQ, Lai TQ, Nguyen ND, Nguyen TV (2012) Vegetarianism, bone  
773 loss, fracture and vitamin D: a longitudinal study in Asian vegans and non-vegans. *Eur J Clin*  
774 *Nutr* 66:75-82
- 775 130. Hu T, Rianon NJ, Nettleton JA, Hyder JA, He J, Steffen LM, Jacobs DR, Criqui MH,  
776 Bazzano LA (2014) Protein intake and lumbar bone density: the Multi-Ethnic Study of  
777 Atherosclerosis (MESA). *Brit J Nutr* 112:1384-1392
- 778 131. Ilich JZ, Brownbill RA, Tamborini L (2003) Bone and nutrition in elderly women:  
779 protein, energy, and calcium as main determinants of bone mineral density. *Eur J Clin Nutr*  
780 *57:554-565*
- 781 132. Iuliano-Burns S, Stone J, Hopper JL, Seeman E (2005) Diet and exercise during  
782 growth have site-specific skeletal effects: a co-twin control study. *Osteoporosis Int* 16:1225-  
783 1232
- 784 133. Libuda L, Alexy U, Remer T, Stehle P, Schoenau E, Kersting M (2008) Association  
785 between long-term consumption of soft drinks and variables of bone modeling and  
786 remodeling in a sample of healthy German children and adolescents. *American Journal of*  
787 *Clinical Nutrition* 88:1670-1677
- 788 134. Libuda L, Wudy SA, Schoenau E, Remer T (2011) Comparison of the effects of  
789 dietary protein, androstenediol and forearm muscle area on radial bone variables in healthy  
790 prepubertal children. *Br J Nutr* 105:428-435
- 791 135. Macdonald HM, New SA, Fraser WD, Campbell MK, Reid DM (2005) Low dietary  
792 potassium intakes and high dietary estimates of net endogenous acid production are associated  
793 with low bone mineral density in premenopausal women and increased markers of bone  
794 resorption in postmenopausal women. *American Journal of Clinical Nutrition* 81:923-933

- 795 136. Meng X, Zhu K, Devine A, Kerr DA, Binns CW, Prince RL (2009) A 5-year cohort  
796 study of the effects of high protein intake on lean mass and BMC in elderly postmenopausal  
797 women. *J Bone Miner Res* 24:1827-1834
- 798 137. Metz JA, Anderson JJ, Gallagher PN, Jr. (1993) Intakes of calcium, phosphorus, and  
799 protein, and physical-activity level are related to radial bone mass in young adult women. *Am*  
800 *J Clin Nutr* 58:537-542
- 801 138. Neville CE, Robson PJ, Murray LJ, Strain JJ, Twisk J, Gallagher AM, McGuinness M,  
802 Cran GW, Ralston SH, Boreham CAG (2002) The effect of nutrient intake on bone mineral  
803 status in young adults: The Northern Ireland Young Hearts project. *Calcified Tissue Int*  
804 70:89-98
- 805 139. Pearce MS, Relton CL, Groom A, Peaston RT, Francis RM (2010) A lifecourse study  
806 of bone resorption in men ages 49-51years: the Newcastle Thousand Families cohort study.  
807 *Bone* 46:952-956
- 808 140. Promislow JH, Goodman-Gruen D, Slymen DJ, Barrett-Connor E (2002) Protein  
809 consumption and bone mineral density in the elderly : the Rancho Bernardo Study. *Am J*  
810 *Epidemiol* 155:636-644
- 811 141. Sahni S, Broe KE, Tucker KL, McLean RR, Kiel DP, Cupples LA, Hannan MT  
812 (2014) Association of total protein intake with bone mineral density and bone loss in men and  
813 women from the Framingham Offspring Study. *Public Health Nutrition* 17:2570-2576
- 814 142. Tanaka M, Itoh K, Abe S, Imai K, Masuda T, Koga R, Itoh H, Kinukawa N,  
815 Matsuyama T, Nakamura M (2001) Relationship between nutrient factors and osteo-sono  
816 assessment index in calcaneus of young Japanese women. *Nutr Res* 21:1475-1482
- 817 143. Tylavsky FA, Anderson JJ (1988) Dietary factors in bone health of elderly  
818 lactoovovegetarian and omnivorous women. *Am J Clin Nutr* 48:842-849
- 819 144. Vatanparast H, Bailey DA, Baxter-Jones ADG, Whiting SJ (2007) The effects of  
820 dietary protein on bone mineral mass in young adults may be modulated by adolescent  
821 calcium intake. *J Nutr* 137:2674-2679
- 822 145. Wang MC, Moore EC, Crawford PB, Hudes M, Sabry ZI, Marcus R, Bachrach LK  
823 (1999) Influence of pre-adolescent diet on quantitative ultrasound measurements of the  
824 calcaneus in young adult women. *Osteoporosis Int* 9:532-535
- 825 146. Weikert C, Walter D, Hoffmann K, Kroke A, Bergmann MM, Boeing H (2005) The  
826 relation between dietary protein, calcium and bone health in women: Results from the EPIC-  
827 Potsdam cohort. *Ann Nutr Metab* 49:312-318
- 828 147. Whiting SJ, Boyle JL, Thompson A, Mirwald RL, Faulkner RA (2002) Dietary  
829 protein, phosphorus and potassium are beneficial to bone mineral density in adult men  
830 consuming adequate dietary calcium. *J Am Coll Nutr* 21:402-409
- 831 148. Yazdanpanah N, Zillikens MC, Rivadeneira F, De Jong R, Lindemans J, Uitterlinden  
832 AG, Pols HAP, Van Meurs JBJ (2007) Effect of dietary B vitamins on BMD and risk of  
833 fracture in elderly men and women: The Rotterdam Study. *Bone* 41:987-994
- 834 149. Zhang Q, Ma GS, Greenfield H, Zhu K, Du XQ, Foo LH, Hu XQ, Fraser DR (2010)  
835 The association between dietary protein intake and bone mass accretion in pubertal girls with  
836 low calcium intakes. *Brit J Nutr* 103:714-723
- 837 150. Abelow BJ, Holford TR, Insogna KL (1992) Cross-cultural association between  
838 dietary animal protein and hip fracture: a hypothesis. *Calcif Tissue Int* 50:14-18
- 839 151. Frassetto LA, Todd KM, Morris RC, Jr., Sebastian A (2000) Worldwide incidence of  
840 hip fracture in elderly women: relation to consumption of animal and vegetable foods. *J*  
841 *Gerontol A Biol Sci Med Sci* 55:M585-592
- 842 152. Key TJ, Appleby PN, Spencer EA, Roddam AW, Neale RE, Allen NE (2007)  
843 Calcium, diet and fracture risk: a prospective study of 1898 incident fractures among 34 696  
844 British women and men. *Public Health Nutr* 10:1314-1320

- 845 153. Sellmeyer DE, Stone KL, Sebastian A, Cummings SR (2001) A high ratio of dietary  
846 animal to vegetable protein increases the rate of bone loss and the risk of fracture in  
847 postmenopausal women. Study of Osteoporotic Fractures Research Group. *Am J Clin Nutr*  
848 73:118-122
- 849 154. Zhong Y, Okoro CA, Balluz LS (2009) Association of total calcium and dietary  
850 protein intakes with fracture risk in postmenopausal women: the 1999-2002 National Health  
851 and Nutrition Examination Survey (NHANES). *Nutrition* 25:647-654
- 852 155. Zhang X, Shu XO, Li H, Yang G, Li Q, Gao YT, Zheng W (2005) Prospective cohort  
853 study of soy food consumption and risk of bone fracture among postmenopausal women.  
854 *Arch Intern Med* 165:1890-1895
- 855 156. Chevalley T, Bonjour JP, van Rietbergen B, Ferrari S, Rizzoli R (2011) Fractures  
856 during childhood and adolescence in healthy boys: relation with bone mass, microstructure,  
857 and strength. *J Clin Endocrinol Metab* 96:3134-3142
- 858 157. Farrin N, Ostadrahimi AR, Mahboob SA, Kolahi S, Ghavami M (2008) Dietary intake  
859 and serum bone related chemistry and their correlations in postmenopausal Iranian women.  
860 *Saudi Med J* 29:1643-1648
- 861 158. Kim J, Lim SY, Kim JH (2008) Nutrient intake risk factors of osteoporosis in  
862 postmenopausal women. *Asia Pac J Clin Nutr* 17:270-275
- 863 159. Park YJ, Lee SJ, Shin NM, Shin H, Kim YK, Cho Y, Jeon S, Cho I (2014) [Bone  
864 mineral density, biochemical bone turnover markers and factors associated with bone health  
865 in young Korean women]. *J Korean Acad Nurs* 44:504-514
- 866 160. Perez Durillo FT, Torio Durantez J, Villarejo Villar AB, Sanchez Vico AB, Cueto  
867 Camarero Mdel M, Durillo JP (2011) [Comparative study of dietary intake and nutritional  
868 status in elderly women with and without hip fracture]. *Aten Primaria* 43:362-368
- 869 161. Preisinger E, Leitner G, Uher E, Alacamlioglu Y, Seidl G, Marktl W, Resch KL  
870 (1995) [Nutrition and osteoporosis: a nutritional analysis of women in postmenopause]. *Wien*  
871 *Klin Wochenschr* 107:418-422
- 872 162. Samieri C, Ginder Coupez V, Lorrain S, Letenneur L, Alles B, Feart C, Paineau D,  
873 Barberger-Gateau P (2013) Nutrient patterns and risk of fracture in older subjects: results  
874 from the Three-City Study. *Osteoporos Int* 24:1295-1305

875



**Table 1: Characteristics of 74 cross-sectional and/or longitudinal correlational studies**

<b>Study</b>	<b>Design</b>	<b>Mean Protein **</b>	<b>Method</b>	<b>Population</b>	<b>n</b>	<b>Outcomes measured</b>
Alexy et al, 2005, Germany[116]	CS	1.4-2.0 +/- 0.3 g/Kg/d	pQCT	Prepubescent and pubescent boys and girls	229	Periosteal Circumference, Cortical Area, BMC, Polar SSI
Alissa et al, 2011, Saudi Arabia[117]	CS	1.03 g/Kg/d	DXA	Postmenopausal women aged 46-70 years old	122	BMD
Alissa et al, 2014, Saudi Arabia[21]	CS	71.4+/-1.55 g/d	DXA	Postmenopausal women, aged 46-88 years	300	BMD
Beasley et al. 2010, USA[118]	CS	TP: 5.7 - 27.6% energy AP:45g/d, VP:19g/d	DXA	Premenopausal women	560	BMD
Beasley et al. 2014, USA[119]	LS	15% total energy	DXA	Postmenopausal women 50-79 years	144,580	BMD
Bounds et al, 2005, USA[51]	CS	55g/d (1.9g/Kg/d)	DXA	6-8 year old children, 25 Boys and 27 Girls	52	BMD/BMC
Budek et al. 2007a, Denmark [120]	CS	2.67 g/Kg/d	BTM	Pubertal boys	81	OC, BAP, CTX
Budek et al. 2007b, Denmark [121]	CS	TP: 1.2 (Girls), 1.3 (Boys), AP: 0.4 (Girls), 0.5 (Boys), DP: 0.4 (Both Girls and Boys) g/Kg/d	DXA	17-year-olds: 63 girls and 46 boys	109	BMC
Chan et al, 2009, Hong Kong/Beijing[22]	CS	65.4-77.5g/d	DXA	Premenopausal women	441	BMD
Chan et al. 2011, Hong Kong[122]	LS	1.3 g/Kg/d	DXA	Older men and women	2217	BMD
Chevalley et al. 2008, Switzerland[52]	CS	47.3 g/d, 1.78 g/Kg/d	DXA	Pre-pubertal boys	232	BMC
Chevalley et al. 2014, Switzerland[123]	LS	Age 7: 1.8; Age 15: 1.1 (g/Kg/d)	DXA; High resolution pQCT	Adolescent boys	176	BMC/BMD/Area at 7.4 and 15.2 years; HR-pQCT distal tibia microstructure and strength

<b>Study</b>	<b>Design</b>	<b>Mean Protein **</b>	<b>Method</b>	<b>Population</b>	<b>n</b>	<b>Outcomes measured</b>
Chiu et al, 1997, Taiwan [23]	CS	1.09 g/Kg/d	DPA (BMD)	Older post F	258	BMD
Coin et al, Italy, 2008[25]	CS	1.02 g/Kg/d	DXA	Males, mean age 73.9+/-5.6 years	136	BMD
Cooper et al, 1996, USA[24]	CS	72 g/d	DPA SPA	Pre (72) and post (218) F	290	BMD, HPO. OC
Dawson-Hughes et al, 2002, USA[124]	CS	79 g/d	DXA	184 men and women(>=65 years old) in placebo (inactive) arm of calcium supplementation trial	184	BMD, OC, NTX
Devine et al, 2005, AUS[26]	CS	1.2 g/Kg/d	DXA , QUS	Elderly F mean age 75y+/-3y Caucasian	1077	BMD, BUA calcaneus
Ekbote et al, 2011, India[53]	CS	18.6g/d-normal and malnourished children combined	DXA	2-3 year old children	71	BMC,BA
Fairweather-Tait et al, 2011, UK[125]	CS	81.3g/d	DXA	Postmenopausal female twin pairs (Monozygotic or dizygotic twins)	2464 pairs	BMD
Freudenheim et al, 1986, USA[27]	CS	1.02 g/Kg/d	SPA	Pre and post F, 35-65y, Caucasian	84	BMC
Geinoz et al, 1993, Switzerland[100]	CS	Mean Intake in g/d by group: 37.8-59.4 g/d	DXA	Elderly M and F Mean age 82y(F); 80y(M)	74	BMD
Genaro et al, 2015, Brazil[126]	CS	66g/d	DXA	Women over 65 years old	200	BMC, BMD
Gregg et al, 1999, USA[29]	CS	0.9 g/Kg/d	QUS	Middle aged (premenopausal) F- mean age= 45.5y	393	BUA Calcaneus, BMD
Gunn et al, 2014, New Zealand[28]	CS	79g/d	BTM, DXA	Postmenopausal women, 60 years of age	142	BMD, CTX,P1NP
Hannan et al. 2000, USA[127]	LS	68g/d (16% of total energy) 0.97 g/kg/d	DXA	224 older men and 391 older women	615	BMD
Henderson et al, 1995 , AUS[30]	CS	1.0 g/Kg/d	DXA	Pre F- mean age=18y	115	BMD
Hernandezavila et al, 1993, USA[128]	CS	76g/d	SPA	Women (50-60 years old)	281	BMD

<b>Study</b>	<b>Design</b>	<b>Mean Protein **</b>	<b>Method</b>	<b>Population</b>	<b>n</b>	<b>Outcomes measured</b>
Hirota et al, 1992, Japan[31]	CS	1.13 g/Kg/d	SPA (BMD)	Young PRE women: 19-25y	161	BMD
Ho et al, 2003, China [56]	CS	1.01 g/Kg/d SP	DXA	<12y POST women (48-62y), Asian	454	BMC, BMD
Ho et al, 2008, China[49]	CS	5.2g/d SP 48.6 g/d TP	DXA	Pre and perimenopausal women 45-55 years old	438	BMC, BMD
Ho-Pham et al, 2009, Vietnam[101]	CS	TP: 35.4-62.6 g/d	DXA	105 Post F Buddhist vegan Nuns and 105 omnivorous women 62+/-10 years old	210	BMD
Ho-Pham et al. 2012, Vietnam[129]	CS/LS	49 g/d; 0.92g/Kg/d	DXA, BTM	Postmenopausal Buddhist Nuns	210	BMD, CTX, PINP
Hoppe et al. 2000, Denmark[54]	CS	73-82g/d	DXA	10 year old children	105	BMC, BA
Horiuchi et al, 2000, Japan[32]	CS	TP: 62.5g/d SP: 12.6g/d	DXA	Post F, 52-83y	85	BMD,OC,BAP,PYD, DPYD
Hu et al, 2014, USA[130]	CS	TP: 11.6-20.4% energy	DXA, QCT	801 women and 857 men (age 62+/-10 years)	1658	BMD
Ilich et al, 2003, USA[131]	CS	1.04 g/kg/d	DXA	Older F, >5 post, 68.7+/-7.1y	136	BMD, BMC
Iuliano-Burns et al, 2005, AUS[132]	CS	76g/d	DXA	7-20 year old Male twins (Monozygotic n=30) and Dizygotic (n=26)	56	BMC, CT, PW, EW
Jaime et al, 2006, Brazil[33]	CS	1.2 g/kg/d	DXA	Men- Over 50y	277	BMD
Jones et al. 2001, Tasmania[55]	CS	83g/d	DXA	Boys and Girls Aged 8 years old	330	BMD
Knurick et al. 2015, USA[34]	CS	69-97g/d	DXA	Adult men and women, 18-50 y	81	BMD
Kumar et al, 2010, India[35]	CS	45.7g/d	DXA	Women aged 20-69 years	225	BMD
Lacey et al, 1991, Japan[36]	CS	1.35 g/kg/d	SPA	Asian pre F(35-40y) and post F (55-60y)	178	BMC
Langsetmo et al., 2015 Canada[62]	LS	0.79g/Kg/d	DXA	Men and women aged over 25 years old	6510	BMD

<b>Study</b>	<b>Design</b>	<b>Mean Protein **</b>	<b>Method</b>	<b>Population</b>	<b>n</b>	<b>Outcomes measured</b>
Lau et al, 1998, China[37]	CS	0.65 g/kg/d	DXA	Vegetarian Post F, 70-89y	76	BMD
Libuda et al. 2008, Germany[133]	CS	1.3 g/Kg/d	pQCT	228 Children and adolescents 8-14 years old		
Libuda et al. 2011, Germany[134]	CS	42.7-46.1 g/d	pQCT	Pre-pubertal children	107	BMC, Cortical Area
Loenekke et al. 2010, USA[50]	CS	72-91g/d	DXA	Males and Females, 22+/-3 years	27	BMD, BMC
MacDonald et al, 2005, UK[135]	CS	79.4g/d	BTM, DXA	45-54y women	5119	PYD, DPYD, BMD
Meng et al. 2009 AUS[136]	LS	80.6 g/d, 1.2 g/kg/d	DXA	Community-dwelling premenopausal women aged 75 +/- 3 years old	862	BMC
Metz et al, 1993, USA[137]	CS	1.24 g/kd/d	SPA	Pre F Caucasian (24-28y)	38	BMD, BMC
Michaelsson et al, 1995, Sweden[38]	CS	59 g/d	DXA	F 28-74y, Caucasian	175	BMD, OC
Nakamura et al, 2004, Japan[39]	CS	1.29 g/kg/d	BTM	Elderly post F, mean age=68.3y , range 43-79	43	OC,BAP,DPYD,NTX
Neville et al, 2002, UK[138]	CS	66-98 g/d	DXA	238 M and 205 F, at both 15 and 20-25 y	443	BMD
New et al, 1997, UK[40]	CS	81+/-22 g/d	DXA	Women aged 44-50 years (Premenopausal)	994	BMD
Oh et al, 2013, Korea[102]	CS	TP 45.0-52.3 g/d	QUS	Men and POM F aged 50-70 years	3330	Calcaneal Bone Stiffness Index
Orozco et al, 1998, Spain[41]	CS	TP: 73.4(17.9) g/d	DXA	Premenopausal women aged 42years old	76	BMD
Orwoll et al, 1987, USA[42]	CS	-	CT,SPA	Men	154	BMC
Pearce et al. 2010, UK[139]	CS	Median: 87.7g/d	Bone Markers	Men aged 49-52 years	412	CTX
Promislow et al, 2002 USA[140]	CS	72.5 g/d	DXA	M/F 55-92y; 572F 388M	960	BMD
Quintas et al, 2003, Spain[43]	CS	1.4-1.7g/d	DPA	Pre F	164	BMD

<b>Study</b>	<b>Design</b>	<b>Mean Protein **</b>	<b>Method</b>	<b>Population</b>	<b>n</b>	<b>Outcomes measured</b>
Rapuri et al, 2003, USA[44]	CS	53.7-71.2 g/d	DXA	Post F- 65-77y	473	BMD,NTX, OC
Rubinacci et al, 1992, Italy[45]	CS	68-83g/d	SPA	Post F	120	BMC
Sahni et al. 2013, USA[141]	CS/LS	81g/d (Men) 77g/d (Women)	DXA	1,280 men and 1,639 women	2919	BMD
Tanaka et al, 2001, Japan[142]	CS	1.3 g/Kg/d	QUS	Pre F- 18-22y	965	Stiffness Index Calcaneus
Teegarden et al, 1998, USA[46]	CS	1.21 g/Kg/d	DXA	Young pre F	215	BMC, BMD
Thorpe et al, 2008, USA[47]	CS	74.7g/d	DXA	Postmenopausal women mean age 68+/-6 years	161	BMD
Tylavsky and Anderson, 1988, USA[143]	CS	1.01 g/Kg/d	SPA	60-98y elderly F	375	BMC, BMD
Vatanparast et al, 2007, Canada[144]	CS	20-25 years: 68-119g/d	DXA	Young adults (59 males, 74 females)	133	BMC, BMD
Wang et al, 1997, USA[48]	CS	0.97 g/Kg/d	DXA	Older post F	125	BMD, BMC
Wang et al. 1999, USA[145]	CS	1.05 g/Kg/d	QUS	63 18-18 year old women	63	BUS, SOS
Weikert et al, 2005, Germany[146]	CS	67.9g/d	QUS	F 35-67y	8178	Os calcis
Whiting et al, 2002, Canada[147]	CS	1.15 g/Kg/d	DXA	M 39-42y	57	BMD
Yazdenpanah et al, 2007, The Netherlands [148]	CS	81.3g/d , 1.1g/Kg/d	DXA	Men and Women aged 55 years and over	5304	BMD
Zhang et al. 2010, China[149]	LS	1.7 g/Kg/d	DXA	757 Girls (Mean age 10 years)	757	BMC

AP, animal protein; BA, Bone area; BAP, Bone Alkaline Phosphatase; BMC, bone mineral content; BMD, bone mineral density; BTM, Bone turnover markers; BUA, Broadband Ultrasound Attenuation; CS, Cross-sectional Study; CT, Computer Tomography; DPA, Dual Photon Absorptiometry; DPYD, Deoxypyridinoline; DXA, Dual energy x-ray absorptiometry; EW, Endosteal Width; HPO, Hydroxyproline; HR-pQCT, High resolution Peripheral Computed Tomography; LS, Longitudinal Study; NTX, N-terminal Telopeptide of Collagen; pQCT, Peripheral Quantitative Computer Tomography; OC, Osteocalcin; P1NP, Procollagen type 1 N-Terminal Propeptide; PW, Periosteal Width; PYD,

Pyridinoline; QUS, Quantitative Ultrasound; SOS, Speed of Sound; SP, soy protein; SPA, Single photon Absorptiometry; SSI, Strength-Strain Index; TP, total protein; VP, vegetable protein.

**Table 2: Characteristics of the 29 studies reporting fracture or osteoporosis diagnosis data (6 of which also in table 1): Cohort and cross-cultural studies.**

Study <sup>a</sup>	Mean Protein	Population	Length	Total n	Fracture/BMD site	Protein type	Parameter	Confounder Adjustments
Abelow et al, 1992, USA cross cultural[150]	10.4g/d-77.8g/d AP	F aged over 50y	-	34 studies, 16 countries	Hip fracture	AP	Fracture Incidence	Age
Beasley et al. 2014, USA[119]	<13.3% to ≥15.6% of energy intake from protein	Women aged 50-79 y at baseline	6y	144,580	Any, Hip, Spine, Forearm	TP	HR	Age, BMI, race-ethnicity, calibrated energy intake, general health, physical activity, history of fracture at age 55 y, history of parental fracture, current smoking, corticosteroid use, glucocorticoid use, treated diabetes, rheumatoid arthritis, and hormone use
Dargent-Molina et al, 2008, France[57]	TP:46(7.5) g/d, AP:29 (8.8) g/d, VP:12(3.0) g/d	POM women	8.37y	36217 (2408 with fracture)	Any low impact fracture	TP AP VP	RR	Adjusted for BMI, physical activity, parity, maternal history of hip fracture, HT use, smoking status, and alcohol intake
Feskanich et al, 1996, USA[59]	79.6 g/d median	F, 35-59y	12y	85,900	Forearm, Hip, Fracture	AP TP VP	RR	Age, BMI, vigorous activity per week; menopausal status, use of postmenopausal hormones; cigarette smoking; use of thyroid hormone medication, thiazide diuretics and alcohol/caffeine.
Frassetto et al, 2000, USA cross cultural[151]	48 to 110.9 g/d	F aged over 50y		33 countries	Hip fracture	TP AP VP	Hip Fracture Incidence	-
Gunn et al, 2014, New Zealand[28]	79 g/d	POM women, 60 years of age DXA	Cross-sectional	142	Osteoporosis diagnosis	TP	Protein intake (g) by BMD diagnosis	-

Study <sup>a</sup>	Mean Protein	Population	Length	Total n	Fracture/BMD site	Protein type	Parameter	Confounder Adjustments
Key et al, 2007, UK[152]	Women: 73.1 (21.6) g/d 77.8(22.6) g/d	Men and Women aged 20-89 years	5.2y	26 749 women and 7947 men	All sites, fractures (including high trauma)	TP	Incident Rate Ratio	Method of recruitment and adjusted for age, smoking, intakes of energy and each other nutrient, alcohol consumption, body mass index, walking, cycling, vigorous exercise, other exercise, physical activity at work, marital status and, for women, parity and use of hormone replacement therapy
Langsetmo et al, 2015, Canada[62]	TP: 0.79(0.60-1.03). AP:17.6(12.8-23)g/d VP:24.3(18.8-31.0)g/d	M and F, aged 25-49 and ≥50 years	5y	6510	Fragility fracture: n=4543  Main fracture: n=4570	TP	HR	Age, height, TEI, centre (women only), education, smoking, alcohol intake, physical activity, sedentary hours, calcium and vitamin D supplement use, hormone therapy (women only), bisphosphonate use (women only), and diagnosis of osteoporosis (women only)
Meyer et al, 1997, Norway[61]	0.8	M/F (mean age 47.1y)	11.4y	19752 F 20035 M	HF- F HF-M	AP	RR	Age at screening, body height, body mass index, self-reported physical activity at work and during leisure time, diabetes, disability pension, marital status, and smoking
Misra et al, 2011, USA[63]	64g/d (energy adjusted)	M/F mean age=75 years	11.6y	946 (n=100 HF)	HF	TP	HR	Age, sex, weight, height and total energy intake
Munger et al, 1999, USA[58]	1.2 units g/Kg/d	Postmenopausal F (55-69y)	1-3y	32 050	HF	AP, TP, VP	RR	Age, body mass index, number of pregnancies, smoking, alcohol use, estrogen use, and physical activity.
Mussolino et al, 1998, USA[60]	<56g/d - >98g/d	Caucasian M (45-74y)	22y	2879	HF	TP	RR	BMI, previous fracture, smoker, physical activity, alcohol, chronic health condition, calcium intake, weight loss.
Sahni et al, 2010, USA[64]	Men TP: 75-79.0g/d	Men and women aged	7 to 14 years	3656	HF	TP, AP VP, AP:VP ratio	HR	Sex and menopause status (group 1: men; group 2: premenopausal women; group 3: postmenopausal women), age(years), weight at baseline (kg), height at baseline (m), physical activity index, intake of energy (MJ/day) and total



Study <sup>a</sup>	Mean Protein	Population	Length	Total n	Fracture/BMD site	Protein type	Parameter	Confounder Adjustments
	AP: 52-54 g/d , VP: 23-25g/d	mean= 55 (9.9)years						vitamin D (IU/day), and smoking status (current versus former/never) and calcium intake
Sellmeyer et al, 2001, USA[153]	49.8g/d	F > 65y old	7.0y +/- 1.5y	1035	HF	VP, AP, AP:VP	RR	Age and body weight
Zhong et al, 2009 USA[154]	Mean(SE)= 61+/-0.8 g/d	POM women 50+ y old	<7y	2006	All fragility fractures	TP	OR	Age, race, body mass index (underweight/normal, overweight, obese), physical activity level, smoking status, alcohol use (heavy, moderate/none), hormone use, general health status, osteoporosis, arthritis, vision impairment, and stroke.
Zhang 2005[155]	SP: 9.6g/d Non Soy: 134g/d	Women aged 40-70 years old	4.5 y	24403	All fractures	SP	RR	Age, body mass index, hours of exercise per week, cigarette smoking, alcohol consumption, history of diabetes mellitus, level of education, family income, season of recruitment, and intakes of total calories, calcium, non-soy protein, fruits, and vegetables

<sup>a</sup>All studies are cohorts unless otherwise stated. AP, Animal Protein; BMD, Bone Mineral Density; HR, Hazard Ratio; OR, Odds Ratio; POM, Postmenopausal; RR, Relative Risk; SP, Soy Protein; TEI, Total Energy Intake; TP, Total Protein; VP, Vegetable Protein.

**Table 3: Characteristics of the 29 studies reporting fracture or osteoporosis diagnosis data (6 of which also in table 1): 13 Case-control studies**

<b>Study</b>	<b>Protein intake</b>	<b>Measure</b>	<b>n</b>	<b>Group/outcome</b>	<b>Confounder Adjustments</b>
Alissa et al, 2011, Saudi Arabia, Non-Prospective [117]	77g/d	DXA	122 POM Women, aged 50-60 years	Normal BMD vs. Osteopenic	Non adjusted for confounders
Chevalley et al. 2011, Switzerland Prospective[156]	47-63 g/d	DXA	176 boys- measured during pre-puberty and adolescence	Fracture vs. No Fracture	Non adjusted for confounders
Chiu et al, 1997, Taiwan Non-Prospective[23]	1.09 g/Kg/d	DPA (BMD)	258 Older post F	OR for Osteopenia diagnosis: LS and FN	Age, BMI, physical activity, calcium intake, non-protein energy intake, long term vegan/vegetarianism
Coin et al, Italy, 2008 Non-Prospective[25]	1.02 g/Kg/d	DXA	136 Males, mean age 73.9+/-5.6 years	OR for low THBMD	BMI
Farrin et al. 2008, Iran Non-prospective[157]	81.4g/d	DXA	58 POM women	LSBMD based diagnosis: Normal/Osteopenic/Osteoporotic	Unadjusted
Kim et al, 2008, Korea Non-prospective[158]	TP= 60g/d, AP= 19g/d, VP= 40g/d	DXA	271 POM women: 134 cases and 137 controls	Osteoporotic vs. Non-Osteoporotic TP, AP, VP	Age, smoking, alcohol drinking, BMI, exercise, family history of osteoporosis, and energy intakes
Martinez-Ramirez et al, 2012, Spain Non-Prospective[65]	TP:105 (1.0) g/d. AP:66-70 (1.3) g/d. VP: 38 (0.63)g/d,	Aged 65 years or over, cases	167 cases and 167 controls	OR : All low energy fractures. TP, AP, VP, AP:VP ratio	Age, sex, energy intake, vegetable protein intake or animal protein intake, serum vitamin C, calcium intake, underlying chronic disease, home access, Katz's index, physical activity, HDL cholesterol and MUFA/PUFA intake.
Nieves et al, 1992, USA Non-prospective[66]	<24g/d to >55g/d	F 50 to 103y	329 (161 cases, 168 controls)	OR Hip fracture	Hospital, age, BMI, oestrogen use, chronic disease status

Park et al, 2014, Korea Non-prospective[159]	81.93+/- 52.31 g/d	DXA	1157 PRE women	Z-Score $\geq 0$ , Z-score $< 0$	Non-adjusted
Perez-Durillo et al, 2011, Spain Non-prospective[160]	Cases 60 (19)g/d; controls 94 (19) g/d	Women > 65 y	44 cases and 42 controls	TP, Hip fracture c	BMI, carbohydrate intake and calcium intake
Preisinger et al, 1995, Austria Non-prospective[161]	15 % total energy, 45- 96 g/d	Osteoporosis	23 POM women 50-70 years old	TP, AP, VP, Osteoporotic vs. Non Osteoporotic	Not required as just intake data
Samieri et al, 2013, France Prospective [162]	70-76 g/d	Men and women >65y	1482	Incident fracture of hip, spine or wrist	Not adjusted for confounders
Wengreen et al, 2004, USA Non-prospective[67]	1.2g/Kg/d	50-89y M/F	2501 (1157 cases, 1334 controls)	Cases Controls, Hip (OR), TP AP VP	BMI, smoking, alcohol, physical activity, oestrogen use, gender, total Calcium and Vitamin D intakes (diet and supplements), potassium intake, age. AP model also adjusted for VP intake, VP model also adjusted for AP intake.

AP, Animal Protein; BMD, Bone Mineral Density; DPA, Dual Photon Absorptiometry; DXA, Dual X-Ray Absorptiometry; LSBMD, Lumbar Spine Bone Mineral Density; OR, Odds Ratio; POM, Postmenopausal; PRE, Premenopausal; THBMD, Total Hip Bone Mineral Density; TP, total protein; VP, vegetable protein

**Table 4: Characteristics of the 30 intervention studies reporting randomised control trials of dietary protein supplementation on bone health outcomes**

<b>Study, Country, Duration</b>	<b>Baseline protein intake</b>	<b>Supplement vs control</b>	<b>Change in protein intake</b>	<b>Subject Total n</b>	<b>Outcomes Measured</b>
Alekel et al, 2000, USA, 24wks[73]	No information in paper	40g/d Soy vs Whey	+ 40g/d	N = 24 SPI+; N = 24 SPI-, N = 21 whey protein (control) PERI F	BMC, BMD, BAP
Aoe et al, 2001, Japan[96]	No information in paper	40mg/d MBP vs Placebo	+ 40mg/d (MBP)	32 PRE F	% change data only: Calcaneal BMD
Aoe et al 2005, Japan, 6mo[69]	No information in paper	40mg/d MBP vs Inactive placebo	+40mg/d (MBP)	27 PERI F	NTX, OC, BMD
Arjmandi et al, 2003, USA, 3mo[87]	Mean (SE): Soy group – 60(6)g/d, MBP group (75(9) g/d	40g/d Soy protein vs MBP	+40g/d	42 POM F	BAP, DPYD
Cao et al, 2011, USA Crossover study 14wks (7 weeks each arm)[1]	No information in paper	61g/d diet ('lower protein control-US daily recommendation) vs. 118g/d ('higher protein' group) diet	+/-57g/d	N=16 40-75 year old POM F,	NTX, DPYD
Ceglia et al, 2009 Crossover study 41 d[82]	Mean (SD): 69.1 (22.1) g/d	Mean (SD)2.1(2.02) Low vs. 96.7(5.7) High	-37.0 g/d (low) +27.6 g/d (high)	M/F 54-82 years old N=10	OC, NTX
Cuneo et al, 2010, Brazil[88]	Mean (SD) : 67(18.8) g/d (placebo), 61.9(24) g/d (collagen)	Hydrolysed collagen (10g/d protein) vs. maltodextrin placebo	+10 g/d	N=36 collagen, N=35 placebo, 45-65 year old POM F	BAP, CTX, OC
Dalais et al, 2003, AUS, 3mo[89]	109(7) g/d Soy group, 112(6)g/d Placebo	40g Soy protein vs casein placebo	+40g/d	106 POM F 50-75 y	PYD, DPYD
Dawson-Hughes et al 2004,USA, 63d[91]	No information in g (17-18% of total energy)	Mean(SD): High: 57.6(8.2) g/d protein , Low: 2.8(0.5) g/d protein	No baseline data in grams	32 Elderly M/F	NTX, OC
Evans et al, 2007, USA Crossover study, 9 months[78]	No information in paper	25.6g Soy protein isolate (I) vs. Milk protein isolate (p),	+25.6g/d	Postmenopausal women N=22, Mean age 63 years	BMD, BAP, CTX

<b>Study, Country, Duration</b>	<b>Baseline protein intake</b>	<b>Supplement vs control</b>	<b>Change in protein intake</b>	<b>Subject Total n</b>	<b>Outcomes Measured</b>
Hunt et al, 2009, USA, 7 wks Cross over study [83]	63(15) g/d LC 69(17) g/d HC	exercise counterbalanced across groups (1/2 in each group exercise, 1/2 in each group no exercise) Protein g/d: LCLP 58, LCHP 112, HCLP 59, HCHP 115	LCLP -5g/d, LCHP +49g/d, HCLP -10g/d, HCHP +46g/d	N=13 in two LC arms, n=14 in two HC arms Post F	DPYD, OC, BAP, TRAP
Ince et al 2004, USA, 2wks Crossover study[94]	1.1 g/kg/d	Change to low (0.8g/Kg/d) protein diet	-0.2g/Kg/d	39 Pre F, 22-39y	NTX, OC
Jenkins et al, 2003, USA, 2mo Crossover study[85]	No information in g (18% of total energy)	Vegetable diet: 189g/d protein, (16% total energy) vs Control diet: 111g/d protein (27% total energy)	+78g/d	20 Middle aged M/F	NTX, BAP
Kenny et al, 2009, USA, 1 year [74]	Mean +/-SD : Soy group- 62.5 (13.7) g/d, Mixed control group- 57.0(21.9)	18g Soy protein (I) vs. 18g Mixed control protein (Casein, Whey and Egg) (p). No isoflavones in these two study arms	+18g/d	Women over 60 years old (mean=71y)	BMD, BAP, NTX
Kerstetter et al, 1999, USA, 4d Crossover study[80]	1.0g/Kg/d	High (2.1g/kg/d)vs low (0.7g/kg/d) protein	+1.1g/Kg/d High -0.3g/Kg/d Low	16 Pre F, 20-40g	OC,BAP, NTX
Kerstetter et al, 2015, USA, 18mo[68]	Mean (SEM): 72.9(1.8) Maltodextrin Group, 73.9(1.9) Whey Group	45g Whey protein or Isocaloric maltodextrin	+45g/d	Men over 70 y and women over 60 years, n=121	BMD, P1NP, CTX, OC
Khalil et al, 2002, US, 3mo[86]	No information in paper	Soy vs Milk protein (40g)	+40g/d	64 M, 59.2+/-17.6y	BAP, DPYD

<b>Study, Country, Duration</b>	<b>Baseline protein intake</b>	<b>Supplement vs control</b>	<b>Change in protein intake</b>	<b>Subject Total n</b>	<b>Outcomes Measured</b>
Lampl et al. 1978, New Guinea, 8 mo[92]	11g/d	Normal diet (11g/d)vs. normal diet plus 20g/d milk protein supplement	31g/d	7-13 year old Bundi children, male and female, with low protein intakes	Periosteal breadth , Endosteal breadth, Compact bone breadth
Martin-Bautista et al. 2011, Spain, 4 mo[93]	Mean(SD): 34.1(26.0)g/d Collagen , 33.0(18.1)g/d Placebo	Drink containing partially hydrolysed collagen vs. placebo drink	+4.25g/d	60 children aged 6-11 years	BAP, OC, TRAP, CTX
Roughead et al, 2003, USA, 8wk Cross over study[81]	No information in paper	High meat diet: 117g/d (20% of energy) versus low meat diet: 68g/d (12% of energy)	+49g/d	15 POM F	HPO, OC, NTX, BAP
Schurch et al, 1998 Switzerland, 6mo[95]	Mean (SD) 45.0 (15.2) g/d protein group vs. 51.0 (19.0) g/d control group	Total protein (20g/d) vs placebo	+20g/d	82 Elderly M/F 80.7y+/- 7.4	%Change data only: DPYD, FSBMD, OC, PFBMD, PYD, BMD
Shapses et al, 1995, USA, crossover study, 5d[84]	0.99 g/Kg/d (no information in g)	LPHC(0.44g/Kg/D protein, p) vs. HPHC (2.71g/kg/d, I) Calcium in both groups=1600mg/d	-0.55g/Kg/d Low +1.70g/Kg/d High	21-42 year old males and females	HPO
Spence et al, 2005, USA, crossover study 28d per phase[79]	No information in paper	Soy protein isolate without isoflavones diet (96g/d) vs. casein-whey protein diet (91g/d)	+0 (as comparing protein type not dose, although 5g/d difference in actuality)	N=15 Post F	BAP ,OC ,NTX
Tkatch et al, 1992, Switzerland, 38days[72]	No information in paper	20.4g/d Protein in nutritional supplement vs. the same nutritional supplement without protein	+20.4g/d	62 M/F elderly, mean age 82y	BMD,OC
Toba et al 2001, Japan, 16d[90]	No information in paper	MBP (300mg/d) vs inactive placebo	+300 mg/d MBP	30 M, 36.2y+/-8.5	NTX,OC

<b>Study, Country, Duration</b>	<b>Baseline protein intake</b>	<b>Supplement vs control</b>	<b>Change in protein intake</b>	<b>Subject Total n</b>	<b>Outcomes Measured</b>
Uenishi et al, 2007, Japan, 6mo[70]	No information in paper	40mg/d MBP vs inactive placebo	+40mg/d MBP	35 Pre F	BMD
Vupadhyahula et al, 2009, USA[75]	Mean (SD): 62.9(1.9) Soy protein group, 61.3(1.6) Milk Protein group	25g soy protein (no isoflavones), 25g milk (casein, whey) protein	+25g/d	203 Post F Mean (SE) age 64 0.6y	BMD, NTX
Yamamura et al, 2002, Japan[77]	No information in paper	MBP(40mg) vs inactive placebo	+40mg/d MBP	33 Pre F	BMD
Zhu et al, 2011, AUS, 2y[76]	Mean(SD): 76(18)g/d High protein group, 76(16) Low protein placebo group	High protein drink 30g vs. low protein drink (placebo) 2.1g	+27.9g/d	219 70-80 year old women	vBMD, BMD
Zou et al 2009, China, 8 mo[71]	No information in paper	Milk with 40mg MBP vs. Milk without MBP	+40mg/d MBP	57 women, 20 years old	BMD

AUS, Australia; BAP, Bone Alkaline Phosphatase; BMC, Bone Mineral Content; BMD, Bone Mineral Density; CTX, C-Terminal Peptide of Collagen; DPYD, Deoxyypyridinoline; FSBMD, Femoral Shaft Bone Mineral Density; HCHP, High Calcium High Protein; HCLP, High Calcium Low Protein; HPO, Hydroxyproline; LCHP, Low Calcium High Protein; LCLP, Low Calcium Low Protein; MBP, Milk Basic Protein; NTX, N-Terminal Peptide of Collagen; OC, Osteocalcin; P1NP, Procollagen type 1 N-Terminal Propeptide; PERI, Perimenopausal; PFBMD, Proximal Femoral Bone Mineral Density; POM, Postmenopausal; PRE, Premenopausal; PYD, Pyridinoline; SPI, Soy Protein Isolate; TRAP, Tartrate Resistant Alkaline Phosphatase; vBMD, volumetric Bone Mineral Density

## Figure Captions

**Fig. 1** PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-analyses flow diagram to illustrate search results

**Fig. 2** Protein intake and all low trauma fractures (lowest intake category: RR=1) Animal (top) Vegetable (middle) Total Protein (bottom)

## Online Resource Caption

Electronic Supplementary File containing further information on study methods and results as well as all Supplementary Figures and Tables.