

Emerging concepts of shear stress in placental development and function

L.C. Morley¹, D.J. Beech¹, J.J. Walker², and N.A.B. Simpson²

¹Leeds Institute of Cardiovascular and Metabolic Medicine, LIGHT Laboratories, University of Leeds, LS2 9JT, UK ²Academic department of Obstetrics and Gynaecology, Level 9, Worsley Building, University of Leeds, LS2 9JT, UK

*Correspondence address. Leeds Institute of Cardiovascular and Metabolic Medicine, LIGHT Laboratories, University of Leeds, LS2 9JT, UK. E-mail: l.c.morley@leeds.ac.uk orcid.org/0000-0002-1109-1396

Submitted on October 15, 2018; resubmitted on March 3, 2019; editorial decision on March 20, 2019

ABSTRACT: Blood flow, and the force it generates, is critical to placental development and function throughout pregnancy. This mechanical stimulation of cells by the friction generated from flow is called shear stress (SS) and is a fundamental determinant of vascular homeostasis, regulating remodelling and vasomotor tone. This review describes how SS is fundamental to the establishment and regulation of the blood flow through the uteroplacental and fetoplacental circulations. Amongst the most recent findings is that alongside the endothelium, embryonic stem cells and the villous trophoblast are mechanically sensitive. A complex balance of forces is required to enable effective establishment of the uteroplacental circulation, while protecting the embryo and placental villi. SS also generates flow-mediated vasodilatation through the release of endothelial nitric oxide, a process vital for adequate placental blood flow. The identification of SS sensors and the mechanisms governing how the force is converted into biochemical signals is a fast-paced area of research, with multiple cellular components under investigation. For example, the Piezo I ion channel is mechanosensitive in a variety of tissues including the fetoplacental endothelium. Enhanced Piezo I activity has been demonstrated in response to the Yoda I agonist molecule, suggesting the possibility for developing tools to manipulate these channels. Whether such agents might progress to novel therapeutics to improve blood flow through the placenta requires further consideration and research.

Key words: placenta / shear stress / mechanosensing / Piezo I / endothelial cells

Introduction

A successful pregnancy outcome is dependent upon effective placentation. This occurs within the first half of pregnancy and describes the development of two distinct but independent circulatory systems involving both the mother (uteroplacental) and baby (fetoplacental). Following implantation, extravillous trophoblast cells migrate out from placental chorionic villi and invade spiral arterioles in the maternal decidua (Kingdom, 1998). The subsequent remodelling transforms these narrow vessels into wider conduits, generating high flow and low resistance. Oxygen and nutrient-rich maternal blood is thus propelled into the evolving IVS (Fig. 1).

To support fetal circulatory and metabolic requirements, blood flow to the uterus increases progressively with advancing gestation. Maternal cardiac output is increasingly directed to the uterus secondary to falling vascular resistance and haemodynamic changes in the systemic circulation (Osol and Moore, 2014). The uterine arteries and endometrial vessels must therefore undergo vascular remodelling to accommodate these profound changes in perfusion (Park *et al.*, 2017). At the same time, the fetoplacental vasculature is develop-

ing through vasculogenesis and branching angiogenesis within the villi (Kingdom *et al.*, 2000). This arborization creates a network of capillaries enabling maximal diffusion of nutrients and gases across the maternal–fetal interface. The circulation is completed by the umbilical arteries transporting deoxygenated blood and waste to the villous tree, to enable replenishment by the maternal supply and returned to baby via the umbilical vein. Effective and responsive gas and nutrient exchange is therefore dependent upon blood flow that can adapt and vary according to circumstance (Kingdom, 1998).

A key element of vascular adaptation in pregnancy is vasodilatation through the production of nitric oxide (NO) and other agonists from the endothelial lining of blood vessels. Within the placenta, fetoplacental vessels lack autonomic innervation and control of vascular tone is therefore dependent upon these vasoactive mediators (Learnmont and Poston, 1996). The most powerful physiological stimulator of NO production is shear stress (SS). This is the mechanical force generated during each cardiac cycle by the haemodynamic force of blood flow. Fluidic SS has long been recognized as being critically important for processes including angiogenesis, vasculogenesis, and control of vascular tone (Ando and Yamamoto, 2013). However, knowledge




Figure 1 Schematic representation of blood flow through the placenta, with contributions from the uteroplacental and fetoplacental circulations. Blue and red arrows show the flow directions of oxygenated (red) and deoxygenated (blue) blood. The white box highlights the villous tree, comprising trophoblast cells. Vascularisation from progressive branching within the fetal circulation forms capillaries in the terminal villi, which are the functional sites of maternal-fetal exchange. Dashed white arrows demonstrate the maternal blood flow through the IVS. The red-to-blue colour gradient represents oxygenation status. Figure from *Slator et al. (2018)*.

of the molecular mechanisms underpinning how cells respond to SS and how this translates into biochemical signals is only now coming to the fore. Failure of these adaptations can lead to pregnancies complicated by gestational disorders including pre-eclampsia and fetal growth restriction (FGR), the pathogenesis of which centres upon impaired spiral artery remodelling, high vascular resistance, and placental hypoperfusion, helpfully reviewed elsewhere (*Kingdom, 1998*). In this New Research Horizons review we present a summary of the most current knowledge of mechanosensing in both placental development and vascular function alongside the key research questions, namely as follows:

- How is SS sensed and transduced in the placenta?
- Does aberrant SS drive or contribute to placental dysfunction?
- Could identifying the molecular complexes responsible for the SS response provide novel targets for therapeutic agents to treat placental dysfunction?

Our focus is on the role of newly identified SS sensors and the opportunities this presents for future research and therapeutic applications.

Flow dynamics and SS

SS is produced by haemodynamic force across the endothelium, and can be quantified using measurements of the inner diameter of the vessel, velocity of flow, and blood viscosity (*Fig. 2*) (*Wareing, 2012*). *In vivo* estimations of SS are complicated by the pulsatile nature of flow, the viscosity of blood contents, and the structural architecture of the vascular tree (*Baratchi et al., 2017*). Depending on the type of vessel, its branching pattern, and variables such as temperature,

flow may be laminar or turbulent and intraluminal force will therefore vary.

Even during embryogenesis SS is involved in the vascular expansion and remodelling required for growth. As the heart starts to beat, haemodynamic force triggers endothelial cells (ECs) to develop a vascular network through vasculogenesis and angiogenesis (*Hyman et al., 2017*). Correspondingly, mice with a disrupted heartbeat show a lack of yolk sac vascularization alongside lethality (*Huang et al., 2003*).

SS is sensed by the endothelium and, through mechanotransduction, activates multiple downstream signalling pathways, resulting in the release of vasoactive mediators such as NO. The endothelium is therefore dynamic, responding to local cues to control vascular homeostasis (*Chatterjee, 2018*). As such, the type and magnitude of SS has a significant impact on the endothelial phenotype (*Malek et al., 1999*).

In the systemic circulation, ECs exposed to physiological SS show increased production of vasodilators and antioxidants, with a reduction in vasoconstrictors and inflammatory mediators (*Malek et al., 1999*). Likewise, disruption of normal SS, for example at sites of turbulent flow, is considered pro-atherosclerotic (*Baratchi et al., 2017*). Indeed, an impaired response to SS has been linked to a variety of cardiovascular disease processes such as aneurysms, thrombosis, and hypertension (*Ando and Yamamoto, 2013*).

Furthermore, in addition to the endothelium, mechanosensing is now being widely reported in multiple cell types that are subjected to fluid flow. This includes the epithelial villous tissue, which has important implications for placentation. It is therefore no surprise that SS and the subsequent cellular response is postulated to have multiple roles in homeostasis from acute modulation of vasomotor tone to angiogenesis and vasculogenesis (*Baratchi et al., 2017*).




Figure 2 Cross-sectional diagram of a vessel showing SS produced by blood flow. SS in force/unit area (dyne/cm²) can be calculated using the equation $SS = (4pQ)/\pi r^3$, where r is the vessel radius (μm), Q is blood flow velocity ($\mu\text{L/s}$), and p is blood viscosity in poise (dyne.s/cm³). 10 dyne/cm² = 1 Pascal. Adapted from Malek *et al.*, 1999, Wareing, 2012.




Figure 3 Components of the EC, each with postulated roles in mechanosensing. G-protein coupled receptors (GPCR) are membrane-spanning receptors e.g. GPR68 (Xu *et al.*, 2018). Integrins are membrane-spanning receptors linking cytoskeletal proteins to the cell matrix. Ion channels may be rapidly activated in response to flow. Examples include members of the TRP family and Piezo I, which are both non-selective cation channels. The glycocalyx is a matrix composed of glycoprotein and glycolipid moieties that covers and protects the cell membrane. Adhesion molecules include proteins on the cell surface (e.g. platelet adhesion cell adhesion molecule), which are expressed on cell-cell junctions. The cytoskeletal scaffold encompasses microtubules, microfilaments and intermediate protein filaments (Chatterjee, 2018). Created using SMART Servier Medical Art (LES LABORATOIRES SERVIER, SAS, France) and adapted from Shihata *et al.* (2016).

SS sensors

There is a growing body of literature dedicated to the identification of mechanosensors, including a variety of proteins, receptors, and transmembrane channels (Fig. 3) (Yamamoto *et al.*, 2006; Shihata *et al.*, 2016; Chatterjee, 2018). The architecture of the cell itself may also be involved in the SS response. The cytoskeleton is composed of protein filaments that scaffold the ECs and may be deformed by shear on the cell surface. This impacts on cellular components, such as integrins (membrane spanning receptors), focal adhesion proteins, and the extracellular matrix, where the force is transduced (Chatterjee, 2018). Integrins themselves may also sense SS, affecting the cytoskeletal filaments directly (Baratchi *et al.*, 2017; Chatterjee, 2018). Force may also be transmitted to the cytoskeleton via the heparan sulfate, chondroitin sulfate, and hyaluronic acid moieties of

the glycocalyx on the EC membrane. As such, deformation of the cytoskeleton by shear may result in EC reorganization and remodelling (Chatterjee, 2018).

It is possible that these putative factors are co-dependent with several SS sensors working in conjunction to form a mechanosome complex (Chatterjee, 2018). One such complex is proposed by Tzima *et al.* (2005). Here the authors suggest that platelet EC adhesion molecule (PECAM-1/CD31), which is expressed on endothelial cell-cell junctions is the direct SS sensor. This results in activation of the junction receptor vascular endothelial cadherin, alongside vascular endothelial growth factor (VEGF) receptor 2 (Tzima *et al.*, 2005). These events may trigger activation of the integrins (Tzima *et al.*, 2005).

The identification of specific fast-acting mechanosensors, which could be pharmacologically manipulated, is of particular interest. For example, the transient receptor potential (TRP) channels are a

major class of calcium ion (Ca^{2+})-permeable and non-selective cationic channel. Ca^{2+} influx through TRP channels produces smooth muscle contractility, alterations in vascular permeability, and remodelling (Baratchi et al., 2017). However, in murine models knock out of the TRP channel, TRPV4, did not affect survival, while TRPP2 homozygous knock-out mice had delayed lethality (Hartmannsgruber et al., 2007; Garcia-Gonzalez et al., 2010).

The ion channel subunit Piezo I was first discovered in 2010 (Hyman et al., 2017). Since then this mechanosensitive membrane protein has risen to prominence as a major player in SS sensing and is considered in more detail below.

Piezo I proteins

Piezo I channels are distinct from conventional ion channels due to their size (900 kDa), structural complexity as a large trimer of Piezo I proteins, and rapid activation/inactivation (Chatterjee, 2018). Cryo-electron microscopy has revealed the structure of Piezo I to be propeller-like, with three highly flexible blades and a central cap enclosing a Ca^{2+} -permeable non-selective cation channel (Fig. 4a and b) (Guo and MacKinnon, 2017; Saotome et al., 2018). The extracellular blade components are thought to act as force sensors within the vessel lumen, regulating the gated channel (Guo and MacKinnon, 2017). The most recent evidence suggests that these channels directly sense and modulate membrane tension via mechanisms, which are currently unknown (Cox et al., 2016; Syeda et al., 2016).

Our group reported that Piezo I responds to SS in human umbilical vein ECs (HUVECs) (Li et al., 2014). We then identified downstream proteins affected by Piezo I in these HUVECs, cultured in both static and flow conditions. After Piezo I depletion using siRNA there was a reduction in endothelial NO synthase (eNOS) activation and furthermore, abolition of VEGF-evoked phosphorylation of eNOS at serine 1177—known to enhance eNOS activity (Li et al., 2014). The mechanism by which SS-activated Piezo I leads to NO release is still under investigation. Wang et al. (2016) suggest that Piezo I is required for the release of endothelial ATP. This leads to activation of the G protein coupled receptor P2Y2 and downstream NO production (Wang et al., 2016). This suggests that Piezo I has a key role in regulating vasodilation through NO.

To further investigate the significance of Piezo I, our lab generated a mouse model with a disrupted endogenous *Piezo1* gene (Li et al., 2014). Inheritance of both the global homozygous and endothelial-specific knock out was lethal in pups at mid-gestation [embryonic days (E) 9.5–11.5]. The phenotype we observed was reduced yolk sac vascularization and FGR prior to *in utero* demise (Fig. 4b). Closer inspection revealed initial endothelial lattice formation preservation, but angiogenesis was significantly inhibited (Li et al., 2014) (Fig. 4c). The finding that Piezo I appears critical for early murine vascular development was also demonstrated by another group, suggesting that Piezo I could be a major player in human embryonic and placental vascular development (Ranade et al., 2014).

In humans, autosomal recessive loss of function mutations in *PIEZO1* causes generalized lymphatic dysplasia (Fotiou et al., 2015, Lukacs et al., 2015). This can present prenatally as non-immune fetal hydrops, which can result in death or pulmonary compromise at birth, secondary to pleural effusions (Martin-Almedina et al., 2018).

Oedema can also recur in childhood and be generalized, affecting the face, limbs, and genitals (Martin-Almedina et al., 2018). The most well-known *PIEZO1* mutation is autosomal dominant, where gain of function results in dehydrated xerocytosis, a disorder of erythrocyte stability (Zarychanski et al., 2012). Interestingly, this can also present with perinatal oedema (Martin-Almedina et al., 2018). The similarity between both phenotypes suggests that Piezo I has a critical role in sensing and developing the lympho-vascular flow system, although this is very much an emerging area of work (Hyman et al., 2017).

Research into Piezo I in other areas of adult disease is rapidly expanding. Recent publications include findings in the pancreas, gastrointestinal tract epithelium, cardiomyocytes, and chondrocytes (Alcaino et al., 2017; Liang et al., 2017; Servin-Vences et al., 2017; Romac et al., 2018). Despite these studies highlighting the pathological relevance of Piezo I, its role in mechanosensing in the placenta remains under researched.

SS sensing in pregnancy

Preimplantation

From the moment of conception, a preimplantation embryo is subjected to the force of fluid flow as it traverses the uterus by peristalsis. To determine if embryos can sense SS, Xie et al., (2006) exposed mouse embryos to either shear or static conditions (Xie et al., 2006). They found no surviving embryos after 24 h of shear at 1.2 dyn/cm² and reduced blastocyst cell numbers. An increase in apoptotic markers and stress-activated protein kinase [mitogen-activated protein kinase (MAPK)] suggested that shear was transduced prior to lethality (Xie et al., 2006). Shear-induced lethality after removal of the zona pellucida (ZP) suggests that the ZP may protect the embryo from mechanical force. This has implications for embryo handling in assisted conception transfer procedures, although transient SS caused by repeated pipetting did not affect development, despite increased MAPK phosphorylation (Xie et al., 2007).

New research has identified Piezo I channels in murine embryonic stem cells (Del Marmol et al., 2018). This raises the question of whether Piezo I activation by SS is important for cell differentiation. The lack of lethality in the knock out mouse models until mid-gestation suggests that Piezo I is not critical for early development but may have fine-tuning roles that are not yet fully identified (Del Marmol et al., 2018).

Post implantation

Transfer of oxygen to the fetus is critically dependent on flow past, and effective diffusion across, the trophoblast (Kingdom, 1998). Despite the importance of SS, extravillous trophoblast cells also form plugs within the spiral arteries, effectively preventing the flow of maternal blood into the IVS (James et al., 2018). It is well known that a hypoxic environment is important for early development. For example, in cases of miscarriage, erythrocytes have been found prematurely in the IVS (Jauniaux et al., 2000). Alongside regulation of oxygen tension, the presence of plugs will also impact on haemodynamics. Until recently, it was thought that plugs prevented any blood from entering the IVS until the end of the first trimester. However, recent work has suggested that capillary-sized channels enable a small, constant influx of blood that increases dramatically after 12 weeks of gestation




Figure 4 Structure and importance of Piezo 1 for murine vascular development. (a) Cryo-electron microscopy Piezo 1 trimeric structure produced using PyMol (Schrodinger Inc. Cambridge, UK) using PDB code 5Z10. Each monomer is represented as ribbons (monomer 1 cyan, monomer 2 magenta, monomer 3 green) (Zhao *et al.*, 2018). (b) Images of sibling yolk sacs (containing embryos) at E 9.5. Scale bar, 1 mm. Piezo 1 wild-type (+/+), haplotype (+/-), and homozygous knock out (-/-) are depicted (Li *et al.*, 2014). (c and d) Dissected yolk sacs stained for CD31, showing wild-type (c) and endothelial-specific *Piezo1*-modified (d) samples. EC: endothelial cell. Both sets of images show reduced vascularization at mid-gestation (Li *et al.*, 2014).

(Roberts *et al.*, 2017). Nevertheless, low SS conditions of <2 dyne/cm² (<0.2 Pa) are produced in the IVS (James *et al.*, 2018). James *et al.* (2012) found that at very low fluidic shear (0.5 and 2 dyne/cm²) trophoblasts did not undergo migration. However, at increased SS (4 and 6 dyne/cm²) migration in the direction of flow occurred. This suggests that high SS at this early stage would damage vascular remodelling by stimulation of migration away from the site of invasion (James *et al.*, 2012). Low SS may therefore be protective of delicate villous tissue through prevention of physical and oxidative stress.

Breakdown of plugs coincides with the end of the first trimester, where increasing blood flow generates SS which encourages trophoblast migration (James *et al.*, 2018).

Placental villi protruding from the trophoblast interact with maternal blood bathing the IVS. Scanning electron microscopy has been used to demonstrate minimal villous growth in BeWo cells and cultured human villous trophoblast under static conditions. By contrast, when exposed to SS, villi formation started at 1 hour. Miura *et al.* (2015) found short villous projections at high SS but these protrusions were

longer (>2 μm) at low SS, increasing over 12 hours. If, however, SS was stopped, the villi decreased (Miura et al., 2015). This therefore suggests that placental villi sense and respond to flow in their fluidic environment. Furthermore, there is a minimum requirement for SS as a “critical external cue” for villous formation (Miura et al., 2015).

When investigating the mechanism behind SS-induced villi formation, Miura et al. (2015) found that application of SS to BeWo cells increased intracellular calcium concentration (Ca^{2+}). Correspondingly, culturing cells in the presence of a Ca^{2+} -chelator inhibited villous growth (Miura et al., 2015). They suggest that the Ca^{2+} channel TRPV6, a member of the TRP family, is a candidate mechanosensor here (Fig. 3). Silencing TRPV6 using siRNA resulted in loss of the SS-induced intracellular Ca^{2+} response. They hypothesize that activation of TRPV6 results in re-localization of Ezrin (a protein linking the membrane and cytoskeleton) within the cell. Rapid Ezrin phosphorylation, but a lack of change in gene expression in response to SS, suggests that trophoblast cells are hyper-responsive to their dynamic fluid environment.

This interplay between the uteroplacental and fetoplacental circulations occurs throughout pregnancy, whereby SS produced by the flow of maternal blood in the IVS continues to affect the villous trophoblast. VEGFs produced by the placenta, including placental growth factor (PlGF), are contributors to angiogenesis and vasodilatation in the placenta. Exposing trophoblast cells from term placentas to SS at 1 dyne/cm² resulted in increased PlGF secretion and intracellular Ca^{2+} influx (Lecarpentier et al., 2016a). Magnetic resonance imaging of term pregnancies has revealed maternal blood flow velocity of 0.94 (+/− 0.14 mm.s^{−1}), which has been used to estimate SS values of <5 dyne/cm² in the IVS (Lecarpentier et al., 2016b). This computer modelling suggests that at term, maintaining low SS at the maternal-fetal interface remains important, and may promote maximal exchange of nutrients and waste.

Regulation of fetoplacental blood flow

Increased synthesis of NO, alongside other vasodilators such as prostacyclin, has been demonstrated in a perfused human placental cotyledon model (Wieczorek et al., 1995). Flow-mediated vasodilatation has also been shown in the perfused placenta model, and in chorionic plate arteries with incremental increases in flow decreasing vascular resistance (Learmont and Poston, 1996; Jones et al., 2015). Increased eNOS expression was also found in ovine fetoplacental ECs (FpECs) after exposure to SS (Li et al., 2004). Correspondingly, NO inhibition increased vascular resistance, a finding which was substantially elevated in FGR samples. Likewise, human placental arterial ECs from the chorionic plate demonstrated increased NO in response to SS (Jones et al., 2015). Wire myography of chorionic plate arteries showed relaxation in response to the NO donor sodium nitroprusside (Mills et al., 2005). Correspondingly, the authors observed increased relaxation in FGR samples (Mills et al., 2005).

This suggests that vessels from dysfunctioning placentae have the capacity to vasodilate over-and-above those from a healthy pregnancy.

In FGR, the anatomy of the placental vascular tree is altered with smaller numbers of immature villi, suggestive of impaired branching angiogenesis. This occurs alongside endothelial dysfunction and creates a hypoxic environment (Kingdom et al., 2000). As such, subsequent increases in transmural pressure in small vessels will heighten placental vascular tone, generating greater SS forces (Krause et al., 2013). This

is supported by the previously mentioned findings of reduced flow-mediated dilatation in FGR (Jones et al., 2015). In addition, total eNOS expression is higher in cells cultured from FGR samples (Myatt et al., 1997; Jones et al., 2015). We propose that mechanotransduction has an important role in regulating placental vascular tone under normal conditions and when SS is elevated in FGR. As such, heightened SS leading to increased production of NO by ECs may act in a compensatory capacity to overcome vascular resistance. When endothelial dysfunction is severe enough to prevent the normal response to SS, mechanosensory compensation will be insufficient and FGR may worsen (Jones et al., 2015; Morley et al., 2018).

A challenge to our hypothesis includes the finding that SS increases PlGF in trophoblast cells in a dose-dependent manner (Lecarpentier et al., 2016a). There are possible considerations:

- As Lecarpentier et al., (2016a) suggest, maternal placental hypoperfusion in FGR may instead ‘decrease’ the SS exerted on the trophoblast. This would result in the low levels of PlGF associated with FGR and pre-eclampsia (Lecarpentier et al., 2016a).
- The failure of PlGF production may represent trophoblast dysfunction and failed compensation, rather than being indicative of the SS response.
- There are microsites with different levels of shear and it is not surprising to find differing responses from the trophoblast versus those observed in uterine, endometrial and FpECs. This is supported by Sprague et al. (2010) who discussed the differences in viscosity secondary to the arrangement of erythrocytes, between capillaries and larger capacity vessels (Sprague et al., 2010).

Piezo1 in the fetoplacental circulation.

‘Stretch-activated non-selective cation channels’ were described in HUVECs in 1998 and it was found that they were involved in Ca^{2+} influx (Kohler et al., 1998; Brakemeier et al., 2002). A doubling in density of these channels was found in HUVECs from pregnancies affected by pre-eclampsia, although their molecular identity remained unknown (Kohler et al., 1998). Given our finding that Piezo1 is present in HUVECs and is critical for vascular development in the murine embryo, we sought to establish if Piezo1 had a role in human FpECs.

Our data revealed consistent Piezo1 gene expression in FpECs, alongside flow-dependent Piezo1 channel activity (26 pS) in fresh arterial ECs extracted from the chorionic plate (Fig. 5a and b). When Piezo1 was depleted in FpECs using siRNA, alignment in response to flow was significantly diminished, suggesting the importance of Piezo1 for SS sensing (Fig. 5c–e). It remains to be determined how SS activates Piezo1 channels in FpECs but our data support the findings in murine endothelium, suggesting that these channels are activated by SS in a membrane-delimited manner (i.e. in excised cell-free membrane patches) (Rode et al., 2017).

Regulation of uteroplacental blood flow

The production of endothelial NO, alongside prostacyclin and endothelial hyperpolarizing factor, has long been recognized as essential for the vasodilatory component of the expansive remodelling in the uteroplacental circulation (Osol and Moore, 2014). This can be demonstrated in uterine arteries, where an 8-fold increase in




Figure 5 Piezo1 is present in placental endothelium and is required for the SS response. (a) Quantitative PCR performed on mRNA isolated from FpECs readily detected *PIEZO1* mRNA expression. (b) Patch clamp recordings from freshly isolated placental arterial ECs showed constitutive Piezo1 channel activity (26 pS). (c) FpECs transfected with scrambled siRNA (Scr.si) after exposure to 48 h of SS caused by an orbital shaker showed alignment (arrow depicts direction of flow, scale 100 μm). (d) Lack of alignment in FpECs exposed to SS after Piezo1 depletion with siRNA (P1.si) (scale 100 μm). (e) Quantification of orientation analysis showing significantly reduced alignment after P1.si versus Scr.si transfection ($P < 0.05$); mean height (SEM) after Scr.si 5674.611 (760.763), P1.si 2185.42 (548.786), static 1728.812 (632.188) ($n = 3/N = 2$). transfection ($P < 0.05$). (f) Changes in intracellular Ca^{2+} occurred in response to increasing concentrations of Yoda1 compared with vehicle control (mean \pm SEM; mean responses to Yoda1 fitted with the Hill equation suggested an approximate EC_{50} of 5.36 μM). Adapted from Morley *et al.* (2018).

eNOS activity was observed in normal pregnancy (Nelson *et al.*, 2000). This has also been shown in animal models where eNOS inhibition prevented uterine artery remodelling (Ko *et al.*, 2018). The augmentation of NO in pregnancy may be secondary to SS generated by the increase in blood flow through the uterine vasculature. This is in conjunction with growth factors such as VEGF and PlGF, and endocrine signals (Osol and Moore, 2014). Successful vascular adaptation normalizes the SS, although definitive values for SS in human uterine vessels both in normal pregnancy and in gestational disorders remains unreported.

The response of uterine artery ECs to SS can be demonstrated by their alignment in the direction of flow (10–20 dyn/cm^2 , based on an assumption of physiological arterial SS in the systemic circulation of 6–40 dyn/cm^2) (de la Paz *et al.*, 2012; Park *et al.*, 2017). Alongside the morphological change in response to SS, Park *et al.* (2017) showed increased expression of VEGF receptor-3 in human uterine artery ECs. This receptor is thought to be involved in angiogenesis and its expression in response to fluidic shear occurred independently of its ligand. Although alterations in flow through the uterine vessels may contribute to vascular remodelling, the mechanisms transducing SS




Figure 6 Schematic illustrating the possible mechanism by which SS activation of Piezo1 results in vasodilatation through the production of NO by phosphorylation of eNOS at serine site I177. NO: nitric oxide, p: phosphorylation, eNOS: endothelial NO synthase. Created using SMART Servier Medical Art (LES LABORATOIRES SERVIER, SAS, France).

into the expression of growth factors remains unknown. Piezo1 has been detected in the uterine artery endothelium of the rat (John et al., 2018). It is therefore possible that Piezo1 activation may lead to NO production, alongside expressions of VEGFs, but further work is required to confirm this.

A low-sodium diet was used by Bigonnesse et al. (2018) to generate a rat model of FGR characterized by reduced uteroplacental perfusion. Impaired blood flow velocity and increased resistance occurred secondary to reduced uterine vessel diameters. In these hypoperfused placentae, increased NO activity was observed, leading the authors to conclude that enhanced production was compensating for vasoconstriction (Bigonnesse et al., 2018). This feeds into our hypothesis that alterations in SS are detected by mechanosensors leading to the release of NO as a compensatory mechanism (Fig. 6).

Analysis of placental bed biopsies showed that increasing intraluminal flow led to dilatation of the small myometrial resistance arteries. By contrast, in women with pre-eclampsia, the SS response was blunted, with absent flow-mediated dilatation (Kublickiene et al., 2000). In contrast, inhibition of eNOS increased myogenic- and norepinephrine-induced tone in arteries from both controls and pre-eclampsia (Kublickiene et al., 2000). As such, NO may still have a role in regulation of vascular tone in pre-eclampsia, but the SS-mediated NO release appears to be absent. Failure of SS-induced vasodilatation may therefore contribute to impaired uteroplacental blood flow (Kublickiene et al., 2000; Osol et al., 2017).

Therapeutic applications

In 2015 a synthetic agonist of Piezo1 was discovered, called Yoda1 (Syeda et al., 2015). This compound activates Piezo1 prolonging the opening of the channel, leading to downstream effects via influx of Ca²⁺ and other cations into the cell.

This exciting step allows us to manipulate Piezo1 function chemically. In our FpECs, Yoda1 caused strong elevation of the intracellular Ca²⁺ concentration with a 50% effect occurring at about 5.4 μM (Morley et al., 2018) (Fig. 5f). Depletion of Piezo1 using siRNA suppressed the Yoda1 response, consistent with it being mediated by Piezo1 channels (Morley et al., 2018). Similar findings have also been demonstrated in iatrogenic pancreatitis secondary to high pressure generated surgically. This showed that Piezo1 channels present in the murine pancreas could be stimulated by Yoda1 (Romac et al., 2018). These findings suggest that the response to mechanical force can indeed be manipulated using drugs.

Although Yoda1 lacks the pharmacological properties of a drug for human use, this is a first step towards developing new compounds with better pharmacokinetic properties. Most recently, a new set of Piezo1 chemical activators (Jedi) has been developed. Interestingly, activation by Jedi required the force-sensing architectural components of the Piezo1 channel to be present to be effective (Wang et al., 2018). Likewise, new inhibitors are also being developed by medicinal chemists, such as Dooku1, which counteracts the effects of Yoda1 (Evans et al., 2018).

Future research

The precise nature of SS and the role of mechanosensors at different stages of placental development still need to be determined. Accurately representing *in vivo* SS conditions in the *in vitro* setting remains a challenge. Sprague et al. (2010) have described the SS variations between vascular beds, type and size of vessel, and even at each bifurcation. It is unlikely that results obtained from pure laminar SS exposure are accurately representative in this context (Sprague et al., 2010). This was recently highlighted where both laminar and disturbed flow were found to activate the same initial pathways involving Piezo1,

P2Y₂, and G protein receptors. However, only disturbed flow resulted in downstream activation of integrins and focal adhesion proteins (Albarran-Juarez *et al.*, 2018).

To overcome this, Sprague *et al.* (2010) suggest using mathematical models and computer simulation to better reflect the complexity of SS and improve the congruency of current experimental models with *in vivo* vasculature. Future studies may also use microfluidic chambers to set targeted shear rates across cell co-cultures, such as placental trophoblast and endothelium (Gnecco *et al.*, 2017).

The impact of *PIEZO1* mutations on the lymphovascular system supports the importance of the channel for human development (Martin-Almedina *et al.*, 2018). However, as far as we are aware, no studies have investigated Piezo1 in the first trimester. Our work has shed new light on mechanosensing in FpECs from healthy term pregnancies (Morley *et al.*, 2018). Urgent work is therefore required to establish differences in SS sensing and mechanosensor expression between normal endothelium and that from pre-eclampsia and FGR. Likewise, the role of specific mechanosensors in the trophoblast remains unexplored. Our working hypothesis links Piezo1 activation by fluid flow to downstream production of NO (Fig. 6). However, it is possible that other extracellular or intracellular proteins may interact with and/or regulate Piezo1 channels and this remains to be determined. The availability of new compounds such as Yoda1, Jedi, and Dooku1, which specifically target the Piezo1 channel, offers opportunities for further fundamental research into this important mechanosensor. The potential for, and consequences of, these mechanosensor modulators crossing the placenta and their effect on the baby remains to be determined. For example, we do not yet know if such agents would have vascular bed specificity within uteroplacental and/or fetoplacental circulations, nor the effect this would have on maternal and fetal wellbeing given the importance of maintaining a balanced pressure gradient between the two circulations.

Conclusion

Previous studies, alongside our own work, have shown that uterine arteries, FpECs, and trophoblast cells are mechanically sensitive. The body of literature describing the role of SS sensing and subsequent signalling in multiple vascular functions within the placenta is growing. It is becoming apparent that SS has implications for a developing pregnancy, from establishing the conceptus through to adequacy of placental blood supply at term. The role of SS appears to be nuanced however, depending upon gestation and potentially, cell type. This paradox stems from the threshold required for survival while preventing damage to the embryo and villi. As pregnancy progresses however, profound vasodilation and adaptation to flow are required to enable adequate placental perfusion. As such, a delicate balance in fluid forces is required for the complex processes of embryo establishment, uteroplacental remodelling, and fetoplacental vascular development. This therefore requires tight control by regulators capable of sensing and transducing force at each stage of development.

Until relatively recently, the molecular mechanisms of mechanosensing have been unknown. Accelerated research has led to the discovery of Piezo1, which is functionally active in FpECs. We are far from piecing together how Piezo1 activation ultimately results in the complex and orchestrated processes of vascular remodelling. However, it is striking

that even *in vitro* Piezo1 activation is necessary for alignment of ECs to SS.

SS mechanotransduction and subsequent signalling, as well as the interplay between these mediators, and clinical and genetic risk factors for placental dysfunction need to be studied for a better understanding of placental haemodynamic regulation. The ability to manipulate mechanosensory complexes and influence maternal-fetal blood flow with new compounds could pave the way for new treatments to prevent the consequences of placental dysfunction.

Acknowledgements

With many thanks to Dr H Gaunt, Dr K McPhillie and Dr J Shi from our lab for assistance with the production of figures in this manuscript. Our thanks, as always, go to the staff and patients at Leeds Teaching Hospitals for the provision of tissue samples for our research.

Authors' roles

L.C.M. wrote the initial version of the manuscript. D.J.B., J.J.W. and N.A.B.S. provided expertise, supervision and editorial feedback.

Funding

Clinical Research Training Fellowship from the Medical Research Council and the Royal College of Obstetricians and Gynaecologists (MR/P002099/1 to L.C.M.) and Wellcome Trust Institutional Strategic Support Fund (to L.C.M.).

Conflict of interest

None to declare.

References

- Albarran-Juarez J, Iring A, Wang S, Joseph S, Grimm M, Strlic B, Wettschureck N, Althoff TF, Offermanns S. Piezo1 and Gq/G11 promote endothelial inflammation depending on flow pattern and integrin activation. *J Exp Med* 2018;**215**:2655–2672.
- Alcaino C, Farrugia G, Beyder A. Mechanosensitive Piezo channels in the gastrointestinal tract. *Curr Top Membr* 2017;**79**:219–244.
- Ando J, Yamamoto K. Flow detection and calcium signalling in vascular endothelial cells. *Cardiovasc Res* 2013;**99**:260–268.
- Baratchi S, Khoshmanesh K, Woodman OL, Potocnik S, Peter K, McIntyre P. Molecular sensors of blood flow in endothelial cells. *Trends Mol Med* 2017;**23**:850–868.
- Bigonnesse E, Sicotte B, Brochu M. Activated NO pathway in uterine arteries during pregnancy in an IUGR rat model. *Am J Physiol Heart Circ Physiol* 2018;**315**:H415–H422.
- Brakemeier S, Eichler I, Hopp H, Kohler R, Hoyer J. Up-regulation of endothelial stretch-activated cation channels by fluid shear stress. *Cardiovasc Res* 2002;**53**:209–218.
- Chatterjee S. Endothelial mechanotransduction, redox signaling and the regulation of vascular inflammatory pathways. *Front Physiol* 2018;**9**:524.

- Cox CD, Bae C, Ziegler L, Hartley S, Nikolova-Krstevski V, Rohde PR, Ng CA, Sachs F, Gottlieb PA, Martinac B. Removal of the mechanoprotective influence of the cytoskeleton reveals PIEZO1 is gated by bilayer tension. *Nat Commun* 2016;**7**:10366.
- Del, Marmol JI, Touhara KK, Croft G, MacKinnon R. Piezo1 forms a slowly-inactivating mechanosensory channel in mouse embryonic stem cells. *Elife* 2018;**7**:1–18.
- dela Paz NG, Walshe TE, Leach LL, Saint-Geniez M, D'Amore PA. Role of shear-stress-induced VEGF expression in endothelial cell survival. *J Cell Sci* 2012;**125**:831–843.
- Evans EL, Cuthbertson K, Endesh N, Rode B, Blythe NM, Hyman AJ, Hall SJ, Gaunt HJ, Ludlow MJ, Foster R et al. Yoda1 analogue (Dooku1) which antagonizes Yoda1-evoked activation of Piezo1 and aortic relaxation. *Br J Pharmacol* 2018;**175**:1744–1759.
- Fotiou E, Martin-Almedina S, Simpson MA, Lin S, Gordon K, Brice G, Atton G, Jeffery I, Rees DC, Mignot C et al. Novel mutations in PIEZO1 cause an autosomal recessive generalized lymphatic dysplasia with non-immune hydrops fetalis. *Nat Commun* 2015;**6**:8085.
- Garcia-Gonzalez MA, Outeda P, Zhou Q, Zhou F, Menezes LF, Qian F, Huso DL, Germino GG, Piontek KB, Watnick T. Pkd1 and Pkd2 are required for normal placental development. *PLoS One* 2010;**5**:1–12.
- Gnecco JS, Pensabene V, Li DJ, Ding T, Hui EE, Bruner-Tran KL, Osteen KG. Compartmentalized Culture of Perivascular Stroma and Endothelial Cells in a Microfluidic Model of the Human Endometrium. *Ann Biomed Eng* 2017;**45**:1758–1769.
- Guo YR, MacKinnon R. Structure-based membrane dome mechanism for Piezo mechanosensitivity. *Elife* 2017;**6**.
- Hartmannsgruber V, Heyken WT, Kacik M, Kaistha A, Grgic I, Hartneck C, Liedtke W, Hoyer J, Kohler R. Arterial response to shear stress critically depends on endothelial TRPV4 expression. *PLoS One* 2007;**2**:e827.
- Huang C, Sheikh F, Hollander M, Cai C, Becker D, Chu PH, Evans S, Chen J. Embryonic atrial function is essential for mouse embryogenesis, cardiac morphogenesis and angiogenesis. *Development* 2003;**130**:6111–6119.
- Hyman AJ, Tumova S, Beech DJ. Piezo1 Channels in Vascular Development and the Sensing of Shear Stress. *Curr Top Membr* 2017;**79**:37–57.
- James JL, Cartwright JE, Whitley GS, Greenhill DR, Hoppe A. The regulation of trophoblast migration across endothelial cells by low shear stress: consequences for vascular remodelling in pregnancy. *Cardiovasc Res* 2012;**93**:152–161.
- James JL, Saghian R, Perwick R, Clark AR. Trophoblast plugs: impact on utero-placental haemodynamics and spiral artery remodelling. *Hum Reprod* 2018:1430–1441.
- Jauniaux E, Watson AL, Hempstock J, Bao YP, Skepper JN, Burton GJ. Onset of maternal arterial blood flow and placental oxidative stress. A possible factor in human early pregnancy failure. *Am J Pathol* 2000;**157**:2111–2122.
- John L, Ko NL, Gokina A, Gokina N, Mandala M, Osol G. The Piezo1 cation channel mediates uterine artery shear stress mechanotransduction and vasodilation during rat pregnancy. *Am J Physiol Heart Circ Physiol* 2018:1019–1026.
- Jones S, Bischof H, Lang I, Desoye G, Greenwood SL, Johnstone ED, Wareing M, Sibley CP, Brownbill P. Dysregulated flow-mediated vasodilatation in the human placenta in fetal growth restriction. *J Physiol* 2015;**593**:3077–3092.
- Kingdom J. Adriana and Luisa Castellucci Award Lecture 1997. Placental pathology in obstetrics: adaptation or failure of the villous tree? *Placenta* 1998;**19**:347–351.
- Kingdom J, Huppertz B, Seaward G, Kaufmann P. Development of the placental villous tree and its consequences for fetal growth. *Eur J Obstet Gynecol Reprod Biol* 2000;**92**:35–43.
- Ko NL, Mandala M, John L, Gelinne A, Osol G. Venoarterial communication mediates arterial wall shear stress-induced maternal uterine vascular remodeling during pregnancy. *Am J Physiol Heart Circ Physiol* 2018;**315**:H709–H717.
- Kohler R, Schonfelder G, Hopp H, Distler A, Hoyer J. Stretch-activated cation channel in human umbilical vein endothelium in normal pregnancy and in preeclampsia. *J Hypertens* 1998;**16**:1149–1156.
- Krause BJ, Carrasco-Wong I, Caniuguir A, Carvajal J, Farias M, Casanello P. Endothelial eNOS/arginase imbalance contributes to vascular dysfunction in IUGR umbilical and placental vessels. *Placenta* 2013;**34**:20–28.
- Kublickiene KR, Lindblom B, Kruger K, Nisell H. Preeclampsia: evidence for impaired shear stress-mediated nitric oxide release in uterine circulation. *Am J Obstet Gynecol* 2000;**183**:160–166.
- Learmont JG, Poston L. Nitric oxide is involved in flow-induced dilation of isolated human small fetoplacental arteries. *Am J Obstet Gynecol* 1996;**174**:583–588.
- Lecarpentier E, Atallah A, Guibourdenche J, Hebert-Schuster M, Vieillefosse S, Chissey A, Haddad B, Pidoux G, Evain-Brion D, Barakat A et al. Fluid shear stress promotes placental growth factor upregulation in human syncytiotrophoblast through the cAMP-PKA signaling pathway. *Hypertension* 2016a;**68**:1438–1446.
- Lecarpentier E, Bhatt M, Bertin GI, Deloison B, Salomon LJ, Deloron P, Fournier T, Barakat AI, Tsatsaris V. Computational fluid dynamic simulations of maternal circulation: wall shear stress in the human placenta and its biological implications. *PLoS One* 2016b;**11**:e0147262.
- Li J, Hou B, Tumova S, Muraki K, Bruns A, Ludlow MJ, Sedo A, Hyman AJ, McKeown L, Young RS et al. Piezo1 integration of vascular architecture with physiological force. *Nature* 2014;**515**:279–282.
- Li Y, Zheng J, Bird IM, Magness RR. Mechanisms of shear stress-induced endothelial nitric-oxide synthase phosphorylation and expression in ovine fetoplacental artery endothelial cells. *Biol Reprod* 2004;**70**:785–796.
- Liang J, Huang B, Yuan G, Chen Y, Liang F, Zeng H, Zheng S, Cao L, Geng D, Zhou S. Stretch-activated channel Piezo1 is up-regulated in failure heart and cardiomyocyte stimulated by AngII. *Am J Transl Res* 2017;**9**:2945–2955.
- Lukacs V, Mathur J, Mao R, Bayrak-Toydemir P, Procter M, Cahalan SM, Kim HJ, Bandell M, Longo N, Day RW et al. Impaired PIEZO1 function in patients with a novel autosomal recessive congenital lymphatic dysplasia. *Nat Commun* 2015;**6**:8329.
- Malek AM, Alper SL, Izumo S. Hemodynamic shear stress and its role in atherosclerosis. *JAMA* 1999;**282**:2035–2042.
- Martin-Almedina S, Mansour S, Ostergaard P. Human phenotypes caused by PIEZO1 mutations; one gene, two overlapping phenotypes? *J Physiol* 2018;**596**:985–992.
- Mills TA, Wareing M, Bugg GJ, Greenwood SL, Baker PN. Chorionic plate artery function and Doppler indices in normal pregnancy and intrauterine growth restriction. *Eur J Clin Invest* 2005;**35**:758–764.

- Miura S, Sato K, Kato-Negishi M, Teshima T, Takeuchi S. Fluid shear triggers microvilli formation via mechanosensitive activation of TRPV6. *Nat Commun* 2015;**6**:8871.
- Morley LC, Shi J, Gaunt HJ, Hyman AJ, Webster PJ, Williams C, Forbes K, Walker JJ, Simpson NAB, Beech DJ. Piezo1 Channels Are Mechanosensors in Human Fetoplacental Endothelial Cells. *Mol Hum Reprod* 2018;510–250.
- Myatt L, Eis AL, Brockman DE, Greer IA, Lyall F. Endothelial nitric oxide synthase in placental villous tissue from normal, pre-eclamptic and intrauterine growth restricted pregnancies. *Hum Reprod* 1997;**12**: 167–172.
- Nelson SH, Steinsland OS, Wang Y, Yallampalli C, Dong YL, Sanchez JM. Increased nitric oxide synthase activity and expression in the human uterine artery during pregnancy. *Circ Res* 2000;**87**:406–411.
- Osol G, Ko NL, Mandala M. Altered endothelial nitric oxide signaling as a paradigm for maternal vascular maladaptation in preeclampsia. *Curr Hypertens Rep* 2017;**19**:82.
- Osol G, Moore LG. Maternal uterine vascular remodeling during pregnancy. *Microcirculation* 2014;**21**:38–47.
- Park YG, Choi J, Jung HK, Song IK, Shin Y, Park SY, Seol JW. Fluid shear stress regulates vascular remodeling via VEGFR-3 activation, although independently of its ligand, VEGF-C, in the uterus during pregnancy. *Int J Mol Med* 2017;**40**:1210–1216.
- Ranade SS, Qiu Z, Woo SH, Hur SS, Murthy SE, Cahalan SM, Xu J, Mathur J, Bandell M, Coste B *et al*. Piezo1, a mechanically activated ion channel, is required for vascular development in mice. *Proc Natl Acad Sci U S A* 2014;**111**:10347–10352.
- Roberts VHJ, Morgan TK, Bednarek P, Morita M, Burton GJ, Lo JO, Frias AE. Early first trimester uteroplacental flow and the progressive disintegration of spiral artery plugs: new insights from contrast-enhanced ultrasound and tissue histopathology. *Hum Reprod* 2017;**32**:2382–2393.
- Rode B, Shi J, Endesh N, Drinkhill MJ, Webster PJ, Lotteau SJ, Bailey MA, Yuldasheva NY, Ludlow MJ, Cubbon RM *et al*. Piezo1 channels sense whole body physical activity to reset cardiovascular homeostasis and enhance performance. *Nat Commun* 2017;**8**:350.
- Romac JM, Shahid RA, Swain SM, Vigna SR, Liddle RA. Piezo1 is a mechanically activated ion channel and mediates pressure induced pancreatitis. *Nat Commun* 2018;**9**:1715.
- Saotome K, Murthy SE, Kefauver JM, Whitwam T, Patapoutian A, Ward AB. Structure of the mechanically activated ion channel Piezo1. *Nature* 2018;**554**:481–486.
- Servin-Vences MR, Moroni M, Lewin GR, Poole K. Direct measurement of TRPV4 and PIEZO1 activity reveals multiple mechanotransduction pathways in chondrocytes. *Elife* 2017;**6**.
- Shihata WA, Michell DL, Andrews KL, Chin-Dusting JP. Caveolae: a role in endothelial inflammation and mechanotransduction? *Front Physiol* 2016;**7**:628.
- Slator PJ, Hutter J, McCabe L, Gomes ADS, Price AN, Panagiotaki E, Rutherford MA, Hajnal JV, Alexander DC. Placenta microstructure and microcirculation imaging with diffusion MRI. *Magn Reson Med* 2018;**80**:756–766.
- Sprague B, Chesler NC, Magness RR. Shear stress regulation of nitric oxide production in uterine and placental artery endothelial cells: experimental studies and hemodynamic models of shear stresses on endothelial cells. *Int J Dev Biol* 2010;**54**: 331–339.
- Syeda R, Florendo MN, Cox CD, Kefauver JM, Santos JS, Martinac B, Patapoutian A. Piezo1 channels are inherently mechanosensitive. *Cell Rep* 2016;**17**:1739–1746.
- Syeda R, Xu J, Dubin AE, Coste B, Mathur J, Huynh T, Matzen J, Lao J, Tully DC, Engels IH *et al*. Chemical activation of the mechanotransduction channel Piezo1. *Elife* 2015;**4**:1–11.
- Tzima E, Irani-Tehrani M, Kiosses WB, Dejana E, Schultz DA, Engelhardt B, Cao G, DeLisser H, Schwartz MA. A mechanosensory complex that mediates the endothelial cell response to fluid shear stress. *Nature* 2005;**437**:426–431.
- Wang S, Chennupati R, Kaur H, Iring A, Wettschureck N, Offermanns S. Endothelial cation channel PIEZO1 controls blood pressure by mediating flow-induced ATP release. *J Clin Invest* 2016;**126**:4527–4536.
- Wang Y, Chi S, Guo H, Li G, Wang L, Zhao Q, Rao Y, Zu L, He W, Xiao B. A lever-like transduction pathway for long-distance chemical- and mechano-gating of the mechanosensitive Piezo1 channel. *Nat Commun* 2018;**9**:1300.
- Wareing M. Effects of oxygenation and luminal flow on human placenta chorionic plate blood vessel function. *J Obstet Gynaecol Res* 2012;**38**:185–191.
- Wieczorek KM, Brewer AS, Myatt L. Shear stress may stimulate release and action of nitric oxide in the human fetal-placental vasculature. *Am J Obstet Gynecol* 1995;**173**:708–713.
- Xie Y, Wang F, Puscheck EE, Rappolee DA. Pipetting causes shear stress and elevation of phosphorylated stress-activated protein kinase/jun kinase in preimplantation embryos. *Mol Reprod Dev* 2007;**74**:1287–1294.
- Xie Y, Wang F, Zhong W, Puscheck E, Shen H, Rappolee DA. Shear stress induces preimplantation embryo death that is delayed by the zona pellucida and associated with stress-activated protein kinase-mediated apoptosis. *Biol Reprod* 2006;**75**:45–55.
- Xu J, Mathur J, Vessieres E, Hammack S, Nonomura K, Favre J, Grimaud L, Petrus M, Francisco A, Li J *et al*. GPR68 senses flow and is essential for vascular physiology. *Cell* 2018;**173**:762–75.e16.
- Yamamoto K, Sokabe T, Matsumoto T, Yoshimura K, Shibata M, Ohura N, Fukuda T, Sato T, Sekine K, Kato S *et al*. Impaired flow-dependent control of vascular tone and remodeling in P2X4-deficient mice. *Nat Med* 2006;**12**:133–137.
- Zarychanski R, Schulz VP, Houston BL, Maksimova Y, Houston DS, Smith B, Rinehart J, Gallagher PG. Mutations in the mechanotransduction protein PIEZO1 are associated with hereditary xerocytosis. *Blood* 2012;**120**:1908–1915.
- Zhao Q, Zhou H, Chi S, Wang Y, Wang J, Geng J, Wu K, Liu W, Zhang T, Dong MQ *et al*. Structure and mechanogating mechanism of the Piezo1 channel. *Nature* 2018;**554**: 487–492.