

This is a repository copy of Latitudinal constraints in responsiveness of plants to arbuscular mycorrhiza:the 'sun-worshipper' hypothesis.

White Rose Research Online URL for this paper: <a href="https://eprints.whiterose.ac.uk/146047/">https://eprints.whiterose.ac.uk/146047/</a>

Version: Accepted Version

#### Article:

Veresoglou, Stavros, Chen, Baodong, Fischer, Matthias et al. (5 more authors) (2019) Latitudinal constraints in responsiveness of plants to arbuscular mycorrhiza:the 'sunworshipper' hypothesis. New Phytologist. ISSN 1469-8137

https://doi.org/10.1111/nph.15918

### Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

### **Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.





# Latitudinal constraints in responsiveness of plants to arbuscular mycorrhiza: The 'sun-worshipper' hypothesis.

Journal:	New Phytologist
Manuscript ID	NPH-L-2019-29648.R1
Manuscript Type:	L - Letter
Date Submitted by the Author:	n/a
Complete List of Authors:	Veresoglou, Stavros; Freie Universitat Berlin, Plant Ecology Chen, Baodong; Research Centre for Eco-Environmental Sciences, Environmental Sciences Fischer, Matthias; Freie Universitat Berlin, Plant Ecology Helgason, Thorunn; University of York, Department of Biology Mamolos, Andreas; Aristotle University of Thessaloniki, Faculty of Agriculture Rillig, Matthias; Freie Universitat Berlin Fachbereich Biologie Chemie Pharmazie, Institut für Biologie Roldán, Antonio; CSIC, Department of Soil and Water Conservation Johnson, David; University of Manchester, School of Earth and Environmental Sciences
Key Words:	Arbuscular mycorrhiza, eco-evolutionary processes, environmental filtering, Glomeromycotina, mycorrhizal responsiveness, phenotypic adaptation, plant biogeography, the sun-worshipper hypothesis

SCHOLARONE™ Manuscripts

34

```
1
      Letter
 2
      Number of Words (summary): 0
      Number of Words (main text): 1496
 3
      Number of references: 31
 4
      Display items: 2 figures no tables, no text boxes
 5
 6
      Latitudinal constraints in responsiveness of plants to arbuscular mycorrhiza: The
 7
      'sun-worshipper' hypothesis.
 8
 9
      Stavros D. Veresoglou<sup>1,2*</sup>, Baodong Chen<sup>3,4</sup>, Matthias M. Fischer<sup>1</sup>, Thorunn Helgason<sup>5</sup>,
10
      Andreas P. Mamolos<sup>2</sup>, Matthias C. Rillig<sup>1,6</sup>, Antonio Roldán<sup>7</sup>, David Johnson<sup>8</sup>
11
12
      <sup>1</sup> Freie Universität Berlin, Institut für Biologie, Plant Ecology, D-14195 Berlin, Germany
13
      <sup>2</sup> Faculty of Agriculture, Laboratory of Ecology and Environmental Protection, Aristotle
14
      University of Thessaloniki, 541 24 Thessaloniki, Greece.
15
16
      <sup>3</sup> State Key Laboratory of Urban and Regional Ecology, Research Center for Eco-
      Environmental Sciences, Chinese Academy of Sciences, Beijing, China
17
18
      <sup>4</sup> University of Chinese Academy of Sciences, Beijing, China
      <sup>5</sup> Department of Biology, University of York, York, YO10 5DD, UK
19
      <sup>6</sup> Berlin-Brandenburg Institute of Advanced Biodiversity Research (BBIB), D-14195 Berlin,
20
      Germany
21
22
      <sup>7</sup> CSIC-Centro de Edafología y Biología Aplicada del Segura, Department of Soil and Water
      Conservation, P.O. Box 164, Campus de Espinardo, 30100 Murcia, Spain
23
24
      <sup>8</sup> School of Earth and Environmental Sciences, The University of Manchester, Manchester,
      M13 9PT. UK.
25
      * Email address: sveresoglou@zedat.fu-berlin.de, telephone: +49 30 838-53172
26
27
      Keywords
28
      Arbuscular mycorrhiza; eco-evolutionary processes; environmental filtering;
29
      Glomeromycotina; mycorrhizal responsiveness; phenotypic adaptation; plant biogeography;
30
31
      the sun-worshipper hypothesis
32
33
```

Most terrestrial plants depend strongly on associations with arbuscular mycorrhizal (AM) 35 fungi (Subphylum: Glomeromycotina) to establish and survive (Bever, 2002; van der Heijden 36 et al., 1998; Klironomos et al., 2011; Veresoglou et al., 2017), and have evolved a nutritional 37 mutualism. In this mutualism, the plant provides carbon to the fungus, usually subject to the 38 availability of light (Hayman 1970; Heinemeyer et al., 2003; Shi et al. 2014; Konvalinkova & 39 Jansa, 2016), and the fungus provides the plant with mineral nutrients acquired from soil. 40 Because of light constraints, it is expected that latitude exerts a strong influence on reciprocal 41 42 exchange of resources between mycorrhizal plants and fungi, and this could have consequences on the responsiveness of plants to mycorrhizal fungi. Latitude induces changes 43 in the amount of solar energy and the timing when this is made available to primary producers 44 during the year and in the day. At the same time, there is a strong negative relationship 45 between latitude and temperature that may also impact the functioning of the mycorrhizal 46 symbiosis, and in some cases (e.g. in north temperate systems), a general relationship between 47 latitude and several edaphic factors (Read and Perez-Moreno 2003). 48 49 There is compelling evidence that the alpha-diversity of Glomeromycotan fungi, which form AM symbioses, decreases with latitude (Davison et al., 2015). This finding can be 50 partially explained by the transition from ecosystems dominated by AM host plants in the 51 tropics, to ectomycorrhizal and ericoid mycorrhizal dominated ecosystems at higher latitudes 52 (Smith &Read, 2008). We know less about the extent to which latitude impacts the 53 functioning of AM symbioses, which could occur either through latitudinal differences in 54 solar radiation or resulting changes in temperature (Clarke & Gaston, 2006; Schluter, 2016). 55 Here, we propose the 'sun-worshipper' hypothesis that discriminates three different types of 56 underlying responses of latitudinal gradient-induced changes in plant host mycorrhizal 57 responsiveness (Fig. 1). Changes in abiotic conditions may allow plants to derive more 58 benefits from the symbiosis at lower latitudes through phenotypic plasticity (mechanism one – 59 Fig.1). A likely example of phenotypic plasticity might involve changes in the expression of 60 genes that allow crosstalk with AM fungi when light availability is low as has been shown for 61 62 drought (Li et al., 2016). Abiotic conditions more favourable for the symbiosis close to the tropics could further exclude, via competition, species less dependent on AM fungi, resulting 63 in distinct plant communities from a perspective of AM fungal-associating behaviour; such a 64 process can be described as environmental filtering (mechanism two – Fig. 1). Finally, we 65 know that AM plants at high latitudes encounter a less diverse pool of potential symbiotic 66 partners (e.g. because of the observed latitudinal gradient in Glomeromycotan diversity; 67 68 Davison et al., 2015) and at the same time communities at high latitudes are in general

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

dominated by plants that associate with ectomycorrhizal and ericoid mycorrhizal fungi. Plants distant from the tropics could thus form less profitable AM symbioses (but also support fewer AM partners) because of a more limited pool of suitable AM fungal partners, and this mechanism is analogous to the indirect eco-evolutionary causes (Pärtel, 2002) (mechanism three - Fig. 1). Here, we use the term eco-evolutionary processes to describe combined effects of latitude on phenotypic plasticity, environmental filtering and eco-evolutionary adaptation of the host plant trait mycorrhizal dependency (e.g. Thuiller *et al.*, 2013). Even though these different mechanisms are not mutually exclusive, it is important to disentangle how each of them influences how host plants respond to mycorrhizal symbioses along gradients of latitude and solar radiation.

The benefits that plants receive from the symbiosis in relation to the carbon costs vary considerably depending on abiotic growth conditions (Johnson et al., 1997; Hoeksema et al. 2010; Grman and Robinson, 2013), compatibility of the plant host with the local AM fungal community (Klironomos, 2003) and the degree to which a plant can take advantage of nonnutritional functions of mycorrhiza such as protection from pathogens (Veresoglou & Rillig, 2013). Resource stoichiometry of phosphorus (P), nitrogen (N) and light, in particular, represents a proven tool explaining variance in growth responses of mycorrhizal hosts at various spatial scales (Johnson, 2010). Latitude-related predictions could complement such existing tools in understanding why mycorrhizal growth responses differ at large scales. It may additionally illuminate systematic differences in mycorrhizal responsiveness such as those explained by the life history of the hosts (Boerner, 1992; Roumet et al., 2005). This would be the case if the latitudinal effects are mediated through differences in solar radiation. Annual terrestrial plants may never experience light-duration stress during winter, whereas the opposite is the case for perennials that represent the majority of terrestrial plants (for example over 70% of species in the LEDA database are perennials – Kleyer et al. 2008). There is good evidence that perennial AM fungi can survive over winter in the roots of their hosts (Buwalda et al., 1985; Dodd & Jeffries, 1986 but see Hetrick et al., 1984; Mohammed et al., 1988), which could affect the carbon economy of their plant hosts. During winter, plant requirements for nutrients are limited and photoassimilates are in short supply; therefore, plants that can confine the activity of their mycorrhizal partners may benefit through improved survival rates. As a result, we expect that there is evolutionary pressure for perennials to further confine mycorrhizal responsiveness when growing outside the tropics, compared to annuals.

102

We undertook three synthesis activities to establish whether the expectations outlined above are plausible for mycorrhizal systems (see Methods in supplementary material). We first compiled a database on crop plant responses to mycorrhiza to identify phenotypic responses to latitude (Fig. S1). To assess environmental filtering due to AM fungi with regards to latitude, we synthesized data from a common garden experiment on comparative mycorrhizal responsiveness of North-American annual and perennial herbaceous plants (Wilson & Hartnett, 1998) with plant distribution data for the specific plants from USDA (USDA, 2016). We also tested for differences in mycorrhizal responsiveness across genotypes of *Zea mays* (maize) that are routinely used either in temperate or tropical systems, despite that genetic variability could effectively be attributed to breeding. These syntheses activities were not sufficiently robust to address the mechanistic constituents of the sun worshipper hypothesis but were carried out to support the over-arching concept and encourage larger syntheses or experiments exploring the hypothesis in the future.

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

Variance in the database on crop plant responses could be best explained, in our models consisting of a single predictor, by photosynthetic radiation (Fig 2.a). The optimal model had an intercept of -0.5 (F=9.7, P<0.001) and a slope per MJ.m<sup>-2</sup>.day<sup>-1</sup> radiation of 0.23 (F=6.1, P=0.019). Fitted intercepts for the different plant species shared a standard deviation of 0.048 (Fig. 2a). Analysis of maize lines demonstrated that eco-evolutionary processes also drive latitude-dependencies on mycorrhizal responsiveness. The Mann-Whitney test between temperate- (i.e. middle two quartiles) and tropical-climate adapted lines of maize revealed higher responsiveness for temperate lines (U = 24.5, P=0.034). When we repeated this analysis for tropical lines through maintaining the two middle quartiles, the differences became even more apparent (U = 4, P < 0.001, Fig. 2b). Mycorrhizal responsiveness of species in Wilson and Hartnett (1998) could be predicted by latitude of their distribution, which suggests that mycorrhizal responsiveness might induce an environmental filtering. We raised latitude to the fourth power to address fitting issues and obtained an intercept of 0.65; latitude slope of -2.44  $10^{-8}$  (F<sub>lat</sub>=23.32; P < 0.001;  $R^2_{adj} = 0.19$ ; Fig. 2c). We subsequently fitted an additional parameter that differentiated between annual and perennial plants. Inclusion of the categorical variable perennial was significant (F=23.36, P<0.001) and there was a significant interaction between this parameter and latitude (F=9.47, P=0.003) suggesting that slopes also differed. When we analyzed annual and perennial plants separately, we found a significant relationship only for perennials (Kendall Tau was -0.32 - P=0.004, whereas for annuals the Kendall Tau was 0.03 - P = 0.87).

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

The sun-worshipper hypothesis predicts that latitude impacts mycorrhizal responsiveness in plants via three complementary mechanisms, namely phenotypic responses, eco-evolutionary processes and environmental filtering (Fig. 1). Even though the analyses we report have limited resolution, they were supportive of the sun worshipper hypothesis. We found evidence that phenotypic plasticity (Fig. 2a), eco-evolutionary processes (Fig. 2b) and environmental filtering (Fig. 2c) might be operational for all three different types of latitude related differences in mycorrhizal responsiveness (Thuiller et al., 2013). Despite the findings from our analyses, it is important to highlight additional factors that may also influence mycorrhizal responsiveness, and which have the potential to confound our findings. For example, our observations may correlate with systematic differences in soil fertility; high weathering rates generally lead to poorer fertility, as occurs in many parts of the tropics (Read and Perez-Moreno, 2003). Therefore, disentangling the specific role of light versus other edaphic and environmental factors in driving mycorrhizal responsiveness likely requires additional experimentation. Testing competing hypotheses could be done with carefully designed common garden experiments or synthesizing evidence from altitudinal experiments. Nevertheless, our analysis and associated hypothesis prompts further mechanistic analyses to test how resource stoichiometry and other critical functions undertaken by mycorrhizal fungi are influenced by latitudinal gradients.

154155

156

## Acknowledgements

- SDV is partly supported by the DFG project Metacorrhiza (VE 736/2-1). DJ is partly
- supported by the N8 AgriFood programme and NERC. TH is partly supported by the BBSRC
- grant BB/L026007/1. The letter has benefited considerably from comments and suggestions
- 160 from Ian Dickie and three anonymous reviewers.

161

162

## Contributions

- SDV: Conceived the project and carried out the analyses; SDV TH, APM, MCR, AR, DJ:
- Discussed and further developed the idea; SDV, DJ: Wrote the manuscript; All authors
- 165 commented to the manuscript and approved the final version of it.

166

167

## **Supplementary Information**

- 168 Appendix 1: Mycorrhizal dependency as proxy of investment
- 169 Appendix 2: Meta-analysis on phenotypic plasticity

170	Appendix 3: Eco-evolutionary processes on LGMR
171	Appendix 4: Environmental filtering on LGMR
172	Fig. S1: Locations of the 40 studies included in the meta-analysis
173	Table S1. Primary data on the 42 studies which we considered in our meta-analysis.
174	
175	References
176	Bever JD. 2002. Negative feedback within a mutualism: host-specific growth of mycorrhizal
177	fungi reduces plant benefit. Proceedings of the Royal Society B 269, 2595-2601.
178	Boerner REJ. 1992. Plant life span and response to inoculation with vesicular-arbuscular
179	mycorrhizal fungi. Mycorrhiza 1, 153-161.
180	Buwalda JG, Stribley DP, Tinker PB. 1985. VA-mycorrhiza of winter and spring cereals.
181	Journal of Agricultural Science 105, 649-657.
182	Clarke A, Gaston KJ. 2006. Climate, energy and diversity. Proceedings of the Royal Society
183	B <b>273</b> , 2257-2266.
184	Davison J, Moora M, Öpik M, Adholeya A, Ainsaar L, Bâ A, et al. 2015. Global
185	assessment of arbuscular mycorrhizal fungus diversity reveals very low endemism.
186	Science <b>349</b> : 970-973.
187	Dodd JC, Jeffries P. 1986. Early development of VAM in autumn-sown cereals. Soil Biology
188	and Biochemistry 18, 149-154
189	Grman E, Robinson TMP. 2013. Resource availability and imbalance affect plant-
190	mycorrhizal interactions: a field test of three hypotheses. Ecology 94, 62-71.
191	Hayman DS. 1974. Plant growth responses to vesicular-arbuscular mycorrhiza .6 effect of
192	light and temperature. New Phytologist 73, 71-&
193	Heinemeyer A, Ringway KP, Edwards EJ, Benham DG, Young PW, Fitter AH. 2003.
194	Impact of soil warming and shading on colonization and community structure of
195	arbuscular mycorrhizal fungi in roots of a native grassland community. Global
196	Change Biology 10, 52-64.
197	Hetrick BAD, Bockus WW, Bloom J. 1984. The role of vesicular-arbuscular mycorrhizal
198	fungi in the growth of Kansas winter wheat. Canadian Journal of Botany 62, 735-740.
199	Hoeksema JD, Chaudhary VB, Gehring CA, Johnson NC, Karst J, Koide RT, Pringle A,
200	Zabiski C, Bever JD, Moore JC. et al. 2010. A meta-analysis of context-dependency
201	in plant response to inoculation with mycorrhizal fungi. Ecology Letters 13, 394-407.
202	Johnson NC, Graham JH, Smith FA. 1997. Functioning of mycorrhizal associations along
203	the mutualism-parasitism continuum. New Phytologist 135, 575–585.

204	<b>Johnson NC. 2010.</b> Resource stoichiometry elucidates the structure and function of
205	arbuscular mycorrhizas across scales. New Phytologist 185, 631-647.
206	Kleyer M, Bekker R M, Knevel I C, Bakker J P, Thompson K, Sonnenschein M and et al.
207	2008. The LEDA Traitbase: A database of life-history traits of Northwest European
208	flora. Journal of Ecology 96, 1266-1274.
209	Klironomos JN, Zobel M, Tibbett M, Stock WD, Rillig MC, Parrent JL, Moora M, Koch
210	AM, Facelli JM, Facelli E, Dickie IA, Bever JD. 2011. Forces that structure plant
211	communities: quantifying the importance of the mycorrhizal symbiosis. New
212	Phytologist 189, 366-370.
213	Konvalinkova T, Jansa J. 2016. Lights off for arbuscular mycorrhiza: On its symbiotic
214	functioning under light deprivation. Frontiers in Plant Science 7, 782.
215	Li T, Sun Y, Ruan Y, Xu L, Hu Y, Hao Z, Zhang X, Li H, Wang Y, Yang L, Chen B.
216	2016. Potential role of D-myo-inositol-3-phosphate synthase and 14-3-3 genes in the
217	crosstalk between Zea mays and Rhizophagus intraradices under drought stress.
218	Mycorrhiza <b>26</b> , 879-893.
219	Mohammad MJ, Pan WL, Kennedy AC. 1998. Seasonal mycorrhizal colonization of winter
220	wheat and its effect on wheat growth under dryland field conditions. Mycorrhiza 8,
221	139-144.
222	Pärtel M. 2002. Local plant diversity patterns and evolutionary history at the regional scale.
223	Ecology <b>83</b> , 2361-2366.
224	Read DJ, Perez-Moreno J. 2003. Mycorrhizas and nutrient cycling in ecosystems – a
225	journey towards relevance? New Phytologist 157, 475-492.
226	Roumet C, Urcelay C, Díaz S. 2005. Suites of root traits differ between annual and perennial
227	species growing in the field. New Phytologist 170, 357-368.
228	Schluter D. 2016. Speciation, Ecological Opportunity, and Latitude. American Naturalist 187,
229	1-18.
230	Shi G, Liu Y, Johnson NC, Olsson PA, Mao L, Cheng G, Jiang S, An L, Du G, Feng H.
231	2014. Interactive influence of light intensity and soil fertility on root-associated
232	arbuscular mycorrhizal fungi. Plant and Soil 378, 173-188.
233	Smith SE, Read DJ. 2008. Mycorrhizal symbiosis, 33d ed. Academic Press, London, UK.
234	Tedersoo L, Nara K. 2010. General latitudinal gradient of biodiversity is reversed in
235	ectomycorrhizal fungi. New Phytologist 185, 351-354.

236	Thuiller W, Münkemüller T, Lavergne S, Mouillot S, Mousquet N, Schiffers K, Gravel
237	D. 2013. A road map for integrating eco-evolutionary processes into biodiversity
238	models. Ecology Letters 16, 94-105.
239	USDA, NRCS. 2016. The PLANTS Database (http://plants.usda.gov, 28 January 2017).
240	National Plant Data Team, Greensboro, NC 27401-4901 USA.
241	van der Heijden MGA, Klironomos JN, Ursic M, Moutoglis P, Streitwolf-Engel R,
242	Boller T, Wiemken A, Sanders IR. 1998. Mycorrhizal fungal diversity determines
243	plant biodiversity, ecosystem variability and productivity. Nature 396, 72-75.
244	Veresoglou SD, Rillig MC. 2013 Accounting for the adaptation deficit of non-mycorrhizal
245	plants in experiments. Plant and Soil 366, 33-34.
246	Veresoglou SD, Wulf M, Rillig MC. 2017. Facilitation between woody and herbaceous
247	plants that associate with arbuscular mycorrhizal fungi in temperate European forests
248	Ecology and Evolution 7, 1181-1189.
249	Wilson GWT, Hartnett DC. 1998. Interspecific variation in plant responses to mycorrhizal
250	colonization in tallgrass prairie. American Journal of Botany 85, 1732-1738.
251	colonization in tallgrass prairie. American Journal of Botany 85, 1732-1738.

**Legends to Figures** 253 254 Fig. 1. The 'sun-worshipper' hypothesis: Light availability declines with latitude (x-axis) to 255 which plant sprecies that associate with AM fungi respond via reducing their mycorrhizal 256 responsiveness (LGMR – y-axis right). This is depicted in the figure with the black 257 continuous line describing phenotypic plasticity (Mechanism 1). Eco-evolutionary responses 258 partially offset the decline in LGMR with latitude (the resulting black discontinuous line has a 259 lower slope - Mechanism 2). Reduced mycorrhizal responsiveness further induces a decline in 260 species that associate with AM fungi (red line) with species that either associate with 261 ectomycorrhizal (ECM) or ericoid (ERM) mycorrhizal fungi (green continuous line – y-axis 262 left). This represents Mechanism 3. As a result, latitude is a key factor that determining plant 263 responsiveness to mycorrhizal fungi 264 265 266 Fig. 2 (a) Scatterplot of mycorrhizal responsiveness plotted against photosynthetic radiation 267 during the growth period for the annual crops that we included in our crop dataset. The 268 relationship is positive; (b) beehive plot of mycorrhizal responsiveness of maize lines adapted 269 to growth under temperate vs tropical conditions. In the plot, we only used the two central 270 quartiles of the two groups of data. The differences are significant (U = 4, P < 0.001); (c) 271 maximum latitude of the distribution of North American herbaceous plants plotted against the 272 mycorrhizal responsiveness as assayed in a common garden experiment. Each cross describes 273 the maximum latitude of a single species and the red line is the best fit line for the relationship 274 with mycorrhizal responsiveness. We found few plants having a high mycorrhizal 275 responsiveness occurring at high latitudes. 276



