



UNIVERSITY OF LEEDS

This is a repository copy of *Synthesis and bioluminescence of electronically modified and rotationally restricted colour-shifting infraluciferin analogues*.

White Rose Research Online URL for this paper:

<http://eprints.whiterose.ac.uk/146116/>

Version: Supplemental Material

Article:

Anderson, JC, Chang, C-H, Jathoul, AP et al. (1 more author) (2019) Synthesis and bioluminescence of electronically modified and rotationally restricted colour-shifting infraluciferin analogues. *Tetrahedron*, 75 (3). pp. 347-356. ISSN 0040-4020

<https://doi.org/10.1016/j.tet.2018.11.061>

(c) 2019, Elsevier Ltd. This manuscript version is made available under the CC BY-NC-ND 4.0 license <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Synthesis and bioluminescence of electronically modified and rotationally restricted colour-shifting infraluciferins

James C. Anderson, Amit P. Jathoul and Aisha J. Syed

Department of Chemistry, University College London, 20 Gordon Street, London, WC1H 0AJ, UK.

and


School of Biosciences, University of Cardiff, Sir Martin Evans Building, Museum Avenue, Cardiff, CF10 3AX, UK

Supplementary Information


Copies of ^1H and ^{13}C NMR spectra

Graphs of emission kinetics of enzymes with different compounds acquired in the PIO


¹H – NMR: 4-((tert-butyldimethylsilyl)oxy)-2-nitroaniline (15)




¹H – NMR: 6-((tert-butyldimethylsilyl)oxy)-1H-benzo[d]imidazole-2-carbonitrile (16)




¹³C – NMR: 6-((tert-butyldimethylsilyl)oxy)-1H-benzo[d]imidazole-2-carbonitrile (**16**)




¹H – NMR: 6-hydroxy-1H-benzo[d]imidazole-2-carbonitrile (**17**)




¹H – NMR: 2-(6-hydroxy-1*H*-benzo[*d*]imidazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid (**10**)




¹H – NMR: 8-hydroxybenzo[4,5]imidazo[1,2-*a*]pyridine-3-carbonitrile (**20**)




¹³C-NMR: 8-hydroxybenzo[4,5]imidazo[1,2-*a*]pyridine-3-carbonitrile (**20**)




¹H – NMR: 2-(8-hydroxybenzo[4,5]imidazo[1,2-*a*]pyridin-3-yl)-4,5-dihydrothiazole-4-carboxylic acid (**12**)




¹³C-NMR: 2-(8-hydroxybenzo[4,5]imidazo[1,2-*a*]pyridin-3-yl)-4,5-dihydrothiazole-4-carboxylic acid (**12**)




¹H – NMR: 2-methyl-6-((triisopropylsilyl)oxy)-1*H*-benzo[*d*]imidazole (**22**)




¹³C-NMR: 2-methyl-6-((triisopropylsilyl)oxy)-1*H*-benzo[*d*]imidazole (**22**)




¹H – NMR: 2-methyl-6-((triisopropylsilyl)oxy)-1-((2- (trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazole (**23a**)




¹³C-NMR: 2-methyl-6-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazole (**23a**)




¹H – NMR: 2-methyl-5-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazole (**23b**)




¹³C-NMR: 2-methyl-5-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazole (**23b**)




¹H – NMR: 6-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazole-2-carbaldehyde (**24a**)




¹³C-NMR: 6-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazole-2-carbaldehyde (**24a**)




¹H – NMR: 5-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl) ethoxy) methyl)-1*H*-benzo[*d*]imidazole-2-carbaldehyde (**24b**)





¹³C-NMR: 5-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl) ethoxy) methyl)-1*H*-benzo[*d*]imidazole-2-carbaldehyde (**24b**)




¹H – NMR: methyl (2-(diethoxyphosphoryl)ethanethioyl)serinate (**25**)




¹H – NMR: methyl (*E*)-(3-(6-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)prop-2-enethioyl)serinate (**26a**)




¹H – NMR: methyl(*E*)-(3-(5-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)prop-2-enethioyl)serinate (**26b**)




¹³C-NMR: methyl(*E*)-(3-(5-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)prop-2-enethioyl)serinate (**26b**)




¹H – NMR: methyl (*E*)-(3-(6-hydroxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)prop-2-enethioyl)serinate (**27a**)




¹³C-NMR: methyl (*E*)-(3-(6-hydroxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)prop-2-enethioyl)serinate (**27a**)




¹H – NMR: methyl (*E*)-(3-(5-hydroxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)prop-2-enethioyl)serinate (**27b**)




¹³C-NMR: methyl (*E*)-(3-(5-hydroxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)prop-2-enethioyl)serinate (**27b**)




¹H – NMR: methyl (*E*)-2-(2-(6-hydroxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)vinyl)-4,5-dihydrothiazole-4-carboxylate (**28a**)




¹³C-NMR: methyl (*E*)-2-(2-(6-hydroxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)vinyl)-4,5-dihydrothiazole-4-carboxylate (**28a**)




¹H – NMR: methyl (*E*)-2-(2-(5-hydroxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)vinyl)-4,5-dihydrothiazole-4-carboxylate (**28b**)




¹³C-NMR: methyl (*E*)-2-(2-(5-hydroxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)vinyl)-4,5-dihydrothiazole-4-carboxylate (**28b**)




¹H – NMR: methyl (*E*)-2-(2-(6-hydroxy-1*H*-benzo[*d*]imidazol-2-yl)vinyl)-4,5-dihydrothiazole-4-carboxylate




¹³C-NMR: methyl (*E*)-2-(2-(6-hydroxy-1*H*-benzo[*d*]imidazol-2-yl)vinyl)-4,5-dihydrothiazole-4-carboxylate




HMBC spectrum: methyl (*E*)-2-(2-(6-hydroxy-1*H*-benzo[*d*]imidazol-2-yl)vinyl)-4,5-dihydrothiazole-4-carboxylate




^1H – NMR: (*E*)-2-(2-(6-hydroxy-1*H*-benzo[*d*]imidazol-2-yl)vinyl)-4,5-dihydrothiazole-4-carboxylic acid (**13**)




¹³C-NMR: (E)-2-(2-(6-hydroxy-1*H*-benzo[*d*]imidazol-2-yl)vinyl)-4,5-dihydrothiazole-4-carboxylic acid (**13**)




Emission kinetics of enzymes with different compounds acquired in the PIO
SI Figure:
A




B



C



D



E

