

This is a repository copy of *Estimation of rate coefficients and branching ratios for reactions of organic peroxy radicals for use in automated mechanism construction*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/147281/>

Version: Published Version

Article:

Jenkin, Michael E., Valorso, Richard, Aumont, Bernard et al. (1 more author) (2019) Estimation of rate coefficients and branching ratios for reactions of organic peroxy radicals for use in automated mechanism construction. *Atmospheric Chemistry and Physics*. pp. 7691-7717. ISSN 1680-7324

<https://doi.org/10.5194/acp-19-7691-2019>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Estimation of rate coefficients and branching ratios for reactions of organic peroxy radicals for use in automated mechanism construction

Michael E. Jenkin^{1,2}, Richard Valorso³, Bernard Aumont³, and Andrew R. Rickard^{4,5}

¹ Atmospheric Chemistry Services, Okehampton, Devon, EX20 4QB, UK

² School of Chemistry, University of Bristol, Cantock's Close, Bristol, BS8 1TS, UK

³ LISA, UMR CNRS 7583, Université Paris-Est Créteil, Université de Paris, Institut Pierre Simon Laplace (IPSL), Créteil, France

⁴ Wolfson Atmospheric Chemistry Laboratories, Department of Chemistry, University of York, York, YO10 5DD, UK

⁵ National Centre for Atmospheric Science, University of York, York, YO10 5DD, UK

Correspondence: Michael E. Jenkin (atmos.chem@btinternet.com)

Received: 18 January 2019 – Discussion started: 5 February 2019

Revised: 27 April 2019 – Accepted: 12 May 2019 – Published: 7 June 2019

Abstract. Organic peroxy radicals (RO₂), formed from the degradation of hydrocarbons and other volatile organic compounds (VOCs), play a key role in tropospheric oxidation mechanisms. Several competing reactions may be available for a given RO₂ radical, the relative rates of which depend on both the structure of RO₂ and the ambient conditions. Published kinetics and branching ratio data are reviewed for the bimolecular reactions of RO₂ with NO, NO₂, NO₃, OH and HO₂; and for their self-reactions and cross-reactions with other RO₂ radicals. This information is used to define generic rate coefficients and structure–activity relationship (SAR) methods that can be applied to the bimolecular reactions of a series of important classes of hydrocarbon and oxygenated RO₂ radicals. Information for selected unimolecular isomerization reactions (i.e. H-atom shift and ring-closure reactions) is also summarized and discussed. The methods presented here are intended to guide the representation of RO₂ radical chemistry in the next generation of explicit detailed chemical mechanisms.

generate ozone (O₃), secondary organic aerosol (SOA) and other secondary pollutants (e.g. Lightfoot et al., 1992; Jenkin and Clemitshaw, 2000; Tyndall et al., 2001; Archibald et al., 2009; Orlando and Tyndall, 2012; Ehn et al., 2017), and rigorous representation of their chemistry is therefore essential for chemical mechanisms used in chemistry-transport models. As discussed in the preceding papers in this series (Jenkin et al., 2018a, b), they are formed rapidly and exclusively from the reactions of O₂ with the majority of carbon-centred organic radicals (R) (Reaction R1), these in turn being produced from the reactions that initiate VOC degradation (e.g. reaction with OH radicals), or from other routes such as decomposition of larger oxy radicals (*M* denotes a third body, most commonly N₂ or O₂ under atmospheric conditions):



Under tropospheric conditions, a given RO₂ radical may have several competing reactions available, the relative rates of which are dependent both on the prevailing ambient conditions and on the structure of RO₂. These include a series of bimolecular reactions (i.e. with NO, NO₂, NO₃, OH and HO₂; and the self-reaction and cross-reactions with the multitude of other RO₂ radicals present in the atmosphere), which are generally available for all RO₂ radicals; and specific unimolecular isomerization reactions (i.e. H-atom shift

1 Introduction

Organic peroxy radicals (RO₂) are important intermediates in the tropospheric degradation of hydrocarbons and other volatile organic compounds (VOCs). It is well established that their chemistry plays a key role in the mechanisms that

or ring-closure reactions) that are potentially available for some classes of RO₂.

The propagating channel of the reaction of RO₂ with NO (Reaction R2a) plays a key role in tropospheric O₃ formation, through oxidizing NO to NO₂, and also usually represents the major reaction for RO₂ radicals under comparatively polluted conditions:



The efficiency of this reaction is influenced by the relative importance of the other reactions available for a given RO₂ radical. The contribution of the terminating channel of the reaction of RO₂ with NO (forming an organic nitrate product, RONO₂) depends on the structure and size of RO₂; and the reaction of NO₂ with selected RO₂ radicals forms stable peroxyxynitrate products, ROONO₂. The formation, transport and degradation of these oxidized organic nitrogen reservoirs from the RO₂ + NO and RO₂ + NO₂ reactions has potential impacts in a number of ways, ranging from the inhibition of O₃ formation on local/regional scales to influencing the global budget and distribution of NO_x and O₃ (e.g. Perring et al., 2013). The reactions of RO₂ radicals with NO₃ primarily play a role during the nighttime in moderately polluted air, providing a radical propagation route that potentially supplements nighttime chain oxidation processes (e.g. Carslaw et al., 1997; Bey et al., 2001a, b; Geyer et al., 2003; Walker et al., 2015).

The reactions with OH, HO₂ and the pool of RO₂ radicals gain in importance as the availability of NO_x becomes more limited, and therefore also inhibit O₃ formation by competing with Reaction (R2a). In many cases, the reactions are significantly terminating and collectively make a major contribution to controlling atmospheric free radical concentrations under NO_x-limited conditions, although the branching ratios for the propagating and terminating reaction channels depend on the structure of RO₂. For some classes of RO₂, unimolecular isomerization reactions can compete with (or dominate over) the bimolecular reactions. These reactions therefore potentially play an important role in HO_x radical recycling under NO_x-limited conditions, and in rapid chain oxidation mechanisms generating highly oxidized multifunctional molecules, HOMs (e.g. Peeters et al., 2009, 2014; Crouse et al., 2013; Ehn et al., 2014, 2017; Jokinen et al., 2014; Rissanen et al., 2015; Bianchi et al., 2019). The relative contributions of the various reactions available for RO₂ thus influence the distribution and functional group content of the oxidized products formed, and their physicochemical properties (e.g. volatility and solubility), and therefore the SOA formation propensity of the chemistry.

In this paper, published data on the kinetics and branching ratios for the above bimolecular reactions of hydrocarbon and oxygenated RO₂ radicals are reviewed and discussed. Preliminary information is also presented for selected unimolecular isomerization reactions, which continue to be considered in ongoing work. The information on bimolecular

reactions is used to define and document a set of rules and structure–activity relationship (SAR) methods (a chemical protocol) to guide the representation of the RO₂ reactions in future detailed chemical mechanisms (Vereecken et al., 2018). In particular, the methods presented below are being used to design the next generation of explicit mechanisms based on the Generator for Explicit Chemistry and Kinetics of Organics in the Atmosphere, GECKO-A (Aumont et al., 2005), and the Master Chemical Mechanism, MCM (Saunders et al., 2003). Application of the methods is illustrated with examples in the supporting information provided in the Supplement.

2 Bimolecular reactions of RO₂ radicals

2.1 The reactions of RO₂ with NO

2.1.1 Kinetics

Rate coefficients for the reactions of NO with a variety of specific hydrocarbon and oxygenated RO₂ radicals have been reported, as summarized in Table 1. For the vast majority of the RO₂ radicals formed in detailed mechanisms, however, kinetic data are unavailable and it is therefore necessary to assign generic rate coefficients based on the reported data.

For acyl peroxy radicals (i.e. of structure RC(O)O₂), a generic rate coefficient (k_{APNO}) is applied:

$$k_{\text{APNO}} = 7.5 \times 10^{-12} \exp(290/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}. \quad (1)$$

This is based on the IUPAC Task Group¹ recommendation for the reaction of NO with CH₃C(O)O₂. As shown in Table 1, this is also close to the rate coefficients recommended for the less studied acyl peroxy radicals, C₂H₅C(O)O₂ and CH₂=CH(CH₃)C(O)O₂. The 298 K value reported for C(O)(OOH)CH₂CH₂CH₂CH(OOH)C(O)O₂ (Berndt et al., 2015) is also broadly consistent with k_{APNO} , although further studies of highly oxygenated acyl peroxy radicals would help to establish the effects of additional substituent groups.

For other classes of hydrocarbon and oxygenated peroxy radical, a generic rate coefficient ($k_{\text{RO}_2\text{NO}}$) is applied:

$$k_{\text{RO}_2\text{NO}} = 2.7 \times 10^{-12} \exp(360/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}. \quad (2)$$

The value of $k_{\text{RO}_2\text{NO}}$ at 298 K ($9.0 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$) is based on a rounded average of the 298 K rate coefficients listed for the $\geq \text{C}_2$ alkyl, cycloalkyl, hydroxyalkyl, hydroxyalkenyl, oxoalkyl, hydroxy-oxyalkyl and hydroxy-dioxa-bicyclo RO₂ radicals in Table 1, which show no significant trends related to the identity and structure of R. The temperature dependence is

¹The “IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation” is abbreviated to “IUPAC Task Group” for simplicity. The evaluation is available at <http://iupac.pole-ether.fr/> (last access: January 2019).

Table 1. Kinetic data for the reactions of hydrocarbon and oxygenated peroxy radicals with NO. Where available, the temperature dependence is given by $k = A \exp(-E/RT)$.


Peroxy radical	A ($10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)	E/R (K)	$k_{298\text{K}}$ ($10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)	Comment
Alkyl and cycloalkyl				
CH ₃ O ₂	2.30	−360	7.7	a
C ₂ H ₅ O ₂	2.55	−380	9.1	a
<i>n</i> -C ₃ H ₇ O ₂	2.90	−350	9.4	a,b
<i>i</i> -C ₃ H ₇ O ₂	2.70	−360	9.0	a,b
<i>t</i> -C ₄ H ₉ O ₂			8.3	a
2-C ₅ H ₁₁ O ₂			8.0	b
<i>c</i> -C ₅ H ₉ O ₂			10.9	b
Allylic (alk-2-enyl)				
CH ₂ =CHCH ₂ O ₂			10.5	b
β -Hydroxyalkyl				
HOCH ₂ CH ₂ O ₂			8.7	c
propene-derived			9.5	c,d
but-1-ene-derived			9.6	c,e
CH ₃ CH(OH)CH(O ₂)CH ₃			9.4	c
methylpropene-derived			9.6	c,f
Hydroxyalkenyl				
buta-1,3-diene-derived			8.8	c,g
isoprene-derived			8.8	a,c,h
Oxoalkyl				
CH ₃ C(O)CH ₂ O ₂			8.0	a,i
Hydroxy-oxyalkyl				
methacrolein-derived			9.3	j,k
methyl vinyl ketone-derived			8.4	j,l
Hydroxy-dioxa-bicyclo				
1,3,5-trimethylbenzene-derived			7.7	m
Acyl				
CH ₃ C(O)O ₂	7.5	−290	20	a
C ₂ H ₅ C(O)O ₂	6.7	−340	21	a
CH ₂ =CH(CH ₃)C(O)O ₂	8.7	−290	23	n
C(O)(OOH)CH ₂ CH ₂ CH(OOH)C(O)O ₂			34	o

Comments: ^a IUPAC Task Group recommendation (<http://iupac.pole-ether.fr/>, last access: January 2019). ^b Based on Eberhard and Howard (1996, 1997), Eberhard et al. (1996). ^c Based on Miller et al. (2004). ^d Mixture of CH₂(OH)CH(O₂)CH₃ and CH₂(O₂)CH(OH)CH₃. ^e Mixture of CH₂(OH)CH(O₂)C₂H₅ and CH₂(O₂)CH(OH)C₂H₅. ^f Mixture of CH₂(OH)C(O₂)(CH₃)₂ and CH₂(O₂)C(OH)(CH₃)₂. ^g Mixture of CH₂(OH)CH(O₂)CH=CH₂, CH₂(OH)CH=CHCH₂O₂ and CH₂(O₂)CH(OH)CH=CH₂. ^h Mixture of CH₂(OH)C(O₂)(CH₃)CH=CH₂, CH₂(OH)C(CH₃)=CHCH₂O₂, CH₂(O₂)C(CH₃)(OH)CH=CH₂, CH₂(OH)CH(O₂)C(CH₃)=CH₂, CH₂(OH)CH=C(CH₃)CH₂O₂ and CH₂(O₂)CH(OH)C(CH₃)=CH₂. ⁱ Based on Sehested et al. (1998). ^j Based on Hsin and Elrod (2007). ^k Mixture of CH₂(OH)C(O₂)(CH₃)CHO and CH₂(O₂)C(OH)(CH₃)CHO. ^l Mixture of CH₂(OH)CH(O₂)C(=O)CH₃ and CH₂(O₂)CH(OH)C(=O)CH₃. ^m Elrod (2011). Mixture of two complex radicals of molecular formula HOC₉H₁₂[OO]O₂, although with one isomer likely dominant. ⁿ De Gouw and Howard (1997). ^o Berndt et al. (2015). Inferred to be the complex oxo-di-hydroperoxy acyl peroxy radical shown, on the basis of its molecular mass and a proposed mechanism.

similarly based on the rounded average of the available values within this group, which are limited to those for $C_2H_5O_2$, $n-C_3H_7O_2$ and $i-C_3H_7O_2$. In practice, the preferred values for all the $\geq C_2$ (non-acyl) RO_2 radicals in Table 1 are also equivalent to k_{RO_2NO} within the reported uncertainties, such that the generic rate coefficient can reasonably be applied for simplicity in all cases except CH_3O_2 . Although derived from a more extensive dataset, the expression for k_{RO_2NO} in Eq. (2) is identical to that recommended previously by Atkinson (1997).

2.1.2 Product branching ratios

The following channels are considered for the reactions of RO_2 with NO :



It is well established that the branching ratio for alkyl peroxy radicals depends on temperature, pressure, and the size and degree of substitution of the peroxy radical (e.g. Carter and Atkinson, 1989; Arey et al., 2001; Yeh and Ziemann, 2014a). The branching ratio has also been reported to be influenced by the presence of oxygenated substituents, with most systematic information reported for β - and δ -hydroxy groups (e.g. O'Brien et al., 1998; Matsunga and Ziemann, 2009, 2010; Yeh and Ziemann, 2014b; Teng et al., 2015).

The fraction of the reaction forming a nitrate product ($RONO_2$) via the terminating channel, $R_{2b} = k_{2b}/(k_{2a} + k_{2b})$, is calculated following the method originally reported for secondary alkyl peroxy radicals by Carter and Atkinson (1989), and subsequently updated by Arey et al. (2001) and Teng et al. (2015). Based on this method, the reference branching ratio for secondary alkyl peroxy radicals, $R^\circ = (k_{2b}/k_{2a})^\circ$, is calculated as follows.

$$R^\circ = [A/(1 + (A/B))]F^z \quad (3)$$



with $A = 2 \times 10^{-22} \exp(n_{CON})[M]$, $B = 0.43(T/300)^{-8}$, $F = 0.41$ and $z = (1 + (\log_{10}(A/B))^2)^{-1}$. Variable n_{CON} is the number of carbon, oxygen and nitrogen atoms in the organic group (R) of the peroxy radical (i.e. excluding the peroxy radical oxygen atoms and equivalent to the carbon number in alkyl peroxy radicals), T is the temperature (in kelvin) and $[M]$ is the gas density (in molecule cm^{-3}).

The fractions of the reaction proceeding via the terminating channel, R_{2b} , and the propagating channel, $R_{2a} (= 1 - R_{2b})$, for a specific peroxy radical are then given by

$$R_{2b} = f_a f_b (R^\circ / (1 + R^\circ)). \quad (4)$$

The effect of the degree of substitution (i.e. whether the radical is primary, secondary or tertiary) is described by f_a , with a unity value applied to secondary peroxy radicals by definition. A further scaling factor, f_b , is used to describe systematic variations in the yields of $RONO_2$ resulting from the

Table 2. Values of the scaling factor, f_a , applied to the branching ratio calculation for the reaction of RO_2 with NO .

Class	Substitution	f_a	Comment
default	primary	0.65	^a
	secondary	1.0	
	tertiary	1.0	
	secondary	1.0	^b
	tertiary	0.13	
	secondary	0.43	^b
	tertiary	0.06	

Comments: ^a Applied in all cases, except for those covered by comment ^b. $f_a = 1$ for secondary peroxy radicals by definition. The equivalent value for tertiary peroxy radicals, and the lower value for primary peroxy radicals, is based on a consensus of information from Cassanelli et al. (2007), Orlando and Tyndall (2012) and Teng et al. (2015) and on previous consideration of the OH + isoprene system (Jenkin et al., 2015); ^b Inhibition of nitrate formation has been reported for complex hydroxy-dioxa-bicyclo peroxy radicals derived from aromatics, relative to comparably sized alkyl peroxy radicals by Rickard et al. (2010) and Elrod (2011), with a particular impact from the presence of alkyl substituents reported by Elrod (2011). The reduced values of f_a for tertiary peroxy radicals, and the general reduction in f_a for peroxy radicals with a neighbouring alkyl substituent (as shown), is inferred from the trend in nitrate yields reported for benzene, toluene, *p*-xylene and 1,3,5-trimethylbenzene by Elrod (2011).

presence of oxygenated substituents (e.g. the effect of hydroxyl substituents, as indicated above), or for specific peroxy radical classes, with a value of f_b being required to account for the effect of each relevant substituent. The applied values of f_a and f_b are summarized in Tables 2 and 3, and example calculations are provided in Sect. S1 in the Supplement.

It is also recognized that reaction channel (R2a) is significantly exothermic, such that prompt decomposition or isomerization of a fraction of the initially formed chemically activated oxy radicals has been reported to occur in some cases; with the remainder being collisionally deactivated to form thermalized RO (e.g. Orlando et al., 2003; Calvert et al., 2015). This is particularly important for β -hydroxy-oxy radicals (e.g. Orlando et al., 1998; Vereecken et al., 1999; Vereecken and Peeters, 1999; Caralp et al., 2003) and some other oxygenated oxy radicals (e.g. Christensen et al., 2000; Orlando et al., 2000a; Wallington et al., 2001). The contributions and treatment of these reactions is summarized in Sect. S2.

2.2 The reaction of RO_2 with NO_2

The reactions of RO_2 with NO_2 have generally been reported to proceed via a reversible association reaction in each case to form a peroxy nitrate ($ROONO_2$):






Table 3. Values of the scaling factor, f_b , applied to the branching ratio calculation for the reaction of RO_2 with NO^a .

Class	f_b	Comment
$\text{C}_n\text{H}_{2n+1}\text{OO}$ (alkyl peroxy)	1.0	^b
$\text{OO}-\text{C}-\text{C}(\text{OH})<$, $\text{OO}-\text{C}-\text{C}(\text{OR})<$, $\text{OO}-\text{C}(\text{OH})<$, $\text{OO}-\text{C}(\text{OR})<$, $\text{OO}-\text{C}-\text{C}(\text{ONO}_2)<$, $\text{OO}-\text{C}-\text{C}(\text{OOH})<$	0.65	^c
δ -hydroxy peroxy	0.8	^d
$\text{OO}-\text{C}-\text{C}(=\text{O})-$, $\text{OO}-\text{C}-\text{C}(=\text{O})-\text{O}-$	0.3	^e
$\text{OO}-\text{C}(=\text{O})-$, $\text{OO}-\text{C}-\text{O}-\text{C}(=\text{O})-$	0.0	^{e,f}
	0.33	^g
	0.0	^h

Comments: ^a A value of f_b needs to be applied to account for the effect of each relevant substituent (see Sect. S1 in the Supplement for further information). ^b $f_b = 1$ for alkyl peroxy radicals, by definition, and also used as a default in all cases other than those covered by comments ^{c–h}. ^c Based on a compromise of information from Matsunaga and Ziemann (2009, 2010), Yeh and Ziemann (2014b) and Teng et al. (2015) for β -hydroxy substituents, but also taking account of information reported for a number of other oxygenated systems (e.g. Tuazon et al., 1998a; Crounse et al., 2012; Lee et al., 2014) and previous consideration of the OH + isoprene system (Jenkin et al., 2015). $\text{OO}-\text{C}-\text{C}(\text{OOH})<$ assumed to be in this category by analogy. ^d Based on the relative impacts of β -OH and δ -OH substituents reported by Yeh and Ziemann (2014a) and previous consideration of the OH + isoprene system (Jenkin et al., 2015). ^e f_b value for $\text{OO}-\text{C}-\text{C}(=\text{O})-$ informed by reported studies of ketone oxidation (Lightfoot et al., 1992; Praske et al., 2015); f_b values for $\text{OO}-\text{C}-\text{C}(=\text{O})-\text{O}-$ and $\text{OO}-\text{C}-\text{O}-\text{C}(=\text{O})-$ informed by reported studies of ester and dibasic ester oxidation (Tuazon et al., 1998b, 1999; Cavalli et al., 2001; Picquet-Varrault et al., 2001, 2002; Pimentel et al., 2010). ^f $f_b = 0$ for $\text{OO}-\text{C}(=\text{O})-$ is based on the general lack of observation of acyl nitrate products in systems where acyl peroxy radicals are formed. ^g Value set to recreate the hydroxy-dioxa-bicyclo nitrate yield reported for benzene by Elrod (2011); In conjunction with the values of f_a in Table 2, this also allows a consistent representation of the yields in the toluene, *p*-xylene and 1,3,5-trimethylbenzene systems (Elrod, 2011; Rickard et al., 2010); ^h $f_b = 0$ for phenyl (and other aryl) peroxy radicals is based on the general lack of observation of aryl nitrate products during the oxidation of aromatic hydrocarbons.

Rate coefficients for the forward and reverse reactions for a number of RO_2 radicals are summarized in Table 4. Those for CH_3O_2 and $\text{C}_2\text{H}_5\text{O}_2$ – and for the two simplest acyl peroxy radicals, $\text{CH}_3\text{C}(\text{O})\text{O}_2$ and $\text{C}_2\text{H}_5\text{C}(\text{O})\text{O}_2$ – are based on (or informed by) the IUPAC Task Group recommendations, and describe the pressure and temperature dependences of the reactions. In all other cases, the reactions are assumed to be at the high-pressure limit under atmospheric conditions, and generic parameters are applied. The parameters $k_{f\text{PN}}$ and $k_{b\text{PN}}$ (given in Table 4) can reasonably be applied to reactions involving non-acyl peroxy radicals, being based on the high-pressure limiting rate coefficients (k_∞) for the forward and reverse reactions of $\text{C}_2\text{H}_5\text{O}_2$ and those reported for a number of higher alkyl peroxy radicals at close to atmospheric pressure (see Table 4 comments). This assumption is also broadly consistent with the limited information available for the forward or reverse reactions of other non-acyl oxygenated peroxy radicals (e.g. Orlando and Tyndall, 2012). In practice, however, these reactions are often omitted from atmospheric chemical mechanisms, owing to the instability of the ROONO_2 products under lower tropospheric conditions (lifetime ≈ 0.2 s at 298 K). As a result, only the formation

and decomposition of methyl peroxy nitrate, CH_3OONO_2 , from the most abundant non-acyl peroxy radical, CH_3O_2 , have previously been represented in the MCM (Saunders et al., 2003). This approach remains advocated here for application for lower tropospheric conditions.

The reactions are generally represented for acyl peroxy radicals, for which the product peroxyacyl nitrates, $\text{RC}(\text{O})\text{OONO}_2$, are particularly stable (lifetime ≈ 40 – 50 min at 298 K). The generic parameters, $k_{f\text{PAN}}$ and $k_{b\text{PAN}}$, are applied in the majority of cases (see Table 4). As shown in Fig. 1, larger acyl peroxy radicals have been reported to be slightly more stable than those derived from $\text{CH}_3\text{C}(\text{O})\text{O}_2$ and $\text{C}_2\text{H}_5\text{C}(\text{O})\text{O}_2$ (Roberts and Bertman, 1992; Kabir et al., 2014), and the assigned value of $k_{b\text{PAN}}$ is consistent with the data for the larger species.

Reported data for $\text{CH}_3\text{OC}(\text{O})\text{O}_2$, $\text{C}_6\text{H}_5\text{OC}(\text{O})\text{O}_2$ and $\text{C}_2\text{H}_5\text{OC}(\text{O})\text{O}_2$ (Kirchner et al., 1999; Bossolasco et al., 2011) indicate a reduced thermal stability of peroxyacyl nitrates derived from formate esters, and an increased decomposition rate ($2 \times k_{b\text{PAN}}$) is therefore applied to $\text{ROC}(\text{O})\text{OONO}_2$ species in general.

Table 4. Rate coefficients for the reactions of hydrocarbon and oxygenated RO₂ radicals with NO₂ and for the reverse decomposition of the RO₂NO₂ products. Generic rate coefficients for specified RO₂ classes are shown in bold font.

Peroxy radical	k_0	k_∞	F_c	$k_{298\text{ K}, 760\text{ Torr}}$	Comment
Forward reaction, k_f (cm ³ molecule ⁻¹ s ⁻¹)					
CH ₃ O ₂	$1.2 \times 10^{-30} (T/300)^{-6.9} [M]$	1.8×10^{-11}	0.36	4.2×10^{-12}	a,b
C ₂ H ₅ O ₂	$1.3 \times 10^{-29} (T/300)^{-6.2} [M]$	8.8×10^{-12}	0.31	5.1×10^{-12}	a,c
<i>n</i> - and <i>sec</i> -C ₄ H ₉ O ₂	–	9.6×10^{-12}	–	9.6×10^{-12}	d
RO₂	–	9.0×10^{-12} (= $k_{f\text{ PAN}}$)	–	9.0×10^{-12}	e
CH ₃ C(O)O ₂	$3.28 \times 10^{-28} (T/300)^{-6.87} [M]$	$1.125 \times 10^{-11} (T/300)^{-1.105}$	0.3	8.9×10^{-12}	a,c
C ₂ H ₅ C(O)O ₂	$1.05 \times 10^{-27} (T/300)^{-6.87} [M]$	$1.125 \times 10^{-11} (T/300)^{-1.105}$	0.36	8.9×10^{-12}	a,f
RC(O)O₂, ROC(O)O₂	–	$1.125 \times 10^{-11} (T/300)^{-1.105}$ (= $k_{f\text{ PAN}}$)	–	1.1×10^{-11}	e,g
Reverse reaction, k_b (s ⁻¹)					
CH ₃ O ₂	$9.0 \times 10^{-5} \exp(-9690/T) [M]$	$1.1 \times 10^{16} \exp(-10560/T)$	0.36	1.5	a,c
C ₂ H ₅ O ₂	$4.8 \times 10^{-4} \exp(-9285/T) [M]$	$8.8 \times 10^{15} \exp(-10440/T)$	0.31	3.4	a,c
<i>n</i> - and <i>sec</i> -C ₄ H ₉ O ₂	–	$8.3 \times 10^{15} \exp(-10368/T)$	–	6.4	h
C ₆ H ₁₃ O ₂ isomers	–	$7.5 \times 10^{15} \exp(-10368/T)$	–	5.8	h
C ₈ H ₁₇ O ₂ isomers	–	$4.8 \times 10^{15} \exp(-10368/T)$	–	3.7	h
RO₂	–	$7.6 \times 10^{15} \exp(-10400/T)$ (= $k_{b\text{ PAN}}$)	–	5.3	e,i
CH ₃ C(O)O ₂	$1.1 \times 10^{-5} \exp(-10100/T) [M]$	$1.9 \times 10^{17} \exp(-14100/T)$	0.3	4.3×10^{-4}	a,c
C ₂ H ₅ C(O)O ₂	$1.7 \times 10^{-3} \exp(-11280/T) [M]$	$8.3 \times 10^{16} \exp(-13940/T)$	0.36	3.6×10^{-4}	a,c
CH ₂ =C(CH ₃)C(O)O ₂	–	$1.6 \times 10^{16} \exp(-13500/T)$	–	3.5×10^{-4}	c,j
RC(O)O₂	–	$5.2 \times 10^{16} \exp(-13850/T)$ (= $k_{b\text{ PAN}}$)	–	3.4×10^{-4}	e,k
ROC(O)O₂	–	$2 \times k_{b\text{ PAN}}$	–	6.8×10^{-4}	e,l

Comments: ^a Rate coefficient for a pressure-dependent reaction is calculated using the expression: $k = F k_0 k_\infty / (k_0 + k_\infty)$, where $\log_{10} F = \log_{10} (F_c) / (1 + [\log_{10} (k_0/k_\infty) / N]^2)$ and $N = [0.75 - 1.27 \log_{10} (F_c)]$ (see <http://iupac.pole-ether.fr/>, last access: January 2019); ^b Based on the evaluation of Golden (2005); ^c Recommended by the IUPAC Task Group (<http://iupac.pole-ether.fr/>, last access: January 2019); ^d Reported by McKee et al. (2016) for a mixture of *n*-C₄H₉O₂ and *sec*-C₄H₉O₂ formed from reaction of Cl with butane; ^e Pressure-independent generic rate coefficient. ^f k_∞ assumed equivalent to that for CH₃C(O)O₂ + NO₂ reaction. k_0 scaled relative to that for CH₃C(O)O₂ to preserve the C₂H₅C(O)O₂ + NO₂ ⇌ C₂H₅C(O)OONO₂ equilibrium constant, k_f/k_b , over the pressure range 100–760 Torr. F_c is equivalent to that recommended for k_b . ^g Forward reaction rate coefficient, $k_{f\text{ PAN}}$, is based on k_∞ for the CH₃C(O)O₂ + NO₂ reaction. ^h Based on Zabel et al. (1989), as recommended by Lightfoot et al. (1992), for isomeric mixtures formed from reactions of Cl with butane, hexane or octane. Assumed to be at high-pressure limit at 800 Torr. ⁱ $k_{b\text{ PAN}}$ is rounded average of the reported rate coefficients for C₂–C₈ alkyl peroxy radicals. ^j Based on Roberts and Bertman (1992). Assumed to be at high-pressure limit at 760 Torr. ^k $k_{b\text{ PAN}}$, is based on a value of $3.4 \times 10^{-4} \text{ s}^{-1}$ at 298 K, which is the average of those reported for *n*-C₃H₇C(O)OONO₂, *n*-C₄H₉C(O)OONO₂ and *n*-C₅H₁₁C(O)OONO₂ (Kabir et al., 2014) and CH₂=C(CH₃)C(O)O₂ (Roberts and Bertman, 1992) (see Fig. 1). E/R is based on the average of the high-pressure limit values for CH₃C(O)O₂NO₂, C₂H₅C(O)O₂NO₂ and CH₂=C(CH₃)C(O)O₂, and also consistent with the approximate value for *n*-C₅H₁₁C(O)OONO₂ (Kabir et al., 2014). ^l Pressure-independent generic rate coefficient for thermal decomposition of ROC(O)O₂ is a factor of 2 greater, based on data for CH₃OC(O)O₂ and C₆H₅OC(O)O₂ (Kirchner et al., 1999), with reduced thermal stability also consistent with data for C₂H₅OC(O)O₂ (Bossolasco et al., 2011).


In a limited number of cases, the reaction of RO₂ with NO₂ has been reported to oxidize NO₂ to NO₃ in an irreversible reaction:



These cases include HC(O)C(O)O₂ (Orlando and Tyndall, 2001), CH₃C(O)C(O)O₂ (Jagiella and Zabel, 2008) and the phenylperoxy radical, C₆H₅O₂ (Jagiella and Zabel, 2007). Reaction (R3b) is therefore applied generally to HC(O)C(O)O₂, RC(O)C(O)O₂, C₆H₅O₂ and other aryl peroxy radicals, using the generic rate coefficient $k_{f\text{ PAN}}$.

2.3 The reaction of RO₂ with NO₃

On the basis of reported information for CH₃O₂ and C₂H₅O₂ (e.g. Biggs et al., 1995; Kukui et al., 1995), the reactions of RO₂ with NO₃ are assumed to proceed via a single channel in each case as follows:



Reported rate coefficients are summarized in Table 5. The reaction of C₂H₅O₂ with NO₃ is the most studied, with consistent 298 K rate coefficients reported in a number of studies (Biggs et al., 1995; Ray et al., 1996; Vaughan et al., 2006; Laversin et al., 2016) and with the temperature dependence systematically investigated (Laversin et al., 2016). The corresponding parameters in Table 5 therefore form the basis of a generic rate coefficient for the reactions of non-acyl peroxy radicals with NO₃:

$$k_{\text{RO}_2\text{NO}_3} = 8.9 \times 10^{-12} \exp(-390/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}. \quad (5)$$

Within the reported uncertainties, the value of the rate coefficient at 298 K is consistent with that for *c*-C₆H₁₁O₂ and with the approximate value for *c*-C₅H₉O₂ reported by Vaughan et al. (2006); and the temperature dependence expression for $k_{\text{RO}_2\text{NO}_3}$ is consistent with those reported for the oxygenated primary peroxy radicals, (CH₃)₂C(OH)CH₂O₂, CH₃OCH₂O₂ and CH₃C(O)CH₂O₂, by Kalalian et al. (2018). $k_{\text{RO}_2\text{NO}_3}$ is therefore currently con-

Table 5. Kinetic data for the reactions of alkyl and oxygenated peroxy radicals with NO₃. Where available, the temperature dependence is given by $k = A \exp(-E/RT)$.

Peroxy radical	A (10 ⁻¹² cm ³ molecule ⁻¹ s ⁻¹)	E/R (K)	$k_{298\text{K}}$ (10 ⁻¹² cm ³ molecule ⁻¹ s ⁻¹)	Comment
Alkyl and cycloalkyl				
CH ₃ O ₂			1.2	a
C ₂ H ₅ O ₂	8.9	390	2.4	b
<i>c</i> -C ₅ H ₉ O ₂			~ 1.2	c
<i>c</i> -C ₆ H ₁₁ O ₂			1.9	c
β -Hydroxyalkyl				
(CH ₃) ₂ C(OH)CH ₂ O ₂	16	480	3.2	d
Alkoxyalkyl				
CH ₃ OCH ₂ O ₂	13.6	435	3.2	d
Oxoalkyl				
CH ₃ C(O)CH ₂ O ₂	5.47	282	2.1	d
Acyl				
CH ₃ C(O)O ₂			3.2	e

Comments: ^a IUPAC Task Group recommendation (<http://iupac.pole-ether.fr/>, last access: January 2019). ^b $k_{298\text{K}}$ based on an average of the values reported by Biggs et al. (1995), Ray et al. (1996), Vaughan et al. (2006) and Laversin et al. (2016). E/R taken from Laversin et al. (2016). ^c Taken from Vaughan et al. (2006). ^d Taken from Kalalian et al. (2018). ^e Taken from Doussin et al. (2003). Canosa-Mas et al. (1996) reported a comparable value of $k = (4.0 \pm 1.0) \times 10^{-12}$ cm³ molecule⁻¹ s⁻¹ over the range 403–443 K.




Figure 1. Reported thermal decomposition rates of selected peroxyacyl nitrates at 298 K and 760 Torr. Values for PAN, PPN and MPAN (defined in the figure) are the IUPAC Task Group recommendations (<http://iupac.pole-ether.fr/>, last access: January 2019). The other values are taken from the systematic study of Kabir et al. (2014), which also reports consistent values for PAN and PPN. The broken line is the generic rate coefficient, $k_{b,\text{PAN}}$, for the decomposition of RC(O)OONO₂ structures (see Sect. 2.2 and Table 4).

sidered appropriate for application to all $\geq \text{C}_2$ non-acyl peroxy radicals. For CH₃O₂, the reaction has been well studied at 298 K, and the value in Table 5 is applied in conjunction with the $k_{\text{RO}_2\text{NO}_3}$ pre-exponential factor, leading to

$$k(\text{CH}_3\text{O}_2 + \text{NO}_3) = 8.9 \times 10^{-12} \exp(-600/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}. \quad (6)$$

The generic rate coefficient for acyl peroxy radicals is based on data for CH₃C(O)O₂, which have been shown to react slightly more rapidly with NO₃ (Canosa-Mas et al., 1996; Doussin et al., 2003). The value at 298 K in Table 5 (based on that reported by Doussin et al., 2003) is once again applied in conjunction with the $k_{\text{RO}_2\text{NO}_3}$ pre-exponential factor, leading to

$$k_{\text{APNO}_3} = 8.9 \times 10^{-12} \exp(-305/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}. \quad (7)$$

The resultant weak temperature dependence yields a value of k_{APNO_3} in the range 403–443 K that is fully consistent with that reported by Canosa-Mas et al. (1996).



2.4 The reaction of RO₂ with OH

Kinetics determinations have been reported for the reactions of OH with C₁–C₄ alkyl peroxy radicals. As shown in Table 6, these reactions are reported to occur rapidly at room temperature, with the rate coefficients for all the reactions being essentially equivalent at 298 K, within the reported uncertainties. Based on the study by Yan et al. (2016), a weak temperature dependence is recommended for the reaction of CH₃O₂ with OH, and the resultant expression,

$$k_{\text{RO}_2\text{OH}} = 3.7 \times 10^{-11} \exp(350/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}, \quad (8)$$

is also adopted in the present work as a generic rate coefficient for the reactions of RO₂ with OH.

The following product channels are considered, but with their branching ratios being strongly dependent on the size of R:



In their theoretical studies of the reaction of CH₃O₂ with OH, Bian et al. (2015), Müller et al. (2016) and Assaf et al. (2018) calculated reaction channel (R5a) to be the most favourable, with experimental confirmation of a dominant contribution from this channel reported for CH₃O₂ by Assaf et al. (2017a, 2018). A number of alternative channels have been considered in modelling assessments (e.g. Archibald et al., 2009), including formation of CH₂O₂ and H₂O or CH₃OH and O₂. However, no evidence for formation of CH₂O₂ and H₂O has been observed at room temperature, indicating that this product channel is at most minor (< 5 %) (Yan et al., 2016; Assaf et al., 2017a; Caravan et al., 2018); this is also consistent with theoretical data (e.g. Müller et al., 2016). The formation of CH₃OH and O₂ via channel (R5b) has been shown to make a minor contribution (6 %–9 %) in the experimental study of Caravan et al. (2018), consistent with the theoretical estimate of ~ 7 % by Müller et al. (2016). It is noted that Caravan et al. (2018) also reported evidence for minor CH₃OOH formation at atmospheric pressure, via channel (R5c); although this was calculated to be formed with a low yield (1.7 %) by Assaf et al. (2018). As a result, values of $k_{5a}/k_5 = 0.93$ and $k_{5b}/k_5 = 0.07$ are currently assigned to the reaction of CH₃O₂ with OH in the present work.

The experimental and theoretical study of Assaf et al. (2018) for a series of C₁–C₄ alkyl peroxy radicals has demonstrated that the reaction can more generally be regarded as proceeding by either channel (R5a) or (R5c). Formation of the thermalized hydrotrioxide, ROOOH, via channel (R5c) was found to be increasingly important for the larger RO₂. Based approximately on their theoretical calculations for 298 K and 1 atm of pressure, k_{5c}/k_5 is thus currently assigned a value of 0.0 for CH₃O₂, 0.8 for RO₂ for which

$n_{\text{CON}} = 2$ (e.g. C₂H₅O₂ and HOCH₂O₂) and 1.0 for all other RO₂ radicals. In the $n_{\text{CON}} = 2$ case, the balance of the reaction is assigned to channel (R5a), i.e. with $k_{5b}/k_5 = 0$. As discussed by Assaf et al. (2018), detailed experimental and theoretical studies of the atmospheric fate of ROOOH are therefore clearly required for the effect of the RO₂ + OH reaction to be fully assessed and represented. A provisional treatment is provided in Sect. S3, based mainly on rate coefficients reported in the theoretical studies of Müller et al. (2016), Assaf et al. (2018) and Anglada and Solé (2018).

2.5 The reaction of RO₂ with HO₂

2.5.1 Kinetics

Rate coefficients for the reactions of HO₂ with a variety of specific hydrocarbon and oxygenated RO₂ radicals have been reported, as summarized in Table 7. For the vast majority of the RO₂ radicals formed in detailed mechanisms, however, kinetic data are unavailable, and it is therefore necessary to assign generic rate coefficients based on the reported data.

As discussed previously (Jenkin et al., 1997; Saunders et al., 2003; Boyd et al., 2003a; Orlando and Tyndall, 2012; Wennberg et al., 2018), the 298 K rate coefficients tend to increase with the size of the organic group. Figure 2 shows the data plotted as a function of n_{CON} . The data for alkyl peroxy radicals and β -hydroxyalkyl peroxy radicals (the most systematically studied groups) show comparable values across the n_{CON} range. Based on optimization to these data, the following expression is derived for application to non-acyl peroxy radicals:

$$k_{\text{RO}_2\text{HO}_2} = 2.8 \times 10^{-13} \exp(1300/T) [1 - \exp(-0.23n_{\text{CON}})] \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1} \quad (9)$$

The temperature dependence is typical of that reported for > C₂ alkyl and β -hydroxy RO₂ radicals and remains unchanged from that applied previously by Saunders et al. (2003).

Based on the limited data for acyl peroxy radicals (see Fig. 2 and Table 7), and specifically that for CH₃C(O)O₂, the 298 K rate coefficients are assigned values that are almost a factor of 2 greater than those defined by Eq. (9). The temperature dependences reported for acyl peroxy radicals appear to be weaker than those for similar-sized radicals in other classes, and the temperature coefficient is again based on that recommended for CH₃C(O)O₂. The following expression is therefore assigned to acyl peroxy radicals:

$$k_{\text{APHO}_2} = 3.5 \times 10^{-12} \exp(730/T) [1 - \exp(-0.23n_{\text{CON}})] \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}. \quad (10)$$

Table 6. Kinetic data for the reactions of peroxy radicals with OH. Where available, the temperature dependence is given by $k = A \exp(-E/RT)$.

Peroxy radical	A ($10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)	E/R (K)	$k_{298\text{K}}$ ($10^{-10} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)	Comment
CH_3O_2	3.7	-350	1.2	^a
$\text{C}_2\text{H}_5\text{O}_2$			1.2	^b
(<i>n</i> - and <i>i</i> -) $\text{C}_3\text{H}_7\text{O}_2$			1.4	^c
(<i>n</i> - and <i>sec</i> -) $\text{C}_4\text{H}_9\text{O}_2$			1.5	^c

Comments: ^a IUPAC Task Group recommendation (<http://iupac.pole-ether.fr/>, last access: January 2019) based on Assaf et al. (2016) and Yan et al. (2016) with an uncertainty factor of 2 assigned to $k_{298\text{K}}$; ^b IUPAC Task Group recommendation (<http://iupac.pole-ether.fr/>, last access: January 2019) based on Faragó et al. (2015) with an uncertainty factor of 1.6 assigned to $k_{298\text{K}}$. A consistent value of $k_{298\text{K}} = (1.3 \pm 0.3) \times 10^{-10} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ has more recently been reported by Assaf et al. (2017b); ^c Global rate coefficients, $k_{298\text{K}} = (1.4 \pm 0.3) \times 10^{-10}$ and $(1.5 \pm 0.3) \times 10^{-10} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, reported by Assaf et al. (2017b) for isomeric mixtures of peroxy radicals formed from reactions of Cl atoms with propane and butane, respectively.





Figure 2. Rate coefficients for the reactions of RO_2 radicals with HO_2 as a function of n_{CON} at 298 K. The heavy broken line is the best fit to the data for alkyl and β -hydroxyalkyl RO_2 on the basis of the assumed function $k = A(1 - \exp(Bn_{\text{CON}}))$. The light broken line is the same function with the 298 K value of k increased by a factor of 1.84 (see Sect. 2.5).

2.5.2 Product branching ratios

On the basis of reported information, the following channels are considered for the reactions of RO_2 with HO_2 :



Formation of a hydroperoxide product (ROOH) and O_2 via terminating channel (R6a) is reported to be dominant for reactions of alkyl peroxy radicals, and this is also taken to be the default where no information is available (see Table 8). However, the reactions of HO_2 with oxygenated peroxy radicals have received considerable attention and evidence has been reported for several additional channels leading to both

radical termination, Reactions (R6b) and (R6c), and radical propagation, Reactions (R6d) and (R6e). Table 8 summarizes the 298 K branching ratios that are applied to several classes of oxygenated peroxy radical based on reported information.

The temperature dependences of the reaction channels have generally not been studied, and the branching ratios in Table 8 are thus applied independent of temperature in most cases. The only exception is the reaction of HO_2 with (non-aryl) acyl peroxy radicals. This class of reaction (in particular the reaction of HO_2 with $\text{CH}_3\text{C}(\text{O})\text{O}_2$) has received the most attention and is also a class for which radical propagation is reported to be particularly important at temperatures near 298 K. As shown in Table 8, channels (R6a), (R6b) and (R6d) are reported to contribute. The temperature dependence of k_{6d}/k is based on the recent study of the $\text{CH}_3\text{C}(\text{O})\text{O}_2 + \text{HO}_2$ reaction reported by Hui et al. (2019). The contributions and temperature dependences of k_{6a}/k and k_{6b}/k also take account of the wider database for the same reaction, in particular the experimental characterization of k_{6a}/k_{6b} reported by Horie and Moortgat (1992). This procedure (described in detail in Sect. S4) results in the following fitted Arrhenius expressions for the individual channel rate coefficients:

$$k_{6a\text{APHO}_2} = 3.00 \times 10^{-12} \exp(480/T) [1 - \exp(-0.23n_{\text{CON}})], \quad (11)$$

$$k_{6b\text{APHO}_2} = 8.83 \times 10^{-15} \exp(1910/T) [1 - \exp(-0.23n_{\text{CON}})], \quad (12)$$

$$k_{6d\text{APHO}_2} = 9.35 \times 10^{-12} \exp(235/T) [1 - \exp(-0.23n_{\text{CON}})]. \quad (13)$$

The corresponding temperature dependences of the channel rate coefficients, derived from the $\text{CH}_3\text{C}(\text{O})\text{O}_2$ data, are thus applied to all (non-aryl) acyl peroxy radicals. The variation of the branching ratios and channel rate coefficients are illustrated for the $\text{CH}_3\text{C}(\text{O})\text{O}_2 + \text{HO}_2$ reaction in Figs. S2 and S3 for the 230–300 K temperature range. Summation of the channel rate coefficients given in Eqs. (11)–(13) reproduces the values of k_{APHO_2} calculated for the overall reaction us-

Table 7. Kinetic data for the reactions of hydrocarbon and oxygenated peroxy radicals with HO₂. Where available, the temperature dependence is given by $k = A \exp(-E/RT)$.

Peroxy radical	A ($10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)	E/R (K)	$k_{298 \text{ K}}$ ($10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)	Comment
Alkyl and cycloalkyl				
CH ₃ O ₂	3.8	−780	5.2	a
C ₂ H ₅ O ₂	6.4	−710	6.9	a
<i>neo</i> -C ₅ H ₁₁ O ₂	1.4	−1380	14.4	b
<i>c</i> -C ₅ H ₉ O ₂	2.1	−1323	17.8	b
<i>c</i> -C ₆ H ₁₁ O ₂	2.6	−1245	17.0	b
decane-derived			19.5	c,d
tetradecane-derived			21.1	c,e
Allylic (alk-2-enyl)				
CH ₂ =CHCH ₂ O ₂	~ 10	−700	~ 10	f
β -aryl				
C ₆ H ₅ CH ₂ O ₂	1.5	−1310	12.0	a
α -Hydroxyalkyl				
HOCH ₂ O ₂	0.056	−2300	12.0	a
β -Hydroxyalkyl				
HOCH ₂ CH ₂ O ₂			13.0	a
CH ₃ CH(OH)CH(O ₂)CH ₃			15.0	a,g
(CH ₃) ₂ C(OH)CH ₂ O ₂	0.56	−1650	14.0	a
(CH ₃) ₂ C(OH)C(O ₂)(CH ₃) ₂			15.0	c
HO- <i>c</i> -C ₆ H ₁₀ O ₂			22.4	c,h
α -pinene-derived			20.9	c,i
γ -terpinene-derived			19.7	c,i
<i>d</i> -limonene-derived			21.2	c,i
Hydroxyalkenyl				
isoprene-derived			17.4	c,j
Alkoxyalkyl				
CH ₃ OCH ₂ O ₂			~ 10	k
Oxoalkyl				
CH ₃ C(O)CH ₂ O ₂			9.0	a,l
Acyl				
CH ₃ C(O)O ₂	17.3	−730	20.0	m
C ₆ H ₅ C(O)O ₂	110	−364	37.0	a,n

Comments: ^a IUPAC Task Group recommendation (<http://iupac.pole-ether.fr/>, last access: January 2019). ^b Based on Rowley et al. (1992a, b) and Boyd et al. (2003a). ^c Taken from Boyd et al. (2003a). ^d Mixture of C₁₀H₂₁O₂ radicals derived from the reaction of OH with decane. ^e Mixture of C₁₄H₂₉O₂ radicals derived from the reaction of OH with tetradecane. ^f Approximate value from Boyd et al. (1996a), based on extrapolation of higher temperature data (393–426 K) using assumed value of $E/R = -700$ K. ^g Taken from Jenkin and Hayman (1995). ^h Derived from the reaction of OH with cyclohexene. RO₂ population dominated by β -hydroxy peroxy radical, HO-*c*-C₆H₁₀-O₂, formed from OH addition. ⁱ RO₂ population dominated by hydroxy peroxy radicals formed from OH addition to the given monoterpene. ^j Mixture of HOC₅H₈O₂ radicals derived from the reaction of OH with isoprene. ^k Approximate value from Jenkin et al. (1993a), based on steady state concentration of HO₂ formed from the self-reaction of CH₃OCH₂O₂ during modulated photolysis. ^l Based on Bridier et al. (1993). ^m $k_{298 \text{ K}}$ based on Groß et al. (2014), Winiberg et al. (2016) and Hui et al. (2019). E/R based on Hui et al. (2019) (see Sect. S4). ⁿ Based on Roth et al. (2010).

Table 8. Branching ratios assigned to reaction channels (R6a)–(R6e) for reactions of hydrocarbon and oxygenated peroxy radical classes with HO₂ at 298 K.






Peroxy radical class	Channel branching ratios					Comment
	k_{6a}/k_6	k_{6b}/k_6	k_{6c}/k_6	k_{6d}/k_6	k_{6e}/k_6	
Alkyl (and default)	1.00	–	–	–	–	a
Acyl (R ≠ aryl)	0.37	0.13	–	0.50	–	b
Acyl (R = aryl)	0.65	0.15	–	0.20	–	c
β-Oxoalkyl (prim.)	0.82	–	–	0.18	–	d
β-Oxoalkyl (sec., tert.)	0.52	–	–	0.48	–	e
α-Alkoxyalkyl (prim., sec.)	0.54	–	0.26	–	0.20	f
α-Alkoxyalkyl (tert.)	1.00	–	–	–	–	g
β-Hydroxyallylic	0.92	–	–	0.08	–	h
β-Nitrooxyallylic	0.50	–	–	0.50	–	i

Comments: ^a Based on studies of CH₃O₂ and C₂H₅O₂ (as summarized by Orlando and Tyndall, 2012), and also used as a default in all cases other than those covered by comments ^{b–i}. ^b Based on studies of CH₃C(O)O₂ (Niki et al., 1985; Horie and Moortgat, 1992; Hasson et al., 2004; Jenkin et al., 2007; Dillon and Crowley, 2008; Groß et al., 2014; Winiberg et al., 2016; Hui et al., 2019); see Sect. S4. Hasson et al. (2012) also reported broadly comparable branching ratios for C₂H₅C(O)O₂ and C₃H₇C(O)O₂. ^c Based on studies of C₆H₅C(O)O₂. k_{6d}/k_6 based on Dillon and Crowley (2008) and Roth et al. (2010), with other branching ratios based on those reported by Roth et al. (2010). ^d Based on studies of CH₃C(O)CH₂O₂ (Jenkin et al., 2008; Dillon and Crowley, 2008; Hasson et al., 2012). ^e Based on studies of CH₃C(O)CH(O₂)CH₃ (Dillon and Crowley, 2008; Hasson et al., 2012) and of CH₃C(O)CH(O₂)CH₂OH Praskie et al. (2015). Praskie et al. (2015) also reported possible minor contribution of channel (R6e) and/or (R6c) for CH₃C(O)CH(O₂)CH₂OH. ^f Based on studies of HOCH₂O₂ (Jenkin et al., 2007) and CH₃OCH₂O₂ (Jenkin et al., 2010). Contribution of OH formation in those studies was originally attributed to channel (R6d), but is allocated here to channel (R6e) on the basis of the theoretical study of Nguyen et al. (2010). ^g Full reaction is assigned to channel (R6a), because channels (R6c) and (R6e) are unavailable for tertiary radicals owing to the absence of an α-H atom. ^h Based on study of hydroxyallylic peroxy radicals formed in isoprene system by Liu et al. (2013), with support from the studies of Paulot et al. (2009) and Navarro et al. (2013). ⁱ Based on study of nitrooxyallylic radicals formed in isoprene system by Schwantes et al. (2015).

ing Eq. (10) to within 5 % over this temperature range (see Sect. S4 for further details).

2.6 The permutation reactions of RO₂

The “permutation” reactions of a given RO₂ radical are its self-reactions (R7) and its cross-reactions (R8) with other peroxy radicals, R'O₂, for which a number of product channels may occur:



In view of the large number of RO₂ radicals generated in a detailed chemical mechanism, however, it is unrealistic to represent these reactions explicitly, and the use of simplified parameterizations is essential (Madronich and Calvert, 1990). As described in detail previously (Jenkin et al., 1997), a very simplified approach has traditionally been adopted in the MCM, in which each peroxy radical is assumed to react

with all other peroxy radicals (i.e. the peroxy radical “pool”) at a single, collective rate. This is achieved by defining a parameter “ $\sum[\text{RO}_2]$ ” which is the sum of the concentrations of all peroxy radicals, excluding HO₂. The collective rate of all the permutation reactions of a particular peroxy radical is then represented by a single pseudo-unimolecular reaction, which has an assigned rate coefficient equal to $k_9 \times \sum[\text{RO}_2]$,



with the value of k_9 depending on the structure of the reacting RO₂ radical. A similar, but more detailed, approach has been applied in GECKO-A, in which the peroxy radical population is divided into a number of reactivity classes (Aumont et al., 2005). This requires the inclusion of a pseudo-unimolecular reaction (analogous to Reaction R9) for reaction of a given peroxy radical with each peroxy radical class, but has the advantage that differential reactivity with each of those classes can be represented, as appropriate. The following subsections describe the basis for assigning rate parameters to the single parameterized permutation reactions (Reaction R9) for each peroxy radical in the more simplified MCM approach. Extension of the method to reactions with a number of reactivity classes (as traditionally applied with GECKO-A) is described in Sect. S5.

2.6.1 Kinetics of self-reactions

Rate coefficients for the self-reactions and cross-reactions of a variety of specific hydrocarbon and oxygenated RO₂ radicals have been reported (as summarized in Tables 9–11), and these form the basis of assigning rate parameters to the parameterized permutation reaction (Reaction R9) for each peroxy radical. The data show that the self-reaction reactivity, relative to that of alkyl peroxy radicals, is activated by the presence of numerous functional groups (including allyl-, benzyl-, hydroxy-, alkoxy-, oxo- and acyl-groups), and that the rate coefficients follow the general trend of decreasing reactivity, primary > secondary > tertiary, for peroxy radicals containing otherwise similar functionalities. It also appears that reactivity tends to increase with the size of the organic group towards a “plateau” value, as most clearly demonstrated by the systematic study of secondary alkyl peroxy radicals reported by Boyd et al. (1999). Based on optimization to the complete secondary alkyl peroxy radical dataset, an expression almost identical to that recommended by Boyd et al. (1999) is thus derived as a reference rate coefficient for secondary peroxy radicals at 298 K, as illustrated in Fig. 3 (units of k are cm³ molecule⁻¹ s⁻¹):

$$\log_{10}(k^{\circ}_{\text{RO}_2\text{RO}_2(\text{sec})}) = -12.9 - (3.2 \times \exp[-0.64(n_{\text{CON}} - 2.3)]). \quad (14)$$

The data for primary alkyl peroxy radicals are more limited. Those for C₂H₅O₂, *n*-C₃H₇O₂, *i*-C₄H₉O₂ and *neo*-C₅H₁₁O₂ suggest a similar trend for primary alkyl peroxy radicals, and an analogous expression to Eq. (14) is therefore derived as a reference rate coefficient at 298 K (see Fig. 3):

$$\log_{10}(k^{\circ}_{\text{RO}_2\text{RO}_2(\text{prim})}) = -11.7 - (3.2 \times \exp[-0.55(n_{\text{CON}} - 0.52)]). \quad (15)$$

It is noted, however, that rate coefficients for the self-reactions of *n*-C₄H₉O₂ and *n*-C₅H₁₁O₂ are reported to be comparable to that for *n*-C₃H₇O₂, and a factor of 2 to 3 lower than those for *i*-C₄H₉O₂ and *neo*-C₅H₁₁O₂ (see Table 9), suggesting that there may be sensitivity to whether the alkyl group is linear or branched. In the absence of additional data (and noting that the kinetics of *neo*-C₅H₁₁O₂ were the most directly determined of the set of C₄ and C₅ primary alkyl peroxy radicals), the above (stronger) size dependence is provisionally applied here.

Data for tertiary alkyl peroxy radicals are currently limited to *t*-C₄H₉O₂, and the corresponding rate coefficient is currently applied as the reference rate coefficient at 298 K, independent of radical size (see Fig. 3):

$$k^{\circ}_{\text{RO}_2\text{RO}_2(\text{tert})} = 2.1 \times 10^{-17} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}. \quad (16)$$

Figure 3 also shows data for allylic and β -hydroxyalkyl RO₂, demonstrating that the presence of both these functionalities




Figure 3. Rate coefficients for the self-reactions of alkyl (filled points), β -hydroxyalkyl (open points) and allylic (\times) RO₂ at 298 K as a function of n_{CON} . Grey filled points indicate where the reported rate coefficient has not been corrected for secondary chemistry. Where available, data are shown for primary, secondary and tertiary radicals containing the given functionalities. Primary, secondary and tertiary alkyl and β -hydroxyalkyl radicals are shown as diamonds, circles and triangles, respectively. The “allylic” peroxy radical group contains only primary radicals and includes “ δ -hydroxyallylic” peroxy radicals. The lines represent the calculated rate coefficients fitted to the data using the methods described in Sect. 2.6.

has an activating effect on self-reaction reactivity. The allylic peroxy radical category includes two δ -hydroxyallylic peroxy radicals, and the assumption is made here that the δ -hydroxy group is too remote to have an influence. Table 12 summarizes a series of activation factors (defined in terms of the parameters α and β) for allylic-, β -aryl-, hydroxy-, alkoxy- and oxo-groups, optimized on the basis of the data in Tables 9 and 10. These are used in conjunction with the reference rate coefficients in Eqs. (14)–(16) to calculate the self-reaction rate coefficient for a given peroxy radical at 298 K, $k_{\text{RO}_2\text{RO}_2}$, as follows:

$$k_{\text{RO}_2\text{RO}_2} = k^{\circ}_{\text{RO}_2\text{RO}_2} \times \alpha / (k^{\circ}_{\text{RO}_2\text{RO}_2})^{\beta}, \\ = \alpha \times (k^{\circ}_{\text{RO}_2\text{RO}_2})^{1-\beta}. \quad (17)$$

Here, $k^{\circ}_{\text{RO}_2\text{RO}_2}$ represents the appropriate reference rate coefficient (i.e. for primary, secondary or tertiary RO₂, as appropriate) as defined by Eqs. (14)–(16); and the term $\alpha / (k^{\circ}_{\text{RO}_2\text{RO}_2})^{\beta}$ describes the level of activation from the given substituent. The inclusion of $k^{\circ}_{\text{RO}_2\text{RO}_2}$ within this activation term is required because the relative enhancement of reactivity resulting from a given substituent appears to decrease as the reactivity increases, as illustrated for the β -hydroxyalkyl group data in Fig. 3. Based on this method, the estimated rate coefficients correlate well with those observed for the series of peroxy radicals for which data are currently available (summarized in Tables 9 and 10), as shown in Fig. 4. It is emphasized, however, that the parameters for

Table 9. Kinetic data for the self-reactions of hydrocarbon peroxy radicals. Where available, the temperature dependence is given by $k = A \exp(-E/RT)$.

Peroxy radical	A ($10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)	E/R (K)	$k_{298\text{K}}$ ($10^{-14} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)	Comment
CH ₃ O ₂	1.03	-365	35	a
Alkyl				
Primary				
C ₂ H ₅ O ₂	0.76	0	7.6	a
<i>n</i> -C ₃ H ₇ O ₂			30	a
<i>n</i> -C ₄ H ₉ O ₂			40	b*
<i>i</i> -C ₄ H ₉ O ₂			100	b*
<i>n</i> -C ₅ H ₁₁ O ₂			39	c
<i>neo</i> -C ₅ H ₁₁ O ₂	0.017	-1960	120	d
Secondary				
<i>i</i> -C ₃ H ₇ O ₂	16	2200	0.1	a
<i>sec</i> -C ₄ H ₉ O ₂			7	b*
<i>sec</i> -C ₅ H ₁₁ O ₂			3.3	c,e
<i>sec</i> -C ₁₀ H ₂₁ O ₂			9.4	c,f
<i>sec</i> -C ₁₂ H ₂₅ O ₂			~ 14	c,f
<i>c</i> -C ₅ H ₉ O ₂	2.9	555	4.5	g*
<i>c</i> -C ₆ H ₁₁ O ₂	0.77	184	4.2	g
Tertiary				
<i>t</i> -C ₄ H ₉ O ₂	100	3900	0.0021	a
Allylic (alk-2-enyl)				
CH ₂ =CHCH ₂ O ₂ (primary)	0.54	-760	69	h
β -aryl				
C ₆ H ₅ CH ₂ O ₂ (primary)	0.24	-1620	550	a

Comments: * Reported rate coefficient not corrected for the effects of secondary chemistry, which can lead to either an overestimate or underestimate of the rate coefficient. ^a IUPAC Task Group recommendation (<http://iupac.pole-ether.fr/>, last access: January 2019). ^b Taken from Glover and Miller (2005). ^c Taken from Boyd et al. (1999). ^d Based on Lightfoot et al. (1990). ^e Mixture of 2-pentyl and 3-pentyl peroxy radicals. ^f Mixture of secondary peroxy radicals of given formula. ^g Based Rowley et al. (1991, 1992c). ^h Based on Jenkin et al. (1993b) and Boyd et al. (1996a).

several of the substituent groups are based on data for very limited sets of peroxy radicals, and additional data would be valuable to test and constrain the method.

Information on the effects of multiple substituents is limited to the data for the secondary and tertiary β -hydroxyallylic peroxy radicals, HOCH₂CH(O₂)CH=CH₂ and HOCH₂C(CH₃)(O₂)C(CH₃)=CH₂, given in Table 10. The reported rate coefficients are consistent with the activating impacts of the β -hydroxy and allylic substituents being approximately cumulative, suggesting that an activation factor should be applied for each relevant organic substituent. However, this would lead to unreasonably large estimated values of $k_{\text{RO}_2\text{RO}_2}$ for secondary and tertiary peroxy radicals containing two or three of the most activating substituents, such that the impact needs to be limited. In multifunctional peroxy radicals, therefore, an activating factor is only applied for the most activating oxygenated substituent in a given per-

oxy radical, with an additional factor also applied only for the specific cases of an allylic or a β -aryl substituent, again limited to one (i.e. the most activating) factor if the peroxy radical contains more than one allylic or β -aryl group. In these specific cases, therefore,

$$\alpha = (\alpha_1 \times \alpha_2), \quad (18)$$

$$\beta = (\beta_1 + \beta_2), \quad (19)$$

where α_1 and β_1 refer to the oxygenated substituent, and α_2 and β_2 refer to either the allylic substituent or the β -aryl substituent. Further information is required to allow the impacts of multiple substituents to be defined more rigorously.

2.6.2 Parameterized representation

The rate coefficients for cross-reactions of peroxy radicals (Reaction R8) have often been inferred from those for the

Table 10. Kinetic data for the self-reactions of oxygenated peroxy radicals. Where available, the temperature dependence is given by $k = A \exp(-E/RT)$.

Peroxy radical	A ($10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)	E/R (K)	$k_{298 \text{ K}}$ ($10^{-14} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)	Comment
α -Hydroxyalkyl				
HOCH ₂ O ₂ (primary)			620	a
β -Hydroxyalkyl				
Primary				
HOCH ₂ CH ₂ O ₂	0.78	-1000	220	a
(CH ₃) ₂ C(OH)CH ₂ O ₂	0.14	-1740	480	a
Secondary				
CH ₃ CH(OH)CH(O ₂)CH ₃	0.077	-1330	67	a
HO- <i>c</i> -C ₆ H ₁₀ O ₂			120	b
Tertiary				
(CH ₃) ₂ C(O ₂)CH ₂ OH			1.5	b
(CH ₃) ₂ C(OH)C(O ₂)(CH ₃) ₂	5.9	1420	0.5	c
HO- <i>c</i> -C ₆ H ₈ (CH ₃) ₂ O ₂			2.0	b
Hydroxyallylic (hydroxyalk-2-enyl)				
Primary				
HOCH ₂ CH=CHCH ₂ O ₂ (δ -hydroxy)			280	d
HOCH ₂ C(CH ₃)=C(CH ₃)CH ₂ O ₂ (δ -hydroxy)			390	d
Secondary				
HOCH ₂ CH(O ₂)CH=CH ₂ (β -hydroxy)			570	d
Tertiary				
HOCH ₂ C(CH ₃)(O ₂)C(CH ₃)=CH ₂ (β -hydroxy)			6.9	d
Alkoxyalkyl				
CH ₃ OCH ₂ O ₂ (primary)			210	a
β -Oxoalkyl				
CH ₃ C(O)CH ₂ O ₂ (primary)			800	a
γ -Oxoalkyl				
CH ₃ C(O)C(CH ₃) ₂ CH ₂ O ₂ (primary)			480	e
<i>t</i> -C ₄ H ₉ C(O)C(CH ₃) ₂ CH ₂ O ₂ (primary)			460	e
Acyl				
CH ₃ C(O)O ₂	29	-500	1600	a
C ₂ H ₅ C(O)O ₂			1700	a
(CH ₃) ₂ CHC(O)O ₂			1440	f
(CH ₃) ₃ CC(O)O ₂			1440	f
C ₆ H ₅ C(O)O ₂	3.4	-1110	1400	a
Alkoxyacyl				
CH ₃ OC(O)O ₂ (and HC(O)OCH ₂ O ₂)			2300	g

Comments: ^a IUPAC Task Group recommendation (<http://iupac.pole-ether.fr/>, last access: January 2019). ^b Taken from on Boyd et al. (2003b). ^c Based on Jenkin and Hayman (1995) and Boyd and Lesclaux (1997). ^d Taken from Jenkin et al. (1998). ^e Based on Le Crâne and Lesclaux (2006). ^f Based on Tomas and Lesclaux (2000) and Le Crâne et al. (2004). ^g Taken from Hansen et al. (2003). The kinetics of the two peroxy radicals formed from the reaction of Cl or F with methyl formate reported to possess indistinguishable kinetics.

Table 11. Kinetic data for the cross-reactions of hydrocarbon or oxygenated peroxy radicals at 298 K. Where available, the temperature dependence expression is given in the comments.

Peroxy radical 1	Peroxy radical 2	$k_{298\text{K}}$ ($10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)	Comment
CH ₃ O ₂	C ₂ H ₅ O ₂	2	a
	<i>neo</i> -C ₅ H ₁₁ O ₂	15	b
	<i>c</i> -C ₆ H ₁₁ O ₂	1.0	c
	<i>t</i> -C ₄ H ₉ O ₂	0.031	a,d
	CH ₂ =CHCH ₂ O ₂	17	b,e
	C ₆ H ₅ CH ₂ O ₂	< 20	b
	CH ₃ C(O)CH ₂ O ₂	38	a
	CH ₃ C(O)O ₂	110	a,f
C ₂ H ₅ O ₂	<i>neo</i> -C ₅ H ₁₁ O ₂	5.6	b
	<i>c</i> -C ₆ H ₁₁ O ₂	0.4	b
	CH ₃ C(O)O ₂	100	b
	C ₂ H ₅ C(O)O ₂	120	a
CH ₃ C(O)O ₂	CH ₃ C(O)CH ₂ O ₂	50	a
	<i>c</i> -C ₆ H ₁₁ O ₂	104	g
	<i>t</i> -C ₄ H ₉ O ₂	111	g
	<i>sec</i> -C ₁₀ H ₂₁ O ₂	109	g
	<i>sec</i> -C ₁₂ H ₂₅ O ₂	105	g
HO- <i>c</i> -C ₆ H ₉ (CH ₃)O ₂ (secondary)	HO- <i>c</i> -C ₆ H ₉ (CH ₃)O ₂ (tertiary)	6.2	h
HOCH ₂ CH=CHCH ₂ O ₂	HOCH ₂ CH(O ₂)CH=CH ₂	39	i
HOCH ₂ C(CH ₃)=C(CH ₃)CH ₂ O ₂	HOCH ₂ C(CH ₃)(O ₂)C(CH ₃)=CH ₂	25	i

Comments: ^a IUPAC Task Group recommendation (<http://iupac.pole-ether.fr/>, last access: January 2019). ^b Taken from Villenave and Lesclaux (1996).

^c Based on Villenave and Lesclaux (1996) and Nozière and Hanson (2017). ^d Temperature dependence expression is $3.8 \times 10^{-13} \exp(-1430/T)$.

^e Temperature dependence expression is $2.8 \times 10^{-13} \exp(515/T)$. ^f Temperature dependence expression is $2.0 \times 10^{-12} \exp(500/T)$. ^g Taken from Villenave et al. (1998). ^h Taken from Boyd et al. (2003b). The structures refer to the isomeric secondary and tertiary peroxy radicals formed from the addition of OH to 1-methylcyclohexene. ⁱ Taken from Jenkin et al. (1998). Presented values are limited to those reported for the cross-reactions of the major radicals formed from the terminal addition of OH to buta-1,3-diene and the terminal addition of OH to 2,3-dimethyl-buta-1,3-diene.

Table 12. Substituent activation factors applied to self-reaction rate coefficients, based on Eq. (17).

Substituent	α	β	Comment
alkyl	1.00	0	a
β -hydroxy	8.0×10^{-5}	0.4	b
allylic (alk-2-enyl)	4.0×10^{-2}	0.15	c
β -aryl	5.8×10^{-2}	0.15	d,e
α -alkoxy	7.0×10^{-5}	0.4	f,g
β -oxo	1.6×10^{-4}	0.4	h,g
γ -oxo	5.3×10^{-5}	0.4	i,g

Comments: ^a $\alpha = 1.00$ and $\beta = 0$ by definition for alkyl peroxy radicals. These are also used as a default for peroxy radical classes not covered by comments ^b to ⁱ, with the exception of acyl peroxy radicals (discussed in Sect. 2.6); ^b Based on data for β -hydroxyalkyl peroxy radicals in Table 10; ^c Based on data for allylic and δ -hydroxyallylic peroxy radicals in Tables 9 and 10; ^d Based on data for C₆H₅CH₂O₂ (Table 9); ^e β assumed equivalent to that for allylic substituent; ^f Based on data for CH₃OCH₂O₂ (Table 10); ^g β assumed equivalent to that for β -hydroxy substituent; ^h Based on data for CH₃C(O)CH₂O₂ (Table 10); ⁱ Based on data for γ -oxoalkyl peroxy radicals in Table 10.

self-reactions of the participating peroxy radicals, using a geometric mean rule as first suggested by Madronich and Calvert (1990), i.e.

$$k_8 = 2 \times (k_7 \times k_{7'})^{0.5}, \quad (20)$$

where k_8 is the cross-reaction rate coefficient, and k_7 and $k_{7'}$ are the self-reaction rate coefficients for the participating peroxy radicals, RO₂ and R'O₂. Figure 5 shows that such a correlation provides a reasonable guide in many cases (although a clear deviation from the rule occurs for the particular case of reactions involving acyl peroxy radicals). In the very simplified MCM approach, the rate coefficient for the single parameterized permutation reaction of a given peroxy radical (Reaction R9) is based on that estimated for the cross-reaction of the peroxy radical with CH₃O₂. This is regarded as a logical choice because CH₃O₂ is the most abundant organic peroxy radical in the atmosphere (and therefore most commonly the major reaction partner), and also possesses a self-reaction rate coefficient that is in the middle of the range of reported values (see Tables 9 and 10). Taking account of the correlations in Fig. 5, the rate coefficients (in $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$) for the parameterized permutation reactions at 298 K are defined as follows.




Figure 4. Scatter plot of estimated rate coefficients ($k_{\text{RO}_2\text{RO}_2}$) for peroxy radical self-reactions with those reported (k_7), as listed in Tables 9 and 10. Those shown with reduced size symbols are where the reported value of k_7 was not corrected for secondary chemistry (see Table 9 comments). The broken lines show the factor of 3 range.

$$\text{For acyl RO}_2, k_{\text{AP}(298\text{ K})} = 1.1 \times 10^{-11}. \quad (21)$$

$$\begin{aligned} \text{For other RO}_2 \text{ (except CH}_3\text{O}_2), k_{\text{RO}_2(298\text{ K})} \\ = f_{\text{RO}_2} \times 2 \times (k_{\text{RO}_2\text{RO}_2} \times k_{298}(\text{CH}_3\text{O}_2 + \text{CH}_3\text{O}_2))^{0.5}. \end{aligned} \quad (22)$$

Here, $k_{298}(\text{CH}_3\text{O}_2 + \text{CH}_3\text{O}_2)$ is the rate coefficient for the self-reaction of CH_3O_2 at 298 K ($= 3.5 \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$) and $k_{\text{RO}_2\text{RO}_2}$ is the 298 K self-reaction rate coefficient, estimated as described above (Sect. 2.6.1). f_{RO_2} is a scaling factor that is introduced to describe systematic deviations from the geometric mean rule, if required. Based on the correlations in Fig. 5, a unity value of f_{RO_2} is considered acceptable for primary and secondary peroxy radicals (i.e. no deviation from the geometric mean rule), whereas a value of $f_{\text{RO}_2} = 2$ is applied to tertiary peroxy radicals. This elevated scaling factor is based on observation of Jenkin et al. (1998) for complex tertiary RO_2 cross-reactions.

Based on the reported temperature dependences of peroxy radical self- and cross-reactions (see Tables 9 and 10, and Table 11 comments), k_{AP} and k_{RO_2} are assigned respective pre-exponential factors of 2.0×10^{-12} and $1.0 \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. For acyl peroxy radicals, this is consistent with the temperature dependence reported for the reaction of $\text{CH}_3\text{C}(\text{O})\text{O}_2$ with CH_3O_2 , and results in the following temperature-dependent expression in all cases:

$$k_{\text{AP}} = 2.0 \times 10^{-12} \exp(508/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}. \quad (23)$$

For k_{RO_2} , the pre-exponential factor is a rounded value, based on the geometric mean of those for the self-reactions of non-acyl peroxy radicals given in Tables 9 and 10. This results




Figure 5. Scatter plot of rate coefficients for peroxy radical cross-reactions (k_8) with the geometric mean of the self-reaction rate coefficients (denoted k_7 and k_7') for the participating peroxy radicals, RO_2 and $\text{R}'\text{O}_2$. Open circles are reactions involving an acyl peroxy radical and a non-acyl peroxy radical; closed circles are reactions involving combinations of primary and secondary peroxy radicals; open diamonds are reactions involving a tertiary peroxy radical and a primary or secondary peroxy radical. The heavy broken line is a 1 : 1 relationship; the light broken line is a 2 : 1 relationship; the dot-dash line is $k_8 = 1.1 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$.

in the following temperature dependence expression for non-acyl peroxy radicals (except CH_3O_2),

$$\begin{aligned} k_{\text{RO}_2} = 1.0 \times 10^{-13} \\ \exp(-(E_{\text{RO}_2}/R)/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}, \end{aligned} \quad (24)$$

with E_{RO_2}/R having a case-dependent value of $-298 \times \ln(k_{\text{RO}_2(298\text{ K})}/10^{-13})$, where $k_{\text{RO}_2(298\text{ K})}$ is defined by Eq. (22). Examples of specific rate coefficients estimated using this method are given in Sect. S5 for the peroxy radicals formed from the sequential addition of OH and O_2 to isoprene. As indicated above, the collective rate of all the permutation reactions of a particular peroxy radical is then represented by a pseudo-unimolecular reaction (Reaction R9), which has an assigned rate coefficient equal to $k_{\text{AP}} \times \sum[\text{RO}_2]$ for acyl peroxy radicals, and $k_{\text{RO}_2} \times \sum[\text{RO}_2]$ for all other peroxy radicals (except CH_3O_2). For the specific case of CH_3O_2 , the applied rate coefficient ($k_{\text{CH}_3\text{O}_2}$) is twice the self-reaction rate coefficient given in Table 9,

$$\begin{aligned} k_{\text{CH}_3\text{O}_2} = 2.06 \times 10^{-13} \\ \exp(365/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}, \end{aligned} \quad (25)$$

with the pseudo-unimolecular reaction rate coefficient equal to $k_{\text{CH}_3\text{O}_2} \times \sum[\text{RO}_2]$. This representation is therefore consistent with CH_3O_2 being lost via its self-reaction with the recommended rate coefficient when it is the dominant radical.

Each reaction potentially has up to four product channels, the branching ratios of which depend on the structure of the radical, as shown in Table 13:



Channels (R9a)–(R9c) have been considered previously in the MCM (Jenkin et al., 1997; Saunders et al., 2003). They are the pseudo-unimolecular representation of the self-reaction channels (R7a) and (R7b) and the cross-reaction channels (R8a)–(R8c), which are reported to account for most of the reaction, particularly for smaller peroxy radicals (e.g. Lightfoot et al., 1992; Orlando and Tyndall, 2012). As shown in Table 13, channels (R9a)–(R9c) continue to represent the complete reaction in the current parameterized methodology.

Although not currently included in the parameterized representation, channel (R9d) is listed to acknowledge the potential formation of peroxide products (i.e. Reactions R7c and R8d). Although these channels have generally been reported to be minor for small peroxy radicals (e.g. Lightfoot et al., 1992; Orlando and Tyndall, 2012), recent studies suggest that they may be more significant for larger peroxy radicals containing oxygenated substituents, and they have been reported to play a role in the formation of low-volatility products in a number of studies (Ziemann, 2002; Ng et al., 2008; Ehn et al., 2014; Jokinen et al., 2014; Mentel et al., 2015; Rissanen et al., 2015; Berndt et al., 2015, 2018a, b; Zhang et al., 2015; McFiggans et al., 2019). These reactions may therefore play a potentially important role in particle formation and growth in the atmosphere. The product denoted “ $\text{RO}_{(\text{peroxide})}$ ” in the pseudo-unimolecular approach represents the monomeric contribution the given peroxy radical makes to the total formation of (dimeric) peroxide products, but is not an independent species for which subsequent gas-phase chemistry can be rigorously defined. In principle, channel (R9d) can be included for the permutation reactions of a subset of larger peroxy radicals, with the $\text{RO}_{(\text{peroxide})}$ product assumed to transfer completely to the condensed phase (i.e. not participating in gas-phase reactions). However, there is currently insufficient information on the structural dependence of the contributions of channels (R7c) or (R8d) to the overall self- and cross-reactions to allow the branching ratio of channel (R9d) to be defined reliably. Further systematic studies of these channel contributions are therefore required as a function of peroxy radical size and functional group content.

3 Unimolecular reactions of RO_2 radicals

Unimolecular isomerization reactions are potentially available for some classes of RO_2 . These generally fall into the category of either ring-closure reactions (where the peroxy radical adds intra-molecularly to an unsaturated linkage to form a peroxide-bridged radical product) or reactions involving the migration of a hydrogen atom to the peroxy radical group (e.g. forming a hydroperoxy-substituted organic radical product when abstraction from a C–H bond occurs). For some RO_2 structures, these reactions have been shown to compete with (or dominate over) the bimolecular reactions under some atmospheric conditions, as discussed further below in Sect. 3.1 and 3.2. Evidence for the operation of peroxy radical isomerization reactions has been reported in numerous theoretical and laboratory studies (e.g. Vereecken and Peeters, 2004; Peeters et al., 2009; 2014; Crouse et al., 2013; Ehn et al., 2014, 2017; Jokinen et al., 2014; Rissanen et al., 2015; Jørgensen et al., 2016; Praske et al., 2017, 2019; Otkjær et al., 2018; Mohammed et al., 2018), and new information is constantly emerging on this important aspect of peroxy radical chemistry (e.g. Bianchi et al., 2019; Xu et al., 2019; Møller et al., 2019). The present section provides a summary of selected classes of isomerization reactions that are currently being considered and represented in ongoing mechanism development work. However, it does not currently attempt to provide a full treatment of unimolecular reactions of RO_2 radicals, which will be considered further in future work as more new information becomes available.

3.1 Ring-closure reactions of RO_2

Table 14 shows representative rate coefficients for selected template ring-closure reactions. The first entry relates to the β -hydroxy cyclohexadienylperoxy radicals formed from the addition of O_2 to OH-aromatic hydrocarbon adducts. As discussed in the companion paper on the OH-initiated oxidation of aromatic VOCs (Jenkin et al., 2018b), these peroxy radicals are represented to undergo rapid and exclusive ring closure to produce a hydroxy-dioxa-bicyclo or “peroxide-bicyclic” radical. This reaction has been calculated to dominate over alternative bimolecular reactions of the peroxy radicals under atmospheric conditions (see Table 14), although evidence for competitive loss via bimolecular reactions has been characterized in experimental studies using high concentrations of NO and/or RO_2 (e.g. Birdsall et al., 2010; Birdsall and Elrod, 2011).

The remaining reactions in Table 14 are based on information presented by Vereecken and Peeters (2004) for specific peroxy radicals formed from the sequential addition of OH and O_2 to isoprene, α -pinene and β -pinene. That information has been used to assign or infer representative rate coefficients to the series of related template peroxy radical structures presented in Table 14. In these cases, the reactions are expected to occur at rates that can compete to varying extents

Table 13. Branching ratios assigned to parameterized permutation reactions of RO₂ (see text).

Peroxy radical class	Channel branching ratios			Comment
	k_{9a}/k_9	k_{9b}/k_9	k_{9c}/k_9	
CH ₃ O ₂	$7.2 \times \exp(-885/T)$	$(1 - (k_{9a}/k_9))/2$	$(1 - (k_{9a}/k_9))/2$	a
Primary and secondary	0.6	0.2	0.2	b
Tertiary and acyl	0.8	–	0.2	c

Comments: ^a Based on IUPAC Task Group recommendation for the CH₃O₂ self-reaction (<http://iupac.pole-ether.fr/>, last access: January 2019). An alternative representation using temperature-dependent channel rate coefficients is provided in Sect. S5. ^b Based on a rounded mean of the reported 298 K branching ratios for the self-reactions of C₂H₅O₂, i-C₃H₇O₂, HOCH₂CH₂O₂, (CH₃)₂C(OH)CH₂O₂, CH₃C(O)CH₂O₂, CH₃OCH₂O₂ and C₆H₅CH₂O₂ based on IUPAC Task Group recommendations (<http://iupac.pole-ether.fr/>, last access: January 2019); *neo*-C₅H₁₁O₂, *c*-C₆H₁₁O₂ and CH₂=CHCH₂O₂ based on Lightfoot et al. (1990), Rowley et al. (1991), Jenkin et al. (1993a, b) and Boyd et al. (1996a); and for the self- and cross-reactions of primary and secondary RO₂ formed from reactions of OH with conjugated dienes (Jenkin et al., 1998). ^c Based on a rounded mean of the reported 298 K branching ratios for the following cross-reactions: CH₃C(O)O₂ + CH₃O₂, C₂H₅C(O)O₂ + C₂H₅O₂ and CH₃C(O)O₂ + CH₃C(O)CH₂O₂ based on IUPAC Task Group recommendations (<http://iupac.pole-ether.fr/>, last access: January 2019); and HOCH₂C(CH₃)(O₂)C(CH₃)=CH₂ + HOCH₂C(CH₃)=C(CH₃)CH₂O₂ formed from reaction of OH with 2,3-dimethyl-but-1,3-diene (Jenkin et al., 1998).

with loss via bimolecular reactions (or other unimolecular reactions discussed below) under atmospheric conditions. It is noted that Xu et al. (2019) have also very recently reported information for a series of isomerization reactions (including ring-closure reactions) for the α - and β -pinene systems, which are being considered in ongoing work.

3.2 Hydrogen atom migration reactions of RO₂













Table 15 shows selected hydrogen atom migration reactions that are currently considered. The rate coefficient assigned generally to the 1,4-formyl H-shift reaction of α -formyl peroxy radicals is based on that determined for the methacrolein-derived peroxy radical, HOCH₂C(CH₃)(O₂)C(=O)H, in the experimental study of Crouse et al. (2012). It is noted that this is slightly higher than, but comparable with, the range of values reported for α -formyl peroxy radicals in the preliminary calculations of Peeters and Nguyen (2012).

The rate coefficients assigned to the 1,4-hydroxyl H-shift reactions of (thermalized) α -hydroxy peroxy radicals are based on those estimated for secondary, tertiary and cyclic peroxy radicals in the theoretical study of Hermans et al. (2005). As discussed in the companion paper on the OH-initiated oxidation of aliphatic VOCs (Jenkin et al., 2018a), thermalized α -hydroxy peroxy radicals are represented to be increasingly formed from the reactions of O₂ with larger α -hydroxy organic radicals (i.e. those with $n_{\text{CON}} > 5$). At the assigned rates, the 1,4-hydroxyl H-shift reaction is likely to be the major fate of the majority of thermalized α -hydroxy peroxy radicals under atmospheric conditions, and therefore indistinguishable from that of the chemically activated α -hydroxy peroxy radical adducts that are formed predominantly from the reactions of O₂ with small α -hydroxy organic radicals (see Sect. 6.2 of Jenkin et al., 2018a). However, the rates of the 1,4-hydroxyl H-shift reactions are formalized in the present work to allow for the representation of

competing rapid isomerization reactions for specific structurally complex peroxy radicals (e.g. the 1,6-enol H-shift reaction discussed below), or with bimolecular reactions under appropriate conditions. It is noted that evidence for competitive loss via bimolecular reactions has been characterized in experimental studies using high concentrations of NO (e.g. Orlando et al., 2000b; Jenkin et al., 2005; Aschmann et al., 2010), leading to the formation of organic acids.

The remaining reactions in Table 15 are inferred from information reported for specific unsaturated peroxy radicals formed during the OH-initiated oxidation of isoprene, taking particular account of the work of Peeters et al. (2009, 2014) on the Leuven isoprene mechanism (LIM1), which has been largely verified by experimental study (e.g. Wennberg et al., 2018; and references therein). The rate coefficients for the 1,5-hydroxyl H-shift reactions are those reported by Peeters et al. (2014) for the corresponding unsaturated secondary and tertiary β -hydroxy peroxy radicals formed from the sequential addition of OH and O₂ to isoprene, with these also being generally consistent with those reported by da Silva et al. (2010). The rate coefficient assigned to the 1,6-hydroxyalkyl H-shift reaction is the geometric mean of rate coefficients applied to (Z)-CH₂(OH)C(CH₃)=CHCH₂O₂ (CISOPAO2) and (Z)-CH₂(OH)CH=C(CH₃)CH₂O₂ (CISOPCO2) in MCM v3.3.1. As discussed by Jenkin et al. (2015), those rate coefficients are derived from the LIM1 calculations of Peeters et al. (2014), but with some scaling to recreate the observations of Crouse et al. (2011, 2014). The generic rate coefficient is applied generally to unsaturated δ -hydroxy peroxy radicals containing the substructure shown, but with the exceptions of CISOPAO2 and CISOPCO2 themselves, for which the species-specific rate coefficients are applied (see Sect. S6 and Table S5). Similarly, the rate coefficient for the rapid 1,6-enol H-shift reaction is the geometric mean of those calculated for (Z)-HOCH=C(CH₃)CH(O₂)CH₂OH and (Z)-HOCH=CHC(CH₃)(O₂)CH₂OH by Peeters and

Table 14. Representative rate coefficients for template ring-closure reactions of peroxy radicals^a.

Radical	Product		<i>A</i> (s ⁻¹)	<i>E/R</i> (K)	<i>k</i> _{298 K} (s ⁻¹)	Comment
			–	–	(3.6–2500) × 10 ²	^b
		sec. ^c	1.0 × 10 ¹⁰	8140	0.014	^d
		tert.	1.0 × 10 ¹⁰	7740	0.053	^e
		sec.	1.0 × 10 ¹⁰	7740	0.053	^e
		tert.	1.0 × 10 ¹⁰	7340	0.20	^f
		sec.	4.8 × 10 ¹⁰	7850	0.17	^e
		tert.	4.8 × 10 ¹⁰	7450	0.67	^g
		sec.	4.8 × 10 ⁹	6750	0.70	^e
		tert.	4.8 × 10 ⁹	6350	2.7	^h
		tert.	1.4 × 10 ¹⁰	7100	0.63	ⁱ

Comments: ^a Temperature dependence of rate coefficient given by $k = A \exp(-E/R/T)$. Rapid reaction of the product radical with O₂ dominates over the reverse ring-opening reaction under atmospheric conditions. Entries in bold font are based on reported data for the specific or closely related structures, with other entries inferred using assumptions given in the following comments. ^b Range of 298 K values based on the calculations of Raoult et al. (2004), Glowacki et al. (2009) and Olivella et al. (2009) for the dominant conformer of the example peroxy radical, formed during the oxidation of benzene. Based on these data, and data for other aromatic systems, analogous ring-closure reactions are assumed to be the exclusive fates of corresponding peroxy radicals formed during the oxidation of aromatic hydrocarbons (Jenkin et al., 2018b). ^c Denotes substitution of product radical. ^d Based on information reported by Vereecken and Peeters (2004) for calculations for the given peroxy radical. ^e *E/R* for formation of a tertiary radical assumed to be 400 K lower than for formation of a secondary radical, corresponding to a difference in *E* of ≈ 3.3 kJ mol⁻¹. This is consistent with differences in energy barriers reported for formation of secondary and tertiary radicals (Vereecken and Peeters, 2004). ^f Based on the calculations of Vereecken and Peeters (2004) for a relevant tertiary peroxy radical formed during the oxidation of isoprene. ^g Based on the calculations of Vereecken and Peeters (2004) for a relevant tertiary peroxy radical formed during the oxidation of α-pinene. Applies specifically to *anti*-conformers, when the OH and peroxy radical groups on the opposite sides of the ring (as shown), which were calculated to account for 60 % of the *anti*- + *syn*-population (Vereecken and Peeters, 2004). ^h Based on the calculations of Vereecken and Peeters (2004) for a relevant tertiary peroxy radical formed during the oxidation of α-pinene. Applies specifically to *syn*-conformers, when the OH and peroxy radical groups on the same side of the ring (as shown), which were calculated to account for 40 % of the *anti*- + *syn*-population (Vereecken and Peeters, 2004). ⁱ Based on the calculations of Vereecken and Peeters (2004) for the a relevant tertiary peroxy radical, formed during the oxidation of β-pinene.

Nguyen (2012). Once again, the 1,6-enol H-shift reaction is likely to be the major fate of the majority of peroxy radicals containing the relevant substructure (see Table 15) under atmospheric conditions, but the rate is formalized in the present work to allow for the representation of competing rapid isomerization reactions for specific structurally complex peroxy radicals, e.g. the 1,4-hydroxyl H-shift reaction discussed above, or other reactions that may be considered and represented in future work.

As indicated above, the present paper does not attempt to provide a full treatment of unimolecular reactions of RO₂ radicals, which ideally require systematic information on the rates of a series of 1, *n*-H-shift reactions from C–H and O–H bonds in different environments. In this respect, it is noted that the systematic influence of a series of neighbouring functional groups and transition state sizes have been considered in theoretical studies of a number of model systems (e.g. Crouse et al., 2013; Jørgensen et al., 2016; Praske

Table 15. Representative rate coefficients for selected H-shift isomerization reactions of peroxy radicals.

Radical	Product(s)	$k(T)$ (s^{-1})	$k_{298\text{K}}$ (s^{-1})	Comment
1,4-formyl H-shift reaction				
		$3.0 \times 10^7 \exp(-5300/T)$	0.57	a,b
1,4-hydroxyl H-shift reaction				
		$3.6 \times 10^{12} \exp(-6310/T)$	2.3×10^3	b,c
		$6.7 \times 10^{12} \exp(-5780/T)$	2.5×10^4	b,d
		$5.6 \times 10^{12} \exp(-6010/T)$	9.8×10^3	b,e
1,5-hydroxyl H-shift reaction				
		$1.9 \times 10^{11} \exp(-9750/T)$	1.2×10^{-3}	b,f
		$1.0 \times 10^{11} \exp(-9750/T)$	6.2×10^{-4}	b,f
1,6-hydroxyalkyl H-shift reaction				
		$1.3 \times 10^{10} \exp(-8380/T) \times \exp(10^8/T^3)$	3.5×10^{-1}	g
1,6-enol H-shift reaction				
		$2.4 \times 10^{-1} T^{4.1} \exp(-2700/T)$	3.9×10^5	h

Comments: ^a Based on rate coefficient reported for the methacrolein-derived peroxy radical, $\text{HOCH}_2\text{C}(\text{CH}_3)(\text{O}_2)\text{C}(=\text{O})\text{H}$ by Crouse et al. (2012). Applied to primary, secondary and tertiary α -formyl peroxy radicals. ^b The initially formed hydroperoxy-substituted product radical decomposes spontaneously to produce the displayed products. ^c Based on the rate coefficient estimated for $\text{CH}_3\text{CH}(\text{OH})\text{O}_2$ by Hermans et al. (2005); applied to secondary α -hydroxyl peroxy radicals. ^d Based on the rate coefficient estimated for $(\text{CH}_3)_2\text{C}(\text{OH})\text{O}_2$ by Hermans et al. (2005); applied to tertiary α -hydroxy peroxy radicals. ^e Based on the rate coefficient estimated for *cyclo*- $\text{C}_6\text{H}_{10}(\text{OH})\text{O}_2$ by Hermans et al. (2005); applied generally to cyclic α -hydroxy peroxy radicals (i.e. where the OH and OO groups are substituents to a ring). ^f Based on rate coefficients reported by Peeters et al. (2014) for corresponding unsaturated secondary and tertiary β -hydroxy peroxy radicals formed in isoprene oxidation. Applied generally to unsaturated β -hydroxy peroxy radicals containing the substructures shown. ^g Based on the geometric mean of rate coefficients applied to (Z)- $\text{CH}(\text{OH})\text{C}(\text{CH}_3)=\text{CHCH}_2\text{O}_2$ (CISOPAO2) and (Z)- $\text{CH}(\text{OH})\text{CH}=\text{C}(\text{CH}_3)\text{CH}_2\text{O}_2$ (CISOPCO2) in MCM v3.3.1 (Jenkin et al., 2015) based on the calculations of Peeters et al. (2014) and observations of Crouse et al. (2011). Applied generally to unsaturated δ -hydroxy peroxy radicals containing the substructure shown, except for CISOPAO2 and CISOPCO2 themselves for which the species-specific rate coefficients are applied (see Table S5). Rapid reaction of the product radical with O_2 dominates over the reverse isomerization reaction under atmospheric conditions. ^h Based on the geometric mean of rate coefficients reported for (Z)- $\text{HOCH}=\text{C}(\text{CH}_3)\text{CH}(\text{O}_2)\text{CH}_2\text{OH}$ and (Z)- $\text{HOCH}=\text{CHC}(\text{CH}_3)(\text{O}_2)\text{CH}_2\text{OH}$ in the calculations of Peeters and Nguyen (2012). Applied to peroxy radicals containing the substructure shown. Rapid reaction of the product radical with O_2 dominates over the reverse isomerization reaction under atmospheric conditions.

et al., 2017; Otkjær et al., 2018). Such studies provide the basis for defining systematic structure–activity methods for a wide range of RO_2 radicals and their potential isomerization reactions, and are being considered in ongoing work. A further consideration, highlighted in those studies, is that the rates of the reverse isomerization reactions are sometimes sufficiently rapid that the product radical may not be fully trapped by onward reaction (e.g. addition of O_2) under atmo-

spheric conditions. It is noted that the explicit representation of a very large number of rapid reversible reactions in detailed mechanisms can have implications for computational efficiency, and needs to be considered carefully in method development and implementation.

4 Conclusions

Published kinetics and branching ratio data have been reviewed for the bimolecular reactions of organic peroxy radicals (RO_2), with information for selected unimolecular isomerization reactions also summarized and discussed. The information has been used to define generic rate coefficients and structure–activity relationship (SAR) methods for the reactions of a series of important classes of hydrocarbon and oxygenated RO_2 radicals, for application in the next generation of explicit detailed chemical mechanisms, based on GECKO-A and the MCM.

The availability of kinetic and mechanistic data for peroxy radical reactions has increased substantially since the appraisals of Saunders et al. (2003) and Aumont et al. (2005), on which the previous treatments of peroxy radical chemistry in the MCM and GECKO-A were mainly based. These advances have allowed improved and updated methods to be defined and summarized in the present work for an extended set of peroxy radical reactions. Nevertheless, there are still a number of specific areas (commented on in Sects. 2 and 3) where information is lacking and further studies would be beneficial. These include the following.

- Kinetics studies of the reactions with NO have only been reported for a limited number of acyl peroxy radicals. Further studies, particularly for larger and highly oxygenated acyl peroxy radicals, would help to establish whether size and/or the presence of additional substituent groups has an effect on reactivity.
- Further systematic data on RONO_2 yields from the reactions with NO are required to help improve branching ratio parameterizations. These include additional data for a variety of acyclic and cyclic oxygenated RO_2 as a function of size and structure.
- For the reactions with NO_3 , studies for $\geq \text{C}_2$ (non-acyl) RO_2 are dominated by primary peroxy radicals. Further studies are therefore required for secondary and tertiary radicals, and product information is generally required for a variety of peroxy radical classes to test assumption that the reaction proceeds via a single channel forming RO, NO_2 and O_2 .
- The reactions of $\geq \text{C}_2$ hydrocarbon RO_2 with OH are believed to produce a thermalized hydrotrioxide, ROOOH , as the major product. Detailed experimental and theoretical studies are therefore required to establish the atmospheric fate of these ROOOH species. Studies of the reactions of oxygenated RO_2 with OH are also required.
- The reactions of HO_2 with several oxygenated RO_2 classes have been shown to proceed via multiple channels, although the temperature dependences of the product channels have generally not been studied. Addi-

tional studies of their temperature dependences would therefore be valuable, in addition to information for larger sets of oxygenated RO_2 within some classes. Kinetics studies have only been reported for a limited number of acyl peroxy radicals. Further studies, particularly for larger and highly oxygenated acyl peroxy radicals, would help to establish whether size and/or the presence of additional substituent groups has an effect on reactivity.

- For the self- and cross-reactions of peroxy radicals, further information is required to allow the impacts of multiple substituents on the kinetics to be defined more rigorously. Further systematic studies of the formation of $\text{ROOR} + \text{O}_2$ (from the self-reaction of RO_2) and $\text{ROOR}' + \text{O}_2$ (from the cross-reaction of RO_2 with $\text{R}'\text{O}_2$) are also required as a function of peroxy radical size and functional group content.
- For unimolecular isomerization reactions, further systematic studies are required for the rates of 1, n -H-shift reactions from C–H and O–H bonds in different chemical environments, and for the effect of ring size and substituents on ring-closure reactions, to build upon recently reported data for these reaction classes.

Data availability. All relevant data have been tabulated or provided as supporting information in the Supplement.

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/acp-19-7691-2019-supplement>.

Author contributions. All authors defined the scope of the work. MEJ developed and revised the estimation methods and drafted the manuscript. The estimation methods were tested, evaluated and refined by all authors. The manuscript was reviewed and revised by all authors.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This work was performed as part of the MAGNIFY project, with funding from the UK Natural Environment Research Council (NERC) and the French National Research Agency (ANR). It was also partially funded by the European Commission through EUROCHAMP-2020. Paul Wennberg (California Institute of Technology), Luc Vereecken (Forschungszentrum Jülich), Geoff Tyndall (NCAR, Boulder), John Crouse (California Institute of Technology) and an anonymous referee are gratefully acknowledged for review and discussion comments that helped to improve the paper.

Financial support. This research has been supported by the UK Natural Environment Research Council (NERC) (grant no. NE/M013448/1), the French National Research Agency (ANR) (grant no. ANR-14-CE01-0010), and the European Commission through EUROCHAMP-2020 (grant no. 730997).

Review statement. This paper was edited by Dwayne Heard and reviewed by Paul O. Wennberg, Geoffrey Tyndall, and one anonymous referee.

References

- Anglada, J. M. and Solé, A.: Tropospheric oxidation of methyl hydrotrioxide (CH_3OOOH) by hydroxyl radical, *Phys. Chem. Chem. Phys.*, 20, 27406–27417, 2018.
- Archibald, A. T., Petit, A. S., Percival, C. J., Harvey, J. N., and Shallcross, D. E.: On the importance of the reaction between OH and RO_2 radicals, *Atmos. Sci. Lett.*, 10, 102–108, 2009.
- Arey, J., Aschmann, S. M., Kwok, E. S. C., and Atkinson, R.: Alkyl nitrate, hydroxyalkyl nitrate, and hydroxycarbonyl formation from the NO_x -air photooxidations of C_5 – C_8 n-alkanes, *J. Phys. Chem. A*, 105, 1020–1027, <https://doi.org/10.1021/jp003292z>, 2001.
- Aschmann, S. M., Tuazon, E. C., Arey, J., and Atkinson, R.: Products and mechanisms of the gas-phase reactions of OH radicals with 1-octene and 7-tetradecene in the presence of NO, *Environ. Sci. Technol.*, 44, 3825–3831, 2010.
- Assaf, E., Song, B., Tomas, A., Schoemaeker, C., and Fittschen, C.: Rate constant of the reaction between CH_3O_2 radicals and OH radicals revisited, *J. Phys. Chem. A*, 120, 8923–8932, 2016.
- Assaf, E., Sheps, L., Whalley, L., Heard, D., Tomas, A., Schoemaeker, C., and Fittschen, C.: The reaction between CH_3O_2 and OH radicals: product yields and atmospheric implications, *Environ. Sci. Technol.*, 51, 2170–2177, 2017a.
- Assaf, E., Tanaka, S., Kajii, Y., Schoemaeker, C., and Fittschen, C.: Rate constants of the reaction of C_2 – C_4 peroxy radicals with OH radicals, *Chem. Phys. Lett.*, 684, 245–249, 2017b.
- Assaf, E., Schoemaeker, C., Vereecken, L., and Fittschen, C.: Experimental and theoretical investigation of the reaction of RO_2 radicals with OH radicals: Dependence of the HO_2 yield on the size of the alkyl group, *Int. J. Chem. Kinet.*, 50, 670–680, 2018.
- Aumont, B., Szopa, S., and Madronich, S.: Modelling the evolution of organic carbon during its gas-phase tropospheric oxidation: development of an explicit model based on a self-generating approach, *Atmos. Chem. Phys.*, 5, 2497–2517, <https://doi.org/10.5194/acp-5-2497-2005>, 2005.
- Berndt, T., Richters, S., Kaethner, R., Voigtländer, J., Stratmann, F., Sipilä, M., Kulmala, M. and Herrmann, H.: Gas-phase ozonolysis of cycloalkenes: formation of highly oxidized RO_2 radicals and their reactions with NO, NO_2 , SO_2 , and other RO_2 radicals, *J. Phys. Chem. A*, 119, 10336–10348, 2015.
- Berndt, T., Scholz, W., Mentler, B., Fischer, L., Herrmann, H., Kulmala, M., and Hansel, A.: Accretion product formation from self- and cross-reactions of RO_2 radicals in the atmosphere, *Angew. Chem. Int. Edit.*, 57, 3820–3824, 2018a.
- Berndt, T., Mentler, B., Scholz, W., Fischer, L., Herrmann, H., Kulmala, M. and Hansel, A.: Accretion product formation from ozonolysis and OH radical reaction of α -pinene: mechanistic insight and the influence of isoprene and ethylene, *Environ. Sci. Technol.*, 52, 11069–11077, 2018b.
- Bian, H., Zhang, S. and Zhang, H.: Theoretical study on the atmospheric reaction of CH_3O_2 with OH, *Int. J. Quantum. Chem.*, 115, 1181–1186, 2015.
- Bianchi, F., Kurtén, T., Riva, M., Mohr, C., Rissanen, M. P., Roldin, P., Berndt, T., Crouse, J. D., Wennberg, P. O., Mentel, T. F., Wildt, J., Junninen, H., Jokinen, T., Kulmala, M., Worsnop, D. R., Thornton, J. A., Donahue, N., Kjaergaard, H. G., and Ehn, M.: Highly oxygenated organic molecules (HOM) from gas-phase autoxidation involving peroxy radicals: A key contributor to atmospheric aerosol, *Chem. Rev.*, 119, 3472–3509, 2019.
- Birdsall, A. W. and Elrod, M. J.: Comprehensive NO-dependent study of the products of the oxidation of atmospherically relevant aromatic compounds, *J. Phys. Chem. A*, 115, 5397–5407, 2011.
- Birdsall, A. W., Andreoni, J. F., and Elrod, M. J.: Investigation of the role of bicyclic peroxy radicals in the oxidation mechanism of toluene, *J. Phys. Chem. A*, 114, 10655–10663, 2010.
- Bey, I., Aumont, B., and Toupance, G.: A modeling study of the nighttime radical chemistry in the lower continental troposphere: 1 – Development of a detailed chemical mechanism including nighttime chemistry, *J. Geophys. Res.*, 106, 9959–9990, 2001a.
- Bey, I., Aumont, B. and Toupance, G.: A modeling study of the nighttime radical chemistry in the lower continental troposphere: 2 – Origin and evolution of HO_x , *J. Geophys. Res.*, 106, 9990–10001, 2001b.
- Biggs, P., Canosa-Mas, C., Fracheboud, J. M., Shallcross, D. E., and Wayne, R. P.: Rate constants for the reactions of C_2H_5 , $\text{C}_2\text{H}_5\text{O}$ and $\text{C}_2\text{H}_5\text{O}_2$ radicals with NO_3 at 298 K and 2.2 Torr, *J. Chem. Soc. Faraday T.*, 91, 817–825, 1995.
- Bossolasco, A. G., Malanca, F. E., and Argüello, G. A.: Peroxy ethoxyformyl nitrate, $\text{CH}_3\text{CH}_2\text{OC}(\text{O})\text{OONO}_2$. Spectroscopic and thermal characterization, *J. Photochem. Photobiol. A*, 221, 58–63, 2011.
- Boyd, A. A. and Lesclaux, R.: The temperature dependence of the rate coefficients for β -hydroxyperoxy radical self reactions, *Int. J. Chem. Kinet.*, 29, 323–331, 1997.
- Boyd, A. A., Nozière, B., and Lesclaux, R.: Kinetic studies of the allylperoxy radical self-reaction and reaction with HO_2 , *Chem. Soc. Faraday T.*, 92, 201–206, 1996a.
- Boyd, A. A., Lesclaux, R., Jenkin, M. E., and Wallington, T. J.: A spectroscopic, kinetic, and product study of the $(\text{CH}_3)_2\text{C}(\text{OH})\text{CH}_2\text{O}_2$ radical self reaction and reaction with HO_2 , *J. Phys. Chem.*, 100, 6594–6603, 1996b.
- Boyd, A. A., Villenave, E., and Lesclaux, R.: Structure-reactivity relationships for the self-reactions of linear secondary alkylperoxy radicals: an experimental investigation, *Int. J. Chem. Kinet.*, 31, 37–46, 1999.
- Boyd, A. A., Flaud, P.-M., Daugey, N., and Lesclaux, R.: Rate constants for $\text{RO}_2 + \text{HO}_2$ reactions measured under a large excess of HO_2 , *J. Phys. Chem. A*, 107, 818–821, 2003a.
- Boyd, A. A., Villenave, E., and Lesclaux, R.: Self- and cross-reactions of β -hydroxyperoxy radicals of relevance to tropospheric monoterpene oxidation: structure-activity relationships for rate coefficients, *Atmos. Environ.*, 37, 2751–2760, 2003b.
- Bridier, I., Veyret, B., Lesclaux, R., and Jenkin, M. E.: Flash-photolysis study of the uv spectrum and kinetics of reactions of

- the acetylperoxy radical, *J. Chem. Soc. Faraday T.*, 89, 2993–2997, 1993.
- Calvert, J. G., Orlando, J. J., Stockwell, W. R., and Wallington, T. J.: The mechanisms of reactions influencing atmospheric ozone, Oxford University Press, Oxford, ISBN 978-0-19-023301-0, 2015.
- Canosa-Mas, C. E., King, M. D., Lopez, R., Percival, C. J., Wayne, R. P., Shallcross, D. E., Pyle, J. A., and Daele, V.: Is the reaction between $\text{CH}_3\text{C}(\text{O})\text{O}_2$ and NO_3 important in the night-time troposphere?, *J. Chem. Soc. Faraday T.*, 92, 2211–2222, 1996.
- Caralp, F., Forst, W., and Rayez, M. T.: Chemical activation in OH radical-oxidation of 1-n-alkenes, *Phys. Chem. Chem. Phys.*, 5, 476–486, 2003.
- Caravan, R. L., Khan, M. A. H., Zádor, J., Sheps, L., Antonov, I. O., Rotavera, B., Ramasesha, K., Kendrew, A., Chen, M., Rösch, D., Osborn, D. L., Fittschen, C., Schoemaeker, C., Duncianu, M., Grira, A., Dusanter, S., Tomas, A., Percival, C., Shallcross, D. E., and Taatjes, C. A.: The reaction of hydroxyl and methylperoxy radicals is not a major source of atmospheric methanol, *Nat. Commun.*, 9, 4343, <https://doi.org/10.1038/s41467-018-06716-x>, 2018.
- Carslaw, N., Carpenter, L. J., Plane, J. M. C., Allan, B. J., Burgess, R. A., Clemitshaw, K. C., Coe, H., and Penkett, S. A.: Simultaneous measurements of nitrate and peroxy radicals in the marine boundary layer, *J. Geophys. Res.*, 102, 18917–18933, 1997.
- Carter, W. P. L. and Atkinson, R.: Alkyl nitrate formation from the atmospheric photooxidation of alkanes; a revised estimation method, *J. Atmos. Chem.*, 8, 165–173, 1989.
- Cassanelli, P., Fox, D. J., and Cox, R. A.: Temperature dependence of pentyl nitrate formation from the reaction of pentyl peroxy radicals with NO, *Phys. Chem. Chem. Phys.*, 9, 4332–4337, 2007.
- Cavalli, F., Barnes, I., and Becker, K.-H.: FT-IR kinetic and product study of the OH radical and Cl-atom-initiated oxidation of dibasic esters, *Int. J. Chem. Kinet.*, 33, 431–439, 2001.
- Christensen, L. K., Ball, J. C., and Wallington, T. J.: Atmospheric oxidation mechanism of methyl acetate, *J. Phys. Chem. A*, 104, 345–351, 2000.
- Crouse, J. D., Paulot, F., Kjaergaard, H. G., and Wennberg, P. O.: Peroxy radical isomerization in the oxidation of isoprene, *Phys. Chem. Chem. Phys.*, 13, 13607–13613, 2011.
- Crouse, J. D., Knap, H. C., Ørnsø, K. B., Jørgensen, S., Paulot, F., Kjaergaard, H. G., and Wennberg, P. O.: Atmospheric fate of methacrolein. 1. Peroxy radical isomerization following addition of OH and O_2 , *J. Phys. Chem. A*, 116, 5756–5762, 2012.
- Crouse, J. D., Nielsen, L. B., Jørgensen, S., Kjaergaard, H. G., and Wennberg, P. O.: Autoxidation of organic compounds in the atmosphere, *J. Phys. Chem. Lett.*, 4, 3513–3520, 2013.
- Crouse, J. D., Teng, A., and Wennberg, P. O.: Experimental constraints on the distribution and fate of peroxy radicals formed in reactions of isoprene + OH + O_2 , presented at Atmospheric Chemical Mechanisms: Simple Models – Real World Complexities, University of California, Davis, USA, 10–12 December 2014.
- Da Silva, G., Graham, C., and Wang, Z.-F.: Unimolecular β -hydroxyperoxy radical decomposition with OH recycling in the photochemical oxidation of isoprene, *Environ. Sci. Technol.*, 44, 250–256, 2010.
- De Gouw, J. A. and Howard, C. J.: Direct measurement of the rate coefficient for the $\text{CH}_2\text{C}(\text{CH}_3)\text{C}(\text{O})\text{O}_2 + \text{NO}$ reaction using chemical ionization mass spectrometry, *J. Phys. Chem. A*, 101, 8662–8667, 1997.
- Dillon, T. J. and Crowley, J. N.: Direct detection of OH formation in the reactions of HO_2 with $\text{CH}_3\text{C}(\text{O})\text{O}_2$ and other substituted peroxy radicals, *Atmos. Chem. Phys.*, 8, 4877–4889, <https://doi.org/10.5194/acp-8-4877-2008>, 2008.
- Doussin, J. F., Picquet-Varrault, B., Durand-Jolibois, R., Loirat, H. and Carlier, P.: A visible and FTIR spectrometric study of the nighttime chemistry of acetaldehyde and PAN under simulated atmospheric conditions, *J. Photochem. Photobiol. A*, 157, 283–293, 2003.
- Eberhard, J. and Howard, C. J.: Temperature-dependent kinetics studies of the reactions of $\text{C}_2\text{H}_5\text{O}_2$ and $n\text{-C}_3\text{H}_7\text{O}_2$ radicals with NO, *Int. J. Chem. Kinet.*, 28, 731–740, 1996.
- Eberhard, J. and Howard, C. J.: Rate coefficients for the reactions of some C_3 to C_5 hydrocarbon peroxy radicals with NO, *J. Phys. Chem. A*, 101, 3360–3366, 1997.
- Eberhard, J., Villalta, P. W., and Howard, C. J.: Reaction of isopropyl peroxy radicals with NO over the temperature range 201–401 K, *J. Phys. Chem.*, 100, 993–997, 1996.
- Ehn, M., Thornton, J. A., Kleist, E., Sipilä, M., Junninen, H., Pullinen, I., Springer, M., Rubach, F., Tillmann, R., Lee, B., Lopez-Hilfiker, F., Andres, S., Acir, I.-H., Rissanen, M., Jokinen, T., Schobesberger, S., Kangasluoma, J., Kontkanen, J., Nieminen, T., Kurtén, T., Nielsen, L. B., Jørgensen, S., Kjaergaard, H. G., Canagaratna, M., Maso, M. D., Berndt, T., Petäjä, T., Wahner, A., Kerminen, V.-M., Kulmala, M., Worsnop, D. R., Wildt, J., and Mentel, T. F.: A large source of low-volatility secondary organic aerosol, *Nature*, 506, 476–479, <https://doi.org/10.1038/nature13032>, 2014.
- Ehn, M., Berndt, T., Wildt, J., and Mentel, T.: Highly oxygenated molecules from atmospheric autoxidation of hydrocarbons: a prominent challenge for chemical kinetics studies, *Int. J. Chem. Kinet.*, 49, 821–831, 2017.
- Elrod, M. J.: Kinetics study of the aromatic bicyclic peroxy radical + NO Reaction: overall rate constant and nitrate product yield measurements, *J. Phys. Chem. A*, 115, 8125–8130, 2011.
- Farágó, E. P., Schoemaeker, C., Viskolcz, B., and Fittschen, C.: Experimental determination of the rate constant of the reaction between $\text{C}_2\text{H}_5\text{O}_2$ and OH radicals, *Chem. Phys. Lett.*, 619, 196–200, 2015.
- Geyer, A., Bächmann, K., Hofzumahaus, A., Holland, F., Konrad, S., Klüpfel, T., Pätz, H.W., Perner, D., Mihelcic, D., Schäfer, H. J., Volz-Thomas, A., and Platt, U.: Nighttime formation of peroxy and hydroxyl radicals during the BERLIOZ campaign: Observations and modeling studies, *J. Geophys. Res.*, 108, 8249, <https://doi.org/10.1029/2001JD000656>, 2003.
- Glover, B. G. and Miller, T. J.: Near-IR cavity ringdown spectroscopy and kinetics of the isomers and conformers of the butyl peroxy radical, *J. Phys. Chem. A*, 109, 11191–11197, 2005.
- Glowacki, D. R., Wang, L., and Pilling, M. J.: Evidence of formation of bicyclic species in the early stages of atmospheric benzene oxidation, *J. Phys. Chem. A*, 113, 5385–5396, 2009.
- Groß, C. B. M., Dillon, T. J., Schuster, G., Lelieveld, J., and Crowley, J. N.: Direct kinetic study of OH and O_3 formation in the reaction of $\text{CH}_3\text{C}(\text{O})\text{O}_2$ with HO_2 , *J. Phys. Chem. A*, 118, 974–985, <https://doi.org/10.1021/jp412380z>, 2014.

- Hansen, J. C., Li, Y., Rosado-Reyes, C. M., Francisco, J. S., Szente, F. J., and Maricq, M. M.: Theoretical and experimental investigation of the UV cross section and kinetics of the methyl formate peroxy radical, *J. Phys. Chem. A*, 107, 5306–5316, 2003.
- Hasson, A. S., Tyndall, G. S., and Orlando, J. J.: A product yield study of the reaction of HO₂ Radicals with ethyl peroxy (CH₃CH₂O₂), acetyl peroxy (CH₃C(O)O₂), and acetonyl peroxy (CH₃C(O)CH₂O₂) radicals, *J. Phys. Chem. A*, 108, 5979–5989, 2004.
- Hasson, A. S., Tyndall, G. S., Orlando, J. J., Singh, S., Hernandez, S. Q., Campbell, S., and Ibarra, Y.: Branching ratios for the reaction of selected carbonyl-containing peroxy radicals with hydroperoxy radicals, *J. Phys. Chem.*, 116, 6264–6281, 2012.
- Hermans, I., Muller, J. F., Nguyen, T. L., Jacobs, P. A., and Peeters, J.: Kinetics of α -hydroxy-alkylperoxyl radicals in oxidation processes. HO₂-initiated oxidation of ketones/aldehydes near the tropopause, *J. Phys. Chem. A*, 109, 4303–4311, 2005.
- Horie, O. and Moortgat, G. K.: Reactions of CH₃C(O)O₂ radicals with CH₃O₂ and HO₂ between 263 K and 333 K, *J. Chem. Soc. Faraday T.*, 88, 3305–3312, 1992.
- Hsin, H. Y. and Elrod, M. J.: Overall rate constant measurements of the reaction of hydroxy- and chloroalkylperoxyradicals derived from methacrolein and methyl vinyl ketone with nitric oxide, *J. Phys. Chem. A*, 111, 613–619, 2007.
- Hui, A. O., Fradet, M., Okumura, M., and Sander, S. P.: Temperature dependence study of the kinetics and product yields of the HO₂ + CH₃C(O)O₂ reaction by direct detection of OH and HO₂ radicals using 2f-IR wavelength modulation spectroscopy, *J. Phys. Chem. A*, 123, 3655–3671, <https://doi.org/10.1021/acs.jpca.9b00442>, 2019.
- Jagiella, S. and Zabel, F.: Reaction of phenylperoxy radicals with NO₂ at 298 K, *Phys. Chem. Chem. Phys.*, 9, 5036–5051, 2007.
- Jagiella, S. and Zabel, F.: Thermal stability of carbonyl radicals. Part II. Reactions of methylglyoxyl and methylglyoxylperoxy radicals at 1 bar in the temperature range 275–311 K, *Phys. Chem. Chem. Phys.*, 10, 1799–1808, 2008.
- Jenkin, M. E. and Clemitshaw, K. C.: Ozone and other secondary photochemical pollutants: chemical processes governing their formation in the planetary boundary layer, *Atmos. Environ.*, 34, 2499–2527, 2000.
- Jenkin, M. E. and Hayman, G. D.: Kinetics of reactions of primary, secondary and tertiary β -hydroxy peroxy radicals: application to isoprene degradation, *J. Chem. Soc. Faraday T.*, 91, 433–446, 1995.
- Jenkin, M. E., Hayman, G. D., Wallington, T. J., Hurley, M. D., Ball, J. C., Nielsen, O. J., and Ellerman, T.: Kinetic and mechanistic study of the self reaction of CH₃OCH₂O₂ radicals at room temperature, *J. Phys. Chem.*, 97, 11712–11723, 1993a.
- Jenkin, M. E., Murrells, T. P., Shalliker, S. J., and Hayman, G. D.: Kinetics and product study of the self-reactions of allyl and allyl peroxy radicals at 296 K, *J. Chem. Soc. Faraday T.*, 89, 433–446, 1993b.
- Jenkin, M. E., Saunders, S. M., and Pilling M. J.: The tropospheric degradation of volatile organic compounds: a protocol for mechanism development, *Atmos. Environ.*, 31, 81–104, 1997.
- Jenkin, M. E., Boyd, A. A., and Lesclaux, R.: Peroxy radical kinetics resulting from the OH-initiated oxidation of 1,3-butadiene, 2,3-dimethyl-1,3-butadiene and isoprene, *J. Atmos. Chem.*, 29, 267–298, 1998.
- Jenkin, M. E., Sulbaek Andersen, M. P., Hurley, M. D., Wallington, T. J., Taketani, F., and Matsumi, Y.: A kinetics and mechanistic study of the OH and NO₂ initiated oxidation of cyclohexa-1,3-diene in the gas phase, *Phys. Chem. Chem. Phys.*, 7, 1194–1204, 2005.
- Jenkin, M. E., Hurley, M. D., and Wallington, T. J.: Investigation of the radical product channel of the CH₃C(O)O₂ + HO₂ reaction in the gas phase, *Phys. Chem. Chem. Phys.*, 9, 3149–3162, 2007.
- Jenkin, M. E., Hurley, M. D., and Wallington, T. J.: Investigation of the radical product channel of the CH₃C(O)CH₂O₂ + HO₂ reaction in the gas phase, *Phys. Chem. Chem. Phys.*, 10, 4274–4280, 2008.
- Jenkin, M. E., Hurley, M. D., and Wallington, T. J.: Investigation of the radical product channel of the CH₃OCH₂O₂ + HO₂ reaction in the gas phase, *J. Phys. Chem.*, 114, 408–416, 2010.
- Jenkin, M. E., Young, J. C., and Rickard, A. R.: The MCM v3.3.1 degradation scheme for isoprene, *Atmos. Chem. Phys.*, 15, 11433–11459, <https://doi.org/10.5194/acp-15-11433-2015>, 2015.
- Jenkin, M. E., Valorso, R., Aumont, B., Rickard, A. R., and Wallington, T. J.: Estimation of rate coefficients and branching ratios for gas-phase reactions of OH with aliphatic organic compounds for use in automated mechanism construction, *Atmos. Chem. Phys.*, 18, 9297–9328, <https://doi.org/10.5194/acp-18-9297-2018>, 2018a.
- Jenkin, M. E., Valorso, R., Aumont, B., Rickard, A. R., and Wallington, T. J.: Estimation of rate coefficients and branching ratios for gas-phase reactions of OH with aromatic organic compounds for use in automated mechanism construction, *Atmos. Chem. Phys.*, 18, 9329–9349, <https://doi.org/10.5194/acp-18-9329-2018>, 2018b.
- Jokinen, T., Sipilä, M., Richters, S., Kerminen, V.-M., Paasonen, P., Stratmann, F., Worsnop, D., Kulmala, M., Ehn, M., Herrmann, H. and Berndt, T.: Rapid autoxidation forms highly oxidized RO₂ radicals in the atmosphere, *Angew. Chem. Int. Ed.*, 53, 14596–14600, <https://doi.org/10.1002/anie.201408566>, 2014.
- Jørgensen, S., Knap, H. C., Otkjær, R. V., Jensen, A. M., Kjeldsen, M. L. H., Wennberg, P. O., and Kjaergaard, H. G.: Rapid hydrogen shift scrambling in hydroperoxy-substituted organic peroxy radicals, *J. Phys. Chem. A*, 120, 266–275, <https://doi.org/10.1021/acs.jpca.5b06768>, 2016.
- Kabir, M., Jagiella, S., and Zabel, F.: Thermal stability of n-acyl peroxy nitrates, *Int. J. Chem. Kinet.*, 46, 462–469, 2014.
- Kalalian, C., Roth, E., and Chakir, A.: Atmospheric reactivity of nitrate radicals: reaction with peroxy radicals, *Atmos. Environ.*, 190, 308–316, 2018.
- Kirchner, K., Mayer-Figge, A., Zabel, F., and Becker, K.-H.: Thermal stability of peroxy nitrates, *Int. J. Chem. Kinet.*, 31, 127–144, 1999.
- Kukui, A. S., Jungkamp, T. P. W., and Schindler, R. N.: Aldehyde formation in the reaction of methoxy radicals with NO₃, *Ber. Bunsen. Phys. Chem.*, 99, 1565–1567, 1995.
- Laversin, H., Cousin, J., Joly, L., Roth, E., Durry, G., and Chakir, A.: Kinetic study of the reaction of nitrate radicals with ethylperoxy radicals between 277 and 358 K, *Chem. Phys. Lett.*, 644, 14–19, 2016.
- Le Crâne, J. P. and Lesclaux, R.: UV absorption spectra and self-reaction rate constants for primary peroxy radicals arising from

- the chlorine-initiated oxidation of carbonyl compounds, *Int. J. Chem. Kinet.*, 38, 276–283, 2006.
- Le Crâne, J. P., Villenave, E., Hurley, M. D., Wallington, T. J., Nishida, S., Takahashi, K., and Matsumi, Y.: Atmospheric chemistry of pivalaldehyde and isobutyraldehyde: kinetics and mechanisms of reactions with Cl atoms, fate of $(\text{CH}_3)_3\text{CC}(\text{O})$ and $(\text{CH}_3)_2\text{CHC}(\text{O})$ radicals, and self-reaction kinetics of $(\text{CH}_3)_3\text{CC}(\text{O})\text{O}_2$ and $(\text{CH}_3)_2\text{CHC}(\text{O})\text{O}_2$ radicals, *J. Phys. Chem. A*, 108, 795–805, 2004.
- Lee, L., Teng, A. P., Wennberg, P. O., Crouse, J. D., and Cohen, R. C.: On rates and mechanisms of OH and O_3 reactions with isoprene-derived hydroxy nitrates, *J. Phys. Chem. A*, 118, 1622–1637, 2014.
- Lightfoot, P. D., Cox, R. A., Crowley, J. N., Destriau, M., Hayman, G. D., Jenkin, M. E., Moortgat, G. K., and Zabel, F.: Organic peroxy radicals: kinetics, spectroscopy and tropospheric chemistry, *Atmos. Environ.*, 26A, 1805–1964, 1992.
- Liu, Y. J., Herdinger-Blatt, I., McKinney, K. A., and Martin, S. T.: Production of methyl vinyl ketone and methacrolein via the hydroperoxyl pathway of isoprene oxidation, *Atmos. Chem. Phys.*, 13, 5715–5730, <https://doi.org/10.5194/acp-13-5715-2013>, 2013.
- Madronich, S. and Calvert, J. G.: Permutation reactions of organic peroxy radicals in the troposphere, *J. Geophys. Res.*, 95, 5697–5715, 1990.
- Matsunaga, A. and Ziemann, P. J.: Yields of beta-hydroxynitrates and dihydroxynitrates in aerosol formed from OH radical initiated reactions of linear alkenes in the presence of NO_x , *J. Phys. Chem. A*, 113, 599–606, <https://doi.org/10.1021/jp807764d>, 2009.
- Matsunaga, A. and Ziemann, P. J.: Yields of beta-hydroxynitrates, dihydroxynitrates, and trihydroxynitrates formed from OH radical-initiated reactions of 2-methyl-1-alkenes, *P. Natl. Acad. Sci. USA*, 107, 6664–6669, <https://doi.org/10.1073/pnas.0910585107>, 2010.
- McFiggans, G., Mentel, T. F., Wildt, J., Pullinen, I., Kang, S., Kleist, E., Schmitt, S., Springer, M., Tillmann, R., Wu, C., Zhao, D., Hallquist, M., Faxon, C., Le Breton, M., Hallquist, Å. M., Simpson, D., Bergström, R., Jenkin, M. E., Ehn, M., Thornton, J. A., Alfara, M. R., Bannan, T. J., Percival, C. J., Priestley, M., Topping, D., and Kiendler-Scharr, A.: Secondary organic aerosol reduced by mixture of atmospheric vapours, *Nature*, 565, 587–593, 2019.
- McKee, K., Blitz, M. A., and Pilling, M. J.: Temperature and pressure studies of the reactions of CH_3O_2 , HO_2 and 1,2- $\text{C}_4\text{H}_9\text{O}_2$ with NO_2 , *J. Phys. Chem. A*, 120, 1408–1420, 2016.
- Mentel, T. F., Springer, M., Ehn, M., Kleist, E., Pullinen, I., Kurtén, T., Rissanen, M., Wahner, A., and Wildt, J.: Formation of highly oxidized multifunctional compounds: autoxidation of peroxy radicals formed in the ozonolysis of alkenes – deduced from structure-product relationships, *Atmos. Chem. Phys.*, 15, 6745–6765, <https://doi.org/10.5194/acp-15-6745-2015>, 2015.
- Miller, A. M., Yeung, L. Y., Kiep, A. C., and Elrod, M. J.: Over-all rate constant measurements of the reactions of alkene-derived hydroxyalkylperoxy radicals with nitric oxide, *Phys. Chem. Chem. Phys.*, 6, 3402–3407, 2004.
- Mohammed, S. Y., Davis, A. A., Al Rashidi, M. J., and Sarathy, S. M.: High-pressure limit rate rules for α -H isomerization of hydroperoxyalkylperoxy radicals, *J. Phys. Chem. A*, 122, 3626–3639, <https://doi.org/10.1021/acs.jpca.7b11955>, 2018.
- Møller, K. H., Bates, K. H., and Kjaergaard, H. G.: The importance of peroxy radical hydrogen-shift reactions in atmospheric isoprene oxidation, *J. Phys. Chem. A*, 123, 920–932, <https://doi.org/10.1021/acs.jpca.8b10432>, 2019.
- Müller, J. F., Liu, Z., Nguyen, V. S., Stavrou, T., Harvey, J. N., and Peeters, J.: The reaction of methyl peroxy and hydroxyl radicals as a major source of atmospheric methanol, *Nat. Commun.*, 7, 13213, <https://doi.org/10.1038/ncomms13213>, 2016.
- Navarro, N. A., Dusanter, S., and Stevens, P. S.: Temperature dependence of the yields of methacrolein and methyl vinyl ketone from the OH-initiated oxidation of isoprene under NO_x -free conditions, *Atmos. Environ.*, 79, 59–66, 2013.
- Ng, N. L., Kwan, A. J., Surratt, J. D., Chan, A. W. H., Chhabra, P. S., Sorooshian, A., Pye, H. O. T., Crouse, J. D., Wennberg, P. O., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol (SOA) formation from reaction of isoprene with nitrate radicals (NO_3), *Atmos. Chem. Phys.*, 8, 4117–4140, <https://doi.org/10.5194/acp-8-4117-2008>, 2008.
- Nguyen, T. L., Vereecken, L., and Peeters, J.: Theoretical study of the $\text{HOCH}_2\text{OO} + \text{HO}_2$ reaction: detailed molecular mechanisms of the three reaction channels, *Z. Phys. Chem.*, 224, 1081–1093, 2010.
- Niki, H., Maker, P. D., Savage, C. M., and Breitenbach, L. P.: FTIR study of the kinetics and mechanism for Cl-atom-initiated reactions of acetaldehyde, *J. Phys. Chem.*, 89, 588–591, 1985.
- Nozière, B. and Hanson, D.: Speciated monitoring of gas-phase organic peroxy radicals by chemical ionization mass spectrometry: Cross-reactions between CH_3O_2 , $\text{CH}_3(\text{CO})\text{O}_2$, $(\text{CH}_3)_3\text{CO}_2$, and $c\text{-C}_6\text{H}_{11}\text{O}_2$, *J. Phys. Chem. A*, 121, 8453–8464, 2017.
- O'Brien, J. M., Czuba, E., Hastie, D. R., Francisco, J. S., and Shepson, P. B.: Determination of the hydroxy nitrate yields from the reaction of $\text{C}_2\text{--C}_6$ alkenes with OH in the presence of NO, *J. Phys. Chem. A*, 102, 8903–8908, <https://doi.org/10.1021/jp982320z>, 1998.
- Olivella, S., Solé, A., and Bofill, J. M.: Theoretical mechanistic study of the oxidative degradation of benzene in the troposphere: reaction of benzene–HO radical adduct with O_2 , *J. Chem. Theory Comput.*, 5, 1607–1623, 2009.
- Orlando, J. J. and Tyndall, G. S.: The atmospheric chemistry of the $\text{HC}(\text{O})\text{CO}$ radical, *Int. J. Chem. Kinet.*, 33, 149–156, 2001.
- Orlando, J. J. and Tyndall, G. S.: Laboratory studies of organic peroxy radical chemistry: an overview with emphasis on recent issues of atmospheric significance, *Chem. Soc. Rev.*, 41, 6294–6317, 2012.
- Orlando, J. J., Tyndall, G. S., Bilde, M., Ferronato, C., Wallington, T. J., Vereecken, L., and Peeters, J.: Laboratory and theoretical study of the oxy radicals in the OH- and Cl-initiated oxidation of ethene, *J. Phys. Chem. A*, 102, 8116–8123, 1998.
- Orlando, J. J., Tyndall, G. S., Vereecken, L., and Peeters, J.: The atmospheric chemistry of the acetonoxyl radical, *J. Phys. Chem. A*, 104, 11578–11588, 2000a.
- Orlando, J. J., Nozière, B., Tyndall, G. S., and Orzechowska, G. E.: Paulson, S. E., and Rudich, Y.: Product studies of the OH and ozone-initiated oxidation of some monoterpenes, *J. Geophys. Res.*, 105, 11561–11572, 2000b.

- Orlando, J. J., Tyndall, G. S., and Wallington, T. J.: The atmospheric chemistry of alkoxy radicals, *Chem. Rev.*, 103, 4657–4689, 2003.
- Otkjær, R. V., Jakobsen, H. H., Tram, C. M., and Kjaergaard, H. G.: Calculated hydrogen shift rate constants in substituted alkyl peroxy radicals, *J. Phys. Chem. A*, 122, 8665–8673, <https://doi.org/10.1021/acs.jpca.8b06223>, 2018.
- Paulot, F., Crouse, J. D., Kjaergaard, H. G., Kurten, A., St Clair, J. M., Seinfeld, J. H., and Wennberg, P. O.: Unexpected epoxide formation in the gas-phase photooxidation of isoprene, *Science*, 325, 730–733, <https://doi.org/10.1126/science.1172910>, 2009.
- Peeters, J. and Nguyen, T. L.: Unusually fast 1,6-H shifts of enolic hydrogens in peroxy radicals: formation of the first-generation C₂ and C₃ carbonyls in the oxidation of isoprene, *J. Phys. Chem. A*, 116, 6134–6141, 2012.
- Peeters, J., Nguyen, T. L., and Vereecken, L.: HO_x radical regeneration in the oxidation of isoprene, *Phys. Chem. Chem. Phys.*, 28, 5935–5939, 2009.
- Peeters, J., Müller, J.-F., Stavrou, T., and Nguyen, V. S.: Hydroxyl radical recycling in isoprene oxidation driven by hydrogen bonding and hydrogen tunneling: the upgraded LIM1 mechanism, *J. Phys. Chem. A*, 118, 8625–8643, 2014.
- Perring, A. E., Pusede, S. E., and Cohen, R. C.: An observational perspective on the atmospheric impacts of alkyl and multifunctional nitrates on ozone and secondary organic aerosol, *Chem. Rev.*, 113, 5848–5870, 2013.
- Picquet-Varrault, B., Doussin, J.-F., Durand-Jolibois, R., and Carlier, P.: FTIR spectroscopic study of the OH-induced oxidation of two linear acetates: ethyl and n-propyl acetates, *Phys. Chem. Chem. Phys.*, 3, 2595–2606, 2001.
- Picquet-Varrault, B., Doussin, J.-F., Durand-Jolibois, R. and Carlier, P.: FTIR spectroscopic study of the OH-induced oxidation of isopropyl, isobutyl, and tert-butyl acetates, *J. Phys. Chem. A*, 106, 2895–2902, 2002.
- Pimentel, A. S., Tyndall, G. S., Orlando, J. J., Hurley, M. D., Wallington, T. J., Sulbaek Andersen, M. P., Marshall, P., and Dibble, T. S.: Atmospheric chemistry of isopropyl formate and tert-butyl formate, *Int. J. Chem. Kinet.*, 42, 479–498, 2010.
- Praske, E., Crouse, J. D., Bates, K. H., Kurtén, T., Kjaergaard, H. G., and Wennberg, P. O.: Atmospheric fate of methyl vinyl ketone: peroxy radical reactions with NO and HO₂, *J. Phys. Chem. A*, 119, 4562–4572, 2015.
- Praske, E., Otkjær, R. V., Crouse, J. D., Hethcox, J. C., Stoltz, B. M., Kjaergaard, H. G., and Wennberg, P. O.: Atmospheric autoxidation is increasingly important in urban and suburban North America, *P. Natl. Acad. Sci. USA*, 115, 64–69, <https://doi.org/10.1073/pnas.1715540115>, 2017.
- Praske, E., Otkjær, R. V., Crouse, J. D., Hethcox, J. C., Stoltz, B. M., Kjaergaard, H. G., and Wennberg, P. O.: Intramolecular hydrogen shift chemistry of hydroperoxy-substituted peroxy radicals, *J. Phys. Chem. A*, 123, 590–600, <https://doi.org/10.1021/acs.jpca.8b09745>, 2019.
- Raoult, S., Rayez, M.-T., Rayez, J.-C., and Lesclaux, R.: Gas phase oxidation of benzene: Kinetics, thermochemistry and mechanism of initial steps, *Phys. Chem. Chem. Phys.*, 6, 2245–2253, 2004.
- Ray, A., Daele, V., Vassalli, I., Poulet, G., and LeBras, G.: Kinetic study of the reactions of C₂H₅O and C₂H₅O₂ with NO₃ at 298 K, *J. Phys. Chem.*, 100, 5737–5744, 1996.
- Rickard, A. R., Wyche, K. P., Metzger, A., Monks, P. S., Ellis, A. M., Dommen, J., Baltensperger, U., Jenkin, M. E., and Pilling, M. J.: Gas phase precursors to anthropogenic secondary organic aerosol: using the Master Chemical Mechanism to probe detailed observations of 1,3,5-trimethylbenzene photo-oxidation, *Atmos. Environ.*, 44, 5423–5433, 2010.
- Rissanen, M. P., Kurtén, T., Sipilä, M., Thornton, J. A., Kausiala, O., Garmash, O., Kjaergaard, H. G., Petäjä, T., Worsnop, D. R., Ehn, M., and Kulmala, M.: Effects of chemical complexity on the autoxidation mechanisms of endocyclic alkene ozonolysis products: from methylcyclohexenes toward understanding α -pinene, *J. Phys. Chem. A*, 119, 4633–4650, <https://doi.org/10.1021/jp510966g>, 2015.
- Roberts, J. M. and Bertman, S. B.: Measurement of the thermal decomposition of peroxyacetic nitric anhydride (PAN) and peroxy-methacrylic nitric anhydride (MPAN), *Int. J. Chem. Kinet.*, 24, 297–307, 1992.
- Roth, E., Chakir, A., and Ferhati, A.: Study of a peroxybenzoyl radical in the gas phase: ultraviolet spectrum and C₆H₅C(O)O₂+HO₂ reaction between 295 and 357 K, *J. Phys. Chem. A*, 114, 10367–10379, 2010.
- Rowley, D. M., Lightfoot, P. D., Lesclaux, R., and Wallington, T. J.: UV Absorption spectrum and self-reaction of cyclohexylperoxy radicals, *J. Chem. Soc. Faraday T.*, 87, 3221–3226, 1991.
- Rowley, D. M., Lesclaux, R., Lightfoot, P. D., Hughes, K., Rudy, S., Hurley, M. D., and Wallington, T. J.: A kinetic and mechanistic study of the reaction of neopentylperoxy radicals with HO₂, *J. Phys. Chem.*, 96, 7043–7048, 1992a.
- Rowley, D. M., Lesclaux, R., Lightfoot, P. D., Noziere, B., Wallington, T. J., and Hurley, M. D.: Kinetic and mechanistic studies of the reactions of cyclopentylperoxy and cyclohexylperoxy radicals with HO₂, *J. Phys. Chem.*, 96, 4889–4894, 1992b.
- Rowley, D. M., Lightfoot, P. D., Lesclaux, R., and Wallington, T. J.: Ultraviolet absorption spectrum and self-reaction of cyclopentylperoxy radicals, *J. Chem. Soc. Faraday T.*, 88, 1369–1376, 1992c.
- Saunders, S. M., Jenkin, M. E., Derwent, R. G., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-aromatic volatile organic compounds, *Atmos. Chem. Phys.*, 3, 161–180, <https://doi.org/10.5194/acp-3-161-2003>, 2003.
- Schwantes, R. H., Teng, A. P., Nguyen, T. B., Coggon, M. M., Crouse, J. D., St. Clair, J. M., Zhang, X., Schilling, K. A., Seinfeld, J. H., and Wennberg, P. O.: Isoprene NO₃ oxidation products from the RO₂+HO₂ pathway, *J. Phys. Chem. A*, 119, 10158–10171, 2015.
- Sehested, J., Christensen, L. K., Nielsen, O. J., Bilde, M., Wallington, T. J., Schneider, W. F., Orlando, J. J., and Tyndall, G. S.: Atmospheric chemistry of acetone: kinetic study of the CH₃C(O)CH₂O₂+NO/NO₂ reactions and decomposition of CH₃C(O)CH₂O₂NO₂, *Int. J. Chem. Kinet.*, 30, 475–489, 1998.
- Teng, A. P., Crouse, J. D., Lee, L., St. Clair, J. M., Cohen, R. C., and Wennberg, P. O.: Hydroxy nitrate production in the OH-initiated oxidation of alkenes, *Atmos. Chem. Phys.*, 15, 4297–4316, <https://doi.org/10.5194/acp-15-4297-2015>, 2015.
- Tomas, A. and Lesclaux, R.: Self-reaction kinetics of the (CH₃)₂CHC(O)O₂ and (CH₃)₃CC(O)O₂ acylperoxy radicals between 275 and 363 K, *Chem. Phys. Lett.*, 319, 521–528, 2000.

- Tuazon, E. C., Aschmann, S. M., Atkinson, R., and Carter, W. P. L.: The reactions of selected acetates with the OH radical in the presence of NO: novel rearrangement of alkoxy radicals of structure RC(O)OCH'R, *J. Phys. Chem. A*, 102, 2316–2321, 1998a.
- Tuazon, E. C., Aschmann, S. M., and Atkinson, R.: Products of the gas-phase reactions of the OH radical with 1-methoxy-2-propanol and 2-butoxyethanol, *Environ. Sci. Technol.*, 32, 3336–3345, 1998b.
- Tyndall, G. S., Cox, R. A., Granier, C., Lesclaux, R., Moortgat, G. K., Pilling, M. J., Ravishankara, A. R., and Wallington, T. J.: Atmospheric chemistry of small organic peroxy radicals, *J. Geophys. Res.*, D106, 12157–12182, 2001.
- Vaughan, S., Canosa-Mas, C. E., Pfrang, C., Shallcross, D. E., Watson, L., and Wayne, R. P.: Kinetic studies of reactions of the nitrate radical (NO₃) with peroxy radicals (RO₂): an indirect source of OH at night?, *Phys. Chem. Chem. Phys.*, 8, 3749–3760, 2006.
- Vereecken, L. and Peeters, J.: Theoretical investigation of the role of intramolecular hydrogen bonding in β -hydroxyethoxy and β -hydroxyethylperoxy radicals in the tropospheric oxidation of ethene, *J. Phys. Chem. A*, 103, 1768–1775, 1999.
- Vereecken, L. and Peeters, J.: Non-traditional (per)oxy ring-closure paths in the atmospheric oxidation of isoprene and monoterpenes, *J. Phys. Chem. A*, 108, 5197–5204, 2004.
- Vereecken, L., Peeters, J., Orlando, J. J., Tyndall, G. S., and Feronato, C.: Decomposition of β -hydroxypropoxy radicals in the OH-initiated oxidation of propene. A theoretical and experimental study *J. Phys. Chem. A*, 103, 4693–4702, 1999.
- Vereecken, L., Aumont, B., Barnes, I., Bozzelli, J. W., Goldman, M. J., Green, W. H., Madronich, S., McGillen, M. R., Mellouki, A., Orlando, J. J., Picquet-Varrault, B., Rickard, A. R., Stockwell, W. R., Wallington, T. J., and Carter, W. P. L.: Perspective on mechanism development and structure-activity relationships for gas-phase atmospheric chemistry, *Int. J. Chem. Kinet.*, 50, 435–469, 2018.
- Villenave, E. and Lesclaux, R.: Kinetics of the cross reactions of CH₃O₂ and C₂H₅O₂ radicals with selected peroxy radicals, *J. Phys. Chem.*, 100, 14372–14382, 1996.
- Villenave, E., Lesclaux, R., Seefeld, S., and Stockwell, W. R.: Kinetics and atmospheric implications of peroxy radical cross reactions involving the CH₃C(O)O₂ radical, *J. Geophys. Res.*, 103, 25273–25285, 1998.
- Walker, H. M., Stone, D., Ingham, T., Vaughan, S., Cain, M., Jones, R. L., Kennedy, O. J., McLeod, M., Ouyang, B., Pyle, J., Bauguitte, S., Bandy, B., Forster, G., Evans, M. J., Hamilton, J. F., Hopkins, J. R., Lee, J. D., Lewis, A. C., Lidster, R. T., Punjabi, S., Morgan, W. T., and Heard, D. E.: Night-time measurements of HO_x during the RONOCO project and analysis of the sources of HO₂, *Atmos. Chem. Phys.*, 15, 8179–8200, <https://doi.org/10.5194/acp-15-8179-2015>, 2015.
- Wallington, T. J., Hurley, M. D., Maurer, T., Barnes, I., Becker, K. H., Tyndall, G. S., Orlando, J. J., Pimentel, A. S., and Bilde, M.: Atmospheric oxidation mechanism of methyl formate, *J. Phys. Chem. A*, 105, 5146–5154, 2001.
- Wennberg, P. O., Bates, K. H., Crouse, J. D., Dodson, L. G., McVay, R. C., Mertens, L. A., Nguyen, T. B., Praske, E., Schwantes, R. H., Smarte, M. D., St Clair, J. M., Teng A. P., Zhang, X., and Seinfeld, J. H.: Gas-phase reactions of isoprene and its major oxidation products, *Chem. Rev.*, 118, 3337–3390, 2018.
- Winiberg, F. A. F., Dillon, T. J., Orr, S. C., Groß, C. B. M., Bejan, I., Brumby, C. A., Evans, M. J., Smith, S. C., Heard, D. E., and Seakins, P. W.: Direct measurements of OH and other product yields from the HO₂ + CH₃C(O)O₂ reaction, *Atmos. Chem. Phys.*, 16, 4023–4042, <https://doi.org/10.5194/acp-16-4023-2016>, 2016.
- Xu, L., Møller, K. H., Crouse, J. D., Otkjær, R. V., Kjaergaard, H. G., and Wennberg, P. O.: Unimolecular reactions of peroxy radicals formed in the oxidation of α -pinene and β -pinene by hydroxyl radicals, *J. Phys. Chem. A*, 123, 1661–1674, <https://doi.org/10.1021/acs.jpca.8b11726>, 2019.
- Yan, C., Kocevská, S., and Krasnoperov, L. N.: Kinetics of the reaction of CH₃O₂ radicals with OH studied over the 292–526 K temperature range, *J. Phys. Chem. A*, 120, 6111–6121, 2016.
- Yeh, G. K. and Ziemann, P. J.: Alkyl nitrate formation from the reactions of C₈–C₁₄ n-alkanes with OH radicals in the presence of NO_x: measured yields with essential corrections for gas-wall partitioning, *J. Phys. Chem. A*, 118, 8147–8157, <https://doi.org/10.1021/jp500631v>, 2014a.
- Yeh, G. K. and Ziemann, P. J.: identification and yields of 1,4-hydroxynitrates formed from the reactions of C₈–C₁₆ n-alkanes with OH radicals in the presence of NO_x, *J. Phys. Chem. A*, 118, 8797–8806, <https://doi.org/10.1021/jp500631v>, 2014b.
- Zabel, F., Reimer, A., Becker, K. H., and Fink, E. H.: Thermal decomposition of alkyl peroxy nitrates, *J. Phys. Chem.*, 93, 5500–5507, 1989.
- Zhang, X., McVay, R., Huang, D. D., Dalleska, N. F., Aumont, B., Flagan, R. C., and Seinfeld, J. H.: Formation and evolution of molecular products in alpha-pinene secondary organic aerosol, *P. Natl. Acad. Sci. USA*, 112, 14168–14173, 2015.
- Ziemann, P. J.: Evidence for low-volatility diacyl peroxides as a nucleating agent and major component of aerosol formed from reactions of O₃ with cyclohexene and homologous compounds, *J. Phys. Chem. A*, 106, 4390–4402, <https://doi.org/10.1021/jp012925m>, 2002.