



This is a repository copy of *Impact of diabetes Mellitus and Chronic Kidney Disease on cardiovascular outcomes and platelet P2Y12 receptor antagonist effects in patients with acute coronary syndromes: Insights from the PLATO trial.*

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/148157/>

Version: Published Version

---

**Article:**

Franchi, F., James, S.K., Ghukasyan Latic, T. et al. (10 more authors) (2019) Impact of diabetes Mellitus and Chronic Kidney Disease on cardiovascular outcomes and platelet P2Y12 receptor antagonist effects in patients with acute coronary syndromes: Insights from the PLATO trial. *Journal of the American Heart Association*, 8 (6). e011139. ISSN 2047-9980

<https://doi.org/10.1161/jaha.118.011139>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# Impact of Diabetes Mellitus and Chronic Kidney Disease on Cardiovascular Outcomes and Platelet P2Y<sub>12</sub> Receptor Antagonist Effects in Patients With Acute Coronary Syndromes: Insights From the PLATO Trial

Francesco Franchi, MD; Stefan K. James, MD, PhD; Tatevik Ghukasyan Lakic, MSc; Andrzej J. Budaj, MD, PhD; Jan H. Cornel, MD, PhD; Hugo A. Katus, MD; Matyas Keltai, MD, DSc; Frederic Kontny, MD, PhD; Basil S. Lewis, MD; Robert F. Storey, MD, DM; Anders Himmelmann, MD, PhD; Lars Wallentin, MD, PhD; Dominick J. Angiolillo, MD, PhD; on behalf of the PLATO Investigators\*

**Background**—There are limited data on how the combination of diabetes mellitus (DM) and chronic kidney disease (CKD) affects cardiovascular outcomes as well as response to different P2Y<sub>12</sub> receptor antagonists, which represented the aim of the present investigation.

**Methods and Results**—In this post hoc analysis of the PLATO (Platelet Inhibition and Patient Outcomes) trial, which randomized acute coronary syndrome patients to ticagrelor versus clopidogrel, patients (n=15 108) with available DM and CKD status were classified into 4 groups: DM+/CKD+ (n=1058), DM+/CKD− (n=2748), DM−/CKD+ (n=2160), and DM−/CKD− (n=9142). The primary efficacy end point was a composite of cardiovascular death, myocardial infarction, or stroke at 12 months. The primary safety end point was PLATO major bleeding. DM+/CKD+ patients had a higher incidence of the primary end point compared with DM−/CKD− patients (23.3% versus 7.1%; adjusted hazard ratio 2.22; 95% CI 1.88–2.63; *P*<0.001). Patients with DM+/CKD− and DM−/CKD+ had an intermediate risk profile. The same trend was shown for the individual components of the primary end point and for major bleeding. Compared with clopidogrel, ticagrelor reduced the incidence of the primary end point consistently across subgroups (*P*-interaction=0.264), but with an increased absolute risk reduction in DM+/CKD+. The effects on major bleeding were also consistent across subgroups (*P*-interaction=0.288).

**Conclusions**—In acute coronary syndrome patients, a gradient of risk was observed according to the presence or absence of DM and CKD, with patients having both risk factors at the highest risk. Although the ischemic benefit of ticagrelor over clopidogrel was consistent in all subgroups, the absolute risk reduction was greatest in patients with both DM and CKD.

**Clinical Trial Registration**—URL: <http://www.clinicaltrials.gov>. Unique identifier: NCT00391872. (*J Am Heart Assoc.* 2019;8:e011139. DOI: 10.1161/JAHA.118.011139.)

**Key Words:** acute coronary syndrome • chronic kidney disease • clopidogrel • diabetes mellitus • ticagrelor

Patients with diabetes mellitus (DM) are at increased risk of atherothrombotic events.<sup>1</sup> Importantly, DM is a key risk factor for the development of chronic kidney disease

(CKD), a well-known cardiovascular risk factor.<sup>2,3</sup> These observations underscore the importance of antiplatelet therapy for secondary prevention of atherothrombotic

From the University of Florida, College of Medicine-Jacksonville, Jacksonville, FL (F.F., D.J.A.); Department of Medical Sciences, Cardiology (S.K.J., L.W.) and Uppsala Clinical Research Center (S.K.J., T.G.L., L.W.), Uppsala University, Uppsala, Sweden; Postgraduate Medical School, Grochowski Hospital, Warsaw, Poland (A.J.B.); Department of Cardiology, Noordwest Ziekenhuisgroep, Alkmaar, Netherlands (J.H.C.); Medizinische Klinik, Universitätsklinikum Heidelberg, Heidelberg, Germany (H.A.K.); Hungarian Institute of Cardiology, Semmelweis University, Budapest, Hungary (M.K.); Department of Cardiology, Stavanger University Hospital, Stavanger, Norway (F.K.); Lady Davis Carmel Medical Center, Haifa, Israel (B.S.L.); Department of Infection, Immunity and Cardiovascular Disease, University of Sheffield, United Kingdom (R.F.S.); AstraZeneca, Research and Development, Gothenburg, Sweden (A.H.).

An accompanying Appendix S1 is available at <https://www.ahajournals.org/doi/suppl/10.1161/JAHA.118.011139>

\*A complete list of the PLATO Investigators is given in Appendix S1.

**Correspondence to:** Dominick J. Angiolillo, MD, PhD, University of Florida College of Medicine-Jacksonville, 655 West 8th St, Jacksonville, FL 32209. E-mail: [dominick.angiolillo@jax.ufl.edu](mailto:dominick.angiolillo@jax.ufl.edu)

Received October 9, 2018; accepted January 30, 2019.

© 2019 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

## Clinical Perspective

### What Is New?

- Acute coronary syndrome patients with diabetes mellitus and chronic kidney disease are at markedly increased risk for long-term atherothrombotic events compared with patients without these risk factors, as well as with those with only 1 of these.
- Although the ischemic benefit of ticagrelor versus clopidogrel was consistent in all patient subgroups, the magnitude of benefit was enhanced according to the patient risk profile.

### What Are the Clinical Implications?

- There is a need to define the most effective treatment options for these high-risk patients, including strategies to reduce the risk of developing chronic kidney disease in patients with diabetes mellitus.
- Similarly, in patients with established chronic kidney disease, glucose control is also critical to reduce the risk of developing diabetes mellitus.
- Clinicians should use more potent platelet-inhibiting therapy in acute coronary syndrome patients with diabetes mellitus and chronic kidney disease who are often undertreated because of high perceived risk of bleeding.

recurrences in these high-risk patients. Dual antiplatelet therapy with aspirin and a P2Y<sub>12</sub> receptor inhibitor is the standard of care for secondary prevention in acute coronary syndrome (ACS) patients.<sup>4</sup> Guidelines recommend that the more potent P2Y<sub>12</sub> receptor inhibitors (ie, prasugrel or ticagrelor) be preferred over clopidogrel for the treatment of ACS patients because of their greater benefit in reducing the risk of cardiovascular events in these patients, albeit at the expense of increased bleeding.<sup>4,5</sup> Nevertheless, clopidogrel remains widely used in ACS patients.<sup>6,7</sup> DM patients treated with clopidogrel have increased rates of recurrent atherothrombotic events, which may be in part because of reduced platelet inhibitory effects of clopidogrel consistently observed among these subjects.<sup>1,8–11</sup> Although studies assessing the impact of CKD status on clopidogrel-induced antiplatelet effects have yielded conflicting findings, pharmacodynamic assessments conducted among DM patients have shown a greater magnitude of impaired clopidogrel-induced platelet inhibition among those with CKD compared with those without CKD.<sup>12–19</sup>

These observations, as well as those from other small observational studies, suggest that the concomitant presence of DM and CKD status can increase ischemic event rates, underscoring the need for more effective platelet-inhibiting therapies in these high-risk patients.<sup>20,21</sup> However, to date most large-scale studies assessing how the presence of DM

and CKD affects cardiovascular outcomes and the relative impact of specific antiplatelet treatment regimens, in particular P2Y<sub>12</sub> receptor inhibitors, have considered these risk factors separately.<sup>1,2</sup> Indeed, the ever-growing prevalence of CKD in patients with DM underscores the need to better risk stratify these patient cohorts. The aim of this analysis was to assess clinical outcomes in ACS patients from the PLATO (Platelet Inhibition and Patient Outcomes) trial according to the presence or absence of DM and CKD, as well as the differential effects of P2Y<sub>12</sub>-inhibiting therapies (ticagrelor versus clopidogrel) in these populations.

## Methods

The PLATO trial (www.ClinicalTrials.gov NCT00391872) was conducted from October 2006 to February 2009 and randomly assigned 18 624 patients with ST-segment-elevation myocardial infarction (MI), non-ST-segment elevation MI, or unstable angina, treated with an invasive or a noninvasive approach, to receive either ticagrelor or clopidogrel as soon as possible after admission. Details of study design, patients, outcome definitions, and results have been described elsewhere.<sup>22</sup> In brief, ticagrelor was administered as a 180-mg loading dose followed by 90 mg twice daily. Patients assigned to clopidogrel received a maintenance dose of 75 mg daily. Those who were clopidogrel naïve were also administered a 300- to 600-mg loading dose. All patients received aspirin unless intolerant. The randomized treatment continued for a minimum of 6 to a maximum of 12 months (median duration 9.1 months). The primary efficacy end point was a composite of cardiovascular death, MI, or stroke. The primary safety end point was all major bleeding according to PLATO definition. Bleeding events were also defined according to the Thrombolysis In Myocardial Infarction (TIMI) and Global Use of Strategies to Open Occluded Arteries (GUSTO) classifications.<sup>22</sup>

Patients randomized in PLATO with available DM and CKD status at the time of randomization were included in the present analysis. Accordingly, patients were classified into 4 groups: DM+/CKD+, DM+/CKD–, DM–/CKD+, and DM–/CKD–. DM status was defined by the investigators at the time of randomization. Serum glucose and hemoglobin A1c were also measured and used to further characterize the study population, with poor glycemic control defined as levels above the median of serum glucose (6.8 mmol/L) and the median of percentage hemoglobin A1c (6.0%).<sup>23</sup> CKD was defined as a creatinine clearance (CrCl) <60 mL/min according to the Cockcroft-Gault equation.<sup>24</sup> There were no exclusion criteria for renal dysfunction in the PLATO trial except for the requirement of dialysis. In an exploratory analysis, CKD status was also stratified according to the Modification of Diet in Renal Disease and Chronic Kidney Disease Epidemiology

Collaboration equations.<sup>25</sup> In addition, in a subgroup of patients (n=13 688), kidney function was assessed based on cystatin C levels measured on stored samples using the Creatinine-Cystatin C Chronic Kidney Disease Epidemiology Collaboration equation.<sup>26</sup>

The PLATO trial adhered to the Declaration of Helsinki and was approved by the appropriate ethical review boards. All patients provided written informed consent. The data, analytic methods, and study materials will not be made available to other researchers for purposes of reproducing the results or replicating the procedure.

## Statistical Analysis

Categorical baseline variables are presented as frequencies and percentages and compared by DM/CKD group using  $\chi^2$  tests. Continuous baseline variables are presented as medians and 25th to 75th percentiles and compared by DM/CKD group using Kruskal–Wallis tests. Kaplan–Meier estimated event rates from randomization to 12 months were plotted by DM/CKD groups. Cox proportional hazards models were used to assess the associations between CKD-DM status and clinical end points. Multivariable Cox regression models included randomized treatment, age, sex, body mass index, heart rate, prior MI, hypertension, dyslipidemia, smoking status, previous percutaneous coronary intervention or coronary artery bypass graft (CABG), and type of ACS as covariates. The interaction between DM/CKD status and randomized treatment was examined by adding an interaction term to the model. Results are presented as adjusted hazard ratios (HR) with 95% CI. In the comparisons between DM/CKD groups, HRs are reported using DM–/CKD– group as reference. All statistical analyses were performed with SAS 9.4 (SAS Institute, Cary, NC). A 2-sided *P* value of <0.05 was considered statistically significant for differences between groups and treatments.

## Results

### Patients and Outcomes According to CKD and DM Status

Among patients randomized in the PLATO trial, 15 108 had DM and CKD status available and were classified as follows: DM+/CKD+ (n=1058), DM+/CKD– (n=2748), DM–/CKD+ (n=2160), and DM–/CKD– (n=9142). Baseline characteristics are reported in Table 1. After excluding patients who prematurely discontinued because of death, the number of patients who discontinued treatment during follow-up was low (43 in the CKD+DM+ group [0.28%], 71 in the CKD–DM+ group [0.47%], 83 in the CKD+DM– group [0.55%], and 206 in the CKD–DM– group [1.36%]). Patients with DM+/CKD+

more frequently had a prior history of cardiovascular disease, including MI, stroke, and peripheral arterial disease; were more frequently diagnosed with non-ST-elevation ACS rather than ST-elevation MI; and were more frequently treated with a noninvasive approach.

Patients with DM+/CKD+ had an over 3-fold higher incidence of the primary end point at 12 months compared with DM–/CKD– patients (23.3% versus 7.1%; adjusted HR 2.22; 95% CI 1.88–2.63). Patients with DM+/CKD– (10.7%; adjusted HR 1.34; 95% CI 1.16–1.55) and DM–/CKD+ (15.8%; adjusted HR 1.60; 95% CI 1.37–1.86) had an intermediate risk profile (*P* for trend <0.001; Figure 1). The same trend was shown for the individual components of the primary end point, cardiovascular death, MI, and stroke, as well as for all-cause mortality (Figure 2). Patients with DM+/CKD+ also had the highest risk of PLATO-defined major bleeding compared with DM–/CKD– patients (14.8% versus 8.5%; adjusted HR 1.47; 95% CI 1.21–1.77) and patients with DM+/CKD– (11.7%; adjusted HR 1.34; 95% CI: 1.17–1.54) or DM–/CKD+ (11.8%; adjusted HR 1.13; 95% CI 0.96–1.33) (Figure 3A). Non-CABG-related major bleeding rates were higher in patients with DM+/CKD+ and DM–/CKD+ compared with patients with DM+/CKD– and DM–/CKD– (Figure 3B). Major bleeding defined according to TIMI and GUSTO criteria showed a similar trend (Figure 4). Results were consistent when measures of poor glycemic control and alternative definitions of CKD were considered (Table 2).

### Outcomes of Ticagrelor Versus Clopidogrel According to CKD and DM Status

Compared with clopidogrel, ticagrelor significantly reduced the incidence of the primary end point consistently across subgroups (*P* interaction=0.3). However, the absolute risk reduction (ARR) with ticagrelor versus clopidogrel was considerably higher in DM+/CKD+ patients (11.26%; adjusted HR 0.78; 95% CI 0.61–1.01) compared with DM–/CKD– (1.37%; adjusted HR 0.86; 95% CI 0.73–1.00) (Figures 5 and 6). Consistent findings were shown on all the components of the primary end point (Table 3). In particular, ticagrelor led to a 5.8% ARR in cardiovascular death in patients with DM+/CKD+ compared with a 0.2% reduction in DM–/CKD– patients. Accordingly, the number-needed-to-treat for the primary end point was 8.9 in DM+/CKD+ and 73 in DM–/CKD–, and for cardiovascular death 17.2 in DM+/CKD+ and 500 in DM–/CKD–.

The effects of ticagrelor versus clopidogrel on PLATO-defined major bleeding were consistent across subgroups (*P* interaction=0.3). In particular, there was no increased risk of major bleeding with ticagrelor compared with clopidogrel in the subgroup of patients with DM+/CKD+ (27.4% versus 26.9%; HR 1.02; 95% CI 0.75–1.40). Accordingly, the effects

**Table 1.** Baseline Characteristics by DM/CKD Status


Group of Characteristics	Characteristic (at Baseline)	DM+/CKD+ (n=1058)	DM+/CKD- (n=2748)	DM-/CKD+ (n=2160)	DM-/CKD- (n=9142)	P Value
Demographics	Age (y), median (Q1–Q3)	72 (66–78)	61 (55–68)	74 (68–79)	59 (52–66)	<0.0001
	Age ≥75 y	429 (40.5%)	233 (8.5%)	1060 (49.1%)	604 (6.6%)	<0.0001
	Female sex	456 (43.1%)	851 (31.0%)	823 (38.1%)	2176 (23.8%)	<0.0001
	Weight (kg), median (Q1–Q3)	75 (65–84)	84 (74–95)	72 (62–80)	80 (70–90)	<0.0001
	Weight <60 kg	107 (10.1%)	120 (4.4%)	349 (16.2%)	498 (5.4%)	<0.0001
	Height (cm), median (Q1–Q3)	165 (160–172)	170 (163–175)	167 (160–173)	171 (165–177)	<0.0001
	BMI (kg/m <sup>2</sup> ), median (Q1–Q3)	26.9 (24.6–30.2)	29.3 (26.4–32.9)	25.4 (23.2–28.1)	27.4 (24.8–30.2)	<0.0001
	Waist circumference (cm), median (Q1–Q3)	99 (91–108)	103 (94–112)	95 (86–102)	97 (90–105)	<0.0001
Race, n (%)	White	922 (87.1)	2515 (91.5)	1928 (89.3)	8553 (93.6)	<0.0001
	Black	22 (2.1)	46 (1.7)	28 (1.3)	71 (0.8)	
	Asian	84 (7.9)	160 (5.8)	160 (7.4)	457 (5.0)	
	Other	30 (2.8)	27 (1.0)	44 (2.0)	61 (0.7)	
Cardiovascular risk factors, n (%)	Habitual smoker	130 (12.3)	800 (29.1)	413 (19.1)	4061 (44.4)	<0.0001
	Hypertension	925 (87.4)	2162 (78.7)	1574 (72.9)	5187 (56.7)	<0.0001
	Dyslipidemia	622 (58.8)	1629 (59.3)	916 (42.4)	3816 (41.7)	<0.0001
History, n (%)	Angina pectoris	651 (61.5)	1423 (51.8)	1137 (52.6)	3647 (39.9)	<0.0001
	Myocardial infarction	360 (34.0)	676 (24.6)	556 (25.7)	1507 (16.5)	<0.0001
	Congestive heart failure	176 (16.6)	188 (6.8)	229 (10.6)	255 (2.8)	<0.0001
	PCI	217 (20.5)	462 (16.8)	290 (13.4)	1025 (11.2)	<0.0001
	CABG	139 (13.1)	236 (8.6)	155 (7.2)	350 (3.8)	<0.0001
	TIA	48 (4.5)	75 (2.7)	81 (3.8)	191 (2.1)	<0.0001
	Nonhemorrhagic stroke	96 (9.1)	129 (4.7)	117 (5.4)	242 (2.6)	<0.0001
	Peripheral arterial disease	149 (14.1)	210 (7.6)	163 (7.5)	422 (4.6)	<0.0001
Medications on arrival, n (%)	Aspirin	1007 (95.2)	2618 (95.3)	2033 (94.1)	8756 (95.8)	0.01
	β-Blockade	842 (79.6)	2257 (82.1)	1613 (74.7)	6739 (73.7)	<0.0001
	ACE-inhibition and/or ARB	806 (76.2)	2049 (74.6)	1397 (64.7)	5361 (58.6)	<0.0001
	Statin	823 (77.8)	2230 (81.1)	1651 (76.4)	7350 (80.4)	<0.0001
	Ca-inhibitor	276 (26.1)	539 (19.6)	352 (16.3)	1054 (11.5)	<0.0001
	Diuretic	497 (47.0)	793 (28.9)	758 (35.1)	1449 (15.8)	<0.0001
	Insulin treatment before admission	282 (26.7)	572 (20.8)			0.0001
Medications index event to discharge, n (%)	GP 2b/3a inhibitor	177 (16.7)	734 (26.7)	413 (19.1)	2686 (29.4)	<0.0001
	Unfractionated heparin	524 (49.5)	1591 (57.9)	1195 (55.3)	5473 (59.9)	<0.0001

Continued

Table 1. Continued

Group of Characteristics	Characteristic (at Baseline)	DM+/CKD+ (n=1058)	DM+/CKD- (n=2748)	DM-/CKD+ (n=2160)	DM-/CKD- (n=9142)	P Value
	Low-molecular-weight heparin	590 (55.8)	1460 (53.1)	1199 (55.5)	4734 (51.8)	0.003
	Fondaparinux	34 (3.2)	74 (2.7)	74 (3.4)	249 (2.7)	0.3
	Bivalirudin	25 (2.4)	90 (3.3)	34 (1.6)	158 (1.7)	<0.0001
Intended approach	Invasive	603 (57.0%)	1912 (69.6%)	1311 (60.7%)	6915 (75.6%)	<0.0001
	Noninvasive	455 (43.0%)	836 (30.4%)	849 (39.3%)	2227 (24.4%)	
Final ACS diagnosis	ST-elevation MI	244 (23.1%)	863 (31.4%)	638 (29.6%)	3980 (43.6%)	<0.0001
	Non-ST-elevation MI	559 (52.9%)	1259 (45.8%)	1038 (48.2%)	3622 (39.6%)	
	Unstable angina	224 (21.2%)	566 (20.6%)	427 (19.8%)	1336 (14.6%)	
	Other	29 (2.7%)	60 (2.2%)	50 (2.3%)	199 (2.2%)	
Randomized treatment	Delay from start of pain (h), median (Q1–Q3)	14.2 (6.8–21.2)	12.7 (5.7–20.4)	14.0 (5.8–21.1)	10.2 (4.3–19.0)	<0.0001
	Treatment duration (d), median (Q1–Q3)	258 (55–361)	276 (179–365)	265 (73–363)	284 (184–366)	<0.0001
Biomarkers	Creatinine ( $\mu\text{mol/L}$ ), median (Q1–Q3)	115.0 (106.0–141.0)	80.0 (70.7–88.0)	106.0 (97.0–124.0)	80.0 (71.0–88.0)	<0.0001
	Glucose (mmol/L), median (Q1–Q3)	9.9 (7.2–13.5)	9.7 (7.2–13.2)	6.5 (5.6–7.9)	6.4 (5.6–7.7)	<0.0001
	HbA1c (mmol/mol), median (Q1–Q3)	7.5 (6.6–8.7)	7.6 (6.7–9.1)	5.9 (5.6–6.2)	5.8 (5.6–6.1)	<0.0001
	Hemoglobin (mmol/mol), median (Q1–Q3)	128.0 (116.0–140.0)	139.0 (128.0–149.0)	134.0 (123.0–145.0)	142.0 (132.0–151.0)	<0.0001
	NT-proBNP (pmol/L), median (Q1–Q3)	1734 (610.0–4071)	395.0 (146.0–953.0)	1002 (320.0–2544)	277.0 (99.0–721.0)	<0.0001
	Troponin I $\mu\text{g/L}$ , median (Q1–Q3)	1.10 (0.12–6.00)	0.95 (0.11–4.30)	1.00 (0.11–5.70)	0.90 (0.12–4.70)	0.01
	Creatinine (mg/dL), median (Q1–Q3)	1.3 (1.2–1.6)	0.9 (0.8–1.0)	1.2 (1.1–1.4)	0.9 (0.8–1.0)	<0.0001
	CrCl (mL/min), median (Q1–Q3)	48.4 (38.9–55.1)	86.7 (73.2–104.5)	50.3 (42.7–55.9)	87.7 (74.5–104.0)	<0.0001

ACE indicates angiotensin converting enzyme; ACS, acute coronary syndrome; ARB, angiotensin receptor blocker; BMI, body mass index; CABG, coronary artery bypass graft; CKD, chronic kidney disease; CrCl, creatinine clearance by Cockcroft-Gault equation; DM, diabetes mellitus; GP, glycoprotein; HbA1c, hemoglobin A1c; MI, myocardial infarction; NT-proBNP, N-terminal pro-brain natriuretic peptide; PCI, percutaneous coronary intervention; TIA, transient ischemic attack.



**Figure 1.** Kaplan–Meier event rate curves for the cumulative incidence of the primary composite end point of cardiovascular (CV) death, myocardial infarction (MI), and stroke stratified by DM/CKD status. *P* value represents the overall comparison among groups according to DM/CKD status. The model is adjusted for age, sex, body mass index, heart rate, prior myocardial infarction, hypertension, dyslipidemia, angina pectoris, smoking status, previous percutaneous coronary intervention or coronary artery bypass graft, type of acute coronary syndrome and randomized treatment. CKD indicates chronic kidney disease; DM, diabetes mellitus.


on non-CABG-related major bleeding were also consistent regardless of CKD/DM status, although the increase in bleeding risk with ticagrelor was numerically higher in patients with CKD (both DM+/CKD+ and DM−/CKD+) (Table 3). The number-needed-to-harm for all major bleeding was 208 in DM+/CKD+ and 49 in DM−/CKD− and for non-CABG-related major bleeding was 73 in DM+/CKD+ and 105 in DM−/CKD−. Major bleeding defined according to TIMI and GUSTO criteria followed the same trend (Table 4).

Results were consistent when measures of poor glycemic control and alternative definitions of CKD were considered. In particular, with poor glycemic control defined by hemoglobin A1c and CKD defined by the Creatinine-Cystatin C Chronic Kidney Disease Epidemiology Collaboration equation, the effects of ticagrelor versus clopidogrel on ischemic and bleeding events were consistent across subgroups (Table 5). In patients with DM+/CKD+, ticagrelor led to a 14% ARR in the primary end point and a 9% ARR in cardiovascular death compared with clopidogrel with no significant increase in major bleeding.

## Discussion

The data from the present post hoc analysis of the PLATO trial represent the largest exploring the impact of having DM, CKD,

or both, on clinical outcomes in ACS patients. Our study showed that (1) the concomitant presence of CKD and DM is not uncommon in patients with ACS, representing 7% of the overall study population; (2) patients with CKD and DM are more likely to already have established atherosclerotic disease, more frequently present with a non-ST-elevation ACS and are more likely to be treated with a noninvasive approach; (3) patients with either DM or CKD are at increased risk of ischemic events compared with patients without these risk factors; and the combination of DM and CKD status is associated with an over 3-fold increased risk of ischemic events compared with patients without these risk factors, including a 6-fold increase in cardiovascular death; (4) the presence of DM and CKD is associated with a significant increase in major bleeding and non-CABG-related major bleeding, but not in CABG-related bleeding; (5) the benefit of ticagrelor over clopidogrel on ischemic outcomes is consistent across DM and CKD status, but the magnitude of absolute benefit is enhanced in higher-risk patients; in particular, in patients with DM and CKD ticagrelor led to a 22% relative risk reduction and an 11% ARR in the primary end point compared with clopidogrel, including a 21% relative risk reduction and an 5.8% ARR in cardiovascular death; and (6) there was no signal of increased risk of bleeding with ticagrelor in patients with CKD and DM as compared with the other subgroups.




**Figure 2.** Kaplan–Meier event rate curves for the cumulative incidence of (A) cardiovascular (CV) death, (B) myocardial infarction (MI), (C) stroke, and (D) all-cause mortality stratified by DM/CKD status. *P* value represents the overall comparison among groups according to DM/CKD status. The model is adjusted for age, sex, body mass index, heart rate, prior myocardial infarction, hypertension, dyslipidemia, angina pectoris, smoking status, previous percutaneous coronary intervention or coronary artery bypass graft, type of acute coronary syndrome, and randomized treatment. CKD indicates chronic kidney disease; DM, diabetes mellitus.

DM and CKD have both been independently associated with an increased risk of cardiovascular events, which may be attributed to abnormalities specific to these patients favoring a prothrombotic and pro-inflammatory status.<sup>1,2</sup> Among patients with DM, impaired clopidogrel-induced antiplatelet effects leading to high levels of platelet reactivity has been largely attributed to an attenuation of clopidogrel's pharmacokinetic profile, characterized by lower active metabolite levels, and in part to dysregulation of the P2Y<sub>12</sub> receptor signaling pathway.<sup>9,10,27</sup> Subgroup analysis of major clinical trials have shown a reduced benefit of clopidogrel in CKD patients.<sup>2</sup> Patients with CKD are characterized by upregulation of the P2Y<sub>12</sub> signaling pathway induced by dinucleoside polyphosphates and impaired hepatic function, which can potentially impact clopidogrel metabolism.<sup>28–32</sup> However, while pharmacodynamic studies have consistently shown DM to be associated with impaired clopidogrel-induced antiplatelet effects, results have been conflicting when assessing how CKD affects clopidogrel

response. These observations may be attributed to confounders within the heterogeneous study populations in which these studies have been performed.<sup>12–19</sup> Pharmacodynamic assessments specifically conducted among DM patients who also have CKD have shown these patients to have greater impairment of clopidogrel-induced platelet inhibition compared with those without CKD.<sup>13,15,16</sup> However, in the absence of DM, renal function has not always been shown to affect clopidogrel's antiplatelet effects.<sup>12,13,17–19</sup> Overall, these findings suggest that there may be some level of synergism of DM and CKD on platelet reactivity in clopidogrel-treated patients, which would be in line with the clinical observations of the present investigation.<sup>16</sup>

A post hoc analysis of the FREEDOM (Comparison of Two Treatments for Multivessel Coronary Artery Disease in Individuals With Diabetes) trial assessing revascularization strategies (surgical versus percutaneous) among DM patients (n=1843) with multivessel coronary artery disease






**Figure 3.** Kaplan–Meier event rate curves for the cumulative incidence of (A) major bleeding, and (B) non-CABG-related major bleeding stratified by DM/CKD status. *P* value represents the overall comparison among groups according to DM/CKD status. Bleeding is defined according to PLATO criteria. The model is adjusted for age, sex, body mass index, heart rate, prior myocardial infarction, hypertension, dyslipidemia, angina pectoris, smoking status, previous percutaneous coronary intervention, or coronary artery bypass graft, type of acute coronary syndrome, and randomized treatment. CABG indicates coronary artery bypass graft; CKD, chronic kidney disease; DM, diabetes mellitus; PLATO, Platelet Inhibition and Patient Outcomes.

evaluated the impact of CKD status on clinical outcomes.<sup>20</sup> In this analysis, CKD affected clinical outcomes irrespective of the strategy used for revascularization, leading to a

nearly 2-fold risk increase in all-cause mortality, cardiovascular death, and stroke and a 1.5-fold risk increase in major bleeding.<sup>20</sup> Our analysis represents the largest data set to



**Figure 4.** Kaplan–Meier event rate curves for the cumulative incidence of major/severe bleeding according to (A) TIMI, and (B) GUSTO criteria stratified by DM/CKD status. *P* value represents the overall comparison among groups according to DM/CKD status. The model is adjusted for age, sex, body mass index, heart rate, prior myocardial infarction, hypertension, dyslipidemia, angina pectoris, smoking status, previous percutaneous coronary intervention or coronary artery bypass graft, type of acute coronary syndrome, and randomized treatment. CKD indicates chronic kidney disease; DM, diabetes mellitus; GUSTO, Global Use of Strategies to Open Occluded Arteries; TIMI, thrombolysis in myocardial infarction.

unravel the contributing role of DM and CKD on cardiovascular outcomes. We extend the findings from the FREEDOM analysis to ACS patients receiving dual antiplatelet therapy

undergoing different treatment strategies (invasive or non-invasive), showing that the presence of either DM or CKD increases long-term cardiovascular events to a similar

**Table 2.** Ischemic and Bleeding Outcomes According to DM/CKD Subgroup, With Poor Glycemic Control Defined by HbA1c and CKD Defined by the Creatinine-Cystatin C CKD-EPI Equation


DM/CKD Subgroup	No. of Events	No. of Patients	Event Rate (%) <sup>*</sup>	HR (95% CI) <sup>†</sup>	P Value <sup>‡</sup>
<b>Cardiovascular death/MI/stroke</b>					
DM−/CKD−	392	1264	6.9		<0.0001
DM+/CKD−	580	5726	10.1	1.33 (1.16–1.52)	
DM−/CKD+	123	734	16.8	1.72 (1.39–2.13)	
DM+/CKD+	263	1264	20.8	2.09 (1.76–2.49)	
<b>Cardiovascular death</b>					
DM−/CKD−	121	5673	2.1		<0.0001
DM+/CKD−	215	5726	3.8	1.54 (1.23–1.94)	
DM−/CKD+	65	734	8.9	2.50 (1.81–3.44)	
DM+/CKD+	155	1264	12.3	3.44 (2.64–4.48)	
<b>MI</b>					
DM−/CKD−	258	5673	4.5		<0.0001
DM+/CKD−	357	5726	6.2	1.24 (1.05–1.47)	
DM−/CKD+	69	734	9.4	1.60 (1.21–2.12)	
DM+/CKD+	130	1264	10.3	1.66 (1.32–2.10)	
<b>All-cause death</b>					
DM−/CKD−	145	5673	2.6		<0.0001
DM+/CKD−	238	5726	4.2	1.45 (1.17–1.79)	
DM−/CKD+	72	734	9.8	2.21 (1.63–2.99)	
DM+/CKD+	174	1264	13.8	3.19 (2.49–4.08)	
<b>Stroke</b>					
DM−/CKD−	46	5673	0.8		0.1679
DM+/CKD−	74	5726	1.3	1.43 (0.98–2.08)	
DM−/CKD+	11	734	1.5	1.15 (0.58–2.29)	
DM+/CKD+	27	1264	2.1	1.67 (0.99–2.81)	
<b>Major bleeding</b>					
DM−/CKD−	484	5673	8.5		0.0039
DM+/CKD−	629	5726	11.0	1.26 (1.11–1.42)	
DM−/CKD+	86	734	11.7	1.14 (0.90–1.45)	
DM+/CKD+	148	1264	11.7	1.14 (0.94–1.39)	
<b>Non-CABG-related major bleeding</b>					
DM−/CKD−	161	5673	2.8		0.0070
DM+/CKD−	180	5726	3.1	1.00 (0.81–1.25)	
DM−/CKD+	44	734	6.0	1.34 (0.94–1.91)	
DM+/CKD+	88	1264	7.0	1.55 (1.16–2.07)	
<b>CABG-related major bleeding</b>					
DM−/CKD−	367	5628	6.5		0.1678
DM+/CKD−	366	5673	6.5	1.02 (0.88–1.18)	
DM−/CKD+	44	727	6.1	0.96 (0.69–1.32)	
DM+/CKD+	96	1250	7.7	1.29 (1.01–1.65)	

The model is adjusted for age, sex, BMI, heart rate, prior myocardial infarction, hypertension, dyslipidemia, angina pectoris, smoking status, previous PCI or CABG, type of ACS define and randomized treatment. BMI indicates body mass index; CABG, coronary artery bypass graft; CKD, chronic kidney disease; CKD-EPI, chronic kidney disease epidemiology collaboration; DM, diabetes mellitus; HbA1c, hemoglobin A1c; HR, hazard ratio; MI, myocardial infarction; PCI, percutaneous coronary intervention.

<sup>\*</sup>The crude event rate, (no. events/no. of subjects)×100%.

<sup>†</sup>Subgroup DM−/CKD− is the reference category.

<sup>‡</sup>P value for the effect of DM/CKD subgroup.




**Figure 5.** Hazard ratios (HR) with 95% CI for the primary composite end point (cardiovascular death, myocardial infarction, and stroke) of ticagrelor (T) vs clopidogrel (C) stratified by DM/CKD status. The model is adjusted for age, sex, body mass index, heart rate, prior myocardial infarction, hypertension, dyslipidemia, angina pectoris, smoking status, previous percutaneous coronary intervention or coronary artery bypass graft, type of acute coronary syndrome, and randomized treatment. CKD indicates chronic kidney disease; DM, diabetes mellitus.

extent but when these risk factors are combined, this risk is further amplified. Notably, this was consistent using multiple definitions of DM and CKD, supporting the validity of our study findings. The ever-rising prevalence of both DM and CKD underscore the relevance of these observations. In fact, both clinical disorders are pandemic public health problems. CKD has a prevalence of 13% in the United States and up to 17% in Europe.<sup>3,33</sup> Importantly, DM is a key risk factor for the development of CKD, and about one third of DM patients are found to have CKD.<sup>3</sup> Therefore, with the increasing prevalence of DM, which is expected to double over the next 20 years, the prevalence of CKD is also expected to rise.<sup>34</sup> These observations underscore the need for defining the most effective treatment options for these high-risk patients, including strategies to reduce the risk of developing CKD in patients with DM. To this extent, sodium-glucose cotransporter-2 inhibitors are new antihyperglycemic therapies known to reduce long-term decline in kidney function.<sup>35,36</sup> Similarly, in patients with established CKD, glucose control is also critical to reduce the risk of developing DM.

Ticagrelor is characterized by more potent and predictable antiplatelet effects compared with clopidogrel, which

translate into better clinical outcomes in ACS patients, albeit at the expense of an increased risk of major bleeding.<sup>22,37</sup> Pharmacodynamic assessments have shown that the enhanced potency of ticagrelor over clopidogrel persists in patients with DM,<sup>38,39</sup> and in the DM subgroup of PLATO, compared with clopidogrel, ticagrelor was associated with a 2.1% ARR in the primary end point, a finding that was consistent with the overall trial results (*P*-interaction: 0.49).<sup>23</sup> In patients with CKD, ticagrelor led to a 4.7% ARR of the primary ischemic end point, which was also consistent with the overall trial results (*P*-interaction: 0.13).<sup>24</sup> However, there are limited data on the pharmacodynamic effects of ticagrelor in CKD patients.<sup>40,41</sup> The present study findings show that, although the benefit of ticagrelor over clopidogrel is consistent across subgroups (*P*-interaction: 0.264), the enhanced benefit of ticagrelor in patients with CKD is even greater in patients who also have DM (11% ARR), including a 5.8% ARR in cardiovascular mortality. Indeed, the higher event rates that characterize these patients can contribute to the greater magnitude of the treatment effect associated with more potent platelet P2Y<sub>12</sub> inhibition induced by ticagrelor. In addition, prior investigations supporting impaired clopidogrel-induced platelet inhibition in DM patients, in particular those



**Figure 6.** Kaplan–Meier event rate curves for the cumulative incidence of the primary composite end point of cardiovascular (CV) death, myocardial infarction, and stroke stratified by treatment group and DM/CKD status. C indicates clopidogrel; CKD, chronic kidney disease; DM, diabetes mellitus; T, ticagrelor.

also with CKD, may contribute to these findings.<sup>9–12,14,16,17</sup> However, because DM and CKD patients are characterized by enhanced vascular inflammation and endothelial dysfunction, it cannot be excluded that they could be more susceptible to the off-target effects of ticagrelor. In fact, ticagrelor increases adenosine levels by inhibiting its reuptake by erythrocytes and adenosine may modulate inflammatory response and favor vasodilation.<sup>42</sup>

Patients with CKD and DM are overall at increased risk of bleeding. This may explain why in some studies these patients are less commonly treated with more potent platelet-inhibiting therapies.<sup>43,44</sup> The increased risk for bleeding among DM and CKD patients was also confirmed in this analysis. However, there was no increased risk of major bleeding with ticagrelor versus clopidogrel in the subgroup of patients with DM+/CKD+. The increase in non-CABG-related major bleeding events was numerically higher in patients with DM+/CKD+, but the relative risk was similar and the effect was overall consistent across groups, also using different bleeding definitions. These findings were also consistent using multiple definitions of DM and CKD.

## Study Limitations

The results of the present study should be interpreted in light of some limitations. Patients with end-stage renal disease requiring hemodialysis were excluded from the trial; therefore,

our results are not applicable to this setting. Although we used different definitions to define CKD status, we did not measure albumin–creatinine ratio and therefore may have underestimated the true prevalence of CKD. Accordingly, the number of patients with CKD+ in our study population was relatively small. CKD was defined according to baseline creatinine levels at the time of ACS presentation. Therefore, creatinine clearance may not be reflective of steady-state kidney function. Indeed, it may be argued that the results of our study pertain to a cohort of CKD patients with mostly moderate (stage 3) degree of renal impairment and the results cannot be extrapolated to those with more advanced stages of renal disease. Moreover, the present investigation does not provide any mechanistic insights for the enhanced rates of adverse outcomes and the inconsistent response to different classes of P2Y<sub>12</sub> inhibiting therapies among patients with concomitant DM and CKD, which is a topic of ongoing investigation (NCT02539160). It may be argued that there are large baseline differences between the DM/CKD groups that might not be possible to fully account for by covariate adjustment. Although an age/sex/comorbid matched analysis could have represented an option, this typically leads to loss of information when not all subjects can be matched, and a similar analysis would have resulted in smaller patient cohorts and ultimately not reflective of risk profile of this patient population in real-world clinical practice. Finally, our results derive from a post hoc subgroup analysis and should

**Table 3.** Outcomes of Ticagrelor Versus Clopidogrel According to DM/CKD Status

DM/CKD Subgroup	Ticagrelor Patients (N)	Clopidogrel Patients (N)	Ticagrelor Event Rate, N (%)	Clopidogrel Event Rate, N (%)	HR (95% CI)	P Value Interaction
<b>Cardiovascular death</b>						0.3
DM+/CKD+	521	537	55 (13.60)	77 (19.40)	0.79 (0.55–1.11)	
DM–/CKD+	1043	1117	69 (8.33)	111 (12.80)	0.68 (0.51–0.92)	
DM+/CKD–	1363	1385	59 (5.30)	62 (5.38)	1.00 (0.70–1.44)	
DM–/CKD–	4621	4521	98 (2.51)	104 (2.71)	0.93 (0.70–1.22)	
<b>MI</b>						0.2
DM+/CKD+	521	537	52 (13.79)	72 (19.66)	0.76 (0.53–1.09)	
DM–/CKD+	1043	1117	77 (9.77)	100 (12.29)	0.83 (0.62–1.12)	
DM+/CKD–	1363	1385	93 (8.76)	84 (7.58)	1.13 (0.84–1.52)	
DM–/CKD–	4621	4521	195 (5.16)	233 (6.33)	0.82 (0.67–0.99)	
<b>All-cause death</b>						0.5
DM+/CKD+	521	537	63 (15.58)	82 (20.66)	0.85 (0.61–1.18)	
DM–/CKD+	1043	1117	80 (9.66)	125 (14.41)	0.70 (0.53–0.93)	
DM+/CKD–	1363	1385	64 (5.75)	69 (5.98)	0.98 (0.70–1.37)	
DM–/CKD–	4621	4521	112 (2.87)	123 (3.21)	0.90 (0.69–1.16)	
<b>Stroke</b>						0.6
DM+/CKD+	521	537	13 (3.26)	18 (4.66)	0.78 (0.38–1.59)	
DM–/CKD+	1043	1117	23 (2.81)	20 (2.33)	1.24 (0.68–2.26)	
DM+/CKD–	1363	1385	22 (1.99)	16 (1.40)	1.42 (0.75–2.71)	
DM–/CKD–	4621	4521	40 (1.03)	31 (0.81)	1.28 (0.80–2.04)	
<b>Major bleeding</b>						0.3
DM+/CKD+	521	537	78 (27.37)	79 (26.89)	1.02 (0.75–1.40)	
DM–/CKD+	1043	1117	129 (21.73)	125 (19.42)	1.13 (0.88–1.44)	
DM+/CKD–	1363	1385	150 (17.61)	171 (18.88)	0.91 (0.73–1.13)	
DM–/CKD–	4621	4521	420 (13.23)	355 (11.19)	1.16 (1.01–1.34)	
<b>Non-CABG-related major bleeding</b>						0.7
DM+/CKD+	521	537	39 (12.87)	32 (10.18)	1.32 (0.82–2.10)	
DM–/CKD+	1043	1117	75 (12.15)	62 (9.14)	1.34 (0.96–1.88)	
DM+/CKD–	1363	1385	48 (5.30)	50 (5.13)	1.03 (0.69–1.52)	
DM–/CKD–	4621	4521	129 (3.88)	97 (2.93)	1.30 (1.00–1.69)	

The model is adjusted for age, sex, body mass index, heart rate, prior myocardial infarction, hypertension, dyslipidemia, angina pectoris, smoking status, previous percutaneous coronary intervention or CABG, type of acute coronary syndrome and randomized treatment. CABG indicates coronary artery bypass graft; CKD, chronic kidney disease; DM, diabetes mellitus; HR, hazard ratio; MI, myocardial infarction.

as such be considered as hypothesis-generating and requiring confirmation in prospectively designed studies.

## Conclusions

In conclusion, the results of the present analysis showed that ACS patients with DM and CKD are at markedly increased risk for long-term atherothrombotic events compared with patients without these risk factors, as well as with those with only 1 of

these. Although the ischemic benefit of ticagrelor versus clopidogrel was consistent in all patient subgroups, the magnitude of benefit was enhanced according to the patient risk profile. Although patients with DM and CKD are at increased risk of bleeding, there were no signals of increased risk of major bleeding events with ticagrelor. Overall, these data underscore the need for using more potent platelet-inhibiting therapy in ACS patients with DM and CKD who are often undertreated because of high perceived risk of bleeding.

**Table 4.** Bleeding Outcomes of Ticagrelor Versus Clopidogrel According to DM/CKD Status According to TIMI and GUSTO Criteria

DM/CKD Subgroup	Ticagrelor Patients (N)	Clopidogrel Patients (N)	Ticagrelor Event Rate, N (%)	Clopidogrel Event Rate, N (%)	HR (95% CI)	P Value Interaction
<b>TIMI major bleeding</b>						<b>0.049</b>
DM+/CKD+	521	537	48 (16.16)	48 (15.71)	1.02 (0.68–1.52)	
DM–/CKD+	1043	1117	78 (12.67)	81 (12.13)	1.05 (0.77–1.43)	
DM+/CKD–	1363	1385	93 (10.58)	124 (13.32)	0.77 (0.59–1.01)	
DM–/CKD–	4621	4521	308 (9.56)	252 (7.82)	1.21 (1.02–1.42)	
<b>TIMI non-CABG-related major bleeding</b>						<b>0.219</b>
DM+/CKD+	521	537	24 (7.84)	15 (4.71)	1.69 (0.89–3.23)	
DM–/CKD+	1043	1117	38 (6.01)	36 (5.22)	1.16 (0.74–1.83)	
DM+/CKD–	1363	1385	27 (2.95)	34 (3.47)	0.84 (0.51–1.40)	
DM–/CKD–	4621	4521	88 (2.63)	57 (1.71)	1.51 (1.08–2.11)	
<b>GUSTO severe bleeding</b>						<b>0.882</b>
DM+/CKD+	521	537	25 (8.12)	34 (10.88)	0.77 (0.46–1.28)	
DM–/CKD+	1043	1117	36 (5.72)	39 (5.70)	0.99 (0.63–1.56)	
DM+/CKD–	1363	1385	33 (3.63)	40 (4.09)	0.88 (0.55–1.39)	
DM–/CKD–	4621	4521	89 (2.67)	92 (2.78)	0.95 (0.71–1.27)	
<b>GUSTO non-CABG-related severe bleeding</b>						<b>0.545</b>
DM+/CKD+	521	537	20 (6.44)	19 (5.98)	1.08 (0.58–2.03)	
DM–/CKD+	1043	1117	25 (3.93)	25 (3.61)	1.06 (0.61–1.85)	
DM+/CKD–	1363	1385	17 (1.85)	25 (2.54)	0.74 (0.40–1.36)	
DM–/CKD–	4621	4521	54 (1.61)	41 (1.23)	1.28 (0.85–1.91)	

The model is adjusted for age, sex, BMI, heart rate, prior myocardial infarction, hypertension, dyslipidemia, angina pectoris, smoking status, previous PCI or CABG, type of ACS, and randomized treatment. ACS indicates acute coronary syndrome; BMI, body mass index; CABG, coronary artery bypass graft; CKD, chronic kidney disease; DM, diabetes mellitus; GUSTO, Global Use of Strategies to Open Occluded Arteries; HR, hazard ratio; PCI, percutaneous coronary intervention; TIMI, thrombolysis in myocardial infarction.

## Acknowledgments

This article is dedicated to the memory of the late Prof. Steen Husted.

## Sources of Funding

The PLATO study was funded by AstraZeneca. Support for the analysis and interpretation of results and preparation of the manuscript was provided through funds to the Uppsala Clinical Research Center and Duke Clinical Research Institute as part of the Clinical Study Agreement.

## Disclosures

James reports institutional research grant, honoraria, and consultant/advisory board fee from AstraZeneca; institutional research grant and consultant/advisory board fee from Medtronic; institutional research grants and honoraria from The Medicines Company; and consultant/advisory board fees from Janssen and Bayer. Latic reports institutional research grants from AstraZeneca. Budaj reports consulting fees from AstraZeneca, Bayer, Bristol Myers Squibb/Pfizer,

GlaxoSmithKline, Sanofi-Aventis, Bayer, and Novartis; investigator fees from AstraZeneca, Sanofi-Aventis, GlaxoSmithKline, Novartis, Bristol Myers Squibb/Pfizer, and Eisai; and honoraria for lectures from AstraZeneca, Bristol Myers Squibb/Pfizer, GlaxoSmithKline, Sanofi-Aventis, and Novartis. Cornel reports consulting fees from Amgen and AstraZeneca. Katus reports personal fees from AstraZeneca, Bayer Vital, and Roche Diagnostics. Keltai has no potential conflicts to report. Kontny reports consultancy fees/honoraria for lectures, advisory board membership, and fee for research work from AstraZeneca; and advisory board membership and consultancy fees from Merck & Co. Lewis reports departmental grants for performing trials from AstraZeneca and MSD; and honoraria and speaker fees from Pfizer and Bristol-Myers Squibb. Storey reports institutional research grants, consultancy fees, and honoraria from AstraZeneca; institutional research grants and consultancy fees from PlaqueTec; consultancy fees and honoraria from Bayer; and consultancy fees from Actelion, Avacta, Bristol-Myers Squibb/Pfizer, Novartis, Thromboserin, and Idorsia. Himmelmann reports being an employee of AstraZeneca. Wallentin reports institutional research grants from AstraZeneca, Bristol-

**Table 5.** Outcomes of Ticagrelor Versus Clopidogrel According to DM/CKD Status, With Poor Glycemic Control Defined by HbA1c and CKD Defined by the Creatinine-Cystatin C CKD-EPI Equation

DM/CKD Subgroup	Ticagrelor Patients (N)	Clopidogrel Patients (N)	Ticagrelor Event Rate, N (%)	Clopidogrel Event Rate, N (%)	HR (95% CI)	P Value Interaction
<b>Cardiovascular death/MI/stroke</b>						<b>0.265</b>
DM+/CKD+	633	631	105 (22.66)	158 (36.57)	0.68 (0.53–0.88)	
DM–/CKD+	344	390	49 (19.68)	74 (27.22)	0.77 (0.54–1.11)	
DM+/CKD–	2841	2886	267 (11.79)	313 (13.64)	0.87 (0.74–1.03)	
DM–/CKD–	2877	2797	191 (8.31)	201 (8.99)	0.92 (0.76–1.13)	
<b>Cardiovascular death</b>						<b>0.257</b>
DM+/CKD+	633	631	57 (11.52)	98 (20.70)	0.63 (0.45–0.87)	
DM–/CKD+	344	390	25 (9.42)	40 (13.63)	0.74 (0.45–1.23)	
DM+/CKD–	2841	2886	103 (4.34)	112 (4.62)	0.96 (0.73–1.25)	
DM–/CKD–	2877	2797	57 (2.39)	64 (2.75)	0.87 (0.61–1.24)	
<b>MI</b>						<b>0.734</b>
DM+/CKD+	633	631	53 (11.28)	77 (17.53)	0.71 (0.50–1.00)	
DM–/CKD+	344	390	29 (11.57)	40 (14.61)	0.84 (0.52–1.36)	
DM+/CKD–	2841	2886	165 (7.24)	192 (8.31)	0.87 (0.71–1.08)	
DM–/CKD–	2877	2797	124 (5.36)	134 (5.97)	0.89 (0.70–1.14)	
<b>All-cause death</b>						<b>0.481</b>
DM+/CKD+	633	631	68 (13.75)	106 (22.39)	0.70 (0.51–0.95)	
DM–/CKD+	344	390	28 (10.55)	44 (14.99)	0.74 (0.46–1.20)	
DM+/CKD–	2841	2886	113 (4.76)	125 (5.15)	0.94 (0.73–1.21)	
DM–/CKD–	2877	2797	64 (2.68)	81 (3.48)	0.77 (0.56–1.07)	
<b>Stroke</b>						<b>0.293</b>
DM+/CKD+	633	631	15 (3.08)	12 (2.57)	1.33 (0.62–2.85)	
DM–/CKD+	344	390	5 (1.90)	6 (2.06)	0.95 (0.29–3.12)	
DM+/CKD–	2841	2886	33 (1.40)	41 (1.70)	0.82 (0.52–1.30)	
DM–/CKD–	2877	2797	29 (1.22)	17 (0.73)	1.67 (0.92–3.05)	
<b>Major bleeding</b>						<b>0.143</b>
DM+/CKD+	633	631	74 (20.61)	74 (20.51)	1.03 (0.75–1.42)	
DM–/CKD+	344	390	43 (23.41)	43 (19.59)	1.12 (0.74–1.71)	
DM+/CKD–	2841	2886	307 (16.44)	322 (16.72)	0.97 (0.83–1.14)	
DM–/CKD–	2877	2797	272 (14.15)	212 (10.97)	1.28 (1.07–1.54)	
<b>Non-CABG-related major bleeding</b>						<b>0.782</b>
DM+/CKD+	633	631	48 (12.89)	40 (10.69)	1.29 (0.84–1.96)	
DM–/CKD+	344	390	23 (12.06)	21 (9.15)	1.36 (0.75–2.45)	
DM+/CKD–	2841	2886	93 (4.71)	87 (4.23)	1.12 (0.84–1.50)	
DM–/CKD–	2877	2797	95 (4.71)	66 (3.29)	1.39 (1.02–1.91)	

The model is adjusted for age, sex, BMI, heart rate, prior myocardial infarction, hypertension, dyslipidemia, angina pectoris, smoking status, previous PCI or CABG, type of ACS, and randomized treatment. ACS indicates acute coronary syndrome; BMI, body mass index; CABG, coronary artery bypass graft; CKD, chronic kidney disease; CKD-EPI, chronic kidney disease epidemiology collaboration; DM, diabetes mellitus; HR, hazard ratio; MI, myocardial infarction; PCI, percutaneous coronary intervention.

Myers Squibb/Pfizer, Boehringer Ingelheim, GlaxoSmithKline, Merck & Co, and Roche Diagnostics; consultancy fees from Abbott; and holds 2 patents involving GDF-15 licensed to Roche Diagnostics (EP2047275B1 and US8951742B2). Angiolillo

reports payments as an individual for (1) consulting fee or honorarium from Amgen, Aralez, AstraZeneca, Bayer, Biosensors, Boehringer Ingelheim, Bristol-Myers Squibb, Chiesi, Daiichi-Sankyo, Eli Lilly, Haemonetics, Janssen, Merck, PLx



Pharma, Pfizer, Sanofi, and The Medicines Company; (2) participation in review activities from CeloNova and St. Jude Medical. He has also received institutional payments for grants from Amgen, AstraZeneca, Bayer, Biosensors, CeloNova, CSL Behring, Daiichi-Sankyo, Eisai, Eli-Lilly, Gilead, Janssen, Matsutani Chemical Industry Co., Merck, Novartis, Osprey Medical, and Renal Guard Solutions; and is the recipient of funding from the Scott R. MacKenzie Foundation and the NIH/NCATS Clinical and Translational Science Award to the University of Florida UL1 TR000064 and NIH/NHGRI U01 HG007269, outside the submitted work. Franchi reports payments as an individual for consulting fee or honorarium from AstraZeneca and Sanofi.

## References

- Ferreiro JL, Angiolillo DJ. Diabetes and antiplatelet therapy in acute coronary syndrome. *Circulation*. 2011;123:798–813.
- Capodanno D, Angiolillo DJ. Antithrombotic therapy in patients with chronic kidney disease. *Circulation*. 2012;125:2649–2661.
- Bonello L, Angiolillo DJ, Aradi D, Sibbing D. P2Y12-ADP receptor blockade in chronic kidney disease patients with acute coronary syndromes. *Circulation*. 2018;138:1582–1596.
- Franchi F, Angiolillo DJ. Novel antiplatelet agents in acute coronary syndrome. *Nat Rev Cardiol*. 2015;12:30–47.
- Capodanno D, Alfonso F, Levine GN, Valgimigli M, Angiolillo DJ. ACC/AHA versus ESC guidelines on dual antiplatelet therapy: JACC guideline comparison. *J Am Coll Cardiol*. 2018;72:2915–2931.
- Sherwood MW, Wiviott SD, Peng SA, Roe MT, Delemos J, Peterson ED, Wang TY. Early clopidogrel versus prasugrel use among contemporary STEMI and NSTEMI patients in the US: insights from the National Cardiovascular Data Registry. *J Am Heart Assoc*. 2014;3:e000849. DOI: 10.1161/JAHA.114.000849.
- Bueno H, Sinnaeve P, Annemans L, Danchin N, Licour M, Medina J, Pocock S, Sanchez-Covisa J, Storey RF, Jukema JW, Zeymer U, Van de Werf F; EPICOR Investigators. Opportunities for improvement in anti-thrombotic therapy and other strategies for the management of acute coronary syndromes: insights from EPICOR, an international study of current practice patterns. *Eur Heart J Acute Cardiovasc Care*. 2016;5:3–12.
- Angiolillo DJ, Bernardo E, Sabate M, Jimenez-Quevedo P, Costa MA, Palazuelos J, Hernandez-Antolin R, Moreno R, Escaned J, Alfonso F, Banuelos C, Guzman LA, Bass TA, Macaya C, Fernandez-Ortiz A. Impact of platelet reactivity on cardiovascular outcomes in patients with type 2 diabetes mellitus and coronary artery disease. *J Am Coll Cardiol*. 2007;50:1541–1547.
- Angiolillo DJ, Jakubowski JA, Ferreiro JL, Tello-Montoliu A, Rollini F, Franchi F, Ueno M, Darlington A, Desai B, Moser BA, Sugidachi A, Guzman LA, Bass TA. Impaired responsiveness to the platelet P2Y12 receptor antagonist clopidogrel in patients with type 2 diabetes and coronary artery disease. *J Am Coll Cardiol*. 2014;64:1005–1014.
- Erlinge D, Varenhorst C, Braun OO, James S, Winters KJ, Jakubowski JA, Brandt JT, Sugidachi A, Siegbahn A, Wallentin L. Patients with poor responsiveness to thienopyridine treatment or with diabetes have lower levels of circulating active metabolite, but their platelets respond normally to active metabolite added ex vivo. *J Am Coll Cardiol*. 2008;52:1968–1977.
- Angiolillo DJ, Fernandez-Ortiz A, Bernardo E, Ramirez C, Sabate M, Jimenez-Quevedo P, Hernandez R, Moreno R, Escaned J, Alfonso F, Banuelos C, Costa MA, Bass TA, Macaya C. Platelet function profiles in patients with type 2 diabetes and coronary artery disease on combined aspirin and clopidogrel treatment. *Diabetes*. 2005;54:2430–2435.
- Gremmel T, Muller M, Steiner S, Seidinger D, Koppensteiner R, Kopp CW, Panzer S. Chronic kidney disease is associated with increased platelet activation and poor response to antiplatelet therapy. *Nephrol Dial Transplant*. 2013;28:2116–2122.
- Baber U, Bander J, Karajgikar R, Yadav K, Hadi A, Theodoropoulos K, Gukathasan N, Roy S, Sayeneni S, Scott SA, Kovacic JC, Yu J, Sartori S, Mehran R, Uribarri J, Badimon JJ, Muntner P, Moreno P, Kini AS, Sharma SK. Combined and independent impact of diabetes mellitus and chronic kidney disease on residual platelet reactivity. *Thromb Haemost*. 2013;110:118–123.
- Franchi F, Rollini F, Angiolillo DJ. Defining the link between chronic kidney disease, high platelet reactivity, and clinical outcomes in clopidogrel-treated patients undergoing percutaneous coronary intervention. *Circ Cardiovasc Interv*. 2015;8:e002760.
- Angiolillo DJ, Bernardo E, Capodanno D, Vivas D, Sabate M, Ferreiro JL, Ueno M, Jimenez-Quevedo P, Alfonso F, Bass TA, Macaya C, Fernandez-Ortiz A. Impact of chronic kidney disease on platelet function profiles in diabetes mellitus patients with coronary artery disease taking dual antiplatelet therapy. *J Am Coll Cardiol*. 2010;55:1139–1146.
- Engwenyu LR, Franchi F, Rollini F, Cho JR, DeGroat C, Bhatti M, Alobaidi Z, Ferrante E, Jakubowski JA, Sugidachi A, Zenni M, Bass TA, Angiolillo DJ. Impact of chronic kidney disease on platelet P2Y12 receptor signalling in patients with type 2 diabetes mellitus. *Thromb Haemost*. 2017;117:201–203.
- Mangiacapra F, Cavallari I, Barbato E, Ricottini E, Patti G, Vizzi V, D'Ambrosio A, De Bruyne B, Wijns W, Di Sciascio G. Impact of chronic kidney disease on platelet reactivity and outcomes of patients receiving clopidogrel and undergoing percutaneous coronary intervention. *Am J Cardiol*. 2014;113:1124–1129.
- Tello-Montoliu A, Ferreiro JL, Kodali MK, Ueno M, Tomasello SD, Rollini F, Capodanno D, Darlington A, Patel R, Desai B, Guzman LA, Bass TA, Angiolillo DJ. Impact of renal function on clopidogrel-induced antiplatelet effects in coronary artery disease patients without diabetes mellitus. *J Thromb Thrombolysis*. 2013;36:14–17.
- Baber U, Mehran R, Kirtane AJ, Gurbel PA, Christodoulidis G, Maehara A, Witzensbichler B, Weisz G, Rinaldi MJ, Metzger DC, Henry TD, Cox DA, Duffy PL, Mazzaferri EL Jr, Xu K, Parise H, Brodie BR, Stuckey TD, Stone GW. Prevalence and impact of high platelet reactivity in chronic kidney disease: results from the Assessment of Dual Antiplatelet Therapy with Drug-Eluting Stents registry. *Circ Cardiovasc Interv*. 2015;8:e001683.
- Baber U, Farkouh ME, Arbel Y, Muntner P, Dangas G, Mack MJ, Hamza TH, Mehran R, Fuster V. Comparative efficacy of coronary artery bypass surgery vs. percutaneous coronary intervention in patients with diabetes and multivessel coronary artery disease with or without chronic kidney disease. *Eur Heart J*. 2016;37:3440–3447.
- Drexler H, Zanolin D, Vonbank A, Rein P, Saely CH. Impaired kidney function is a diabetes risk equivalent in patients with established coronary artery disease. *Diabetes*. 2014;63(supplement 1):A3 [Abstract].
- Wallentin L, Becker RC, Budaj A, Cannon CP, Emanuelsson H, Held C, Horrow J, Husted S, James S, Katus H, Mahaffey KW, Scirica BM, Skene A, Steg PG, Storey RF, Harrington RA; PLATO Investigators, Freij A, Thorsen M. Ticagrelor versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med*. 2009;361:1045–1057.
- James S, Angiolillo DJ, Cornel JH, Erlinge D, Husted S, Kontny F, Maya J, Nicolau JC, Spinar J, Storey RF, Stevens SR, Wallentin L; PLATO Study Group. Ticagrelor vs. clopidogrel in patients with acute coronary syndromes and diabetes: a substudy from the PLATElet inhibition and patient Outcomes (PLATO) trial. *Eur Heart J*. 2010;31:3006–3016.
- James S, Budaj A, Aylward P, Buck KK, Cannon CP, Cornel JH, Harrington RA, Horrow J, Katus H, Keltai M, Lewis BS, Parikh K, Storey RF, Szummer K, Wojdyla D, Wallentin L. Ticagrelor versus clopidogrel in acute coronary syndromes in relation to renal function: results from the Platelet Inhibition and Patient Outcomes (PLATO) trial. *Circulation*. 2010;122:1056–1067.
- Inker LA, Astor BC, Fox CH, Isakova T, Lash JP, Peralta CA, Kurella Tamura M, Feldman HI. KDOQI US commentary on the 2012 KDIGO clinical practice guideline for the evaluation and management of CKD. *Am J Kidney Dis*. 2014;63:713–735.
- Inker LA, Schmid CH, Tighiouart H, Eckfeldt JH, Feldman HI, Greene T, Kusek JW, Manzi J, Van Lente F, Zhang YL, Coresh J, Levey AS; CKD-EPI Investigators. Estimating glomerular filtration rate from serum creatinine and cystatin C. *N Engl J Med*. 2012;367:20–29.
- Ueno M, Ferreiro JL, Tomasello SD, Capodanno D, Tello-Montoliu A, Kodali M, Secheran N, Dharmashankar K, Alissa R, Capranzano P, Desai B, Charlton RK, Bass TA, Angiolillo DJ. Functional profile of the platelet P2Y1(2) receptor signalling pathway in patients with type 2 diabetes mellitus and coronary artery disease. *Thromb Haemost*. 2011;105:730–732.
- Chang H, Yanachkov IB, Michelson AD, Li Y, Barnard MR, Wright GE, Frelinger AL III. Agonist and antagonist effects of diadenosine tetraphosphate, a platelet dense granule constituent, on platelet P2Y1, P2Y12 and P2X1 receptors. *Thromb Res*. 2010;125:159–165.
- Jankowski V, Gunthner T, Hergert-Rosenthal S, Zidek W, Jankowski J. Dinucleoside polyphosphates and uremia. *Semin Dial*. 2009;22:396–399.
- Leblond F, Guevin C, Demers C, Pellerin I, Gascon-Barre M, Pichette V. Downregulation of hepatic cytochrome P450 in chronic renal failure. *J Am Soc Nephrol*. 2001;12:326–332.
- Dreisbach AW, Lertora JJ. The effect of chronic renal failure on drug metabolism and transport. *Expert Opin Drug Metab Toxicol*. 2008;4:1065–1074.
- Jankowski J, Hagemann J, Yoon MS, van der Giet M, Stephan N, Zidek W, Schluter H, Tepel M. Increased vascular growth in hemodialysis patients induced by platelet-derived diadenosine polyphosphates. *Kidney Int*. 2001;59:1134–1141.

33. Bruck K, Stel VS, Gambaro G, Hallan S, Volzke H, Arnlov J, Kastarinen M, Guessous I, Vinhas J, Stengel B, Brenner H, Chudek J, Romundstad S, Tomson C, Gonzalez AO, Bello AK, Ferrieres J, Palmieri L, Browne G, Capuano V, Van Biesen W, Zoccali C, Gansevoort R, Navis G, Rothenbacher D, Ferraro PM, Nitsch D, Wanner C, Jager KJ; European CKD Burden Consortium. CKD prevalence varies across the European general population. *J Am Soc Nephrol*. 2016;27:2135–2147.
34. Geiss LS, Wang J, Cheng YJ, Thompson TJ, Barker L, Li Y, Albright AL, Gregg EW. Prevalence and incidence trends for diagnosed diabetes among adults aged 20 to 79 years, United States, 1980–2012. *JAMA*. 2014;312:1218–1226.
35. Wanner C, Inzucchi SE, Lachin JM, Fitchett D, von Eynatten M, Matthews M, Johansen OE, Woerle HJ, Broedl UC, Zinman B; EMPA-REG OUTCOME Investigators. Empagliflozin and progression of kidney disease in type 2 diabetes. *N Engl J Med*. 2016;375:323–334.
36. Heerspink HJ, Perkins BA, Fitchett DH, Husain M, Cherney DZ. Sodium glucose cotransporter 2 inhibitors in the treatment of diabetes mellitus: cardiovascular and kidney effects, potential mechanisms, and clinical applications. *Circulation*. 2016;134:752–772.
37. Storey RF, Angiolillo DJ, Patil SB, Desai B, Ecob R, Husted S, Emanuelsson H, Cannon CP, Becker RC, Wallentin L. Inhibitory effects of ticagrelor compared with clopidogrel on platelet function in patients with acute coronary syndromes: the PLATO (PLATElet inhibition and patient Outcomes) PLATELET substudy. *J Am Coll Cardiol*. 2010;56:1456–1462.
38. Sweeny JM, Angiolillo DJ, Franchi F, Rollini F, Waksman R, Raveendran G, Dangas G, Khan ND, Carlson GF, Zhao Y, Teng R, Mehran R. Impact of diabetes mellitus on the pharmacodynamic effects of ticagrelor versus clopidogrel in troponin-negative acute coronary syndrome patients undergoing ad hoc percutaneous coronary intervention. *J Am Heart Assoc*. 2017;6:e005650. DOI: 10.1161/JAHA.117.005650.
39. Clavijo LC, Maya J, Carlson G, Angiolillo DJ, Teng R, Caplan R, Price MJ. Platelet inhibition with ticagrelor versus clopidogrel in Hispanic patients with stable coronary artery disease with or without diabetes mellitus. *Cardiovasc Revasc Med*. 2015;16:450–454.
40. Deharo P, Pankert M, Quilici J, Bonnet G, Bassez C, Verdier V, Morange P, Alessi MC, Bonnet JL, Cuisset T. Chronic kidney disease has a significant impact on platelet inhibition of new P2Y12 inhibitors. *Int J Cardiol*. 2015;184:428–430.
41. Barbieri L, Pergolini P, Verdoia M, Rolla R, Nardin M, Marino P, Bellomo G, Suryapranata H, De Luca G; Novara Atherosclerosis Study Group. Platelet reactivity in patients with impaired renal function receiving dual antiplatelet therapy with clopidogrel or ticagrelor. *Vascul Pharmacol*. 2016;79:11–15.
42. Vilahur G, Gutierrez M, Casani L, Varela L, Capdevila A, Pons-Llado G, Carreras F, Carlsson L, Hidalgo A, Badimon L. Protective effects of ticagrelor on myocardial injury after infarction. *Circulation*. 2016;134:1708–1719.
43. Baber U, Chandrasekhar J, Sartori S, Aquino M, Kini AS, Kapadia S, Weintraub W, Muhlestein JB, Vogel B, Faggioni M, Farhan S, Weiss S, Strauss C, Toma C, DeFranco A, Baker BA, Keller S, Efron MB, Henry TD, Rao S, Pocock S, Dangas G, Mehran R. Associations between chronic kidney disease and outcomes with use of prasugrel versus clopidogrel in patients with acute coronary syndrome undergoing percutaneous coronary intervention: a report from the PROMETHEUS study. *JACC Cardiovasc Interv*. 2017;10:2017–2025.
44. Desai RJ, Spoendlin J, Mogun H, Gagne JJ. Contemporary time trends in use of antiplatelet agents among patients with acute coronary syndrome and comorbid diabetes mellitus or chronic kidney disease. *Pharmacotherapy*. 2017;37:1322–1327.

# SUPPLEMENTAL MATERIAL

## Appendix

### PLATO Investigators (A-Z)

Alexander Nikolaevich Parkhomenko (Research Institute of Cardiology, Kiev, UA)

Ali Oto (Hacettepe Univ School of Medicine, Ankara, TR)

Allan Skene (Worldwide Clinical Trials, Inc; Nottingham, GB)

Andrzej Budaj (Grochowski Hospital, Warsaw, PL)

Anneli Freij (AstraZeneca, Mölndal, SE)

Anwar Santoso (Udayana School of Medicine, Denpasar, ID)

Armando (García Castillo, Hospital de Cardiología UMAE, Nuevo Leon, MX)

Basil S Lewis (Lady Davis Carmel Medical Center, Haifa, IL)

Bernhard Meier (University Hospital, Bern, CH)

Cheuk Man Yu (Chinese University of Hong Kong, HK)

Christopher P Cannon (TIMI study group, Brigham and Women's Hospital, Boston, MA, US)

Dato Seri Robaayah Zambahari (National Heart Institute, Kuala Lumpur, MY)

Delon Wu Wu (Chang Gung Memorial Hospital-Linkou, Tao-Yuan, TW)

Diego Ardissino (Ospedale Maggiore di Parma, Parma, IT)

Dimitar Raev (Medical Institute, Ministry of Interior, Sofia, BG)

Dimitrios Kremastinos (Attikon University General Hospital, Athens, GR)

Douglas Weaver (Henry Ford Heart & Vascular Inst, Detroit, MI, US)

Ernesto Paolasso (Estudios Clinicos Latin America, Rosario, AR)

Evangelos Giannitsis (University of Heidelberg, Heidelberg, DE)

Frederic Kontny (Volvat Medical Center, Oslo, NO)

Freek Verheugt (UMC St Radboud, Nijmegen, NL)

Gerald Maurer (Medical University of Vienna, Vienna, AT)

Hugo Katus (Universitätsklinik Heidelberg, Heidelberg, DE)

Håkan Emanuelsson (AstraZeneca, Mölndal, SE)

Jan H Cornel (Medisch Centrum Alkmaar, Alkmaar, NL)

Jay Horrow (AstraZeneca, Wilmington, DE, US)

Jean-Pierre Bassand (University Hospital Jean Minjoz, Besancon, FR)

Jindrich Spinar (University Hospital Brno, Brno, CZ)

João Morais (Santo Andrés Hospital, Leiria, PT)

Jose Lopez Sendon (Hospital Universitario La Paz, Madrid, ES)

José Nicolau (Heart Institute (InCor), São Paulo, BR)

Ki-Bae Seung (Kangnam St. Mary's Hospital, Seoul, KR)

Lars Wallentin (Uppsala Clinical Research Center, Uppsala, SE)

Lim Soo Teik (Heart Centre, Singapore, SG)

Magda Heras (Hospital Clinic of Barcelona, Barcelona, ES)

Marc J Claeys (University Hospital Antwerp, Edegem, BE)

Marc Sabatine (Brigham & Women's Hospital, Boston, MA, US)

Marius Vintila (St Pantelimon University Hospital, Bucharest, RO)

Matyas Keltai (Semmelweis University, Budapest, HU)

Mikhail Ruda (Cardiology Research Center, Moscow, RU)

Mona Thorsén (AstraZeneca, Mölndal, SE)

Neil Kleiman (Baylor College of Medicine, Houston, TX, US)

Noe Babilonia (Philippine Heart Center, Quezon City, PH)

Patrick Commerford (Groote Schuur Hospital, Cape Town, ZA)

Paul Gurbel (Sinai Hospital, Baltimore, MD, US)

Phil Aylward (Flinders Medical Centre, Bedford Park, AU)

Philippe Gabriel Steg (Hospital Bichat-Claude Bernard, Paris, FR)

Pierre Thérioux (Montreal Heart Institute, Montreal, CA)

Piyamitr Sritara (Ramathibodi Hospital, Bangkok, TH)

Prem Pais (St.John's Medical College, Bangalore, IN)

Richard Becker (Duke Clinical Research Institute, Durham, NC, US)

Riitta Lassila (Helsinki University Central Hospital, FI)

Robert A Harrington (Duke Clinical Research Institute, Durham, NC, US)

Robert F Storey (Clinical Science Centre, Northern General Hospital and University of Sheffield, Sheffield, UK)

Runlin Gao (Cardiovascular Inst & Fu Wai Hosp, Beijing, CN)

Steen Husted (Aarhus University Hospital, Aarhus, DK)

Stefan K James (Uppsala Clinical Research Center and Uppsala University Hospital, Uppsala, SE)

Tibor Duris (Hospital Nove Zamky, Nove Zamky, SK)

Zaza Chapichadze (Institute of Cardiology, Tblisi, GE)