



UNIVERSITY OF LEEDS

This is a repository copy of *Tail asymptotic behavior of the supremum of a class of chi-square processes*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/148318/>

Version: Accepted Version

Article:

Ji, L orcid.org/0000-0002-7790-7765, Liu, P and Robert, S (2019) Tail asymptotic behavior of the supremum of a class of chi-square processes. *Statistics and Probability Letters*, 154. 108551. ISSN 0167-7152

<https://doi.org/10.1016/j.spl.2019.07.001>

© 2019, Elsevier B.V. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

TAIL ASYMPTOTIC BEHAVIOR OF THE SUPREMUM OF A CLASS OF CHI-SQUARE PROCESSES

LANPENG JI, PENG LIU, AND STEPHAN ROBERT

Abstract: We analyze in this paper the supremum of a class of chi-square processes over non-compact intervals, which can be seen as a multivariate counterpart of the generalized weighted Kolmogorov-Smirnov statistic. The boundedness and the exact tail asymptotic behavior of the supremum are derived. As examples, the chi-square process generated from the Brownian bridge and the fractional Brownian motion are discussed.

Key Words: chi-square process; exact asymptotics; Brownian bridge; Pickands constants; Piterbarg constants

AMS Classification: Primary 60G15; secondary 60G70

1. INTRODUCTION

Let $X(t), t \geq 0$, be a Gaussian process with almost surely (a.s.) continuous sample paths. For a sequence of constants $\{b_i\}_{i=1}^n$ satisfying

$$1 = b_1 = \cdots = b_k > b_{k+1} \geq \cdots \geq b_n > 0$$

we define the chi-square process as

$$(1) \quad \chi_{\mathbf{b}}^2(t) = \sum_{i=1}^n b_i^2 X_i^2(t), \quad t \geq 0,$$

where X_i 's are independent copies of X . The supremum of chi-square process appears naturally as limiting test statistic in various statistical models; see, e.g., [1, 2, 3, 4]. It also plays an important role in reliability applications in the engineering sciences, see [5, 6, 7, 8] and the references therein.

Of interest in applied probability and statistics is the tail asymptotics of

$$\mathbb{P} \left\{ \sup_{t \in \mathcal{T}} \chi_{\mathbf{b}}^2(t) > u \right\}, \quad u \rightarrow \infty$$

for an interval $\mathcal{T} \subset \mathbb{R}_+$, provided that

$$(2) \quad \sup_{t \in \mathcal{T}} \chi_{\mathbf{b}}^2(t) < \infty \quad a.s..$$

Numerous contributions have been devoted to the study of the tail asymptotics of the supremum of chi-square processes over compact intervals \mathcal{T} ; see, e.g., [6, 9, 10, 11] and the references therein, where the technique used is to transform the supremum of chi-square process into the supremum of a special Gaussian random field. We refer to, e.g., [12, 13, 14, 15, 16, 17, 18] for more discussions on the tail asymptotics (or excursion probability) of Gaussian and related fields.

In this paper, we are interested in the analysis of a class of weighted locally stationary chi-square processes defined by

$$\sup_{t \in \mathcal{T}} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)}, \quad \text{with } \mathcal{T} = (0, 1) \text{ or } (0, 1],$$

where $w(\cdot)$ is some positive continuous function definable on the non-compact set \mathcal{T} , and the generic process X is the locally stationary Gaussian process. More precisely, $X(t), t \in \mathcal{T}$, is a centered Gaussian process with a.s. continuous sample paths, unit variance and correlation function $r(\cdot, \cdot)$ satisfying

$$(3) \quad \lim_{h \rightarrow 0} \frac{1 - r(t, t+h)}{K^2(|h|)} = C(t)$$

uniformly in $t \in I$, for all the compact interval I in \mathcal{T} , where $K(\cdot)$ is a positive regularly varying function at 0 with index $\alpha/2 \in (0, 1]$, and $C(\cdot)$ is a positive continuous function satisfying

$$\lim_{t \rightarrow 0} C(t) = \infty \quad \text{or} \quad \lim_{t \rightarrow 1} C(t) = \infty.$$

We refer to [19] for more discussions on such locally stationary Gaussian processes.

Our motivation for considering the supremum of the weighted locally stationary chi-square processes over the non-compact interval $\mathcal{T} = (0, 1)$ or $(0, 1]$ is from its potential applications in statistics. For instance, in its univariate framework (with $n = 1$) the following generalized weighted Kolmogorov-Smirnov statistic

$$W_w := \sup_{t \in (0,1)} \frac{|\overline{B}(t)|}{w(t)}, \quad \text{with} \quad \overline{B}(t) = \frac{B(t)}{\sqrt{t(1-t)}}, \quad t \in (0, 1),$$

has been discussed in the statistics literature, see, e.g., [20], where B is the standard Brownian bridge with variance function $\text{Var}(B(t)) = t(1-t), t \in [0, 1]$ and w is a suitably chosen weight function such that

$$(4) \quad W_w < \infty \quad a.s..$$

We refer to [20, 21, 22, 23] for further discussions on the generalized weighted Kolmogorov-Smirnov statistic.

An interesting theoretical question is to find sufficient and necessary conditions on w so that the a.s. finiteness of (4) holds. It is shown in [20][Theorem 3.3, Theorem 4.2.3] (see also [21][Theorem 26.3]) that

$$(5) \quad W_w < \infty \quad a.s. \quad \Leftrightarrow \quad \int_0^1 \frac{1}{t(1-t)} e^{-cw^2(t)} dt < \infty \quad \text{for some } c > 0.$$

One of the main results displayed in Theorem 3.1 shows necessary and sufficient conditions on the weight function w under which it holds that

$$(6) \quad \sup_{t \in \mathcal{T}} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} < \infty \quad a.s..$$

This extends the result of (5). Furthermore, for certain w satisfying (6) we derive in Theorem 3.3 the exact asymptotics of

$$(7) \quad \mathbb{P} \left\{ \sup_{t \in \mathcal{T}} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\}, \quad u \rightarrow \infty.$$

As an important application of Theorem 3.3, we obtain in Corollary 3.4 the tail asymptotics of the supremum of the chi-square process generated from the Brownian bridge. It is worth mentioning that this tail asymptotic result is new even for the univariate (i.e., $n = 1$) case. As a second example, the chi-square process generated by the fractional Brownian motion is discussed.

We expect that the derived results will have interesting applications in statistics or beyond.

Organization of the rest of the paper: In Section 2 we present a preliminary result which is a tailored version of Theorem A.1 in [19]. The main results are given in Section 3, followed by examples. All the proofs are displayed in Section 4.

2. PRELIMINARIES

This section concerns a result derived in [19], which is crucial for the derivation of (6). Based on the discussions therein, we shall consider $\int_0^{1/2} (C(s))^{1/\alpha} ds = \infty$ or $\int_{1/2}^1 (C(s))^{1/\alpha} ds = \infty$. For this purpose, of crucial importance is the following function

$$f(t) = \int_{1/2}^t (C(s))^{1/\alpha} ds, \quad t \in (0, 1).$$

We denote by $\overleftarrow{f}(t), t \in (f(0), f(1))$ the inverse function of $f(t), t \in (0, 1)$. Further, for any $d > 0$, let $s_{j,d}^{(1)} = \overleftarrow{f}(jd)$, $j \in \mathbb{N} \cup \{0\}$ if $f(1) = \infty$, and let $s_{j,d}^{(0)} = \overleftarrow{f}(-jd)$, $j \in \mathbb{N} \cup \{0\}$ if $f(0) = -\infty$. Denote $\Delta_{j,d}^{(1)} = [s_{j-1,d}^{(1)}, s_{j,d}^{(1)}], j \in \mathbb{N}$ and $\Delta_{j,d}^{(0)} = [s_{j,d}^{(0)}, s_{j-1,d}^{(0)}], j \in \mathbb{N}$, which give a partition of $[1/2, 1)$ in the case $f(1) = \infty$ and a partition of $(0, 1/2]$ in the case $f(0) = -\infty$, respectively. Moreover, let $q(u) = \overleftarrow{K}(u^{-1/2})$ be the inverse function of $K(\cdot)$ at point $u^{-1/2}$ (assumed to exist asymptotically).

The following (scenario-dependent) restrictions on the positive continuous weight function w^2 and the correlation function $r(\cdot, \cdot)$ of X play a crucial role. Let therefore $S \in \{0, 1\}$.

Condition A(S): The weight function w^2 is monotone in a neighbourhood of S and satisfies $\lim_{t \rightarrow S} w^2(t) = \infty$.

Condition B(S): Suppose that there exists some constant $d_0 > 0$ such that

$$\limsup_{j \rightarrow \infty} \sup_{t \neq s \in \Delta_{j,d_0}^{(S)}} \frac{1 - r(t, s)}{K^2(|f(t) - f(s)|)} < \infty,$$

and when $\alpha = 2$ and $k = 1$, assume further

$$K^2(|t|) = O(t^2), \quad t \rightarrow 0.$$

Condition C(S): Suppose that there exists some constant $d_0 > 0$ such that

$$\liminf_{j \rightarrow \infty} \inf_{t \neq s \in \Delta_{j,d_0}^{(S)}} \frac{1 - r(t, s)}{K^2(|f(t) - f(s)|)} > 0.$$

Moreover, there exist $j_0, l_0 \in \mathbb{N}$, $M_0, \beta > 0$, such that for $j \geq j_0, l \geq l_0$,

$$(8) \quad \sup_{s \in \Delta_{j+l,d_0}^{(S)}, t \in \Delta_{j,d_0}^{(S)}} |r(s, t)| < M_0 l^{-\beta}.$$

For the subsequent discussions we present a tailored version of Theorem A.1 of [19], focusing on $|f(S)| = \infty$. We define

$$I_w(S) = \left| \int_{1/2}^S (C(t))^{1/\alpha} \frac{(w(t))^{k-2}}{q(w^2(t))} e^{-\frac{w^2(t)}{2}} dt \right|.$$

Theorem 2.1. *Let $X(t), t \in (0, 1)$, be a centered locally stationary Gaussian process with a.s. continuous sample paths, unit variance and correlation function $r(\cdot, \cdot)$ satisfying (3) and $r(s, t) < 1$ for $s \neq t \in (0, 1)$. Suppose further that, for $S = 0$ or 1 , we have $|f(S)| = \infty$ and **A(S)**, **B(S)**, **C(S)** are satisfied. Then*

$$(9) \quad \mathbb{P} \left\{ \chi_{\mathbf{b}}^2(t) \leq w^2(t) \text{ ultimately as } t \rightarrow S \right\} = 0, \quad \text{or } 1$$

according to

$$I_w(S) = \infty, \quad \text{or } < \infty.$$

3. MAIN RESULTS

In this section, we first give a criteria for (6) to hold and then display the exact asymptotics of (7) for different types of w such that (6) is valid.

3.1. Analysis of (6). Denote by $E(0) = (0, 1/2]$ and $E(1) = [1/2, 1)$. Recall that $S \in \{0, 1\}$. Under the conditions of Theorem 2.1, we have that if $I_w(S) < \infty$, then

$$\sup_{t \in E(S)} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} < \infty \quad \text{a.s.},$$

however, when $I_w(S) = \infty$ we only see that

$$\sup_{t \in E(S)} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} \geq 1 \quad \text{a.s..}$$

Apparently, the above is not informative for the validity of (6). On the other hand, it is easily shown that

$$\sup_{t \in E(S)} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} < \infty \quad \text{a.s.} \quad \Leftrightarrow \quad \sup_{t \in E(S)} \frac{|X(t)|}{w(t)} < \infty \quad \text{a.s.},$$

which means that, instead of the condition $I_w(S) = \infty$ in Theorem 2.1, a more accurate condition that is independent of n, k should be possible to ensure that (6) holds. Inspired by this fact and given the importance of (6), we provide below a sufficient and necessary condition for (6) to hold.

Define, for any constant $c > 0$ and any positive continuous function w

$$J_{c,w}(S) = \left| \int_{1/2}^S (C(t))^{1/\alpha} e^{-cw^2(t)} dt \right|.$$

Below is our first principal result, a criterion for (6), which is a generalization of (5).

Theorem 3.1. *Under the conditions of Theorem 2.1 we have*

$$\sup_{t \in E(S)} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} < \infty \quad \text{a.s.} \quad \Leftrightarrow \quad J_{c,w}(S) < \infty \quad \text{for some } c > 0.$$

Next we illustrate the criteria presented in Theorem 3.1 by an example of a weighted chi-square process with generic process being the normalized standard Brownian bridge, which further provides us with a clear comparison between $I_w(S)$ and $J_{c,w}(S)$.

Example 3.2. *Let $X(t) = \bar{B}(t)$, $t \in (0, 1)$, and, with $\rho_1 > 0, \rho_2 \in \mathbb{R}$, define*

$$(10) \quad w_{\rho_1, \rho_2}^2(t) = 2\rho_1 \ln \ln \left(\frac{e^2}{t(1-t)} \right) + 2\rho_2 \ln \ln \ln \left(\frac{e^2}{t(1-t)} \right), \quad t \in (0, 1).$$

First note that for the normalized standard Brownian bridge

$$(11) \quad \lim_{h \rightarrow 0} \frac{1 - \mathbb{E}(\bar{B}(t)\bar{B}(t+h))}{|h|} = \frac{1}{2t(1-t)}$$

holds uniformly in $t \in I$, for any compact interval I in $(0, 1)$. This means that \bar{B} is a locally stationary Gaussian process with

$$K(h) = \sqrt{|h|}, \quad \alpha = 1, \quad q(u) = u^{-1}.$$

Furthermore,

$$f(t) = \int_{1/2}^t \frac{1}{2s(1-s)} ds = \frac{1}{2} \ln \left(\frac{t}{1-t} \right)$$

implying that $f(1) = -f(0) = \infty$. Moreover, by the proof of Corollary 2.6 in [19] we have that conditions **B(S)** and **C(S)** are satisfied by $\bar{B}(t), t \in (0, 1)$, and $\mathbb{E}(\bar{B}(t), \bar{B}(s)) < 1$ for $s \neq t, s, t \in (0, 1)$. Thus, all the conditions of Theorem 3.1 are fulfilled.

Next, on one hand, we have

$$\frac{1}{t(1-t)} (w_{\rho_1, \rho_2}(t))^k e^{-\frac{w_{\rho_1, \rho_2}^2(t)}{2}} \sim \frac{Q}{t(1-t) \left(\ln \left(\frac{1}{t(1-t)} \right) \right)^{\rho_1} \left(\ln \ln \left(\frac{e^2}{t(1-t)} \right) \right)^{\rho_2 - k/2}}$$

as $t \rightarrow 0$ or $t \rightarrow 1$, with Q some positive constant. Thus, elementary calculations show that

$$I_w(0) = I_w(1) = \int_{1/2}^1 \frac{(w_{\rho_1, \rho_2}(t))^k}{t(1-t)} e^{-\frac{w_{\rho_1, \rho_2}^2(t)}{2}} dt < \infty$$

holds if and only if

$$(12) \quad \rho_1 > 1, \quad \text{or } \rho_1 = 1 \text{ and } \rho_2 > 1 + k/2.$$

On the other hand, we can show that the functions $w_{\rho_1, \rho_2}(t)$ satisfying that $\exists c > 0$ such that $J_{c,w}(S) < \infty$ are not restricted to the ones satisfying (12). In fact, since for any $\rho_1 > 0$ there exists some c such that $\rho_1 > \frac{1}{2c}$, we have that

$$\begin{aligned} J_{c,w}(0) = J_{c,w}(1) &= \int_{1/2}^1 \frac{1}{t(1-t)} e^{-cw_{\rho_1, \rho_2}^2(t)} dt \\ &\leq \int_{1/2}^1 \frac{1}{t(1-t) \left(\ln \left(\frac{1}{t(1-t)} \right) \right)^{2c\rho_1} \left(\ln \ln \left(\frac{e^2}{t(1-t)} \right) \right)^{2c\rho_2}} dt < \infty \end{aligned}$$

holds for any $\rho_2 \in \mathbb{R}$. Thus, we conclude from Theorem 3.1 that

$$\sup_{t \in (0,1)} \frac{\chi_{\mathbf{b}}^2(t)}{w_{\rho_1, \rho_2}^2(t)} < \infty \quad a.s.$$

holds for any $\rho_1 > 0$ and $\rho_2 \in \mathbb{R}$.

The exact tail asymptotics of $\sup_{t \in (0,1)} \frac{\chi_{\mathbf{b}}^2(t)}{w_{\rho_1, \rho_2}^2(t)}$ will be discussed in next section.

3.2. Asymptotics of (7). For those w such that (6) holds, of interest is the exact tail asymptotic behavior of $\sup_{t \in \mathcal{T}} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)}$. Actually, as we have seen, the behavior of w around 0 and 1 plays a crucial role for the finiteness in (6). However, this does not apply to the tail asymptotics of $\sup_{t \in \mathcal{T}} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)}$. It turns out that only the probability mass in the neighborhood of minimizer of w contribute to the tail asymptotics, indicating that the other part of the process including the part around 0 or 1 can be neglected. As discussed in [24], the weight function is introduced when constructing the Goodness-of-Fit test which is intended to emphasize a specific region of the domain. With these motivations, for the tail asymptotics we shall consider the following two types of w :

Assumption F1: The function w attains its minimum at finite distinct inner points $\{t_i\}_{i=1}^m$ of \mathcal{T} , and

$$(13) \quad w(t_i + t) = w(t_i) + a_i |t_i|^{\beta_i} (1 + o(1)), \quad t \rightarrow t_i$$

holds for some positive constants $a_i, \beta_i > 0, i = 1, 2, \dots, m$.

Assumption F2: The function w attains its minimum at all points on disjoint intervals $[c_i, d_i] \subseteq \mathcal{T}, i = 1, 2, \dots, m$ (i.e., w is a constant on these intervals).

Under assumption **F1**, we need additional conditions which are stated below. Recall $q(u) = \overleftarrow{K}(u^{-1/2})$. It follows that $q(u)$ is a regularly varying function at infinity with index $-1/\alpha$ which can be further expressed as $q(u) =$

$u^{-1/\alpha}L(u^{-1/2})$, with $L(\cdot)$ a slowly varying function at 0. Denote further $\beta = \max_{1 \leq i \leq m} \beta_i$. According to the values of $L(u^{-1/2})$ as $u \rightarrow \infty$, we consider the following three scenarios:

C1(β): $\beta > \alpha$, or $\beta = \alpha$ and $\lim_{u \rightarrow \infty} L(u^{-1/2}) = 0$;

C2(β): $\beta = \alpha$ and $\lim_{u \rightarrow \infty} L(u^{-1/2}) = \mathcal{L} \in (0, \infty)$;

C3(β): $\beta < \alpha$, or $\beta = \alpha$ and $\lim_{u \rightarrow \infty} L(u^{-1/2}) = \infty$.

Before displaying our results, we introduce two important constants. One is the *Pickands constant* defined by

$$\mathcal{H}_{2H} = \lim_{S \rightarrow \infty} \frac{1}{S} \mathbb{E} \left(\exp \left(\sup_{t \in [0, S]} \left(\sqrt{2} B_H(t) - t^{2H} \right) \right) \right),$$

with $B_H(t), t \in \mathbb{R}$, a standard fractional Brownian motion (fBm) defined on \mathbb{R} with Hurst index $H \in (0, 1]$. And the other one is the *Piterbarg constant* defined by

$$\mathcal{P}_{2H}^d = \lim_{\lambda \rightarrow \infty} \mathbb{E} \left(\exp \left(\sup_{t \in [-\lambda, \lambda]} \left(\sqrt{2} B_H(t) - (1+d)|t|^{2H} \right) \right) \right), \quad d > 0.$$

We refer to [13, 25, 26, 27, 28, 29] for the properties and generalizations of the Pickands-Piterbarg type constants. In what follows, α will play a similar role as $2H$. Moreover, We shall use the standard notation for asymptotic equivalence of two functions f and h . Specifically, we write $f(x) \sim h(x)$, if $\lim_{x \rightarrow a} f(x)/h(x) = 1$ ($a \in \mathbb{R} \cup \{\infty\}$), and further, write $f(x) = o(h(x))$, if $\lim_{x \rightarrow a} f(x)/h(x) = 0$.

Let $K = \{1 \leq i \leq m : \beta_i = \beta\}$ and $K^c = \{1 \leq i \leq m : \beta_i < \beta\}$. Below is our second principal result.

Theorem 3.3. *Let $\frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)}, t \in \mathcal{T}$, be the weighted locally stationary chi-square process considered in Theorem 2.1 such that (6) holds. We have:*

(i). *If **F1** is satisfied, then, as $u \rightarrow \infty$,*

$$\mathbb{P} \left\{ \sup_{t \in \mathcal{T}} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\} \sim \left(\prod_{i=k+1}^n (1 - b_i^2)^{-1/2} \right) \mathcal{M}(u) \Upsilon_k(w^2(t_1)u),$$

where (with the convention $\prod_{i=p}^q = 1$ if $q < p$)

$$(14) \quad \Upsilon_k(u) := \mathbb{P} \{ \chi_{k,1}^2(0) > u \} = \frac{2^{(2-k)/2}}{\Gamma(k/2)} u^{k/2-1} \exp \left(-\frac{u}{2} \right), \quad u > 0,$$

and

$$\mathcal{M}(u) = \begin{cases} 2 \left(\sum_{i \in K} a_i^{-1/\beta} (C(t_i))^{1/\alpha} \right) (w(t_1))^{2/\alpha-1/\beta} \Gamma(1/\beta+1) \mathcal{H}_\alpha(q(u))^{-1} u^{-1/\beta}, & \text{for } \mathbf{C1}(\beta), \\ \sum_{i \in K} \mathcal{P}_\alpha^{a_i(w(t_1)C(t_i))^{-1} \mathcal{L}^\alpha} + \#K^c, & \text{for } \mathbf{C2}(\beta), \\ m, & \text{for } \mathbf{C3}(\beta). \end{cases}$$

(ii). *If **F2** is satisfied, then, as $u \rightarrow \infty$,*

$$\mathbb{P} \left\{ \sup_{t \in \mathcal{T}} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\} \sim \left(\prod_{i=k+1}^n (1 - b_i^2)^{-1/2} \right) \left(\sum_{j=1}^m \int_{c_j}^{d_j} (C(t))^{1/\alpha} dt \right) \mathcal{H}_\alpha(q(w^2(c_1)u))^{-1} \Upsilon_k(w^2(c_1)u).$$

We conclude this section with two applications of Theorem 3.3. The first one is on the weighted locally stationary chi-square process discussed in Example 3.2, and the second one concerns the weighted locally stationary chi-square process with generic process X being a normalized standard fBm.

Corollary 3.4. Let $\frac{\chi_{\mathbf{b}}^2(t)}{w_{\rho_1, \rho_2}^2(t)}$, $t \in (0, 1)$, with $\rho_1 > 0$ and $\rho_2 \in \mathbb{R}$, be the weighted locally stationary chi-square process discussed in Example 3.2. We have, as $u \rightarrow \infty$, if $\rho_2 \geq -\rho_1 \ln \ln(4e^2)$, then

$$\mathbb{P} \left\{ \sup_{t \in (0,1)} \frac{\chi_{\mathbf{b}}^2(t)}{w_{\rho_1, \rho_2}^2(t)} > u \right\} \sim \left(\prod_{i=k+1}^n (1 - b_i^2)^{-1/2} \right) \mathcal{M}(u) \Upsilon_k(2A_1 u),$$

where $A_1 = \rho_1 \ln \ln(4e^2) + \rho_2 \ln \ln \ln(4e^2)$ and

$$\mathcal{M}(u) = \begin{cases} 2A_1 \sqrt{\frac{\pi \ln(4e^2) \ln \ln(4e^2)}{\rho_1 \ln \ln(4e^2) + \rho_2}} u^{1/2}, & \text{for } \rho_2 > -\rho_1 \ln \ln(4e^2), \\ 2\Gamma(1/4)A_1 \left(\frac{\ln \ln(4e^2) (\ln(4e^2))^2}{8\rho_1} \right)^{1/4} u^{3/4}, & \text{for } \rho_2 = -\rho_1 \ln \ln(4e^2), \end{cases}$$

and if $\rho_2 < -\rho_1 \ln \ln(4e^2)$, then

$$\mathbb{P} \left\{ \sup_{t \in (0,1)} \frac{\chi_{\mathbf{b}}^2(t)}{w_{\rho_1, \rho_2}^2(t)} > u \right\} \sim 2A_2 \left(\prod_{i=k+1}^n (1 - b_i^2)^{-1/2} \right) \rho_1^{-1} Q \sqrt{-2\pi\rho_2} u^{1/2} \Upsilon_k(2A_2 u),$$

where $A_2 = \rho_2(\ln(-\rho_2) - \ln(\rho_1) - 1)$ and

$$Q = \frac{1}{2t_1 - 1} \ln \left(\frac{e^2}{t_1(1 - t_1)} \right), \quad t_1 = 1/2 + \sqrt{1/4 - e^{2 - e^{-\rho_2/\rho_1}}}.$$

Next, we consider $B_H(t)$, $t \geq 0$, to be the standard fBm with Hurst index $H \in (0, 1)$ and covariance function

$$\text{Cov}(B_H(s), B_H(t)) = \frac{1}{2} (|s|^{2H} + |t|^{2H} - |s - t|^{2H}), \quad s, t \geq 0.$$

Denote by $\bar{B}_H(t) = B_H(t)/t^H$, $t \in (0, 1]$ the normalized standard fBm defined on $(0, 1]$. Further, for any $\rho > 0$ and $\varepsilon \in (0, 1)$, we define

$$(15) \quad w_{\rho, \varepsilon}^2(t) = \begin{cases} \rho \ln \ln(e^2/t), & \text{for } t \in (0, \varepsilon), \\ \rho \ln \ln(e^2/\varepsilon), & \text{for } t \in [\varepsilon, 1]. \end{cases}$$

We have the following result.

Corollary 3.5. Let $\frac{\chi_{\mathbf{b}}^2(t)}{w_{\rho, \varepsilon}^2(t)}$, $t \in (0, 1]$, be a weighted chi-square process with generic process $\bar{B}_H(t)$, $t \in (0, 1]$ and $w_{\rho, \varepsilon}^2$ given in (15). Then, we have, as $u \rightarrow \infty$,

$$\mathbb{P} \left\{ \sup_{t \in (0,1)} \frac{\chi_{\mathbf{b}}^2(t)}{w_{\rho, \varepsilon}^2(t)} > u \right\} \sim \left(\prod_{i=k+1}^n (1 - b_i^2)^{-1/2} \right) (-\ln(\varepsilon)) (\ln \ln(e^2/\varepsilon) \rho/2)^{\frac{1}{2H}} \\ \times \mathcal{H}_{2H} u^{\frac{1}{2H}} \Upsilon_k(\rho \ln \ln(e^2/\varepsilon) u).$$

4. PROOFS

This section is devoted to the proof of all the results presented in Section 3.

Proof of Theorem 3.1: Note that $t^{k/2-1}(q(t))^{-1}$ is a positive regularly varying function at ∞ with index $\kappa = k/2 - 1 + 1/\alpha \geq 0$. Thus, by Potter bound (e.g., [30])

$$c_1 t^{\kappa-1} \leq t^{k/2-1}(q(t))^{-1} \leq c_2 t^{\kappa+1}, \quad t \geq c_3,$$

holds for some constants $c_1, c_2, c_3 > 0$, which, together with the fact that $w^2(t) \rightarrow \infty$ as $t \rightarrow S$, leads to

$$(16) \quad Q_1 e^{-w^2(t)} \leq \frac{(w(t))^{k-2}}{q(w^2(t))} e^{-\frac{w^2(t)}{2}} \leq Q_2 e^{-\frac{w^2(t)}{3}}$$

for all t approaching S , with some positive constants Q_1, Q_2 . Therefore, if $J_{c,w}(S) < \infty$ holds for some $c > 0$, then, by (16),

$$I_{\sqrt{3cw}}(S) < \infty.$$

This together with ii) of Theorem 2.1 yields that

$$\limsup_{t \rightarrow S} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} \leq 3c \quad a.s.$$

showing that

$$\sup_{t \in E(S)} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} < \infty \quad a.s.$$

On the other hand, if $J_{c,w}(S) = \infty$ for all $c > 0$, then, by (16), $I_{\sqrt{cw}}(S) = \infty$. Thus, by iii) of Theorem 2.1

$$\limsup_{t \rightarrow S} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} \geq c \quad a.s.$$

holds for all $c > 0$, implying that

$$\sup_{t \in E(S)} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} = \infty \quad a.s.$$

This completes the proof. \square

We show next a version of the Borell-TIS inequality for chi-square process, which will play a key role in the proof of Theorem 3.3. We refer to, e.g., [12, 31] for discussions on the Borell-TIS inequality for Gaussian random fields. Denote below $\mathcal{S} \subseteq \mathbb{R}$ to be any fixed interval.

Lemma 4.1. *Let $\chi_{\mathbf{b}}^2(t), t \in \mathcal{S}$, be a chi-square process with generic centered Gaussian process X which has a.s. continuous sample paths and variance function denoted by $\sigma_X^2(t)$. If*

$$\sup_{t \in \mathcal{S}} X(t) < \infty \quad a.s.,$$

then there exists some positive constant Q such that for all $u > Q^2$ we have

$$(17) \quad \mathbb{P} \left\{ \sup_{t \in \mathcal{S}} \chi_{\mathbf{b}}^2(t) > u \right\} \leq \exp \left(- \frac{(\sqrt{u} - Q)^2}{2 \sup_{t \in \mathcal{S}} \sigma_X^2(t)} \right).$$

Proof of Lemma 4.1: Using the classical approach when dealing with chi-square processes as, e.g., in [4, 13, 19], we introduce a particular Gaussian random field, namely,

$$(18) \quad Y_{\mathbf{b}}(t, \boldsymbol{\theta}) := \sum_{i=1}^n b_i X_i(t) v_i(\boldsymbol{\theta}), \quad (t, \boldsymbol{\theta}) \in \mathcal{D} := \mathcal{S} \times [-\pi, \pi] \times [-\pi/2, \pi/2]^{n-2},$$

where $\boldsymbol{\theta} = (\theta_2, \theta_3, \dots, \theta_n)$, and $v_n(\boldsymbol{\theta}) = \sin(\theta_n), v_{n-1}(\boldsymbol{\theta}) = \sin(\theta_{n-1}) \cos(\theta_n), \dots, v_1(\boldsymbol{\theta}) = \cos(\theta_n) \cdots \cos(\theta_2)$ are spherical coordinates. In view of [13], for any $u > 0$

$$(19) \quad \mathbb{P} \left(\sup_{t \in \mathcal{S}} \chi_{\mathbf{b}}^2(t) > u \right) = \mathbb{P} \left(\sup_{(t, \boldsymbol{\theta}) \in \mathcal{D}} Y_{\mathbf{b}}(t, \boldsymbol{\theta}) > \sqrt{u} \right).$$

Since the variance function of $Y_{\mathbf{b}}$ satisfies for $u > 0$

$$\mathbb{E} \left((Y_{\mathbf{b}}(t, \boldsymbol{\theta}))^2 \right) = \sigma_X^2(t) \left(1 - (1 - b_n^2) \sin^2(\theta_n) - \sum_{i=k+1}^{n-1} (1 - b_i^2) \left(\prod_{j=i+1}^n \cos^2(\theta_j) \right) \sin^2(\theta_i) \right),$$

we have

$$(20) \quad \sup_{(t, \boldsymbol{\theta}) \in \mathcal{D}} \mathbb{E} \left((Y_{\mathbf{b}}(t, \boldsymbol{\theta}))^2 \right) \leq \sup_{t \in \mathcal{S}} \sigma_X^2(t).$$

Then, by (19) and the Borell-TIS inequality for Gaussian random fields (cf. [12][Theorem 2.1.1]) we conclude that (17) holds with $Q = \mathbb{E} \left(\sup_{(t, \boldsymbol{\theta}) \in \mathcal{D}} Y_{\mathbf{b}}(t, \boldsymbol{\theta}) \right) < \infty$. This completes the proof. \square

The next result concerns an upper bound for the tails of double-sup of the locally stationary chi-square processes, which will also play a key role in the proof of Theorem 3.3.

Lemma 4.2. *Let $\chi_{\mathbf{b}}^2(t), t \in \mathcal{S}$, be a chi-square process with the generic centered locally stationary Gaussian process X which has a.s. continuous sample paths. If further the correlation function of X satisfies*

$$(21) \quad r(s, t) < 1 \text{ for any } s \neq t \in \mathcal{S},$$

then, for any compact intervals $\mathcal{S}_1, \mathcal{S}_2 \subset \mathcal{S}$ such that $\mathcal{S}_1 \cap \mathcal{S}_2 = \emptyset$ we have

$$\mathbb{P} \left\{ \sup_{t \in \mathcal{S}_1} \chi_{\mathbf{b}}^2(t) > u, \sup_{t \in \mathcal{S}_2} \chi_{\mathbf{b}}^2(t) > u \right\} \leq \exp \left(- \frac{(2\sqrt{u} - Q)^2}{2(2 + 2\eta)} \right).$$

for all $u > Q^2$, with some constant $Q > 0$ and $\eta \in (0, 1)$.

Proof of Lemma 4.2: Using the expression of $Y_{\mathbf{b}}(t, \boldsymbol{\theta})$ given in (18), we have

$$\begin{aligned} \mathbb{P} \left\{ \sup_{t \in \mathcal{S}_1} \chi_{\mathbf{b}}^2(t) > u, \sup_{t \in \mathcal{S}_2} \chi_{\mathbf{b}}^2(t) > u \right\} &= \mathbb{P} \left\{ \sup_{(t, \boldsymbol{\theta}) \in \mathcal{D}_1} Y_{\mathbf{b}}(t, \boldsymbol{\theta}) > \sqrt{u}, \sup_{(t, \boldsymbol{\theta}) \in \mathcal{D}_2} Y_{\mathbf{b}}(t, \boldsymbol{\theta}) > \sqrt{u} \right\} \\ &\leq \mathbb{P} \left\{ \sup_{(t, \boldsymbol{\theta}) \in \mathcal{D}_1, (t', \boldsymbol{\theta}') \in \mathcal{D}_2} (Y_{\mathbf{b}}(t, \boldsymbol{\theta}) + Y_{\mathbf{b}}(t', \boldsymbol{\theta}')) > 2\sqrt{u} \right\} \end{aligned}$$

where

$$\mathcal{D}_i = \mathcal{S}_i \times [-\pi, \pi] \times [-\pi/2, \pi/2]^{n-2}, \quad i = 1, 2.$$

By (21) we have that there exists some $\eta \in (0, 1)$ such that

$$\begin{aligned} \mathbb{E} \left((Y_{\mathbf{b}}(t, \boldsymbol{\theta}) + Y_{\mathbf{b}}(t', \boldsymbol{\theta}'))^2 \right) &= \mathbb{E} \left((Y_{\mathbf{b}}(t, \boldsymbol{\theta}))^2 \right) + \mathbb{E} \left((Y_{\mathbf{b}}(t', \boldsymbol{\theta}'))^2 \right) + 2 \sum_{i=1}^n \mathbb{E} (X_i(t) X_i(t')) b_i^2 v_i(\boldsymbol{\theta}) v_i(\boldsymbol{\theta}') \\ &\leq 2 + 2\eta \sum_{i=1}^n b_i^2 v_i(\boldsymbol{\theta}) v_i(\boldsymbol{\theta}') \\ (22) \quad &\leq 2 + 2\eta, \quad (t, \boldsymbol{\theta}) \in \mathcal{D}_1, (t', \boldsymbol{\theta}') \in \mathcal{D}_2. \end{aligned}$$

Consequently, by the Borell-TIS inequality

$$\mathbb{P} \left\{ \sup_{(t, \boldsymbol{\theta}) \in \mathcal{D}_1, (t', \boldsymbol{\theta}') \in \mathcal{D}_2} (Y_{\mathbf{b}}(t, \boldsymbol{\theta}) + Y_{\mathbf{b}}(t', \boldsymbol{\theta}')) > 2\sqrt{u} \right\} \leq \exp \left(- \frac{(2\sqrt{u} - Q)^2}{2(2 + 2\eta)} \right)$$

for all $u > \frac{Q^2}{4}$, with $Q = \mathbb{E} \left(\sup_{(t, \boldsymbol{\theta}) \in \mathcal{D}_1, (t', \boldsymbol{\theta}') \in \mathcal{D}_2} (Y_{\mathbf{b}}(t, \boldsymbol{\theta}) + Y_{\mathbf{b}}(t', \boldsymbol{\theta}')) \right) < \infty$. Thus, the claim follows. This completes the proof. \square

Proof of Theorem 3.3: Without loss of generality, we show the proof only for the case where $\mathcal{T} = (0, 1)$.

(i). Let $\rho > 0$ be a sufficiently small constant such that

$$[t_i - \rho, t_i + \rho] \cap [t_j - \rho, t_j + \rho] = \emptyset, \quad \text{for all } i \neq j.$$

Further, denote $\mathcal{T}_\rho = \mathcal{T} \setminus \bigcup_{i=1}^m [t_i - \rho, t_i + \rho]$. It follows from the Bonferroni inequality (e.g., [32]) that

$$\begin{aligned}
(23) \quad & \sum_{i=1}^m p_i(u) + \mathbb{P} \left\{ \sup_{t \in \mathcal{T}_\rho} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\} \\
& \geq \mathbb{P} \left\{ \sup_{t \in \mathcal{T}} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\} \\
& \geq \sum_{i=1}^m p_i(u) - \sum_{1 \leq i < j \leq m} \mathbb{P} \left\{ \sup_{t \in [t_i - \rho, t_i + \rho]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u, \sup_{t \in [t_j - \rho, t_j + \rho]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\},
\end{aligned}$$

where

$$p_i(u) = \mathbb{P} \left\{ \sup_{t \in [t_i - \rho, t_i + \rho]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\}.$$

We first focus on the asymptotics of $p_i(u)$ as $u \rightarrow \infty$. Denote

$$Y(t) = \frac{w(t_1)}{w(t)} X(t), \quad t \in \mathcal{T}.$$

We have

$$p_i(u) = \mathbb{P} \left\{ \sup_{t \in [t_i - \rho, t_i + \rho]} \sum_{l=1}^n b_l^2 Y_l^2(t) > w^2(t_1) u \right\}, \quad 1 \leq i \leq m,$$

where $\{Y_l\}_{l=1}^n$ is a sequence of independent copies of Gaussian process Y . It can be shown that, by **F1**, for any $i = 1, 2, \dots, m$,

$$\sigma_Y(t) = \sqrt{\mathbb{E}((Y(t))^2)} = \frac{w(t_1)}{w(t)}, \quad t \in [t_i - \rho, t_i + \rho],$$

attains its maximum which is equal to 1 at the unique point t_i , and further

$$\sigma_Y(t_i + t) = 1 - \frac{a_i}{w(t_1)} |t|^{\beta_i} (1 + o(1)), \quad t \rightarrow 0.$$

Moreover, by (3)

$$1 - \text{Corr}(Y(t_i + t), Y(t_i + s)) = C(t_i) K^2(|t - s|)(1 + o(1)), \quad t \rightarrow 0.$$

Consequently, it follows from [8][Theorem 5.2] that, as $u \rightarrow \infty$,

$$(24) \quad p_i(u) \sim \left(\prod_{l=k+1}^n (1 - b_l^2)^{-1/2} \right) \mathcal{M}_i(\beta_i, u) \Upsilon_k(w^2(t_1)u),$$

where $\Upsilon_k(\cdot)$ is given in (14) and

$$\mathcal{M}_i(\beta_i, u) = \begin{cases} 2a_i^{-1/\beta_i} (w(t_1))^{2/\alpha - 1/\beta_i} (C(t_i))^{1/\alpha} \Gamma(1/\beta_i + 1) \mathcal{H}_\alpha(q(u))^{-1} u^{-1/\beta_i}, & \text{for } \mathbf{C1}(\beta_i), \\ \mathcal{P}_\alpha^{a_i(w(t_1)C(t_i))^{-1} \mathcal{L}^\alpha} & \text{for } \mathbf{C2}(\beta_i), \\ 1, & \text{for } \mathbf{C3}(\beta_i). \end{cases}$$

In the sequel, we discuss the three scenarios **C1**(β), **C2**(β), **C3**(β) one-by one.

C1(β). Using the fact that $\beta = \max_{i=1}^m \beta_i$, we have that

$$\mathcal{M}_j(\beta_j, u) = o(\mathcal{M}_i(\beta_i, u)), \quad u \rightarrow \infty$$

for any $i \in K$ and $j \in K^c$. This implies that

$$(25) \quad \sum_{i=1}^m p_i(u) \sim \sum_{i \in K} p_i(u) \sim \left(\prod_{l=k+1}^n (1 - b_l^2)^{-1/2} \right) \mathcal{M}(u) \Upsilon_k(w^2(t_1)u),$$

where $\mathcal{M}(\cdot)$ is given in (15). On the other hand, it follows directly from Lemma 4.1 that

$$\mathbb{P} \left\{ \sup_{t \in \mathcal{T}_\rho} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\} \leq \exp \left(- \frac{\inf_{t \in \mathcal{T}_\rho} w^2(t) (\sqrt{u} - Q)^2}{2} \right)$$

holds for all $u > Q^2$, with Q some positive constant. Since further, by **F1**,

$$\inf_{t \in \mathcal{T}_\rho} w^2(t) > w^2(t_1),$$

we have that

$$(26) \quad \mathbb{P} \left\{ \sup_{t \in \mathcal{T}_\rho} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\} = o(\mathcal{M}(u) \Upsilon_k(w^2(t_1)u)), \quad u \rightarrow \infty.$$

Moreover, since for any $i \neq j$

$$\begin{aligned} & \mathbb{P} \left\{ \sup_{t \in [t_i - \rho, t_i + \rho]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u, \quad \sup_{t \in [t_j - \rho, t_j + \rho]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\} \\ & \leq \mathbb{P} \left\{ \sup_{t \in [t_i - \rho, t_i + \rho]} \chi_{\mathbf{b}}^2(t) > w^2(t_1)u, \quad \sup_{t \in [t_j - \rho, t_j + \rho]} \chi_{\mathbf{b}}^2(t) > w^2(t_1)u \right\}. \end{aligned}$$

we have from Lemma 4.2 that, for all u large,

$$\mathbb{P} \left\{ \sup_{t \in [t_i - \rho, t_i + \rho]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u, \quad \sup_{t \in [t_j - \rho, t_j + \rho]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\} \leq \exp \left(- \frac{(2w(t_1)\sqrt{u} - Q_{i,j})^2}{2(2 + 2\eta)} \right), \quad 1 \leq i < j \leq m,$$

with $Q_{i,j}$'s some positive constants and $\eta \in (0, 1)$. Therefore, as $u \rightarrow \infty$,

$$(27) \quad \sum_{1 \leq i < j \leq m} \mathbb{P} \left\{ \sup_{t \in [t_i - \rho, t_i + \rho]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u, \quad \sup_{t \in [t_j - \rho, t_j + \rho]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\} = o(\mathcal{M}(u) \Upsilon_k(w^2(t_1)u)).$$

Combining (25)–(27) with (23) we establish the claim of **C1**(β).

C2(β). In this case, we have that (24) holds with

$$\mathcal{M}_i(\beta_i, u) = \begin{cases} \mathcal{P}_\alpha^{a_i(w(t_1)C(t_i))^{-1}\mathcal{L}^\alpha}, & i \in K \\ 1, & i \in K^c. \end{cases}$$

Consequently,

$$\sum_{i=1}^m p_i(u) \sim \left(\prod_{l=k+1}^n (1 - b_l^2)^{-1/2} \right) \left(\sum_{i \in K} \mathcal{P}_\alpha^{a_i(w(t_1)C(t_i))^{-1}\mathcal{L}^\alpha} + \#K^c \right) \Upsilon_k(w^2(t_1)u).$$

Note that (26) and (27) still hold. Similarly as the case **C1**(β), we establish the claim of **C2**(β).

C3(β). In this case, we have that (24) holds with

$$\mathcal{M}_i(\beta_i, u) = 1, \quad 1 \leq i \leq m.$$

Consequently,

$$\sum_{i=1}^m p_i(u) \sim m \left(\prod_{l=k+1}^n (1 - b_l^2)^{-1/2} \right) \Upsilon_k(w^2(t_1)u).$$

Similarly as before, the claim of **C3**(β) follows.

(ii). By **F2** we have for any sufficiently small $\varepsilon > 0$ it holds that

$$\inf_{t \in \mathcal{T}_\varepsilon} w(t) > w(c_1), \quad \text{with } \mathcal{T}_\varepsilon = \mathcal{T} \setminus \bigcup_{i=1}^m [c_i - \varepsilon, d_i + \varepsilon].$$

Similarly to (23) we have

$$\begin{aligned}
& \sum_{i=1}^m \mathbb{P} \left\{ \sup_{t \in [c_i - \varepsilon, d_i + \varepsilon]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\} + \mathbb{P} \left\{ \sup_{t \in \mathcal{T}_\varepsilon} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\} \\
(28) \quad & \geq \mathbb{P} \left\{ \sup_{t \in \mathcal{T}} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\} \\
& \geq \sum_{i=1}^m \mathbb{P} \left\{ \sup_{t \in [c_i, d_i]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\} - \sum_{1 \leq i < j \leq m} \mathbb{P} \left\{ \sup_{t \in [c_i, d_i]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u, \sup_{t \in [c_j, d_j]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\}.
\end{aligned}$$

Next, we have from **F2** that for $1 \leq i \leq m$

$$\begin{aligned}
\mathbb{P} \left\{ \sup_{t \in [c_i, d_i]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\} &= \mathbb{P} \left\{ \sup_{t \in [c_i, d_i]} \chi_{\mathbf{b}}^2(t) > w^2(c_1)u \right\}, \\
\mathbb{P} \left\{ \sup_{t \in [c_i - \varepsilon, d_i + \varepsilon]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\} &\leq \mathbb{P} \left\{ \sup_{t \in [c_i - \varepsilon, d_i + \varepsilon]} \chi_{\mathbf{b}}^2(t) > w^2(c_1)u \right\}.
\end{aligned}$$

It is noted that the result in Theorem 2.1 of [19] also holds when $g(t) = 0$. Thus, it follows from that result, as $u \rightarrow \infty$,

$$\begin{aligned}
\mathbb{P} \left\{ \sup_{t \in [c_i, d_i]} \chi_{\mathbf{b}}^2(t) > w^2(c_1)u \right\} &\sim \prod_{j=k+1}^n (1 - b_j^2)^{-1/2} \mathcal{H}_\alpha \int_{c_i}^{d_i} (C(t))^{1/\alpha} dt (q(w^2(c_1)u))^{-1} \Upsilon_k(w^2(c_1)u), \\
\mathbb{P} \left\{ \sup_{t \in [c_i - \varepsilon, d_i + \varepsilon]} \chi_{\mathbf{b}}^2(t) > w^2(c_1)u \right\} &\sim \prod_{j=k+1}^n (1 - b_j^2)^{-1/2} \mathcal{H}_\alpha \int_{c_i - \varepsilon}^{d_i + \varepsilon} (C(t))^{1/\alpha} dt (q(w^2(c_1)u))^{-1} \Upsilon_k(w^2(c_1)u).
\end{aligned}$$

Moreover, since

$$\begin{aligned}
& \mathbb{P} \left\{ \sup_{t \in [c_i, d_i]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u, \sup_{t \in [c_j, d_j]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\} \\
&= \mathbb{P} \left\{ \sup_{t \in [c_i, d_i]} \chi_{\mathbf{b}}^2(t) > w^2(c_1)u, \sup_{t \in [c_j, d_j]} \chi_{\mathbf{b}}^2(t) > w^2(c_1)u \right\},
\end{aligned}$$

we have from Lemma 4.2 that, for all u large,

$$\mathbb{P} \left\{ \sup_{t \in [c_i, d_i]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u, \sup_{t \in [c_j, d_j]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\} \leq \exp \left(-\frac{(2w(c_1)\sqrt{u} - Q_{i,j})^2}{2(2 + 2\eta)} \right), \quad 1 \leq i < j \leq m,$$

with $Q_{i,j}$'s some positive constants and $\eta \in (0, 1)$. This implies that

$$\sum_{1 \leq i < j \leq m} \mathbb{P} \left\{ \sup_{t \in [c_i, d_i]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u, \sup_{t \in [c_j, d_j]} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\} = o \left((q(w^2(c_1)u))^{-1} \Upsilon_k(w^2(c_1)u) \right), \quad u \rightarrow \infty.$$

Moreover, Lemma 4.1 gives that

$$\mathbb{P} \left\{ \sup_{t \in \mathcal{T}_\varepsilon} \frac{\chi_{\mathbf{b}}^2(t)}{w^2(t)} > u \right\} = o \left((q(w^2(c_1)u))^{-1} \Upsilon_k(w^2(c_1)u) \right), \quad u \rightarrow \infty.$$

Consequently, by letting $\varepsilon \rightarrow 0$ we conclude that the claim in (ii) is established. This completes the proof. \square

Proof of Corollary 3.4: We have from Example 3.2, for $\rho_1 > 0$ and $\rho_2 \in \mathbb{R}$,

$$\sup_{t \in (0,1)} \frac{\chi_{\mathbf{b}}^2(t)}{w_{\rho_1, \rho_2}^2(t)} < \infty \quad a.s.$$

Furthermore, for the generic locally stationary Gaussian process $X = \overline{B}$ we have

$$(29) \quad C(t) = \frac{1}{2t(1-t)}, \quad K(h) = \sqrt{|h|}, \quad h \in (0, 1).$$

Next, in order to apply Theorem 3.3 we analyze the function $w_{\rho_1, \rho_2}(t)$.

For simplicity, we define

$$f(x) = 2\rho_1 x + 2\rho_2 \ln x, \quad x(t) = \ln \ln \left(\frac{e^2}{t(1-t)} \right).$$

Apparently,

$$w_{\rho_1, \rho_2}^2(t) = f(x(t)), \quad t \in (0, 1), \quad \{x(t) : t \in (0, 1)\} = [\ln \ln(4e^2), \infty).$$

Since

$$\frac{\partial f(x)}{\partial x} = 2\rho_1 + \frac{2\rho_2}{x}, \quad x \in [\ln \ln(4e^2), \infty),$$

the following three different cases will be discussed separately:

$$a). \rho_2 > -\rho_1 \ln \ln(4e^2); \quad b). \rho_2 = -\rho_1 \ln \ln(4e^2); \quad c). \rho_2 < -\rho_1 \ln \ln(4e^2).$$

a). $\rho_2 > -\rho_1 \ln \ln(4e^2)$: In this case, we have

$$f'(x) = \frac{\partial f(x)}{\partial x} > 0, \quad x \in [\ln \ln(4e^2), \infty),$$

which means that $f(x)$ attains its minimum over $[\ln \ln(4e^2), \infty)$ at the unique point $x_0 = \ln \ln(4e^2)$, and $f'(x_0) > 0$.

Since further

$$x'(t) = \frac{\partial x(t)}{\partial t} = \frac{t(1-t)}{\ln \left(\frac{e^2}{t(1-t)} \right)} (t^{-1}(1-t)^{-2} - t^{-2}(1-t)^{-1}),$$

we conclude that the minimizer of $f(x(t))$ over $(0, 1)$ is unique and equal to $t_1 = 1/2$, and $x(t_1) = x_0$, $x'(t_1) = 0$.

Next, we look at the Taylor expansion of $f(x(t))$, $t \in (0, 1)$ at t_1 . Note that

$$\frac{\partial f(x(t))}{\partial t} \Big|_{t=t_1} = 0, \quad \frac{\partial^2 f(x(t))}{\partial t^2} \Big|_{t=t_1} = \frac{\partial f}{\partial x} \Big|_{x=x_0} \frac{\partial^2 x}{\partial t^2} \Big|_{t=t_1} = \frac{16}{\ln(4e^2)} \left(\rho_1 + \frac{\rho_2}{\ln \ln(4e^2)} \right) =: Q_1 > 0.$$

Thus, by Taylor expansion

$$f(x(t)) - f(x(t_1)) = \frac{Q_1}{2} (t - t_1)^2 (1 + o(1)), \quad t \rightarrow t_1$$

which yields that

$$(30) \quad w_{\rho_1, \rho_2}(t) = w_{\rho_1, \rho_2}(t_1) + \frac{Q_1}{4w_{\rho_1, \rho_2}(t_1)} (t - t_1)^2 (1 + o(1)), \quad t \rightarrow t_1.$$

Moreover, by (29) and (30)

$$C(t_1) = 2, \quad \alpha = 1 < 2 = \beta, \quad q(u) = u^{-1}.$$

Since further $\Gamma(1/2 + 1) = 1/2\sqrt{\pi}$ and $\mathcal{H}_1 = 1$, by applying Theorem 3.3 we conclude that the claim in (a) is established.

b). $\rho_2 = -\rho_1 \ln \ln(4e^2)$: In this case, we have that $f(x)$ attains its minimum over $[\ln \ln(4e^2), \infty)$ at the unique point $x_0 = \ln \ln(4e^2)$, but with $f'(x_0) = 0$. Further, the minimizer of $f(x(t))$ over $(0, 1)$ is unique and equal to $t_1 = 1/2$, and $x(t_1) = x_0$, $x'(t_1) = 0$. Thus, we have

$$\frac{\partial f(x(t))}{\partial t} \Big|_{t=t_1} = 0, \quad \frac{\partial^2 f(x(t))}{\partial t^2} \Big|_{t=t_1} = 0.$$

Next, let us calculate higher-order derivatives of $f(x(t))$. We have

$$\begin{aligned}\frac{\partial^3 f(x(t))}{\partial t^3} &= \frac{\partial^3 f}{\partial x^3} \left(\frac{\partial x}{\partial t} \right)^3 + 3 \frac{\partial^2 f}{\partial x^2} \frac{\partial x}{\partial t} \frac{\partial^2 x}{\partial t^2} + \frac{\partial f}{\partial x} \frac{\partial^3 x}{\partial t^3}, \\ \frac{\partial^4 f(x(t))}{\partial t^4} &= \frac{\partial^4 f}{\partial x^4} \left(\frac{\partial x}{\partial t} \right)^4 + 6 \frac{\partial^3 f}{\partial x^3} \left(\frac{\partial x}{\partial t} \right)^2 \frac{\partial^2 x}{\partial t^2} + 3 \frac{\partial^2 f}{\partial x^2} \left(\frac{\partial^2 x}{\partial t^2} \right)^2 + 4 \frac{\partial^2 f}{\partial x^2} \frac{\partial x}{\partial t} \frac{\partial^3 x}{\partial t^3} + \frac{\partial f}{\partial x} \frac{\partial^4 x}{\partial t^4}.\end{aligned}$$

This implies that

$$\frac{\partial^3 f(x(t))}{\partial t^3} \Big|_{t=t_1} = 0, \quad \frac{\partial^4 f(x(t))}{\partial t^4} \Big|_{t=t_1} = 3 \frac{\partial^2 f}{\partial x^2} \Big|_{x=x_0} \left(\frac{\partial^2 x}{\partial t^2} \Big|_{t=t_1} \right)^2 = \frac{384\rho_1}{\ln \ln(4e^2)(\ln(4e^2))^2} =: Q_2 > 0.$$

Therefore, by Taylor expansion we conclude that

$$(31) \quad w_{\rho_1, \rho_2}(t) = w_{\rho_1, \rho_2}(t_1) + \frac{Q_2}{48w_{\rho_1, \rho_2}(t_1)}(t - t_1)^4(1 + o(1)), \quad t \rightarrow t_1.$$

Similarly as in (a), the claim of (b) follows by applying Theorem 3.3.

c). $\rho_2 < -\rho_1 \ln \ln(4e^2)$: In this case, we have that $f(x)$ attains its minimum over $[\ln \ln(4e^2), \infty)$ at an inner point $x_0 = -\rho_2/\rho_1$, for which $f'(x_0) = 0$. Furthermore, the minimizer of $f(x(t))$ over $(0, 1)$ are two distinct points $t_1 = \frac{1 + \sqrt{1 - 4e^{2-\epsilon^{-\rho_2/\rho_1}}}}{2}$ and $t_2 = \frac{1 - \sqrt{1 - 4e^{2-\epsilon^{-\rho_2/\rho_1}}}}{2}$, for which $x'(t_i) \neq 0, i = 1, 2$. Thus, we have, for $i = 1, 2$,

$$\frac{\partial f(x(t))}{\partial t} \Big|_{t=t_i} = 0, \quad \frac{\partial^2 f(x(t))}{\partial t^2} \Big|_{t=t_i} = \frac{\partial^2 f}{\partial x^2} \Big|_{x=x_0} \left(\frac{\partial x}{\partial t} \Big|_{t=t_i} \right)^2 = \frac{2\rho_1^2}{-\rho_2} Q_3 > 0,$$

where, by symmetry of $x(t), t \in (0, 1)$,

$$Q_3 := \left(\frac{\partial x}{\partial t} \Big|_{t=t_1} \right)^2 = \left(\frac{\partial x}{\partial t} \Big|_{t=t_2} \right)^2 > 0.$$

Consequently, by Taylor expansion we conclude that

$$(32) \quad w_{\rho_1, \rho_2}(t) = w_{\rho_1, \rho_2}(t_i) + \frac{\rho_1^2 Q_3}{-2\rho_2 w_{\rho_1, \rho_2}(t_i)}(t - t_i)^2(1 + o(1)), \quad t \rightarrow t_i.$$

In addition,

$$w_{\rho_1, \rho_2}(t_1) = w_{\rho_1, \rho_2}(t_2) = \sqrt{2\rho_2(\ln(-\rho_2) - \ln(\rho_1) - 1)}.$$

Similarly as in (a), the claim of (c) follows by applying Theorem 3.3. This completes the proof. \square

Proof of Corollary 3.5: First note that

$$\lim_{h \rightarrow 0} \frac{1 - \mathbb{E}(\overline{B}_H(t), \overline{B}_H(t+h))}{|h|^{2H}} = \frac{1}{2t^{2H}}$$

holds uniformly in $t \in I$, for any compact interval I in $(0, 1]$. This means that \overline{B}_H is a locally stationary Gaussian process with

$$C(t) = \frac{1}{2t^{2H}}, \quad K(h) = |h|^H, \quad \alpha = 2H.$$

We shall first discuss the finiteness of $\sup_{t \in (0, 1]} \frac{\chi_{\mathbf{b}}^2(t)}{w_{\rho, \epsilon}^2(t)}$, for which we only need to verify the conditions in Theorem 3.1 for the case where $S = 0$. First, condition **A**(0) is satisfied by $w_{\rho, \epsilon}$, and clearly

$$f(0) = \frac{1}{2^{1/(2H)}} \int_{1/2}^0 t^{-1} dt = -\infty.$$

Further, we have from the calculations in the proof of Corollary 2.7 in [19] that $\mathbb{E}(\overline{B}_H(t), \overline{B}_H(s)) < 1$ for $s \neq t$, $s, t \in (0, 1]$, and conditions **B(0)** and **C(0)** are satisfied by $\overline{B}_H(t)$, $t \in (0, 1]$. Moreover, we have that

$$J_{c, w_{\rho, \varepsilon}}(0) \leq \frac{1}{2^{1/(2H)}} \left(\frac{1}{(\ln(e^2/\varepsilon))^{\rho c}} \int_{\varepsilon \wedge 1/2}^{1/2} \frac{1}{t} dt + \int_0^\varepsilon \frac{1}{t (\ln(e^2/t))^{\rho c}} dt \right) < \infty$$

for any $c > 1/\rho$. Consequently, it follows from Theorem 3.1 that

$$\sup_{t \in (0, 1]} \frac{\chi_{\mathbf{b}}^2(t)}{w_{\rho, \varepsilon}^2(t)} < \infty \quad a.s.$$

Next, since by definition $w_{\rho, \varepsilon}$ attains its minimum over $(0, 1]$ on the interval $[\varepsilon, 1]$, we conclude from (ii) of Theorem 3.3 that the claim follows. \square

Acknowledgement: P. Liu was partially supported by the Swiss National Science Foundation Grant 200021-166274.

REFERENCES

- [1] D. Jarušková, “Testing appearance of polynomial trend,” *Extremes*, vol. 2, no. 1, pp. 25–37, 1999.
- [2] D. Jarušková, “Asymptotic behaviour of a test statistic for detection of change in mean of vectors,” *J. Statist. Plann. Inference*, vol. 140, no. 3, pp. 616–625, 2010.
- [3] J. Albin and D. Jarušková, “On a test statistic for linear trend,” *Extremes*, vol. 6, no. 3, pp. 247–258 (2004), 2003.
- [4] D. Jarušková and V. Piterbarg, “Log-likelihood ratio test for detecting transient change,” *Statist. Probab. Lett.*, vol. 81, no. 5, pp. 552–559, 2011.
- [5] G. Lindgren, “Extreme values and crossings for the χ^2 -process and other functions of multidimensional gaussian processes with reliability applications,” *Adv. Appl. Probab.*, vol. 12, pp. 746–774, 1980.
- [6] G. Lindgren, “Slepian models for χ^2 -processes with dependent components with application to envelope upcrossings,” *J. Appl. Probab.*, vol. 26, no. 1, pp. 36–49, 1989.
- [7] E. Hashorva and L. Ji, “Piterbarg theorems for chi-processes with trend,” *Extremes*, vol. 18, no. 1, pp. 37–64, 2015.
- [8] P. Liu and L. Ji, “Extremes of chi-square processes with trend,” *Probab. Math. Statist.*, vol. 36, no. 1, pp. 1–20, 2016.
- [9] D. Konstantinides, V. Piterbarg, and S. Stamatovic, “Gnedenko-type limit theorems for cyclostationary χ^2 -processes,” *Lithuanian Mathematical Journal*, vol. 44, no. 2, pp. 157–167, 2004.
- [10] Z. Tan and E. Hashorva, “Limit theorems for extremes of strongly dependent cyclo-stationary χ -processes,” *Extremes*, vol. 16, no. 2, pp. 241–254, 2013.
- [11] E. Hashorva and L. Ji, “Extremes of $\alpha(\mathbf{t})$ -locally stationary Gaussian random fields,” *Trans. Amer. Math. Soc.*, vol. 368, no. 1, pp. 1–26, 2016.
- [12] R. Adler and J. Taylor, *Random fields and geometry*. Springer Monographs in Mathematics, New York: Springer, 2007.
- [13] V. Piterbarg, *Asymptotic methods in the theory of Gaussian processes and fields*, vol. 148 of *Translations of Mathematical Monographs*. Providence, RI: American Mathematical Society, 1996. Translated from the Russian by V.V. Piterbarg, revised by the author.
- [14] J. Azaïs and M. Wschebor, *Level sets and extrema of random processes and fields*. Hoboken, NJ: John Wiley & Sons Inc., 2009.
- [15] D. Marinucci and S. Vадlamani, “High-frequency asymptotics for Lipschitz–Killing curvatures of excursion sets on the sphere,” *Ann. Appl. Probab.*, vol. 26, pp. 462–506, 2016.
- [16] D. Cheng and Y. Xiao, “Excursion probability of Gaussian random fields on sphere,” *Bermoulli*, vol. 22, pp. 1113–1130, 2016.
- [17] D. Cheng and Y. Xiao, “The mean Euler characteristic and excursion probability of Gaussian random fields with stationary increments,” *Ann. Appl. A Probab.*, vol. 26, pp. 722–759, 2016.
- [18] E. Bernardino, A. Estrade, and J. R. Leon, “A test of Gaussianity based on the Euler characteristic of excursion sets,” *Electron. J. Statist.*, vol. 11, pp. 843–890, 2017.
- [19] P. Liu and L. Ji, “Extremes of locally stationary chi-square processes with trend,” *Stochastic Process. Appl.*, vol. 127, no. 2, pp. 497–525, 2017.

- [20] M. Csörgö, S. Csörgö, L. Horváth, and D. M. Mason, “Weighted empirical and quantile processes,” *Annals of Probability*, vol. 14, no. 1, pp. 31–85, 1986.
- [21] A. DasGupta, *Asymptotic Theory of Statistics and Probability*. Springer Texts in Statistics, Springer-Verlag New York, 2008.
- [22] V. Piterbarg and V. Taran, “Exact asymptotics for large deviation probabilities of normalized Brownian bridge with applications to empirical processes,” *Computers Math. Applic.*, vol. 22, no. 7, pp. 85–91, 1991.
- [23] L. Dümbgen, P. Kolesnyk, and R. A. Wilke, “Bi-log-concave distribution functions,” *Journal of Statistical Planning and Inference*, vol. 184, pp. 1–17, 2017.
- [24] R. Chicheportiche and J.-P. Bouchaud, “Weighted Kolmogorov-Smirnov test: Accounting for the tails,” *Phys. Rev. E*, vol. 86, pp. 041–115, 2012.
- [25] K. Dębicki, “Ruin probability for Gaussian integrated processes,” *Stochastic Processes and their Applications*, vol. 98, no. 1, pp. 151–174, 2002.
- [26] K. Dębicki and M. Mandjes, “Exact overflow asymptotics for queues with many Gaussian inputs,” *J. Appl. Probab.*, vol. 40, no. 3, pp. 704–720, 2003.
- [27] A. Dieker and B. Yakir, “On asymptotic constants in the theory of extremes for Gaussian processes,” *Bernoulli*, vol. 20, no. 3, pp. 1600–1619, 2014.
- [28] L. Bai, K. Dębicki, E. Hashorva, and L. Luo, “On generalised Piterbarg constants,” *Methodol Comput Appl Probab*, vol. 20, pp. 137–164, 2018.
- [29] K. Dębicki and E. Hashorva, “On extremal index of max-stable stationary processes,” *Probability and Mathematical Statistics*, vol. 37, pp. 299–317, 2017.
- [30] N. Bingham, C. Goldie, and J. Teugels, *Regular variation*, vol. 27 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1989.
- [31] G. Samorodnitsky, “Probability tails of Gaussian extrema,” *Stochastic Process. Appl.*, vol. 38, no. 1, pp. 55–84, 1991.
- [32] Z. Michina, “Remarks on Pickands theorem,” *Probability and Mathematical Statistics*, vol. 37, no. 2, pp. 373–393, 2017.

LANPENG JI, SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, WOODHOUSE LANE, LEEDS LS2 9JT, UNITED KINGDOM
E-mail address: `l.ji@leeds.ac.uk`

PENG LIU, DEPARTMENT OF STATISTICS AND ACTUARIAL SCIENCE, UNIVERSITY OF WATERLOO, CANADA
E-mail address: `peng.liu1@uwaterloo.ca`

STEPHAN ROBERT, INSTITUTE FOR INFORMATION AND COMMUNICATION TECHNOLOGIES, SCHOOL OF BUSINESS AND ENGINEERING VAUD (HEIG-VD), UNIVERSITY OF APPLIED SCIENCES OF WESTERN SWITZERLAND, SWITZERLAND
E-mail address: `Stephan.Robert@heig-vd.ch`