

This is a repository copy of *Analysing the genetic architecture of clubroot resistance variation in Brassica napus by associative transcriptomics*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/148586/>

Version: Published Version

Article:

Hejna, Ondrej, Havlickova, Lenka orcid.org/0000-0002-5874-8615, He, Zhesi orcid.org/0000-0001-8335-9876 et al. (2 more authors) (2019) Analysing the genetic architecture of clubroot resistance variation in Brassica napus by associative transcriptomics. *Molecular Breeding*. 112. ISSN 1380-3743

<https://doi.org/10.1007/s11032-019-1021-4>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Analysing the genetic architecture of clubroot resistance variation in *Brassica napus* by associative transcriptomics

Ondrej Hejna · Lenka Havlickova · Zhesi He · Ian Bancroft  · Vladislav Curn

Received: 6 February 2019 / Accepted: 8 July 2019 / Published online: 20 July 2019
© The Author(s) 2019

Abstract Clubroot is a destructive soil-borne pathogen of Brassicaceae that causes significant recurrent reductions in yield of cruciferous crops. Although there is some resistance in oilseed rape (a crop type of the species *Brassica napus*), the genetic basis of that resistance is poorly understood. In this study, we used an associative transcriptomics approach to elucidate the genetic basis of resistance to clubroot pathotype ECD 17/31/31 across a genetic diversity panel of 245 accessions of *B. napus*. A single nucleotide polymorphism (SNP) association analysis was performed with 256,397 SNPs distributed across the genome of *B. napus* and combined with transcript abundance data of 53,889 coding DNA sequence (CDS) gene models. The SNP association analysis identified two major loci (on chromosomes A2 and A3) controlling resistance and seven minor loci. Within these were a total of 86 SNP markers. Altogether, 392 genes were found in these regions. Another 21 genes were implicated as potentially involved in resistance using gene expression marker

(GEM) analysis. After GO enrichment analysis and InterPro functional analysis of the identified genes, 82 candidate genes were identified as having roles in clubroot resistance. These results provide useful information for marker-assisted breeding which could lead to acceleration of pyramiding of multiple clubroot resistance genes in new varieties.

Keywords Association genetics · *Brassica napus* · Clubroot · Transcriptomics

Introduction

Clubroot disease caused by the soil-borne obligate biotrophic pathogen *Plasmodiophora brassicae* is one of the most serious diseases of cruciferous crops, especially the allotetraploid *Brassica napus* ($2n = 38$, AACC) and its diploid progenitor species *Brassica rapa* ($2n = 20$, AA) and *Brassica oleracea* ($2n = 18$, CC) (Dixon 2009; Fredua-Agyeman and Rahman 2016). Two-phase infection by this parasite leads to formation of galls or clubs on the roots and hypocotyls of diseased plants (Hwang et al. 2012). Creation of these club-shaped roots ultimately leads to the interruption of the uptake and flow of water and minerals in roots. This results in wilting, stunted growth, chlorosis and leaf abscission leading even to the death of infected plants (Kageyama and Asano 2009). The symptoms of clubroot disease result in significant reduction of seed yield as well as decreasing of oil content in seed of susceptible cruciferous plants worldwide (Dixon 2009). The average loss of yield in

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s11032-019-1021-4>) contains supplementary material, which is available to authorized users.

O. Hejna · V. Curn
Biotechnological Centre, Faculty of Agriculture, University of South Bohemia, Studentska, 1668 Ceske Budejovice, Czech Republic

O. Hejna · L. Havlickova · Z. He · I. Bancroft (✉)
Department of Biology, University of York, Heslington, York YO10 5DD, UK
e-mail: ian.bancroft@york.ac.uk

areas with *P. brassicae* exceeds 20% but may lead to total crop failure (Pageau et al. 2006). Indeed, the ever-growing rapeseed production all over the world in the past years also increase areas infested by *P. brassicae*. Dissemination has been observed in Europe, China, India, Canada and Australia (Diederichsen et al. 2014; Chai et al. 2014; Bhattacharya et al. 2014; Rahman et al. 2014; Donald and Porter 2014).

Three main systems have been set to distinguish between pathotypes of *P. brassicae*. The most commonly used is the Williams classification (Williams 1966). This system is based on 4 different resistant varieties and is able to distinguish 16 different pathotypes. Seven different pathotypes have been revealed in the Czech Republic. The most common pathotypes are 7 (43%) and 6 (29%) (Ricarova et al. 2016). Other systems distinguishing pathotypes are the European clubroot differential (ECD) set (Buczacki et al. 1975) and the differential system of Some et al. (1996).

Traditional agricultural practices such as soil liming or agrochemical treatments by fungicides are expensive and not very effective to control this disease (Hwang et al. 2014), which is further complicated due to the broad range of hosts in which the pathogen is able to reproduce and the ability of spores to survive more than 20 years (Wallenhammar 1996). The most effective and economic strategy to eliminate clubroot disease is a combination of conventional disease-management measures including crop rotation, liming, application of fungicides and disinfection of equipment in combination with the use of varieties possessing multi-resistant genes (Faggian and Strelkov 2009).

Intensive breeding of cultivars resistant to clubroot has been ongoing for many years. The main sources of clubroot resistance (CR) genes were found in European fodder turnip cultivars (*Brassica rapa*), namely Gelria R, Siloga, Debra and Milan White (Hirai 2006). These sources of resistance were successfully introgressed into *Brassica napus*, which has led to the release of several resistant rapeseed cultivars (Piao et al. 2009).

The effort of scientists to find resistance genes is increasing every year along with the growing areas affected by *P. brassicae*. Recently, two CR genes in *B. rapa* have been cloned: *CRa* has been mapped on chromosome A3 (Ueno et al. 2012) and *Crr1* on chromosome A8 (Hatakeyama et al. 2013). Furthermore, seven other loci have been finely mapped on the A3 chromosome of *B. rapa*, namely *CRb*, *CRb-kato*, *CRd*, *Crr3*, *Rcr1*, *Rcr2* and *Rcr4* (Saito et al. 2006; Kato et al.

2013; Chu et al. 2014; Zhang et al. 2014; Huang et al. 2017; Yu et al. 2017; Pang et al. 2018). At least ten another CR loci were discovered in the A genome; *Crr2* was mapped on A1 (Suwabe et al. 2003); *CRc* and *Rcr8* on A2 (Sakamoto et al. 2008; Yu et al. 2017); *CRk*, *PbBa3.1* and *PbBa3.3* on A3 (Sakamoto et al. 2008; Chen et al. 2013); *CrrA5* on A5 (Nguyen et al. 2018); *Crr4* on A6 (Suwabe et al. 2006); and *Rcr9* on A8 (Yu et al. 2017).

In contrast with *B. rapa*, less progress has been made towards the identification of CR genes in the *B. oleracea* genome (C genome). Previous studies have presumed that there are much less dominant CR genes in C genome and CR is quantitative under polygenic control there (Zhang et al. 2016). So far, five loci have been described: *CR2a*, *CR2b*, *Pb3*, *Pb4* and *PbBo1* (Landry et al. 1992; Grandclément and Thomas 1996; Voorrips et al. 1997; Rocherieux et al. 2004). Furthermore, at least 10 CR quantitative trait loci (QTL) have been mapped in the C genome: *QTL-LG3* on C1 (Nomura et al. 2005), *Pb-Anju1*, *Pb-Anju2* and *CRQTL-YC* on C2 (Nagaoka et al. 2010; Lee et al. 2016), *Pb-Anju3* on C3 (Nagaoka et al. 2010), *Pb-GC1* and *QTL-LG9* on C5 (Nagaoka et al. 2010; Nomura et al. 2005) and *Pb-Anju4* and *Rcr7* on C7 (Nagaoka et al. 2010; Dakouri et al. 2018).

Currently, more than 30 CR loci and two dominant CR genes have been proposed in the AC genome of *B. napus*. Manzanares-Dauleux et al. (2000) discovered one dominant gene *Pb-Bn1* on A4 and two QTLs on A4 and C5. So far, the most CR loci have been mapped by Werner et al. (2008), in which 19 QTL (most race-specific) were detected across 8 chromosomes.

The first GWAS-based study using *Brassica* 60 K SNP arrays for screening a natural population of 472 *B. napus* accessions in an infected field to detect resistance genes to most predominant pathotype 4 of *P. brassicae* in China mapped 10 loci on A4, A10, C3, C4, C6 and C9 chromosomes (Li et al. 2016). Compared to 60 K SNP array, current associative transcriptomics (AT) platform (Havlickova et al. 2018) offers much better SNP coverage with the added benefit of using transcript abundance data. AT that was first described by Harper et al. (2012) has been previously used to identify genes underlying control of seed glucosinolate content (Lu et al. 2014), anion homeostasis (Koprivova et al. 2014), cell wall polysaccharides (Wood et al. 2017) and leaf calcium and magnesium accumulation (Alcock et al. 2017).

In this study, 245 diverse *B. napus* genotypes were inoculated under controlled conditions by the mixture of most predominant *P. brassicae* pathotype ECD 17/31/31 in the Czech Republic and scored for clubroot resistance. These data were used for AT with aim to elucidate associated regions associated with a source of clubroot resistance.

Materials and methods

Plant material

A panel of 245 accessions of *Brassica napus* used to test clubroot resistance and subsequent association analysis has been previously reported (Havlickova et al. 2018). Based on previous relatedness characterization, accessions were defined as winter oilseed rape (101), winter fodder (4), spring oilseed rape (92), swede (17), kale (2), semi-winter (5) and not assigned crop type (24). This collection is composed of lines all over the world, including varieties from Europe, Asia, North America, Australia and North Africa, with lines released from the 1950s up to today's modern winter Canola type oilseed rape (Supplemental Table 1).

Pathogen isolates

Inoculum used for testing of the resistance to clubroot was composed by the most aggressive pathotypes *P. brassicae* within the Czech Republic. Clubs for inoculum preparation were collected from the hardest hit areas affected by clubroot near Svetla Hora in the Moravian-Silesian Region (Ricarova et al. 2016). According to the identification method called the ECD (European Clubroot Differential set) Buczacki et al. (1975), the pathogen was identified as pathotypes 17/31/31.

Inoculum standing spores were obtained from a solid club of infected plants. Before use, the tumours were stored at $-18\text{ }^{\circ}\text{C}$. The clubs were pureed in distilled water for 3 min at the highest speed to prepare an inoculum. The final suspension was filtered through a muslin cloth and then centrifuged three times for 7 min; the resultant clusters of spores were resuspended in distilled water and adjusted with Bürker chamber to the desired concentration of 100 M spores in 1 ml of inoculum.

Inoculation of spores and plant cultivation

Cultivation trays with a cell size of 4×4 cm were filled with a mixture of coarse perlite and conventional growing medium for vegetables (Forestina, Czech Republic) with a pH 6.5 in the ratio 1:1. The seeds of tested genotypes were sown in each pot on the surface of the growth substrate. On every seed was applied by micropipette 0.5 ml of inoculum at a concentration of 100 M spores ml^{-1} and covered with 1 cm of coarse perlite. As the standard, extremely susceptible variety of Chinese cabbage 'Granaat' has been used. The prepared plant pots were placed with five replications in a randomized design on trays within constantly maintained about 1 cm high water level. Plants were grown in a growth chamber with a 16-h day (80 to $100\ \mu\text{E m}^{-2}\ \text{s}^{-1}$ at $20\text{ }^{\circ}\text{C}$) and an 8-h night ($18\text{ }^{\circ}\text{C}$) photoperiod over a period of 7 weeks. Indication of infection was observed 3 weeks after sowing. The plants were fertilized weekly using a solution of Kristalon Start (AGRO CS a.s., Czech Republic) (0.5 g per 10 l of water).

Evaluation of infestation

Disease severity was assessed 7 weeks after inoculation from roots using a standard 0–3 scale, where 0 = no visual symptoms, 1 = clubs only on the lateral roots, 2 = main root clubs and 3 = deformed entire root system (Buczacki et al. 1975) (Fig. 1a) The disease index (DI) was determined as follows:

$$\text{DI} = [(n_1 + 2n_2 + 3n_3) / (N_T \times 3)] \times 100$$
 where n_1 to n_3 were the number of plants with different disease severity of 1–3 scale and N_T represented the total number of identified plants, respectively (Li et al. 2016). Very sensitive genotypes showed DI above 80%, while high resistant genotypes reveal DI less than 20%.

Transcriptome sequencing, SNP detection, gene expression and associative transcriptomics

Plant material was a subset of the genetically diverse AT panel of 383 rapeseed accessions described by Havlickova et al. (2018). The growth conditions for RNA extraction, transcriptome sequencing and functional genotype determination were reported previously by He et al. (2017). The genotypes reported in Havlickova et al. (2018) were re-used for the present study. Mapping and associative transcriptomics were performed as described by Havlickova et al. (2018). In




Fig. 1 **a** Disease severity was assessed after 7 weeks from inoculation by using a standard 0–3 scale (0 = no visual symptoms, 1 = clubs only on the lateral roots, 2 = main root clubs, 3 = deformed

entire root system). **b** Histogram of disease index (DI) in 245 accessions including colour coded crop types

total, 256,397 SNPs (MAF > 0.01) and 53,889 CDS models (RPKM > 0.4) were used for Manhattan plots. Regions of the genome containing multiple markers showing significant association with trait variation are termed herein association peaks. In the Manhattan plots, simple SNP markers (i.e. polymorphisms between resolved bases) and hemi-SNPs (i.e. polymorphisms involving multiple bases called at the SNP position in one allele of the polymorphism) that have been directly linkage mapped and can be assigned to a genome with confidence are shown as dark points. Hemi-SNP markers that have not been linkage mapped are shown as light points indicating that the polymorphism could be at either the position of the CDS gene model plotted or within the homeologous CDS gene model in the other genomes. Association peaks comprising only light

points are termed herein shadow peaks. Shadow peaks are not expected to represent the positions of trait control loci. To evaluate minor effect loci, we implemented two thresholds for calling association peaks as described by Li et al. (2016). The major peaks were designated as significant associated loci, when they contained at least one SNP with significance $-\log_{10}P < 10^{-5}$ (sSNP). The minor loci were called as a potential associated loci, if a minimum of one SNP in the peak has significance $10^{-5} < -\log_{10}P < 10^{-4}$ (pSNP) and a locus has a distinguishable shape of peak as well.

Candidate gene annotation

Pairwise linkage disequilibrium was calculated, and heat maps were produced for each individual

chromosome as previously described by Havlickova et al. (2018). Potential candidate genes were identified within the range of LD block regarded as region with the most significant SNPs ($-\log_{10}P > 4$) which $r^2 > 0.4$ (Li et al. 2016); when not present, a region of ± 0.2 Mb (Samayoa et al. 2015) from the significant SNPs on the pseudomolecule reference sequence was checked for annotated genes putatively involved in plant response to the club root. To uncover potential candidate genes, GO enrichment analysis, InterPro functional analysis and manual annotation based on the similarity of *A. thaliana* were performed.

Results

Phenotypic variation of clubroot resistance in a diversity panel

Resistance to clubroot was assessed in an association panel of 245 *Brassica napus* accessions by using DI (Figs. 1b and 2; Supplemental Table 1). The measured values showed a high level of phenotypic variation within the panel. The DIs ranged from 0 to 100 with an average value of 45.51 ± 2.32 standard error (SE). The frequency distributions diagrams of DIs indicated a

certain degree of separation between almost fully resistant lines and sensitive ones. Moreover, 35.5% of lines were classified as resistant ($DI < 20$), whereas 19.6% demonstrated extreme susceptibility to the clubroot disease ($DI > 80$) with prevalent amount of lines with DI in the range of 61–70 (24%). Unequal distributions of data may indicate that the resistance is driven by major resistance gene or genes accompanied by multi-loci of weaker effect. To evaluate the effect of the crop type to clubroot resistance, frequency distribution was assessed (Fig. 1b). Both kale lines were found to be resistant ($DI = 0$), followed by swede ($DI = 3.9 \pm 3.8$), spring OSR ($DI = 23.5 \pm 3.3$), semiwinter OSR ($DI = 43.9 \pm 8.8$), fodder with largest variation ($DI = 56.7 \pm 17.2$), not assigned crop types ($DI = 66.2 \pm 5.8$) and winter OSR ($DI = 68.2 \pm 2.4$).

SNP association analysis

AT analysis of DI identified 9 SNP association peaks with trait variation (Fig. 3). They were called according to location on the *B. napus* pseudomolecule (*BnA01_0308*, *BnA02_0265*, *BnA02_0286*, *BnA03_0186*, *BnA03_0263*, *BnA08_0009*, *BnC02_0414*, *BnC07_0238*, *BnC07_0421*).




Fig. 2 Population structure and trait variation across 245 *B. napus* accessions. **a** Relatedness of accessions in the panel based on 355,536 scored SNPs. **b** Main crop types in the panel, colour-coded: orange for spring oilseed rape, green for semi-winter oilseed rape, light blue for swede, dark blue for kale, black for fodder

and red for winter oilseed rape, grey for crop type not assigned. **c** Population structure for highest likelihood $K = 2$. **d** Variation for clubroot resistance by using disease index DI ($DI = 0$, no visual symptoms; $DI = 100$, deformed entire root system)




Fig. 3 Transcriptome SNP association analysis for clubroot resistance. The SNP markers are positioned on the x-axis based in the genomic order of the gene models in which the polymorphism was scored, with the significance of the trait association, as $-\log_{10}P$, on the y-axis. A1 to A10 and C1 to C9 are the chromosomes of *B. napus*, shown in alternating black and blue colours to permit boundaries to be distinguished. Hemi-SNP markers (i.e. polymorphisms involving multiple bases called at the SNP position in one

allele of the polymorphism) for which the genome of the polymorphism cannot be assigned are shown as light points, whereas simple SNP markers (i.e. polymorphisms between resolved bases) and hemi-SNPs that have been directly linkage mapped, both of which can be assigned to a genome, are shown as dark points. The broken blue and red horizontal lines mark significance $-\log_{10}P = 5$ and $-\log_{10}P = 4$, respectively

Five SNP association peaks included sSNPs as shown in Supplemental Table 2. In total, 86 SNPs were highly associated with clubroot resistance, 29 of them were called as sSNP and the remaining 57 as pSNP. Five SNPs were lying alone above $-\log_{10}P < 10^{-4}$ threshold line without any close sSNPs or pSNPs to create a distinguishable peak, therefore excluded from further investigation. The rest of the associated SNPs were clustered in small loci and formed clear peaks. Graphic representation of SNP association analysis in Manhattan plots for individual chromosomes is seen in Supplemental Figure 3. The most significantly associated peak *BnA03_0263* with predominant simple sSNPs ($-\log_{10}P > 7$) that can be assigned with confidence to a genome was discovered on chromosome A03 (Fig. 4). This peak was accompanied by the presence of corresponding shadow peak in homeologous region of chromosome C07 (*BnC07_0421*) shown in Supplemental Figure 3h. Furthermore, other associated peaks in very small regions with simple sSNPs *BnA02_0265* and *BnA03_0186* were found on chromosomes A02 and A03 respectively (Supplemental Figure 3b, d). Both of them were also accompanied by the presence of corresponding shadow peaks with hemi-SNPs in homeologous regions of chromosomes C02 and C07: *BnC02_0414* (Supplemental Figure 3f) and

BnC07_0238 (Supplemental Figure 3g) respectively. The last three potential loci were found on A01: *BnA01_0308* (Supplemental Figure 3a), A02: *BnA02_0286* (Supplemental Figure 3c) and A08: *BnA08_0009* (Supplemental Figure 3e).

Candidate genes from SNP analysis were searched in the total area of 392 genes (in LD blocks containing significant SNPs; Supplemental Figure 4 or at a distance of 0.2 Mb from the pSNPs/sSNPs in locus). In these regions, potential candidate genes directly/indirectly involved in clubroot resistance are identified Supplemental Table 2. Using the results of enrichment analysis, genes containing phrase “defence response”, “ethylene”, “jasmonic acid”, “salicylic acid”, “abscisic acid”, “auxin” and “gibberellin” were considered as potential candidate genes. Based on the InterPro analysis and annotation, the transcription factors with plant defence domains ERF, bZIP, WRKY, MYB, plant defence *cis*-regulatory ET/JA motifs, G-box, GCC-box, W-box and pathogen-related proteins were included to candidate gene group (Singh et al. 2002). The estimated narrow-sense heritability (h^2) of DI 0.469 suggests that genetic variability may play a substantial role in CR resistance. In total, 63 candidate genes have been identified from SNP analysis (Supplemental Table 2).




Fig. 4 SNP association analysis for clubroot resistance focus on part of pseudomolecule with the highest associated locus (around 26 million bases from the beginning of the A03 chromosome). The SNP are positioned on the *x*-axis based on their location (units 10^5), the positions of the candidate genes for this locus are further indicated on *x*-axis. On the *y*-axis are values of the trait association

significance ($-\log_{10}P$). The black signs represent simple SNP and hemi-SNP markers assigned to the corresponding genome and grey hemi-SNP markers for which the genome of the polymorphism cannot be assigned. The dashed blue and red lines mark significance $-\log_{10}P = 5$ and $-\log_{10}P = 4$, respectively

GEMs analysis

Candidate genes were identified on the significance limit of $-\log_{10}P > 3.5$. Graphic representation of the gene expression analysis is shown in Supplemental Figure 1. The gene expression analysis identified 21 genes which exceeded the defined limit of significance (Supplemental Table 3). In the total, we identified 21 genes above this limit (Supplemental Table 3). These genes were located on the chromosomes A05, A07, A09, A10, C02, C03, C04, C05, C06, C07, C08 and C09 (Supplemental Figure 1). GO enrichment analysis, InterPro analysis and annotation were performed at the same way as for SNP candidate genes. Overall, 12 potential candidate genes have been identified from GEM analysis (Supplemental Table 3). These do not correspond to the positions of the SNP associations. This low number of candidates is a consequence of the available transcript abundance data being derived from leaf tissue, whereas the trait was measured in roots.

Discussion

Many studies have reported clubroot resistance loci in *B. napus* and its diploid progenitors *B. rapa* and *B. oleracea*, for example: *Anju1*, *Anju2*, *Anju3*, *Anju4*, *CRa*, *CrrA5*, *CRb*, *CRb^{kato}*, *CRQTL-GN_1*, *CRQTL-GN_2*, *CRc*, *CRd*, *Crr1*, *Crr2*, *Crr3_CrK*, *QTL_LG9*, *MCR-A4*, *MCR-C3*, *MCR-C9*, *PbBA31*, *PbBA32*, *SCR-A10a*, *SCR-A10b*, *SCR-C3*, *SCR-C4a*, *SCR-C4b*, *SCR-C6*, *Rcr1*, *Rcr2*, *Rcr4*, *Rcr7*, *Rcr8* and *Rcr9* (Chen et al. 2013; Chu et al. 2014; Dakouri et al. 2018; Hayashida et al. 2008; Huang et al. 2017; Kato et al. 2013; Lee et al. 2016; Li et al. 2016; Nagaoka et al. 2010; Nguyen et al. 2018; Nomura et al. 2005; Pang et al. 2018; Saito et al. 2006; Sakamoto et al. 2008; Suwabe et al. 2003; Yu et al. 2017; Zhang et al. 2014). In this study, we aimed to use AT to identify further loci and candidate genes playing key roles in clubroot resistance in oilseed rape. The clubroot disease is difficult to control, once the soil is infested with spores of *P. brassicae*, the soil

contamination could last for more than 20 years (Dixon 2009). In the last decade, this disease is spreading rapidly (Ricarova et al. 2017). In the light of seriousness, numerous studies have been conducted to discover resistance genes. However, most studies were carried out on *B. rapa*. In *B. napus*, a source of the major resistance gene has been found in Mendel variety, which shows resistance against certain pathogens. Unfortunately, the nature of resistance is based on only one resistance gene and it was overcome by new pathotype or high pathogen pressure (Diederichsen et al. 2014).

Recent GWAS analysis conducted in *B. napus* for Chinese pathotype 4 identified nine new resistance loci (Li et al. 2016). In this study, we performed GWAS with a mix of European *P. brassicae* pathotypes 17/31/31. We used a large diversity panel of *B. napus* representing genotypes from around the world. For association analysis, we combined SNP markers and transcript abundance from mRNA-Seq to detect new resistance loci and potential candidate genes for resistance against clubroot disease.

In total, we identified 86 SNPs to be highly associated with clubroot resistance. Twenty-nine of them with significance of $P < 10^{-5}$ and 57 SNPs with significance $10^{-5} < P < 10^{-4}$. SNPs were located into nine small loci (*BnA01_0308*, *BnA02_0265*, *BnA02_0286*, *BnA03_0186*, *BnA03_0263*, *BnA08_0009*, *BnC02_0414*, *BnC07_0238*, *BnC07_0421*). This suggests that the clubroot resistance is probably quantitatively inherited trait controlled by multiple loci.

The most prominent association peak was located on chromosome A03, locus *BnA03_0263* (Fig. 4), with few hemi-SNPs (markers with ambiguous genome anchoring) present in homeologous position on chromosome C07, locus *BnC07_0421* (Supplemental Figure 3h). Among the genes containing most of the sSNPs on chromosome A03, 7 candidate genes were found, of which 6 belong to the Disease Resistance (TIR-NBS-LRR class) family. In the corresponding region of *B. rapa* genome or in close proximity, resistance genes *CRa*, *CRb^{kato}*, *Rcr1*, *Rcr2* and *Rcr4* have been identified (Hayashida et al. 2008; Ueno et al. 2012; Kato et al. 2013; Chu et al. 2014; Huang et al. 2017; Yu et al. 2017). The GEM analysis revealed among 6 disease resistance candidates, one with a high correlation with the clubroot DI (*BnaA03g45000D*). This gene shows the greatest similarity with the already cloned *CRa* clubroot resistance gene in the *B. rapa* genome. However, in the study by Zhang et al. (2016) focusing

specifically on *CRa* ortholog in *B. napus*, localized in high proximity of our locus *BnA03_0263*, the same principle of resistance has not been demonstrated and it has been suggested that resistance to *P. brassicae* may be controlled by the combined effect of a new CR gene and *CRa* from *B. rapa* (Zhang et al. 2016).

Another associated locus *BnA03_0186* (Supplemental Figure 3d) has been detected on chromosome A03. We found 7 candidate genes directly containing sSNPs lying in close proximity of a group of 4 leucine-rich repeat transmembrane kinase genes. The ortholog of this gene in *A. thaliana* is localized to the plasma membrane, and it is involved in the regulation of plant innate immunity. Moreover, this gene has the ability to recognize chitin (Le et al. 2014) and its transcription is strongly declined under clubroot infection (Siemens et al. 2006). Another candidate gene containing pSNPs is ERF domain 11; ethylene response factor acts as a negative regulator of JA-responsive defence gene expression, resistance to fungal pathogen *Fusarium oxysporum* and antagonist of JA inhibition of root elongation (Lyons et al. 2013).

The second most prominent association peak was located on chromosome A02, locus *BnA02_0265* (Fig. 5). In this locus, we identified 9 candidate genes. The most promising are Tryptophan RNA-binding attenuator contains sSNPs. *A. thaliana* ortholog is in the direct interaction with PEN3 required for non-host resistance (Campe et al. 2016) and pectin lyase-like superfamily also contains sSNPs. Its role is cell wall modification (Etchells et al. 2012). This locus also included RING U-box superfamily, which is disease resistance protein (TIR-NBS class) with function of signal transduction, apoptosis and innate immune response. Other candidate genes can be seen below.

A second locus on A02 chromosome *BnA02_0286* contains two candidate genes, *A. thaliana* orthologs *cytochrome P450*, family 71, subfamily B, polypeptide 20 (potential genetic target of Whirly transcription factor) and defence response CAP-gly domain linker.

Loci *BnA02_0265* and *BnA02_0286* overlap a relatively large locus designed as *Rcr8* (Yu et al. 2017).

Another associated locus with pSNPs was found on chromosome A01. Within this locus, 11 candidate genes are present and their position is not overlapping with previously described region where *Crr2* gene has been identified in *B. rapa* (Suwabe et al. 2003). The direct gene encompassing pSNP is an ortholog ATPase, AAA-type, *CDC48* and negative regulator of NLR-mediated




Fig. 5 SNP association analysis for clubroot resistance focus on part of pseudomolecule with the second most prominent association peak (around 26 million bases from the beginning of the A02 chromosome). The SNPs are positioned on the *x*-axis based on their location (units 10^5), the positions of the candidate genes for this locus are further indicated on *x*-axis. On the *y*-axis are values of the

trait association significance ($-\log_{10}P$). The black signs represent simple SNP and hemi-SNP markers assigned to the corresponding genome and grey hemi-SNP markers for which the genome of the polymorphism cannot be assigned. The dashed blue and red lines mark significance $-\log_{10}P = 5$ and $-\log_{10}P = 4$, respectively

immunity (Copeland et al. 2016). Next to this gene is another interesting candidate gene ortholog of *PAT-TERN-TRIGGERED IMMUNITY (PTI) COMPROMISED RECEPTOR-LIKE CYTOPLASMIC KINASE 1* and *PCRK1*. This gene is important for immunity induced by damage associated molecular pattern—DAMPs (Sreekanta et al. 2015a, b). Interesting gene in this locus is also phytosulphokine 5 precursor, which is an important signalling in resistance in root infection of *Fusarium oxysporum* (Shen and Diener 2013).

The other associated locus was found on chromosome A8. This does not correspond to resistance loci identified previously on this chromosome, *Rcr9* and *Crr1*, being in different positions on the chromosome (Yu et al. 2017; Suwabe et al. 2003). It contains 6 candidate genes, for example cluster of orthologs of *GDSL-motif lipase 2* and *GDSL-like Lipase Acylhydrolase* superfamily. They are involved in disease resistance and negatively regulate auxin signalling (Lee et al. 2009). *GDSL-like lipase* is also engaged in defence against *Alternaria brassicicola* (Oh et al. 2005).

GEM association revealed 21 genes, exceeding the limit of $-\log_{10}p$ 3.5. These genes were distributed across the whole genome, with slight predominance towards C genome. Performing GO enrichment

analysis, InterPro analysis and using *blastn* for annotation gene with orthologs of *A. thaliana*, we reduced the number of candidate genes to nineteen. The most promising candidates were orthologs of pentacyclic triterpene synthase 1, which is important for *PEN1* and *PEN2* functions in powdery mildew non-host interaction. This synthase mediates transport required for innate immunity and focal accumulation of syntaxin *PEN1* (Nielsen et al. 2012). Another candidate cytokinin response factor 2 plays important role in response to stress condition and auxin regulation (Simackova et al. 2015). *S*-adenosyl-L-methionine-dependent methyltransferase superfamily is another stress candidate gene especially important in drought tolerance (Nir et al. 2014). It is important to recognize that only genes with expression in the source tissue for mRNAseq (i.e. leaves) correlated with the CR trait can be identified as GEM associations; those with root-specific or infection-specific expression cannot be identified by AT with the expression data available. However, AT analysis revealed candidate genes directly or indirectly involved in clubroot resistance, not only as results of the significant association between DI and sequence variation present in our diversity panel, but also as result of difference in

transcript abundance within the panel and therefore provides added value in the association analysis.

Translation of our findings into improved clubroot resistance of new *B. napus* varieties will involve the development of molecular markers to select alleles associated with greater resistance. To aid this, we have compiled (Supplemental Figure 5) shortlists of suitable polymorphisms to underpin the development of high throughput SNP markers. The use of molecular markers to pre-select seedling for trialling will improve the speed and efficiency of breeding for clubroot resistance.

Funding information This work was supported by The Czech National Agency for Agricultural Research (NAZV QJ1510172 and QK1910070) and UK Biotechnology and Biological Sciences Research Council (BB/L002124/1).

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Alcock TD, Havlickova L, He Z, Bancroft I, White PJ, Broadley MR, Graham N (2017) Identification of candidate genes for calcium and magnesium accumulation in *Brassica napus* L. by association genetics. *Front Plant Sci* 8:1968. <https://doi.org/10.3389/fpls.2017.01968>
- Bhattacharya I, Dutta S, Mondal S, Mondal B (2014) Clubroot disease on Brassica crops in India. *Can J Plant Pathol* 36: 154–160. <https://doi.org/10.1080/07060661.2013.875064>
- Buczacki ST, Toxopeus H, Mattusch P, Johnston TD, Dixon GR, Hobolth LA (1975) Study of physiologic specialization in *Plasmodiophora Brassicae*: proposals for attempted rationalization through an international approach. *Trans Br Mycol Soc* 65:295–303. [https://doi.org/10.1016/S0007-1536\(75\)80013-1](https://doi.org/10.1016/S0007-1536(75)80013-1)
- Campe R, Langenbach C, Leissing F, Popescu GV, Popescu SC, Goellner K, Beckers GJM, Conrath U (2016) ABC transporter *PEN3/PDR8/ABCG36* interacts with calmodulin that, like *PEN3*, is required for Arabidopsis nonhost resistance. *New Phytol* 209:294–306. <https://doi.org/10.1111/nph.13582>
- Chai AL, Xie XW, Shi YX, Li BJ (2014) Research status of clubroot (*Plasmodiophora Brassicae*) on cruciferous crops in China. *Can J Plant Pathol* 36:142–153. <https://doi.org/10.1080/07060661.2013.868829>
- Chen J, Jing J, Zhan Z, Zhang T, Zhang C, Piao Z (2013) Identification of novel QTLs for isolate-specific partial resistance to *Plasmodiophora brassicae* in *Brassica rapa*. *PLoS One* 8:e85307. <https://doi.org/10.1371/journal.pone.0085307>
- Chu M, Song T, Falk KC, Zhang X, Liu X, Chang A, Lahlali R, McGregor L, Gossen BD, Yu F, Peng G (2014) Fine mapping of *Rcr1* and analyses of its effect on transcriptome patterns during infection by *Plasmodiophora brassicae*. *BMC Genomics* 15:1166. <https://doi.org/10.1186/1471-2164-15-1166>
- Copeland C, Woloshen V, Huang Y, Li X (2016) *AtCDC48A* is involved in the turnover of an NLR immune receptor. *Plant J* 88:294–305. <https://doi.org/10.1111/tpj.13251>
- Dakouri A, Zhang X, Peng G, Falk KC, Gossen BD, Strelkov SE, Yu F (2018) Analysis of genome-wide variants through bulked segregant RNA sequencing reveals a major gene for resistance to *Plasmodiophora Brassicae* in *Brassica oleracea*. *Sci Rep* 8:17657. <https://doi.org/10.1038/s41598-018-36187-5>
- Diederichsen E, Frauen M, Ludwig-Mueller J (2014) Clubroot disease management challenges from a German perspective. *Can J Plant Pathol* 36:85–98. <https://doi.org/10.1080/07060661.2013.861871>
- Dixon GR (2009) *Plasmodiophora Brassicae* in its environment. *J Plant Growth Regul* 28:212–228. <https://doi.org/10.1007/s00344-009-9098-3>
- Donald EC, Porter IJ (2014) Clubroot in Australia: the history and impact of *Plasmodiophora brassicae* in *Brassica* crops and research efforts directed towards its control. *Can J Plant Pathol* 36:66–84. <https://doi.org/10.1080/07060661.2013.873482>
- Etchells JP, Moore L, Jiang WZ, Prescott H, Capper R, Saunders NJ, Bhatt AM, Dickinson HG (2012) A role for *BELLRINGER* in cell wall development is supported by loss-of-function phenotypes. *BMC Plant Biol* 12:212. <https://doi.org/10.1186/1471-2229-12-212>
- Faggian R, Strelkov SE (2009) Detection and Measurement of *Plasmodiophora brassicae*. *J Plant Growth Regul* 28(3):282–288
- Fredua-Agyeman R, Rahman H (2016) Mapping of the clubroot disease resistance in spring *Brassica napus* canola introgressed from European winter canola Cv. ‘Mendel’. *Euphytica* 211:201–213. <https://doi.org/10.1007/s10681-016-1730-2>
- Grandclément C, Thomas G (1996) Detection and analysis of QTLs based on RAPD markers for polygenic resistance to *Plasmodiophora brassicae* Woron in *Brassica oleracea* L. *Theor Appl Genet* 93:86–90. <https://doi.org/10.1007/BF00225731>
- Harper AL, Trick M, Higgins J, Fraser F, Clissold L, Wells R, Hattori C, Werner P, Bancroft I (2012) Associative transcriptomics of traits in the polyploid crop species *Brassica napus*. *Nat Biotechnol* 30:798–802. <https://doi.org/10.1038/nbt.2302>
- Hatakeyama K, Suwabe K, Tomita RN, Kato T, Nunome T, Fukuoka H, Matsumoto S (2013) Identification and characterization of *Crr1a*, a gene for resistance to clubroot disease (*Plasmodiophora brassicae* Woronin) in *Brassica rapa* L. *PLoS One* 8:e54745. <https://doi.org/10.1371/journal.pone.0054745>
- Havlickova L, He Z, Wang L, Langer S, Harper AL, Kaur H, Broadley MR, Gegas V, Bancroft I (2018) Validation of an updated associative transcriptomics platform for the

- polyploid crop species *Brassica napus* by dissection of the genetic architecture of erucic acid and tocopherol isoform variation in seeds. *Plant J* 93:181–192. <https://doi.org/10.1111/tpj.13767>
- Hayashida N, Takabatake Y, Nakazawa N, Aruga D, Nakanishi H, Taguchi G, Sakamoto K, Matsumoto E (2008) Construction of a practical SCAR marker linked to clubroot resistance in Chinese cabbage, with intensive analysis of HC352b genes. *J Jpn Soc Hortic Sci* 77:150–154. <https://doi.org/10.2503/jjshs1.77.150>
- He Z, Wang L, Harper AL, Havlickova L, Pradhan AK, Parkin IAP, Bancroft I (2017) Extensive homoeologous genome exchanges in allopolyploid crops revealed by mRNAseq-based visualization. *Plant Biotechnol J* 15(5):594–604
- Hirai M (2006) Genetic analysis of clubroot resistance in *Brassica* crops. *Breed Sci* 56:223–229. <https://doi.org/10.1270/jsbbs.56.223>
- Huang Z, Peng G, Liu X, Deora A, Falk KC, Gossen BD, McDonald MR, Yu F (2017) Fine mapping of a clubroot resistance gene in chinese cabbage using snp markers identified from bulked segregant RNA sequencing. *Front Plant Sci* 8. <https://doi.org/10.3389/fpls.2017.01448>
- Hwang SF, Strelkov SE, Feng J, Gossen BD, Howard RJ (2012) *Plasmodiophora brassicae*: a review of an emerging pathogen of the Canadian canola (*Brassica napus*) crop. *Mol Plant Pathol* 13:105–113. <https://doi.org/10.1111/j.1364-3703.2011.00729.x>
- Hwang SF, Ahmed HU, Zhou Q, Strelkov SE, Gossen BD, Peng G, Turnbull GD (2014) Efficacy of Vapam fumigant against clubroot (*Plasmodiophora brassicae*) of canola. *Plant Pathol* 63:1374–1383. <https://doi.org/10.1111/ppa.12207>
- Kageyama K, Asano T (2009) Life cycle of *Plasmodiophora brassicae*. *J Plant Growth Regul* 28:203–211. <https://doi.org/10.1007/s00344-009-9101-z>
- Kato T, Hatakeyama K, Fukino N, Matsumoto S (2013) Fine mapping of the clubroot resistance gene *CRb* and development of a useful selectable marker in *Brassica rapa*. *Breed Sci* 63:116–124. <https://doi.org/10.1270/jsbbs.63.116>
- Koprivova A, Harper AL, Trick M, Bancroft I, Kopriva S (2014) Dissection of the control of anion homeostasis by associative transcriptomics in *Brassica napus*. *Plant Physiol* 166:442–450. <https://doi.org/10.1104/pp.114.239947>
- Landry BS, Hubert N, Crete R, Chang MS, Lincoln SE, Etoh T (1992) A genetic map for *Brassica oleracea* based on RFLP markers detected with expressed DNA sequences and mapping of resistance genes to race 2 of *Plasmodiophora brassicae* (Woronin). *Genome* 35:409–420. <https://doi.org/10.1139/g92-061>
- Le MH, Cao Y, Zhang XC, Stacey C (2014) *LIK1*, A *CERK1*-interacting kinase, regulates plant immune responses in *Arabidopsis*. *PLoS One* 9:e102245. <https://doi.org/10.1371/journal.pone.0102245>
- Lee DS, Kim BK, Kwon SJ, Jin HC, Park OK (2009) *Arabidopsis* GDSL lipase 2 plays a role in pathogen defense via negative regulation of auxin signaling. *Biochem Biophys Res Commun* 379:1038–1042. <https://doi.org/10.1016/j.bbrc.2009.01.006>
- Lee J, Izzah NK, Choi BS, Joh HJ, Lee SC, Perumal S, Seo J, Ahn K, Jo EJ, Choi GJ, Nou IS, Yu Y, Yang TJ (2016) Genotyping-by-sequencing map permits identification of clubroot resistance QTLs and revision of the reference genome assembly in cabbage (*Brassica oleracea* L.). *DNA Res* 23:29–41. <https://doi.org/10.1093/dnares/dsv034>
- Li L, Luo Y, Chen B, Xu K, Zhang F, Li H, Huang Q, Xiao X, Zhang T, Hu J, Li F, Wu X (2016) A genome-wide association study reveals new loci for resistance to clubroot disease in *Brassica napus*. *Front Plant Sci* 7:1483. <https://doi.org/10.3389/fpls.2016.01483>
- Lu G, Harper AL, Trick M, Morgan C, Fraser F, O'Neill C, Bancroft I (2014) Associative transcriptomics study dissects the genetic architecture of seed glucosinolate content in *Brassica napus*. *DNA Res* 21:613–625. <https://doi.org/10.1093/dnares/dsu024>
- Lyons R, Iwase A, Gänsewig T, Sherstnev A, Duc C, Barton GJ, Hanada K, Higuchi-Takeuchi M, Matsui M, Sugimoto K, Kazan K, Simpson GG, Shirasu K (2013) The RNA-binding protein FPA regulates Flg22-triggered defense responses and transcription factor activity by alternative polyadenylation. *Sci Rep* 3:2866. <https://doi.org/10.1038/srep02866>
- Manzanares-Dauleux MJ, Divaret I, Baron F, Thomas G (2000) Evaluation of French *Brassica oleracea* landraces for resistance to *Plasmodiophora brassicae*. *Euphytica* 113:211–218. <https://doi.org/10.1023/A:1003997421340>
- Nagaoka T, Doullah MAU, Matsumoto S, Kawasaki S, Ishikawa T, Hori H, Okazaki K (2010) Identification of QTLs that control clubroot resistance in *Brassica oleracea* and comparative analysis of clubroot resistance genes between *B. rapa* and *B. oleracea*. *Theor Appl Genet* 120:1335–1346. <https://doi.org/10.1007/s00122-010-1259-z>
- Nguyen ML, Monakhos GF, Komakhin RA, Monakhos SG (2018) The new Clubroot resistance locus is located on chromosome A05 in Chinese cabbage (*Brassica rapa* L.). *Russ J Genet* 54:296–304. <https://doi.org/10.1134/S1022795418030080>
- Nielsen ME, Feechan A, Böhlenius H, Ueda T, Thordal-Christensen H (2012) *Arabidopsis* ARF-GTP exchange factor, *GNOM*, mediates transport required for innate immunity and focal accumulation of syntaxin *PEN1*. *Proc Natl Acad Sci U S A* 109:11443–11448. <https://doi.org/10.1073/pnas.1117596109>
- Nir I, Moshelion M, Weiss D (2014) The *Arabidopsis* *GIBBERELLIN METHYL TRANSFERASE 1* suppresses gibberellin activity, reduces whole-plant transpiration and promotes drought tolerance in transgenic tomato. *Plant Cell Environ* 37:113–123. <https://doi.org/10.1111/pce.12135>
- Nomura K, Minegishi Y, Kimizuka-Takagi C, Fujioka T, Moriguchi K, Shishido R, Ikehashi H (2005) Evaluation of F2 and F3 plants introgressed with QTLs for clubroot resistance in cabbage developed by using SCAR markers. *Plant Breed* 124:371–375. <https://doi.org/10.1111/j.1439-0523.2005.01105.x>
- Oh IS, Park AR, Bae MS, Kwon SJ, Kim YS, Lee JE, Kang NY, Lee S, Cheong H, Park OK (2005) Secretome analysis reveals an *Arabidopsis* lipase involved in defense against *Alternaria brassicicola*. *Plant Cell* 17:2832–2847. <https://doi.org/10.1105/tpc.105.034819>
- Pageau K, Reisdorf-Cren M, Morot-Gaudry JF, Masclaux-Daubresse C (2006) The two senescence-related markers, *GSI* (cytosolic glutamine synthetase) and *GDH* (glutamate dehydrogenase), involved in nitrogen mobilization, are differentially regulated during pathogen attack and by stress

- hormones and reactive oxygen species in *Nicotiana tabacum* L. leaves. *J Exp Bot* 57:547–557. <https://doi.org/10.1093/jxb/erj035>
- Pang W, Fu P, Li X, Zhan Z, Yu S, Piao Z (2018) Identification and mapping of the clubroot resistance gene CRd in Chinese cabbage (*Brassica Rapa* ssp. *Pekinensis*). *Front Plant Sci* 9. <https://doi.org/10.3389/fpls.2018.00653>
- Piao Z, Ramchiary N, Lim YP (2009) Genetics of clubroot resistance in *Brassica* species. *J Plant Growth Regul* 28:252–264. <https://doi.org/10.1007/s00344-009-9093-8>
- Rahman H, Peng G, Yu F, Falk KC, Kulkarni M, Selvaraj G (2014) Genetics and breeding for clubroot resistance in canadian spring canola (*Brassica napus* L.). *Can J Plant Pathol* 36: 122–134. <https://doi.org/10.1080/07060661.2013.862571>
- Ricarova V, Kazda J, Baranyk P, Rysanek P (2016) Clubroot – an emerging disease faced by Czech oilseed rape growers. *Sci Agric Bohem* 47:105–112. <https://doi.org/10.1515/sab-2016-0015>
- Ricarova V, Kazda J, Baranyk P, Rysanek P (2017) Greenhouse and field experiments with winter oilseed rape cultivars resistant to *Plasmodiophora brassicae* Wor. *Crop Prot* 92:60–69. <https://doi.org/10.1016/j.cropro.2016.10.009>
- Rocherieux J, Glory P, Giboulot A, Boury S, Barbeyron G, Thomas G, Manzaneres-Dauleux MJ (2004) Isolate-specific and broad-spectrum QTLs are involved in the control of clubroot in *Brassica oleracea*. *Theor Appl Genet* 108: 1555–1563. <https://doi.org/10.1007/s00122-003-1580-x>
- Saito M, Kubo N, Matsumoto S, Suwabe K, Tsukada M, Hirai M (2006) Fine mapping of the clubroot resistance gene, *Crr3*, in *Brassica rapa*. *Theor Appl Genet* 114:81–91. <https://doi.org/10.1007/s00122-006-0412-1>
- Sakamoto K, Saito A, Hayashida N, Taguchi G, Matsumoto E (2008) Mapping of isolate-specific QTLs for clubroot resistance in Chinese cabbage (*Brassica rapa* L. ssp. *pekinensis*). *Theor Appl Genet* 117:759–767. <https://doi.org/10.1007/s00122-008-0817-0>
- Samayoa LF, Malvar RA, Olukolu BA, Holland JB, Butron A (2015) Genome-wide association study reveals a set of genes associated with resistance to the Mediterranean corn borer (*Sesamia nonagrioides* L) in a maize diversity panel. *BMC Plant Biol* 15:350. <https://doi.org/10.1186/s12870-014-0403-3>
- Shen Y, Diener AC (2013) *Arabidopsis thaliana* RESISTANCE TO FUSARIUM OXYSPORUM 2 implicates tyrosine-sulfated peptide signaling in susceptibility and resistance to root infection. *PLoS Genet* 9:e1003525. <https://doi.org/10.1371/journal.pgen.1003525>
- Siemens J, Keller I, Sarx J, Kunz S, Schuller A, Nagel W, Schmülling T, Parniske M, Ludwig-Müller J (2006) Transcriptome analysis of *Arabidopsis* clubroots indicate a key role for cytokinins in disease development. *Mol Plant-Microbe Interact* 19:480–494. <https://doi.org/10.1094/MPMI-19-0480>
- Simackova M, O'Brien JA, Khan M, Noorden GV, Otvos K, Vieten A, De Clercq I, Van Haperen JMA, Cuesta C, Hoyerová K, Vanneste S, Mrhavy P, Wabnik K, Van Breusegem F, Nowack M, Murphy A, Friml J, Weijeres D, Beekman T, Benková E (2015) Cytokinin response factors regulate PIN-FORMED auxin transporters. *Nat Commun* 6: 8717. <https://doi.org/10.1038/ncomms9717>
- Singh K, Foley RC, Onate-Sanchez L (2002) Transcription factors in plant defense and stress responses. *Curr Opin Plant Biol* 5: 430–436. [https://doi.org/10.1016/S1369-5266\(02\)00289-3](https://doi.org/10.1016/S1369-5266(02)00289-3)
- Some A, Manzaneres MJ, Laurens F, Baron F, Thomas G, Rouxel F (1996) Variation for virulence on *Brassica napus* L. amongst *Plasmodiophora brassicae* collections from France and derived single-spore isolates. *Plant Pathol* 45: 432–439. <https://doi.org/10.1046/j.1365-3059.1996.d01-155.x>
- Sreekanta S, Bethke G, Hatsugai N, Tsuda K, Thao A, Wang L, Katagiri F, Glazebrook J (2015a) The receptor-like cytoplasmic kinase *PCRK1* contributes to pattern-triggered immunity against *Pseudomonas syringae* in *Arabidopsis thaliana*. *New Phytol* 207:78–90. <https://doi.org/10.1111/nph.13345>
- Sreekanta S, Haruta M, Minkoff BB, Glazebrook J (2015b) Functional characterization of *PCRK1*, a putative protein kinase with a role in immunity. *Plant Signal Behav* 10: e1063759–e1063751. <https://doi.org/10.1080/15592324.2015.1063759>
- Suwabe K, Tsukazaki H, Iketani H, Hatakeyama K, Fujimura M, Nunome T, Fukuoka H, Matsumoto S, Hirai M (2003) Identification of two loci for resistance to clubroot (*Plasmodiophora brassicae* Woronin) in *Brassica rapa* L. *Theor Appl Genet* 107:997–1002. <https://doi.org/10.1007/s00122-003-1309-x>
- Suwabe K, Tsukazaki H, Iketani H, Hatakeyama K, Kondo M, Fujimura M, Nunome T, Fukuoka H, Hirai M, Matsumoto S (2006) Simple sequence repeat-based comparative genomics between *Brassica rapa* and *Arabidopsis thaliana*: the genetic origin of clubroot resistance. *Genetics* 173:309–319. <https://doi.org/10.1534/genetics.104.038968>
- Ueno H, Matsumoto E, Aruga D, Kitagawa S, Matsumura H, Hayashida N (2012) Molecular characterization of the *Cra* gene conferring clubroot resistance in *Brassica rapa*. *Plant Mol Biol* 80:621–629. <https://doi.org/10.1007/s11103-012-9971-5>
- Voorrips RE, Jongerius MC, Kanne HJ (1997) Mapping of two genes for resistance to clubroot (*Plasmodiophora brassicae*) in a population of doubled haploid lines of *Brassica oleracea* by means of RFLP and AFLP markers. *Theor Appl Genet* 94: 75–82. <https://doi.org/10.1007/s001220050384>
- Wallenhammar AC (1996) Prevalence of *Plasmodiophora brassicae* in a spring oilseed rape growing area in central Sweden and factors influencing soil infestation levels. *Plant Pathol* 45:710–719. <https://doi.org/10.1046/j.1365-3059.1996.d01-173.x>
- Werner S, Diederichsen E, Frauen M, Schondelmaier J, Jung C (2008) Genetic mapping of clubroot resistance genes in oilseed rape. *Theor Appl Genet* 116:363–372. <https://doi.org/10.1007/s00122-007-0674-2>
- Williams PH (1966) A system for the determination of races of *Plasmodiophora brassicae* that infect cabbage and rutabaga. *Phytopathology* 56:624–626
- Wood IP, Pearson BM, Garcia-Gutierrez EG, Havlicková L, He Z, Harper AL, Bancroft I, Waldron KW (2017) Carbohydrate microarrays and their use for the identification of molecular markers for plant cell wall composition. *Proc Natl Acad Sci U S A* 114:6860–6865. <https://doi.org/10.1073/pnas.1619033114>
- Yu F, Zhang X, Peng G, Falk KC, Strelkov SE, Gossen BD (2017) Genotyping-by-sequencing reveals three QTL for clubroot

- resistance to six pathotypes of *Plasmiodiophora Brassicae* in *Brassica rapa*. *Sci Rep* 7:4516. <https://doi.org/10.1038/s41598-017-04903-2>
- Zhang T, Zhao Z, Zhang C, Pang W, Choi SR, Lim YP, Piao Z (2014) Fine genetic and physical mapping of the *CRb* gene conferring resistance to clubroot disease in *Brassica rapa*. *Mol Breed* 34:1173–1183. <https://doi.org/10.1007/s11032-014-0108-1>
- Zhang H, Feng J, Hwang SF, Strelkov SE, Falak I, Huang X, Sun R (2016) Mapping of clubroot (*Plasmiodiophora brassicae*) resistance in canola (*Brassica napus*). *Plant Pathol* 65:435–440. <https://doi.org/10.1111/ppa.12422>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.