



This is a repository copy of *Draft whole-genome sequences of 10 Aeromonas strains from clinical and environmental sources.*

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/149408/>

Version: Published Version

---

**Article:**

Gray, K., Green, L.R. [orcid.org/0000-0002-2836-2041](https://orcid.org/0000-0002-2836-2041), Chaudhuri, R.R. [orcid.org/0000-0001-5037-2695](https://orcid.org/0000-0001-5037-2695) et al. (1 more author) (2019) Draft whole-genome sequences of 10 *Aeromonas* strains from clinical and environmental sources. *Microbiology Resource Announcements*, 8 (30). e00170-19. ISSN 2576-098X

<https://doi.org/10.1128/mra.00170-19>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:  
<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>



# Draft Whole-Genome Sequences of 10 *Aeromonas* Strains from Clinical and Environmental Sources

Katie Gray,<sup>a</sup> Luke R. Green,<sup>a</sup> Roy R. Chaudhuri,<sup>b</sup> Jonathan G. Shaw<sup>a</sup>

<sup>a</sup>Department of Infection, Immunity and Cardiovascular Disease, University of Sheffield, Sheffield, United Kingdom

<sup>b</sup>Department of Molecular Biology and Biotechnology, University of Sheffield, Sheffield, United Kingdom

**ABSTRACT** *Aeromonas* bacteria are able to cause disease in a wide range of animals from humans to fish. In this article, we report the draft whole-genome sequences of 10 *Aeromonas* strains from clinical and environmental sources. These genome sequences will provide a repository of information for further investigations into the pathogenicity of this enigmatic pathogen.

*Aeromonas* species are aquatic bacteria found in brackish water and freshwater and are able to cause disease in both cold-blooded and warm-blooded animals (1). In humans, they can give rise to numerous clinical manifestations but are mostly associated with gastrointestinal disease and wound infections (2). A plethora of factors have been linked with aeromonad pathogenicity, including toxigenic factors, such as aerolysin and type III secretion systems (T3SS), and adhesins, such as type IV pili and both polar and lateral flagella (3–5). However, aeromonad pathogenicity is thought to be multifactorial and not reliant upon one virulence determinant. *Aeromonas caviae* strains Sch3 and Sch29 were isolated from children presenting with gastroenteritis at Sheffield Children's Hospital, and *Aeromonas veronii* biovar *sobria* BC88 was isolated from a child with dysentery in Western Australia (6); all the other aeromonads were environmental strains isolated from the River Don at various locations within South Yorkshire, United Kingdom. *A. caviae* Sch3 has been extensively studied due to its production of lateral flagella (4) and its ability to glycosylate its polar flagella (7, 8). *A. veronii* biovar *sobria* BC88 is a model for aeromonad adhesion, as it produces a type IV bundle-forming pilus (5). The analysis of the genomes presented here will allow a deeper understanding of the biology of aeromonads in relation to both their physiology and pathogenicity.

All environmental strains were isolated from river water by inoculation onto cystine lactose electrolyte-deficient (CLED) agar plus Andrade's indicator (Oxoid) and Brilliance UTI clarity agar (Oxoid), as these were nonselective and provided a differential presumptive identification of organisms. Human clinical strains were isolated by the inoculation of feces onto *Aeromonas* isolation agar (Fluka) containing ampicillin. All strains were identified to the species level by multisequence alignment against the sequences of known aeromonad species. All strains were grown in tryptic soy broth (TSB), and genomic DNA was extracted using a QIAamp DNA minikit (Qiagen). For each of the 10 *Aeromonas* strains, genomic DNA libraries were prepared using the Nextera XT library kit (Illumina, San Diego, CA). The genomes were nucleotide sequenced using an Illumina HiSeq 2500 platform at MicrobesNG (University of Birmingham, UK). Sequencing used 2 × 250-bp paired-end reads that gave from 41 to 150× depth (Table 1). Trimmomatic v0.38 (9) was used to trim the reads with a sliding window quality cutoff of Q15, and read quality analyses were performed with FastQC software v0.11.8 (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>). The genomes were assembled and annotated using SPAdes v3.7 (10) and Prokka v1.12, respectively, using the standard default settings (11). Genome assembly metrics were calculated with

**Citation** Gray K, Green LR, Chaudhuri RR, Shaw JG. 2019. Draft whole-genome sequences of 10 *Aeromonas* strains from clinical and environmental sources. *Microbiol Resour Announc* 8:e00170-19. <https://doi.org/10.1128/MRA.00170-19>.

**Editor** David Rasko, University of Maryland School of Medicine

**Copyright** © 2019 Gray et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Jonathan G. Shaw, [J.G.Shaw@sheffield.ac.uk](mailto:J.G.Shaw@sheffield.ac.uk).

**Received** 19 February 2019

**Accepted** 1 July 2019

**Published** 25 July 2019

**TABLE 1** Genome accession numbers and characteristics of 10 *Aeromonas* strains taken from either a clinical or environmental source

Strain	Species (source)	Mean coverage (x)	No. of reads	No. of contigs	Largest contig (bp)	Total length (bp)	GC content (%)	$N_{50}$ (bp)	$N_{75}$ (bp)	$L_{50}$ (contigs)	$L_{75}$ (contigs)	No. of CDS	ENA accession and assembly no.
Sch29	<i>A. caviae</i> (human clinical diarrhea)	49.94	874,346	195	182,866	4,425,373	61.34	86,948	50,669	21	37	3,953	<a href="#">ERS3090042</a> , <a href="#">GCA_901202955</a>
Sch3N	<i>A. caviae</i> (human clinical diarrhea)	84.83	1,155,915	192	240,555	4,737,922	61.39	90,385	43,401	18	37	4,313	<a href="#">ERS3090043</a> , <a href="#">GCA_901212305</a>
BC88	<i>A. veronii</i> (human clinical dysentery)	111.91	1,381,586	155	1,023,078	4,604,788	58.54	215,763	118,836	6	13	4,065	<a href="#">ERS3090044</a> , <a href="#">GCA_901212295</a>
KLG1	<i>A. hydrophila</i> (environmental water)	53.88	730,744	154	260,972	4,908,673	61.22	103,738	64,788	15	30	4,373	<a href="#">ERS3090045</a> , <a href="#">GCA_901212375</a>
KLG2	<i>A. allosaccharophila</i> (environmental water)	150.39	1,670,538	85	626,394	4,513,594	58.87	292,420	141,273	5	11	4,059	<a href="#">ERS3090046</a> , <a href="#">GCA_901212385</a>
KLG5	<i>A. veronii</i> (environmental water)	70.63	974,322	103	818,983	4,740,061	58.51	280,270	130,857	6	12	4,269	<a href="#">ERS3090047</a> , <a href="#">GCA_901212355</a>
KLG6	<i>A. media</i> (environmental water)	104.31	1,255,304	454	159,134	4,542,840	61.20	36,002	19,277	40	80	4,070	<a href="#">ERS3090048</a> , <a href="#">GCA_901212365</a>
KLG7	<i>A. veronii</i> (environmental water)	53.29	687,242	104	345,454	4,552,893	58.80	139,212	84,196	11	21	4,069	<a href="#">ERS3090049</a> , <a href="#">GCA_901212345</a>
KLG8	<i>A. veronii</i> (environmental water)	54.65	738,573	76	568,763	4,590,381	58.62	198,583	115,053	7	15	4,137	<a href="#">ERS3090050</a> , <a href="#">GCA_901212395</a>
KLG9	<i>A. veronii</i> (environmental water)	40.98	578,468	74	399,442	4,605,689	58.71	180,084	120,803	9	17	4,160	<a href="#">ERS3090051</a> , <a href="#">GCA_901212405</a>

QUAST, and the number of annotated coding sequences (CDS) for each aeromonad genome is shown in Table 1.

The sequences will provide a great resource for further investigations into the physiology and pathogenicity of the *Aeromonas* genus.

**Data availability.** The reads used for assembly of the 10 annotated aeromonad genomes were deposited in the European Nucleotide Archive (ENA) at the European Molecular Biology Laboratory (EMBL) under the accession number [PRJEB31025](https://www.ebi.ac.uk/ena/record/PRJEB31025). The specific accession numbers for each sample are supplied in Table 1. The versions described in this paper are the first versions.

## ACKNOWLEDGMENT

MicrobesNG provided genome sequencing, which was supported by the BBSRC (grant number BB/L024209/1).

## REFERENCES

- Janda JM, Abbott SL. 2010. The genus *Aeromonas*: taxonomy, pathogenicity and infection. *Clin Microbiol Rev* 23:35–73. <https://doi.org/10.1128/CMR.00039-09>.
- Parker JL, Shaw JG. 2011. *Aeromonas* spp. clinical microbiology and disease. *J Infect* 62:109–118. <https://doi.org/10.1016/j.jinf.2010.12.003>.
- Rabaan AA, Gryllos I, Tomas JM, Shaw JG. 2001. Motility and the polar flagellum are required for *Aeromonas caviae* adherence to HEp-2 cells. *Infect Immun* 69:4257–4267. <https://doi.org/10.1128/IAI.69.7.4257-4267.2001>.
- Gavin R, Rabaan AA, Merino S, Tomas JM, Gryllos I, Shaw JG. 2002. Lateral flagella of *Aeromonas* species are essential for epithelial cell adherence and biofilm formation. *Mol Microbiol* 43:383–397. <https://doi.org/10.1046/j.1365-2958.2002.02750.x>.
- Hadi N, Yang Q, Barnett TC, Tabei SMB, Kirov SM, Shaw JG. 2012. Bundle-forming pilus locus of *Aeromonas veronii* bv sobria. *Infect Immun* 80:1351–1369. <https://doi.org/10.1128/IAI.06304-11>.
- Kirov SM, Sanderson K. 1996. Characterization of a type IV bundle forming pilus (SFP) from a gastroenteritis strain of *Aeromonas veronii* biovar sobria. *Microb Pathog* 21:23–34. <https://doi.org/10.1006/mpat.1996.0039>.
- Tabei SMB, Hitchen PG, Day-Williams MJ, Merino S, Vart R, Pang P-C, Horsburgh GJ, Viches S, Wilhelms M, Tomás JM, Dell A, Shaw JG. 2009. An *Aeromonas caviae* genomic island is required for both O-antigen lipopolysaccharide biosynthesis and flagellin glycosylation. *J Bacteriol* 191:2851–2863. <https://doi.org/10.1128/JB.01406-08>.
- Parker JL, Lowry R, Couto NAS, Wright PC, Stafford GS, Shaw JG. 2014. Maf-dependent bacterial flagellin glycosylation occurs before chaperone binding and flagellar T3SS export. *Mol Microbiol* 92:258–272. <https://doi.org/10.1111/mmi.12549>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.