



UNIVERSITY OF LEEDS

This is a repository copy of *Oxic Sediments*.

White Rose Research Online URL for this paper:  
<https://eprints.whiterose.ac.uk/153842/>

Version: Accepted Version

---

**Book Section:**

Tostevin, R and Poulton, SW [orcid.org/0000-0001-7621-189X](https://orcid.org/0000-0001-7621-189X) (2019) *Oxic Sediments*. In: Gargaud, M, Irvine, WM, RicardAmils, R, Cleaves, HJ, Pinti, D, Quintanilla, JC and Viso, M, (eds.) *Encyclopedia of Astrobiology, Living Edition*. Springer , Berlin, Germany . ISBN 978-3-642-27833-4

[https://doi.org/10.1007/978-3-642-27833-4\\_5414-1](https://doi.org/10.1007/978-3-642-27833-4_5414-1)

---

© Springer-Verlag GmbH Germany, part of Springer Nature 2020. This is an author produced version of a paper published in *Encyclopedia of Astrobiology, Living Edition*. Uploaded in accordance with the publisher's self-archiving policy.

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

## **Oxic sediments**

Rosalie Tostevin<sup>1</sup> and Simon W. Poulton<sup>2</sup>

<sup>1</sup>Department of Earth Sciences, University of Oxford, Oxford, OX13AN, UK.

Email: Rosalie.tostevin@earth.ox.ac.uk

<sup>2</sup>School of Earth and Environment, University of Leeds, Leeds, LS2 9JT, UK

Email: S.Poulton@leeds.ac.uk

### **Definition**

Pore waters within oxic sediments contain measureable amounts of dissolved oxygen. In general, pore water oxygen concentrations decline with depth due to organic matter degradation. The depth of oxygen penetration varies depending on the sedimentation rate, mixing processes, the flux and type of organic carbon and the oxygen content of bottom waters.

**Keywords:** Oxygen, redox, diagenesis, organic carbon, microbes, respiration, pore waters.

### **History**

#### **Overview**

Sediments at the seafloor are porous and poorly consolidated, and as such readily allow diffusion of bottom waters and the dissolved gases that they contain into the sediment. With depth, sediments undergo compaction and eventually lose contact with overlying seawater. While sediment pore waters at or close to the sediment-water interface in the global ocean commonly contain dissolved oxygen, the concentration of oxygen declines with depth due to organic matter degradation.

Deposited organic matter is degraded at the seafloor and with burial in the sediments, with less than 1% surviving to become part of the rock record (Emerson and Hedges, 1988). Aerobic respiration is the first pathway used by microbes for organic matter degradation, as it is associated with the largest free energy yield. As such, aerobic respiration is responsible for the majority of organic carbon remineralisation in the oceans and sediments (Bender and Heggie, 1984). But at low dissolved oxygen concentrations (<10  $\mu\text{M}$ ), the rate of aerobic respiration begins to decline, and anaerobic respiration processes may proceed with alternate electron acceptors (Tostevin & Poulton, this volume). Aerobic respiration tends to be most prolific at the sediment-water interface, but oxygen may be present in pore waters for centimeters or even meters below the sediment-water interface.


The depth of oxygen penetration in sediments is highly variable, and is controlled by the chemical, physical and biological setting. One major control is the concentration of oxygen in overlying bottom waters. The concentration of dissolved oxygen in surface waters is set by atmospheric oxygen levels, as well as the solubility of  $\text{O}_2$  gas in water, which is itself controlled by salinity and temperature. At depth in the water column, oxygen is consumed by respiration, but the rate of decline depends on the local carbon cycle and mixing rates. Oxygen levels in oceanic bottom waters vary from up to 300  $\mu\text{M}$  below polar surface waters, to <5  $\mu\text{M}$  below oxygen minimum zones in intense upwelling regions. Higher oxygen levels

at the sediment-water interface increase the average depth to which oxygen can persist in the sediments.

Sedimentation rate exerts another important control on the depth of oxygen penetration, by influencing the amount of time organic matter remains exposed at or near the sediment-water interface (Figure 1). Under the high sedimentation rates common in coastal and shallow shelf environments ( $\sim 1 \text{ mmyr}^{-1}$ ), organic matter rapidly loses contact with overlying bottom waters. Therefore, a relatively large fraction of organic matter is often buried in such sediments, and oxygen in pore waters is rapidly consumed (Figure 1). In contrast, under the low sedimentation rates common in deeper water settings ( $< 0.01 \text{ mmyr}^{-1}$ ), organic matter remains exposed to oxic bottom waters for longer (1000's of years), and so there is less organic carbon leftover to drive anaerobic heterotrophic communities in the sediments.

Mixing in shallow water environments can disturb any stratification, bringing anoxic sediments from a few centimeters depth back into contact with oxic bottom waters. Mixing may occur mechanically, for example during storm events, or biologically, through bioturbation (Kristensen, 2000). More intense mixing results in re-oxidation of reduced species and allows deeper oxygen penetration (Figure 1). The properties of the sediments can influence the degree of contact between bottom waters and pore waters. For example, fine grained sediments such as mud have a lower porosity, and so limit diffusion of oxygen from bottom waters into pore waters, hence facilitating a steeper oxygen gradient than coarse grained sediments such as sand.

Finally, the flux and type of organic matter is an important variable. Large fluxes of organic matter, as are common in coastal environments, can result in more rapid consumption of the available oxygen in shallow sediments. Furthermore, organic matter varies considerably in terms of its reactivity, with phytoplankton-derived carbon easier to digest than terrestrial organic matter which contains tough plant lignins. All of these parameters contribute to the exposure time of marine sediments to oxic conditions.



**Figure 1:** The depth of the oxic zone in sediments is partly controlled by the sedimentation rate, the flux of organic matter, and mixing processes. In general, the oxic zone extends deeper in deeper ocean settings where sedimentation rates and organic carbon fluxes are lower.

#### Cross references

Tostevin, R., Poulton, S. W., Suboxic sediments

#### Further reading

#### References

- Bender, M.L., Heggie, D.T., 1984. Fate of organic carbon reaching the deep sea floor: a status report. *Geochim. Cosmochim. Acta* 48, 977–986.
- Emerson, S., Hedges, J., 1988. Processes controlling the organic carbon content of open ocean sediments. *Paleoceanography* 3, 621–634.
- Kristensen, E., 2000. Organic matter diagenesis at the oxic/anoxic interface in coastal marine sediments, with emphasis on the role of burrowing animals, in: Liebezeit, G., Dittmann, S., Kröncke, I. (Eds.), *Life at Interfaces and Under Extreme Conditions, Developments in Hydrobiology*. Springer Netherlands, pp. 1–24.