

Practice review: Evidence-based and effective management of pain in patients with advanced cancer

Palliative Medicine

1–10

© The Author(s) 2020



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/0269216319896955

journals.sagepub.com/home/pmj

Emma J Chapman¹, Zoe Edwards¹, Jason W Boland²,
Matthew Maddocks³, Lucy Fettes³, Catherine Malia⁴,
Matthew R Mulvey¹ and Michael I Bennett¹

Abstract

Background: Pain of a moderate or severe intensity affects over half of patients with advanced cancer and remains undertreated in at least one-third of these patients.

Aim: The aim of this study was to provide a pragmatic overview of the evidence supporting the use of interventions in pain management in advanced cancer and to identify where encouraging preliminary results are demonstrated but further research is required.

Design: A scoping review approach was used to examine the evidence supporting the use of guideline-recommended interventions in pain management practice.

Data sources: National or international guidelines were selected if they described pain management in adult cancer patients and were written within the last 5 years in English. The Cochrane Database of Systematic Reviews (January 2014 to January 2019) was searched for ‘cancer’ AND ‘pain’ in the title, abstract or keywords. A MEDLINE search was also made.

Results: A strong opioid remains the drug of choice for treating moderate or severe pain. Bisphosphonates and radiotherapy are also effective for cancer-related bone pain. Optimal management requires a tailored approach, support for self-management and review of treatment outcomes. There is likely a role for non-pharmacological approaches. Paracetamol should not be used in patients taking a strong opioid to treat pain. Cannabis-based medicines are not recommended. Weak opioids, ketamine and lidocaine are indicated in specific situations only.

Conclusion: Interventions commonly recommended by guidelines are not always supported by a robust evidence base. Research is required to evaluate the efficacy of non-steroidal anti-inflammatory drugs, anti-convulsants, anti-depressants, corticosteroids, some invasive anaesthetic techniques, complementary therapies and transcutaneous electrical nerve stimulation.

Keywords

Pain, analgesia, evidence-based practice, neoplasms, terminal care, review

What is already known about the topic?

- Pain is not adequately controlled in all patients with advanced cancer.
- There are no universally used guidelines for the treatment of pain in patients with advanced cancer.
- Not all guideline recommendations are evidence-based.

¹Academic Unit of Palliative Care, Leeds Institute of Health Sciences, Leeds, UK

²Wolfson Palliative Care Research Centre, Hull York Medical School, University of Hull, Hull, UK

³Cicely Saunders Institute of Palliative Care, Policy & Rehabilitation, Florence Nightingale Faculty of Nursing, Midwifery & Palliative Care, King’s College London, London, UK

⁴St Gemma’s Hospice, Leeds, UK

Corresponding author:

Emma J Chapman, Academic Unit of Palliative Care, Leeds Institute of Health Sciences, Worsley Building, Clarendon Way, Leeds, LS2 9JT, UK.
Email: e.j.chapman@leeds.ac.uk

What this paper adds?

- We provide a pragmatic overview of the current 'state of the science' supporting pain management practice.
- 'Do' and 'Do not' practice recommendations are made based upon a set of strength recommendations, that is, strong: a large and consistent body of evidence, such as a systematic review, moderate: solid empiric evidence from one or more papers and tentative: limited empiric evidence, a strong recommendation cannot be made.
- Interventions are categorised as 'Don't know' where there is currently a lack or paucity of high-quality empiric evidence to support the approach in management of cancer pain but some evidence to suggest that the approach is effective in a non-cancer context. These are interventions where further research is required before the routine use of the practice can be recommended.

Implications for practice, theory or policy

- An oral strong opioid remains the drug of choice for treating moderate or severe pain in patients with advanced cancer and should be supplemented with bisphosphonates and/or radiotherapy for bone pain. Optimal management requires a tailored approach, support for self-management and regular review of treatment outcomes. There is likely a role for non-pharmacological approaches.
- Oral paracetamol should not routinely be used for patients who are already taking a strong opioid to treat moderate-to-severe pain. Cannabis-based medicines are not recommended. Weak opioids, ketamine and lidocaine are only indicated in specific situations.
- Further research is required into the use of non-steroidal anti-inflammatory drugs, anti-convulsants, anti-depressants, corticosteroids, some invasive anaesthetic techniques, acupuncture, massage and transcutaneous electrical nerve stimulation.

Introduction

Pain is of a moderate or severe intensity in more than half of patients with advanced cancer¹ and is undertreated in around one-third of cases.² Poorly controlled pain is the most common cause for community-based cancer patients to contact out-of-hours primary care services³ and is one of the most frequent reasons for their hospitalisation.⁴ Although home is the preferred place of care and death for many patients,⁵ national surveys show that this is where pain control is poorest.⁶ However, accessing specialist palliative care improves pain management for patients living at home.⁷

There are no standardised guidelines for the management of pain in patients with advanced cancer and not all guideline recommendations are evidence-based. We set out to provide a pragmatic overview of the state of science supporting approaches commonly recommended by key guidelines rather than produce a novel consensus guideline. We also identified where encouraging preliminary results have been demonstrated but further research is required before the routine use of the practice can be recommended.

Methods

This practice review, guided by scoping review methodology,^{8,9} was used to determine the extent of the evidence supporting the use of interventions in the management of pain in patients with advanced cancer. Guidelines from national or international bodies were selected if they described the management of pain in adult cancer patients and were written within the last 5 years in English language (Table 1). The Cochrane Database of Systematic Reviews

(January 2014 to January 2019) was searched for the terms 'cancer' AND 'pain' in the title, abstract or keywords. After independent screening by two researchers, 19 articles were identified as relevant to this review. An additional search of MEDLINE was made for further key publications that provided additional primary data to add to the evidence base for the interventions considered in Table 1 (search date: 31 May 2019). Methods were compliant with the Preferred Reporting Items for Systematic reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) Checklist (Supplementary Material 1). We determined the strength of evidence for commonly recommended interventions plus some non-pharmacological approaches for which there is currently limited evidence but a low potential for harm. Each intervention was then allocated to a practice recommendation category of 'Do', 'Do not' or 'Don't know' based on the direction and strength of the supporting evidence (Table 2). A 'strong' recommendation was made where there was evidence from a large and consistent body of evidence, such as a systematic review, a 'moderate' recommendation was made in light of solid empiric evidence from one or more papers and a 'tentative' recommendation where there was limited empiric evidence. Where 'tentative' evidence supports use of an intervention, the likely increased risk to the patient is considered (Table 3).

Practice recommendations – 'Do'

Screen for pain at every engagement with a healthcare professional. All guidelines examined recommend that patients should be screened for pain^{10–15,17} and at every engagement with a healthcare professional.¹⁰ Meta-analysis shows that

Table 1. Guidelines examined and their position on the interventions recommended by this practice review.

	Screen	Assess	Strong opioids	Opioid switch	Epidural/ intrathecal	Corticosteroids	Radiotherapy/ bisphosphonates	Self-management	Physiotherapy	Non-pharmacological	Regularly review and evaluate
Standards for the management of cancer-related pain across Europe – a position paper from the EFIC Task Force on cancer pain (2019) ¹⁰	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded
National Clinical Practice Guidelines in Oncology (NCCN Guidelines®) – adult cancer pain (Version 2. 2019) ¹¹	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded
Scottish Palliative Care Guidelines – pain management (updated 2019) ¹²	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded
Management of cancer pain in adult patients: European Society for Medical Oncology (ESMO) Clinical Practice Guidelines (2018) ¹³	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded
Pharmacotherapy of pain in cancer patients. Recommendations of the Polish Association for the Study of Wordliczek et al. (2018) ¹⁴	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded
World Health Organization (WHO) Guidelines for the pharmacological and radiotherapeutic management of cancer pain in adults and adolescents (2018) ¹⁵	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded
Spanish Society of Medical Oncology (SEOM, 2017) ¹⁶	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded
Pharmacological management of cancer pain in adults. National Clinical Guideline No. 9 (Ireland, 2015) ¹⁷	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded	Shaded

Shaded box shows where intervention is included in the guideline.

Table 2. How categories of recommendation were decided.

Do	High-quality published evidence for effectiveness in pain management in cancer patients AND Clear favourable benefit to risk ratio
Do not	Evidence that the approach is not effective in pain management in cancer patients OR Lack of high-quality evidence to support its use AND Unfavourable benefit to risk balance
Don't know	High-quality published data to support its use in the management of pain in a context other than advanced cancer AND/OR Limited evidence for effectiveness in pain in cancer patients, further research is required
<i>Strength of recommendations</i>	
Strong	A large and consistent body of evidence, such as a systematic review
Moderate	Solid empiric evidence from one or more papers
Tentative	Limited empiric evidence

using patient-reported measurements of pain reduces pain intensity in cancer patients perhaps by prompting discussion about symptoms between patient and professional.¹⁸

A large, randomised trial found that integration of a systematic bedside pain assessment tool into cancer care improves pain outcomes.¹⁹

Table 3. Summary of recommendations for addressing pain in advanced cancer patients.

	Strength of evidence
Do	
1. Screen for pain	Strong
2. Assess the components of the pain (e.g. neuropathic, nociceptive, inflammatory)	Tentative
3. Agree upon a tailored pain management plan	
(a) Treat moderate or severe pain with a strong opioid, such as morphine	Strong
(b) Consider opioid switching	Moderate
(c) Consider epidural or intrathecal administration of opioids	Moderate
(d) Treat metastatic bone pain with radiotherapy and/or bisphosphonates	Strong
(e) Support patients to use self-management strategies	Strong
(f) Physiotherapy-based interventions	Tentative
(g) Non-pharmacological approaches, for example, music therapy	Tentative
4. Regularly review and evaluate cancer pain treatment outcomes	Moderate
Do not	
1. Routinely use oral paracetamol to treat moderate or severe pain in patients on strong opioids	Moderate
2. Routinely use weak opioids such as codeine	Moderate
3. Routinely use ketamine	Moderate
4. Routinely use intravenous lidocaine	Moderate
5. Use cannabis-based medicines	Strong
Don't know if there is benefit of	
1. Non-steroidal anti-inflammatory drugs	
2. Anti-convulsants, anti-depressants	
3. Corticosteroids	
4. Other anaesthetic interventions, such as coeliac plexus block and spinal cord stimulation	
5. Acupuncture, massage	
6. Transcutaneous electrical nerve stimulation	

Assess the components of the pain. It is widely recommended that assessment should include classification of the type and pathophysiology of the pain.^{10–14} Aetiology, mechanisms and impact of cancer pain should be evaluated and this information used to inform the treatment plan.¹⁰ Identification and grading²⁰ of neuropathic pain are of particular importance as it is a predictor of poor pain management outcome in cancer patients.²¹ A tentative recommendation is made due to the potential benefit and low risk of harm yet a lack of high-quality evidence to support a role for pain assessment in increasing efficacy of treatment.

Agree upon a tailored pain management plan

Treat moderate or severe pain with a strong opioid. Morphine remains the first-line treatment for moderate-to-severe cancer pain; all commonly used oral or transdermal opioids have similar efficacy.^{22,23} However, less than half of patients who die of cancer are prescribed a strong opioid, and this occurs around median 9 weeks before death.^{24–26} Patients with inadequately managed pain may benefit from referral to specialist palliative care at an early stage, as a longer duration of specialist palliative care is associated with increased access to opioids and improved quality of life indicators.²⁷

Consider opioid switching. Switching to an alternative opioid^{28–32} is recommended if there is inadequate pain relief or unacceptable side effects.^{11,12,13,14,16,17} Evidence to support opioid switching is of moderate strength, better pain relief or reduced side effects were reported in 50% of patients who switched due to sub-optimal response to another opioid in a recent four-arm, phase-IV randomised controlled trial (RCT).³³ Of note, methadone has similar analgesic actions as morphine³⁴ but a long and unpredictable plasma half-life, which can lead to problems with accumulation and toxicity. Methadone should only be used for cancer pain under the guidance of specialist palliative care professionals.

Consider epidural or intrathecal opioids. Spinal or epidural administration of opioids or implanted intrathecal pumps is advocated.^{10,11,13,14,16,17} The evidence to support this from two RCTs is judged as moderate, the complexity of running trials in this group of patients is acknowledged.^{35–37} Implantation of a spinal opioid infusion requires access to a specialist pain management team for ongoing review and follow-up and careful consideration of fitness to undergo the procedure in patients with end-stage cancer.

Treat metastatic bone pain with radiotherapy, bisphosphonates or both. Metastatic bone pain can be

treated with radiotherapy or radioablation.^{11,12,13–17} A meta-analysis of RCTs found that around 60% of patients with uncomplicated bone metastases experienced pain relief in response to external beam radiotherapy and up to a quarter of these reported complete pain relief within 3–4 weeks.³⁸ Use of bisphosphonates is recommended^{11,13–17} and this treatment has a role in reducing skeletal morbidity in metastatic disease.³⁹ Bisphosphonates may delay the onset of pain but there was no evidence of an analgesic effect in a systematic review.⁴⁰ However, a more recent randomised multicentre trial (not included in the systematic review) provided good evidence that ibandronate had similar efficacy to radiotherapy.⁴¹

Support patients to use self-management strategies. Supporting self-management involves helping patients make the decision to manage their pain, enhance their self-efficacy by solving problems and incorporate pain-relieving strategies into daily life.⁴² Interventions that support self-management behaviour are infrequently recommended by guidelines but evidence shows that they are effective in pain control.^{43–48} Good communication, in particular, about opioids is key to maximising patient compliance with and therefore efficacy of treatment.⁴⁹

Consider physiotherapy-based interventions. Physiotherapy and rehabilitation medicine approaches include physical exercise (walking and home-based resistance exercises), myofascial release and proprioceptive neuromuscular facilitation techniques and are associated with reduced pain intensity in a pilot RCT, a large randomised trial of telerehabilitation, and as a symptom co-existing with fatigue.^{50–52}

Non-pharmacological approaches, for example, music therapy. Non-pharmacological approaches are recommended.^{10,11,14,16} Music-based interventions had a significant effect on pain intensity based on low-quality evidence from seven studies analysed in a Cochrane systematic review.⁵³ Where no adverse effects have been reported, we make a tentative suggestion to consider non-pharmacological approaches, such as music therapy, as part of a holistic management plan.

Regularly review and evaluate treatment cancer pain treatment outcomes. Multiple guidelines^{10,11,13,14,15,16} recommend that pain intensity and its management plan should be regularly reviewed to assess treatment outcome and plan onward care. Moderate evidence shows that review and evaluation of cancer pain management at the service level improves care and allows comparison between different services for example, hospices, community services and hospitals.^{54,55}

Practice recommendations – ‘Do not’

Routinely use oral paracetamol to treat moderate or severe pain in patients on strong opioids. Paracetamol is currently recommended for mild-to-moderate pain in guidelines based upon World Health Organization (WHO) guidance and the classic ‘cancer pain ladder’.^{11–13,14,15,16,17} However, patients with moderate-to-severe cancer pain that is already being treated with a strong opioid are unlikely to gain any additional benefit. A Cochrane systematic review found that there is no evidence that using paracetamol alone or in combination with opioids is of any benefit in treating cancer pain.⁵⁶ Furthermore, Cochrane systematic reviews also find high-quality evidence of no effect compared to placebo in non-cancer lower back pain⁵⁷ and recently osteoarthritis.⁵⁸ However, single-dose intravenous paracetamol is effective in acute post-operative pain⁵⁹ and may offer benefits in acute cancer pain, but this needs to be substantiated.

Routinely use weak opioids, such as codeine. Moderate-quality evidence from a Cochrane systematic review shows that there is only a little, low-quality evidence for the use of codeine in cancer pain⁶⁰ and the effectiveness of this step only lasts for about a month after which patients need to move to a strong opioid to gain sufficient analgesia.⁶¹ In an open-label RCT of moderate cancer pain, low-dose morphine had an earlier and more adequate analgesic effect than weak opioids and a similar tolerability.⁶² Nevertheless, in low resource countries and/or where morphine use may be limited, a weak opioid could still be a suitable choice for moderate pain.

Routinely use ketamine. Some guidelines include the option to use ketamine^{10,11,14,16} but acknowledge the lack of evidence to support its routine use. A Cochrane systematic review concluded that there was insufficient evidence to determine the benefits and harms of ketamine as an adjuvant to opioids.⁶³ Oral ketamine was neither superior to placebo in a RCT of patients with cancer-related neuropathic pain⁶⁴ nor when used subcutaneously as an adjunct to opioids in a randomised, double-blind, placebo-controlled study in patients with cancer pain.⁶⁵ However, ketamine should not be completely ruled out as a third-line approach because of its effectiveness in non-cancer, acute pain contexts.^{66,67} Ketamine might be of benefit in specific situations, such as patients with central sensitisation.⁶⁸

Routinely use intravenous lidocaine. There is limited data that intravenous lidocaine may reduce pain intensity in some patients^{69,70} but a lack of efficacy has been seen in other trials.⁷¹ Use of lidocaine comes with a risk of frequent adverse effects and therefore specialist supervision

is needed. Lidocaine could therefore be considered as an option for the treatment of opioid-refractory cancer pain as recommended by some guidelines.^{11,17}

Use cannabis-based medicines. Pre-clinical experiments show that tetrahydrocannabinol (THC) enhances the anti-nociceptive effect of morphine,⁷² and preliminary studies indicated a role as an adjuvant treatment for pain in cancer patients.⁷³ However, larger, RCTs of cannabis-based drugs as an adjuvant to standard treatment in advanced cancer patients have not confirmed any benefit.^{74,75} A recent systematic review⁷⁶ also concluded that very weak evidence suggests no effect of THC or oromucosal nabiximols on cancer pain that is not relieved by strong opioids.

'Don't know'

Further research is needed in patients with moderate and severe cancer pain because it is not clear if there is benefit of the following.

Non-steroidal anti-inflammatory drugs. Although non-steroidal anti-inflammatory drugs (NSAIDs) are recommended by guidelines,^{11–15,17} a Cochrane systematic review found that there is no high-quality evidence to determine their place in the management of cancer pain.⁷⁷ There is some evidence for a beneficial effect in other types of pain^{78,79} yet considering the potential for significant adverse effects, it is important to assess and carefully reassess the use of NSAIDs.¹³ RCT's evidence is urgently needed.

Anti-convulsants, anti-depressants. Despite evidence of effectiveness of anti-convulsants and anti-depressants in non-cancer contexts,⁸⁰ there is only very limited evidence of efficacy in cancer patients.⁸¹ A systematic review and meta-analysis found that combining anti-convulsant drugs (gabapentin or pregabalin) with opioids did not improve pain relief in cancer patients compared with opioids alone.⁸² The authors noted that due to heterogeneity of patients, a beneficial effect in some patients with neuropathic pain cannot be discounted. Currently high-quality evidence to support their use is lacking and the benefit to risk balance is unclear.

Corticosteroids. Multiple current guidelines recommend corticosteroids for some types of cancer pain,^{11,12,14–16} particularly where pain is related to inflammation and oedema. Evidence for their efficacy is weak, although a Cochrane systematic review found that some studies showed a beneficial effect on cancer-related pain after 1 week of treatment.⁸³ In light of the many well-documented adverse effects,⁸⁴ the likely benefit to risk ratio for the individual needs careful consideration.

Other anaesthetic interventions. Guidelines recommend that nerve blocks, cordotomy and spinal cord stimulation are considered in specific situations, such as refractory pain^{11,13,16} or as adjuvant therapy.¹² Currently there is not enough good-quality evidence to judge whether spinal cord stimulation is superior to pharmacological treatments in regards to safety and efficacy for management of pain in advanced cancer patients.^{35,85} RCTs are needed but the difficulty of conducting them in the relevant population is acknowledged.

Acupuncture, massage. NCCN recommend that massage and acupuncture should be considered alongside pharmacologic interventions.¹¹ Systematic reviews found that there is insufficient evidence to recommend the use of acupuncture⁸⁶ or massage⁸⁷ for relief of pain in cancer patients. Patients are often keen to explore complementary and integrative therapies and report a beneficial effect. Considering, the low potential for harm, further robust research into the efficacy of complementary therapies should be considered.

Transcutaneous electrical nerve stimulation. Transcutaneous electrical nerve stimulation (TENS) is categorised as an integrative intervention¹¹ or adjuvant therapy.¹² There is currently not enough evidence to recommend that TENS can be used for cancer pain in general or in advanced cancer patients due to a lack of power and undersizing in the studies published.⁸⁸ As a relatively simple, portable, non-invasive intervention with high safety profile, TENS merits further research in effectiveness trials.

Limitations

Limitations are acknowledged in that as this was a practice review produced using scoping review methodology, a systematic review of the evidence base for each intervention was not performed. There may be additional current guidelines and relevant primary literature that we have not examined. The source and direction of the evidence, for example, conclusion of the systematic review, was considered but the quality of evidence for each original study was not re-appraised.

Summary

After comprehensive assessment of the patient, a strong opioid remains the drug of choice for treating moderate or severe pain in patients with advanced cancer. Treatment with bisphosphonates, radiotherapy or both for bone pain is effective. Optimal management requires a tailored approach, support for self-management and regular review of treatment outcomes. Interventions commonly recommended by international guidelines are not always supported by a robust evidence base. Oral paracetamol should not be used in patients already

taking a strong opioid to manage moderate or severe pain. Cannabis-based drugs are not recommended due to evidence of lack of effect. Future research should focus on defining the efficacy of NSAIDs, anti-depressants, anti-convulsants, corticosteroids and the likely role for non-pharmacological approaches in management of pain in patients with advanced cancer. Data are needed to support the appropriate selection of patients who might benefit from third-line approaches, such as lidocaine and ketamine.

Author contributions

Original idea, identification of guidelines and preparation of manuscript was done by E.J.C. and M.I.B.; the Cochrane Database of Systematic Reviews was independently screened by E.J.C. and Z.E; all authors contributed content to the review and revised the final submission.

Notes on contributors

Dr Emma J Chapman is a Senior Research Fellow in Palliative Care; Zoe Edwards is a Research Fellow in Medicines Optimisation and Pharmacist; Dr Jason W Boland is a Senior Clinical Lecturer and Honorary Consultant in Palliative Medicine; Dr Matthew Maddocks is a Senior Lecturer and Specialist Physiotherapist, Ms Lucy Fettes is a Physiotherapist with experience of Pain Management Programmes in Palliative Care; Ms Catherine Malia is a Nurse Consultant at St Gemma's Hospice, Leeds; Dr Matthew R Mulvey is a University Academic Fellow, with an interest in neuropathic pain; Prof. Michael I Bennett is the St Gemma's Professor of Palliative Medicine, Director of the St Gemma's Academic Unit of Palliative Care at the University of Leeds and an internationally recognised expert in Pain Management.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship and/or publication of this article: This work was supported by Yorkshire Cancer Research Programme (Grant No. L412) – 'RESOLVE: Improving health status and symptom experience for people living with advanced cancer'.

ORCID iDs

Emma J Chapman  <https://orcid.org/0000-0003-2859-2020>

Zoe Edwards  <https://orcid.org/0000-0001-6525-8190>

Jason W Boland  <https://orcid.org/0000-0001-5272-3057>

Lucy Fettes  <https://orcid.org/0000-0002-2642-8318>

Matthew R Mulvey  <https://orcid.org/0000-0002-6357-3848>

Supplemental material

Supplemental material for this article is available online.

References

- Breivik H, Cherny N, Collett B, et al. Cancer-related pain: a pan-European survey of prevalence, treatment, and patient attitudes. *Ann Oncol* 2009; 20(8): 1420–1433.
- Greco MT, Roberto A, Corli O, et al. Quality of cancer pain management: an update of a systematic review of under-treatment of patients with cancer. *J Clin Oncol* 2014; 32(36): 4149–4154.
- Adam R, Wassell P and Murchie P. Why do patients with cancer access out-of-hours primary care? A retrospective study. *Br J Gen Pract* 2014; 64(619): e99–e104.
- Numico G, Cristofano A, Mozzicafreddo A, et al. Hospital admission of cancer patients: avoidable practice or necessary care? *PLoS ONE* 2015; 10(3): e0120827.
- Gomes B, Higginson IJ, Calanzani N, et al. Preferences for place of death if faced with advanced cancer: a population survey in England, Flanders, Germany, Italy, The Netherlands, Portugal and Spain. *Ann Oncol* 2012; 23(8): 2006–2015.
- Dixon J, King D, Matosevic T, et al. *Equity in the provision of palliative care in the UK: review of evidence*. London: Personal Social Services Research Unit, London School of Economics and Political Science, 2015, <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/healthcaresystem/bulletins/nationalsurveyofbereavedpeoplevoices/england2015#relief-of-pain-in-the-last-3-months-of-life>
- ElMokhallati Y, Woodhouse N, Farragher T, et al. Specialist palliative care support is associated with improved pain relief at home during the last 3 months of life in patients with advanced disease: analysis of 5-year data from the national survey of bereaved people (VOICES). *BMC Med* 2019; 17(1): 50.
- Munn Z, Peters MDJ, Stern C, et al. Systematic review or scoping review? Guidance for authors when choosing between a systematic or scoping review approach. *BMC Med Res Methodol* 2018; 18(1): 143.
- Tricco AC, Lillie E, Zarin W, et al. PRISMA extension for Scoping Reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med* 2018; 169(7): 467–473.
- Bennett MI, Eisenberg E, Ahmedzai SH, et al. Standards for the management of cancer-related pain across Europe-A position paper from the EFIC Task Force on Cancer Pain. *Eur J Pain* 2019; 23(4): 660–668.
- National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology. In: *Adult cancer pain* (version 2. 2019), [09/05/19], 2019, https://www.nccn.org/professionals/physician_gls/pdf/pain.pdf
- NHS. Scottish palliative care guidelines – pain management 2019, <https://www.palliativecareguidelines.scot.nhs.uk/guidelines/pain/pain-management.aspx>
- Fallon M, Giusti R, Aielli F, et al. Management of cancer pain in adult patients: ESMO Clinical Practice Guidelines. *Ann Oncol* 2018; 29(Suppl. 4): iv166–iv191.
- Wordliczek J, Kotlinska-Lemieszek A, Leppert W, et al. Pharmacotherapy of pain in cancer patients – recommendations of the Polish Association for the Study of Pain, Polish Society of Palliative Medicine, Polish Society of Oncology, Polish Society of Family Medicine, Polish Society of Anaesthesiology and Intensive Therapy and Association of Polish Surgeons. *Pol Przegl Chir* 2018; 90(4): 55–84.
- WHO. *Guidelines for the pharmacological and radiotherapeutic management of cancer pain in adults and adolescents*. Geneva: World Health Organization, 2018, <https://www.ncbi.nlm.nih.gov/books/NBK537492/>

16. Jara C, Del Barco S, Gravalos C, et al. SEOM clinical guideline for treatment of cancer pain (2017). *Clin Transl Oncol* 2018; 20(1): 97–107.
17. National Clinical Effectiveness Committee. *Pharmacological management of cancer pain in adults*. National clinical guideline no. 9, 2015, https://health.gov.ie/wp-content/uploads/2015/11/Pharma-Mgmt-Cancer-Pain_web.pdf
18. Adam R, Burton CD, Bond CM, et al. Can patient-reported measurements of pain be used to improve cancer pain management? A systematic review and meta-analysis. *BMJ Support Palliat Care* 2017; 7(4). DOI: 10.1136/bmjspcare-2016-001137.
19. Fallon M, Walker J, Colvin L, et al. Pain management in cancer center inpatients: a cluster randomized trial to evaluate a systematic integrated approach – the Edinburgh pain assessment and management tool. *J Clin Oncol* 2018; 36(13): 1284–1290.
20. Finnerup NB, Haroutounian S, Kamerman P, et al. Neuropathic pain: an updated grading system for research and clinical practice. *Pain* 2016; 157(8): 1599–1606.
21. Rayment C, Hjermstad MJ, Aass N, et al. Neuropathic cancer pain: prevalence, severity, analgesics and impact from the European Palliative Care Research Collaborative-Computerised Symptom Assessment study. *Palliat Med* 2013; 27(8): 714–721.
22. Wiffen PJ, Wee B, Derry S, et al. Opioids for cancer pain – an overview of Cochrane reviews. *Cochrane Database Syst Rev* 2017; 7: CD012592.
23. Corli O, Floriani I, Roberto A, et al. Are strong opioids equally effective and safe in the treatment of chronic cancer pain? A multicenter randomized phase IV ‘real life’ trial on the variability of response to opioids. *Ann Oncol* 2016; 27(6): 1107–1115.
24. Higginson IJ and Gao W. Opioid prescribing for cancer pain during the last 3 months of life: associated factors and 9-year trends in a nationwide United Kingdom cohort study. *J Clin Oncol* 2012; 30(35): 4373–4379.
25. Gagnon B, Scott S, Nadeau L, et al. Patterns of community-based opioid prescriptions in people dying of cancer. *J Pain Symptom Manage* 2015; 49(1): 36–44.
26. Ziegler L, Mulvey M, Blenkinsopp A, et al. Opioid prescribing for patients with cancer in the last year of life: a longitudinal population cohort study. *Pain* 2016; 157(11): 2445–2451.
27. Ziegler LE, Craigs CL, West RM, et al. Is palliative care support associated with better quality end-of-life care indicators for patients with advanced cancer? A retrospective cohort study. *BMJ Open* 2018; 8(1): e018284.
28. Schmidt-Hansen M, Bennett MI, Arnold S, et al. Oxycodone for cancer-related pain. *Cochrane Database Syst Rev* 2017; 8: CD003870.
29. Hadley G, Derry S, Moore RA, et al. Transdermal fentanyl for cancer pain. *Cochrane Database Syst Rev* 2013; 10: CD010270.
30. Schmidt-Hansen M, Bromham N, Taubert M, et al. Buprenorphine for treating cancer pain. *Cochrane Database Syst Rev* 2015; 3: CD009596.
31. Wiffen PJ, Derry S, Naessens K, et al. Oral tapentadol for cancer pain. *Cochrane Database Syst Rev* 2015; 9: CD011460.
32. Bao YJ, Hou W, Kong XY, et al. Hydromorphone for cancer pain. *Cochrane Database Syst Rev* 2016; 10: CD011108.
33. Corli O, Roberto A, Corsi N, et al. Opioid switching and variability in response in pain cancer patients. *Support Care Cancer* 2019; 27(6): 2321–2327.
34. Nicholson AB, Watson GR, Derry S, et al. Methadone for cancer pain. *Cochrane Database Syst Rev* 2017; 2: CD003971.
35. Vayne-Bossert P, Afsharimani B, Good P, et al. Interventional options for the management of refractory cancer pain – what is the evidence. *Support Care Cancer* 2016; 24(3): 1429–1438.
36. Rauck RL, Cherry D, Boyer MF, et al. Long-term intrathecal opioid therapy with a patient-activated, implanted delivery system for the treatment of refractory cancer pain. *J Pain* 2003; 4(8): 441–447.
37. Smith TJ, Staats PS, Deer T, et al. Randomized clinical trial of an implantable drug delivery system compared with comprehensive medical management for refractory cancer pain: impact on pain, drug-related toxicity, and survival. *J Clin Oncol* 2002; 20(19): 4040–4049.
38. Chow E, Zeng L, Salvo N, et al. Update on the systematic review of palliative radiotherapy trials for bone metastases. *Clin Oncol* 2012; 24(2): 112–124.
39. O’Carrigan B, Wong MH, Willson ML, et al. Bisphosphonates and other bone agents for breast cancer. *Cochrane Database Syst Rev* 2017; 10: CD003474.
40. Porta-Sales J, Garzon-Rodriguez C, Llorens-Torrome S, et al. Evidence on the analgesic role of bisphosphonates and denosumab in the treatment of pain due to bone metastases: a systematic review within the European Association for Palliative Care guidelines project. *Palliat Med* 2017; 31(1): 5–25.
41. Hoskin P, Sundar S, Reczko K, et al. A multicenter randomized trial of ibandronate compared with single-dose radiotherapy for localized metastatic bone pain in prostate cancer. *J Natl Cancer Inst* 2015; 107(10): djv197.
42. Yamanaka M. A concept analysis of self-management of cancer pain. *Asia Pac J Oncol Nurs* 2018; 5(3): 254–261.
43. Adam R, Bond C and Murchie P. Educational interventions for cancer pain. A systematic review of systematic reviews with nested narrative review of randomized controlled trials. *Patient Educ Couns* 2015; 98(3): 269–282.
44. Bennett MI, Bagnall AM and Jose Closs S. How effective are patient-based educational interventions in the management of cancer pain? Systematic review and meta-analysis. *Pain* 2009; 143(3): 192–199.
45. Edwards Z, Ziegler L, Craigs C, et al. Pharmacist educational interventions for cancer pain management: a systematic review and meta-analysis. *Int J Pharm Pract* 2019; 27: 336–345.
46. Boger E, Ellis J, Latter S, et al. Self-management and self-management support outcomes: a systematic review and mixed research synthesis of stakeholder views. *PLoS ONE* 2015; 10(7): e0130990.
47. Johnston BM, Milligan S, Foster C, et al. Self-care and end of life care – patients’ and carers’ experience a qualitative study utilising serial triangulated interviews. *Support Care Cancer* 2012; 20(8): 1619–1627.
48. Oldenmenger WH, Geerling JI, Mostovaya I, et al. A systematic review of the effectiveness of patient-based educational

- interventions to improve cancer-related pain. *Cancer Treat Rev* 2018; 63: 96–103.
49. Ekstedt M and Rustoen T. Factors that hinder and facilitate cancer patients' knowledge about pain management—a qualitative study. *J Pain Symptom Manage* 2019; 57(4): 753–760.
 50. Lopez-Sendin N, Alburquerque-Sendin F, Cleland JA, et al. Effects of physical therapy on pain and mood in patients with terminal cancer: a pilot randomized clinical trial. *J Altern Complement Med* 2012; 18(5): 480–486.
 51. Pyszora A, Budzynski J, Wojcik A, et al. Physiotherapy programme reduces fatigue in patients with advanced cancer receiving palliative care: randomized controlled trial. *Support Care Cancer* 2017; 25(9): 2899–2908.
 52. Chevillat AL, Moynihan T, Herrin J, et al. Effect of collaborative telerehabilitation on functional impairment and pain among patients with advanced-stage cancer: a randomized clinical trial. *JAMA Oncol* 2019; 5: 644–652.
 53. Bradt J, Dileo C, Magill L, et al. Music interventions for improving psychological and physical outcomes in cancer patients. *Cochrane Database Syst Rev* 2016; 8: CD006911.
 54. Eagar K, Watters P, Currow DC, et al. The Australian Palliative Care Outcomes Collaboration (PCOC) – measuring the quality and outcomes of palliative care on a routine basis. *Aust Health Rev* 2010; 34(2): 186–192.
 55. Currow DC, Allingham S, Yates P, et al. Improving national hospice/palliative care service symptom outcomes systematically through point-of-care data collection, structured feedback and benchmarking. *Support Care Cancer* 2015; 23(2): 307–315.
 56. Wiffen PJ, Derry S, Moore RA, et al. Oral paracetamol (acetaminophen) for cancer pain. *Cochrane Database Syst Rev* 2017; 7: CD012637.
 57. Saragiotto BT, Machado GC, Ferreira ML, et al. Paracetamol for low back pain. *Cochrane Database Syst Rev* 2016; 6: CD012230.
 58. Leopoldino AO, Machado GC, Ferreira PH, et al. Paracetamol versus placebo for knee and hip osteoarthritis. *Cochrane Database Syst Rev* 2019; 2: CD013273.
 59. McNicol ED, Ferguson MC, Haroutounian S, et al. Single dose intravenous paracetamol or intravenous propacetamol for postoperative pain. *Cochrane Database Syst Rev* 2016; 5: CD007126.
 60. Straube C, Derry S, Jackson KC, et al. Codeine, alone and with paracetamol (acetaminophen), for cancer pain. *Cochrane Database Syst Rev* 2014; 9: CD006601.
 61. Ventafridda V, Tamburini M, Caraceni A, et al. A validation study of the WHO method for cancer pain relief. *Cancer* 1987; 59(4): 850–856.
 62. Bandieri E, Romero M, Ripamonti CI, et al. Randomized trial of low-dose morphine versus weak opioids in moderate cancer pain. *J Clin Oncol* 2016; 34(5): 436–442.
 63. Bell RF, Eccleston C and Kalso EA. Ketamine as an adjuvant to opioids for cancer pain. *Cochrane Database Syst Rev* 2017; 6: CD003351.
 64. Fallon MT, Wilcock A, Kelly CA, et al. Oral ketamine vs placebo in patients with cancer-related neuropathic pain: a randomized clinical trial. *JAMA Oncol* 2018; 4(6): 870–872.
 65. Hardy J, Quinn S, Fazekas B, et al. Randomized, double-blind, placebo-controlled study to assess the efficacy and toxicity of subcutaneous ketamine in the management of cancer pain. *J Clin Oncol* 2012; 30(29): 3611–3617.
 66. Brinck EC, Tiippana E, Heesen M, et al. Perioperative intravenous ketamine for acute postoperative pain in adults. *Cochrane Database Syst Rev* 2018; 12: CD012033.
 67. Assouline B, Tramer MR, Kreienbuhl L, et al. Benefit and harm of adding ketamine to an opioid in a patient-controlled analgesia device for the control of postoperative pain: systematic review and meta-analyses of randomized controlled trials with trial sequential analyses. *Pain* 2016; 157(12): 2854–2864.
 68. Schug SA and Goddard C. Recent advances in the pharmacological management of acute and chronic pain. *Ann Palliat Med* 2014; 3(4): 263–275.
 69. Peixoto RD and Hawley P. Intravenous lidocaine for cancer pain without electrocardiographic monitoring: a retrospective review. *J Palliat Med* 2015; 18(4): 373–377.
 70. Sharma S, Rajagopal MR, Palat G, et al. A phase II pilot study to evaluate use of intravenous lidocaine for opioid-refractory pain in cancer patients. *J Pain Symptom Manage* 2009; 37(1): 85–93.
 71. Lee JT, Sanderson CR, Xuan W, et al. Lidocaine for cancer pain in adults: a systematic review and meta-analysis. *J Palliat Med* 2019; 22(3): 326–334.
 72. Smith FL, Cichewicz D, Martin ZL, et al. The enhancement of morphine antinociception in mice by delta9-tetrahydrocannabinol. *Pharmacol Biochem Behav* 1998; 60(2): 559–566.
 73. Johnson JR, Burnell-Nugent M, Lossignol D, et al. Multicenter, double-blind, randomized, placebo-controlled, parallel-group study of the efficacy, safety, and tolerability of THC: CBD extract and THC extract in patients with intractable cancer-related pain. *J Pain Symptom Manage* 2010; 39(2): 167–179.
 74. Fallon MT, Albert Lux E, McQuade R, et al. Sativex oromucosal spray as adjunctive therapy in advanced cancer patients with chronic pain unrelieved by optimized opioid therapy: two double-blind, randomized, placebo-controlled phase 3 studies. *Br J Pain* 2017; 11(3): 119–133.
 75. Lichtman AH, Lux EA, McQuade R, et al. Results of a double-blind, randomized, placebo-controlled study of nabiximols oromucosal spray as an adjunctive therapy in advanced cancer patients with chronic uncontrolled pain. *J Pain Symptom Manage* 2018; 55(2): 179–188.
 76. Hauser W, Welsch P, Klose P, et al. Efficacy, tolerability and safety of cannabis-based medicines for cancer pain: a systematic review with meta-analysis of randomised controlled trials. *Schmerz* 2019; 33: 424–436.
 77. Derry S, Wiffen PJ, Moore RA, et al. Oral nonsteroidal anti-inflammatory drugs (NSAIDs) for cancer pain in adults. *Cochrane Database Syst Rev* 2017; 7: CD012638.
 78. Enthoven WT, Roelofs PD, Deyo RA, et al. Non-steroidal anti-inflammatory drugs for chronic low back pain. *Cochrane Database Syst Rev* 2016; 2: CD012087.
 79. McNicol ED, Ferguson MC and Schumann R. Single-dose intravenous diclofenac for acute postoperative pain in adults. *Cochrane Database Syst Rev* 2018; 8: CD012498.
 80. Finnerup NB, Attal N, Haroutounian S, et al. Pharmacotherapy for neuropathic pain in adults: a systematic review and meta-analysis. *Lancet Neurol* 2015; 14(2): 162–173.
 81. Mishra S, Bhatnagar S, Goyal GN, et al. A comparative efficacy of amitriptyline, gabapentin, and pregabalin in neuropathic cancer pain: a prospective randomized double-blind

- placebo-controlled study. *Am J Hosp Palliat Care* 2012; 29(3): 177–182.
82. Kane CM, Mulvey MR, Wright S, et al. Opioids combined with antidepressants or antiepileptic drugs for cancer pain: systematic review and meta-analysis. *Palliative Med* 2018; 32(1): 276–286.
83. Haywood A, Good P, Khan S, et al. Corticosteroids for the management of cancer-related pain in adults. *Cochrane Database Syst Rev* 2015; 4: CD010756.
84. Jaward LR, O’Neil TA, Marks A, et al. Differences in adverse effect profiles of corticosteroids in palliative care patients. *Am J Hosp Palliat Care* 2019; 36(2): 158–168.
85. Peng L, Min S, Zejun Z, et al. Spinal cord stimulation for cancer-related pain in adults. *Cochrane Database Syst Rev* 2015; 6: CD009389.
86. Paley CA, Johnson MI, Tashani OA, et al. Acupuncture for cancer pain in adults. *Cochrane Database Syst Rev* 2015; 10: CD007753.
87. Shin ES, Seo KH, Lee SH, et al. Massage with or without aromatherapy for symptom relief in people with cancer. *Cochrane Database Syst Rev* 2016; 6: CD009873.
88. Hurlow A, Bennett MI, Robb KA, et al. Transcutaneous electric nerve stimulation (TENS) for cancer pain in adults. *Cochrane Database Syst Rev* 2012; 3: CD006276.