



This is a repository copy of *Hypoxia modulates platelet purinergic signalling pathways.*

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/155062/>

Version: Published Version

Article:

Paterson, G.G., Young, J.M., Willson, J.A. et al. (9 more authors) (2019) Hypoxia modulates platelet purinergic signalling pathways. *Thrombosis and Haemostasis*, 120 (2). pp. 253-261. ISSN 0340-6245

<https://doi.org/10.1055/s-0039-3400305>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:
<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>



Hypoxia Modulates Platelet Purinergic Signalling Pathways

Gordon G. Paterson^{1,2} Jason M. Young^{1,2} Joseph A. Willson³ Christopher J. Graham^{1,2}
Rebecca C. Dru^{1,2} Eleanor W. Lee^{1,2} Greig S. Torpey^{1,2} Sarah R. Walmsley³ Melissa V. Chan⁴
Timothy D. Warner⁴ John Kenneth Baillie^{1,5,6} Alfred Arthur Roger Thompson^{1,7}

¹ APEX (Altitude Physiology Expeditions), Edinburgh, United Kingdom

² Edinburgh Medical School, University of Edinburgh, Edinburgh, United Kingdom

³ University of Edinburgh Centre for Inflammation Research, The Queen's Medical Research Institute, University of Edinburgh, Edinburgh, United Kingdom

⁴ Centre for Immunobiology, Blizard Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, United Kingdom

⁵ Division of Genetics and Genomics, The Roslin Institute, University of Edinburgh, Edinburgh, United Kingdom

⁶ Department of Anaesthesia, Critical Care and Pain Medicine, Royal Infirmary of Edinburgh, NHS Lothian, Edinburgh, United Kingdom

⁷ Department of Infection, Immunity and Cardiovascular Disease, University of Sheffield, Sheffield, United Kingdom

Address for correspondence Alfred Arthur Roger Thompson, BSc, MB ChB, MRCP, PhD, Department of Infection, Immunity and Cardiovascular Disease, University of Sheffield, M127a, Royal Hallamshire Hospital, Beech Hill Road, Sheffield, S10 2RX, United Kingdom (e-mail: R.Thompson@sheffield.ac.uk).

Thromb Haemost

Abstract

Background Hypoxia resulting from ascent to high-altitude or pathological states at sea level is known to increase platelet reactivity. Previous work from our group has suggested that this may be adenosine diphosphate (ADP)-specific. Given the clinical importance of drugs targeting ADP pathways, research into the impact of hypoxia on platelet ADP pathways is highly important.

Methods Optimal aggregometry was performed on plasma from 29 lowland residents ascending to 4,700 m, allowing systematic assessment of platelet reactivity in response to several platelet agonists. Aggregometry was also performed in response to ADP in the presence of inhibitors of the two main ADP receptors, P2Y₁ and P2Y₁₂ (MRS2500 and cangrelor, respectively). Phosphorylation of vasodilator-stimulated phosphoprotein (VASP), a key determinant of platelet aggregation, was analysed using the VASPFix assay.

Results Hypobaric hypoxia significantly reduced the ability of a fixed concentration of cangrelor to inhibit ADP-induced aggregation and increased basal VASP phosphorylation. However, in the absence of P2Y receptor inhibitors, we did not find evidence of increased platelet sensitivity to any of the agonists tested and found reduced sensitivity to thrombin receptor-activating peptide-6 amide.

Conclusion Our results provide evidence of increased P2Y₁ receptor activity at high altitude and suggest down-regulation of the P2Y₁₂ pathway through increased VASP phosphorylation. These changes in ADP pathway activity are of potential therapeutic significance to high-altitude sojourners and hypoxic sea level patients prescribed platelet inhibitors and warrant further investigation.

Keywords

- ▶ hypoxia
- ▶ platelet physiology
- ▶ anti-platelet agents
- ▶ ADP receptors
- ▶ high altitude

received
April 24, 2019
accepted after revision
September 29, 2019

DOI <https://doi.org/10.1055/s-0039-3400305>.
ISSN 0340-6245.

© Georg Thieme Verlag KG
Stuttgart · New York

License terms



Introduction

Acute hypobaric hypoxia, such as that induced by ascent to high altitude, has long been considered to produce a thrombogenic phenotype.¹ Consistent with this, epidemiological studies report a markedly increased risk of strokes at high altitude: up to 30 times that at sea level.² Furthermore, these events are reported to occur in younger patients with fewer cardiovascular risk factors.³ Although some studies examining the effect of acute hypoxia on coagulation have reported minimal changes in coagulation, they were investigating the 'economy class syndrome' and thus exposure to hypoxia was brief.^{4,5} On the contrary, studies examining a longer exposure to hypoxia have demonstrated a hypercoagulable state.^{6–8}

Platelets are key to haemostasis^{9,10} and appear to have an important role in the hypercoagulable state induced by hypoxia: expression of soluble P-selectin, an *in vivo* platelet activation marker, was shown to be increased 2.5-fold after ascent to high altitude.¹¹ Furthermore, hypoxia has recently been shown to significantly alter the platelet proteome, including up-regulation of calpain small subunit 1.¹² Calpains are calcium-dependent proteases involved in various physiological processes including platelet activation.¹³ The same study also found increased intracellular calcium concentration in hypoxic platelets¹² which is a common downstream effect of platelet activation by several agonists, including adenosine diphosphate (ADP).¹⁰

Recent work from our group has demonstrated a hypercoagulable state in 63 subjects participating in a controlled, non-exertional sojourn to 5,300 m.⁶ Multiplate aggregation assays using fixed doses of several platelet activators at this altitude suggested that hypoxia-induced platelet hyper-reactivity is specific to ADP as responses to thrombin receptor-activating peptide (TRAP) and collagen were not altered.⁶ It is known that ADP acts on two pro-aggregatory pathways, via P2Y₁ and P2Y₁₂ receptors,¹⁴ and in light of our previous findings we aimed to further investigate these pathways. P2Y₁₂ inhibitors are widely used anti-platelet medications, and effects of hypoxia on their efficacy are of high clinical relevance. This is particularly important given the increasing number of older sojourners to high altitude, who may have cardiovascular co-morbidities.¹⁵

Our study is the first to our knowledge to investigate the effect of hypoxia on the major platelet activation pathways with full concentration–response curves. Optimul aggregometry is a recently developed assay which applies the principles of light-transmission aggregometry to a 96-well format, with platelet agonists lyophilised in each well.¹⁶ This assay requires significantly less plasma volume than traditional aggregometry, thus facilitating systematic assessment of platelet aggregation in response to several agonists. *Ex vivo* use of ADP receptor inhibitors also permitted further detailed study of the purinergic pathways in a group of healthy, lowland volunteers ascending to 4,700 m.

Methods

Data were collected from 29 participants before and during the APEX 5 Expedition. All participants had no known

cardiovascular or respiratory conditions and no pre-existing coagulopathy. Participants were asked to refrain from alcohol and anti-platelet medications during the week prior to sampling. This study was approved by the ACCORD Research Ethics Committee (17-HV-030) and all participants gave informed consent as per the Declaration of Helsinki. Reagents were supplied by Sigma-Aldrich (Irvine, United Kingdom) unless otherwise specified.

Ascent Profile and Sample Collection

Participants were resident at < 250 m above sea level and had not travelled to high altitude (> 2,500 m) in the 2 months prior to the study. The ascent profile and study timeline are summarised in **Fig. 1**. We chose to perform aggregometry assays on day 6 (one day following ascent to 4,700 m to examine effects of sub-acute hypoxia) and day 11 (the time point we had previously observed changes in Multiplate aggregation with ADP⁶). Venepuncture was performed using 21G needles (Williams Medical, Rhymney, United Kingdom) into citrated tubes. Platelet-rich plasma (PRP) was prepared by centrifugation of whole blood at 175 × g for 15 minutes (EBA 280, Hettich, Tuttlingen, Germany). Platelet-poor plasma (PPP) was generated by further centrifugation at 3,624 × g for 2 minutes. Peripheral oxygen saturation (SpO₂) was measured using a pulse oximeter (SM-100, Santamedical, Tustin, United States) at baseline and every day of the expedition.

Optimul Aggregometry

Modified optimul aggregometry plates were prepared as previously described by Chan and Warner.¹⁶ In brief, pre-diluted platelet agonists were added to individual wells of a gelatin-coated 96-well plate and the plates were lyophilised. Plates were vacuum-sealed and protected from light before use. Each 96-well plate contained lyophilised concentration ranges of arachidonic acid (0.03–1 mM), ADP (0.005–40 μM),




Fig. 1 Ascent profile. Baseline testing was performed in April 2017, 2 months prior to the expedition. Subjects landed in La Paz (3,700 m) where they spent four nights before ascending to Huayna Potosi Base Camp (4,700 m) by bus on day 5 where they stayed for the remainder of the study. Optimul aggregometry, VASPFix and full blood count samples were collected at baseline and on day 11. On day 6, only optimul aggregometry was performed.

collagen (0.01–40 µg/mL), epinephrine (0.0004–10 µM), TRAP-6 amide (0.03–40 µM), and U46619 (thromboxane mimetic, 0.005–40 µM).

PRP was aliquoted and incubated for 30 minutes (at 37°C) alone or in the presence of the P2Y₁ inhibitor MRS2500 (1 µM, Tocris Bioscience, Bristol, United Kingdom) or the P2Y₁₂ inhibitor cangrelor (100 nM, The Medicines Company, New Jersey, United States). After incubation, 40 µL of PRP (plus or minus inhibitor) was quickly added to wells containing the lyophilised platelet agonists. PPP was added to four agonist-free wells to provide a signal equivalent to 100% aggregation. PRP was also added to agonist-free wells and wells containing vehicle alone as a 0% aggregation control. Plates were then placed on a thermal shaker (BioShake iQ, QInstruments, Jena, Germany) for 5 minutes (37°C, 1,200 revolutions per minute). Light absorbance was read at 595 nm in a 96-well plate reader (SPECTROstar Nano, BMG LABTECH, Aylesbury, United Kingdom).

Percentage aggregation was calculated using absorbance values for PPP (100%) and PRP (0%) in agonist-free wells as reference values. A visual inspection of concentration–response curves was conducted to remove any clearly erroneous curves (deviating from a sigmoid shape). Data were also removed if they showed no response, with response defined as greater than 30% aggregation in response to two doses of agonist.

Vasodilator-Stimulated Phosphoprotein Phosphorylation

Vasodilator-stimulated phosphoprotein (VASP) is a platelet protein whose phosphorylation is modulated by cyclic adenosine monophosphate (cAMP).¹⁷ cAMP production is inhibited by the action of G_{ai} on adenylate cyclase secondary to activation of P2Y₁₂ receptors.¹⁸ The degree of VASP phosphorylation can thus be used as a marker of P2Y₁₂ activity.¹⁹ The VASPFix assay²⁰ was used to quantify VASP phosphorylation.

Aliquots of PRP were incubated for 6 minutes in one of three conditions: phosphate-buffered saline (PBS), a prostacyclin analogue iloprost (1 nM), or ADP (5 µM) + iloprost (1 nM) and mixed in a 1:5 ratio with VASPFix (Platelet Solutions Ltd., Nottingham, United Kingdom), vortexed, and snap frozen on dry ice. Samples from high altitude were transported to the United Kingdom for analysis on dry ice by a specialist company.

Flow cytometry was performed and the median fluorescein isothiocyanate fluorescence (mf) recorded for each condition. Iloprost induces maximal VASP phosphorylation with PBS acting as a negative control.

If iloprost did not induce phosphorylation (i.e., if mf (iloprost) < mf (saline)), it was assumed that there was a technical failure and these samples removed from the analysis. Similarly, if mf (iloprost + ADP) was greater than mf (iloprost), the sample was also removed from analysis.

Full Blood Count

Three millilitres of blood were collected into an ethylenediaminetetraacetic acid blood tube (Sarstedt Ltd., Leicester, United Kingdom) and samples analysed within 24 hours by clinical haematology laboratories (NHS Lothian Laboratories, Edin-

burgh, United Kingdom, and SELADIS, Universidad Mayor de San Andrés, La Paz, Bolivia).

Statistics

Optimal aggregometry data were fit to Eq. (1) by least squares, non-linear regression using the `scipy.optimize` library (for Python 2.7.2).

Eq. (1): Agonist-response equation where y is the percentage aggregation, x is $\log[\text{agonist}]$, EC_{50} the concentration resulting in half maximal aggregation, $Top/Bottom$ the maximum/minimum aggregation and the Hill Slope the steepness of the curve. For studies examining the dose–response curve of cangrelor, IC_{50} replaces EC_{50} .

$$y = \frac{Bottom + (Top - Bottom)}{1 + 10^{(LogEC_{50} - x) \cdot Hill\ Slope}}$$

The concentration resulting in half-maximal aggregation (EC_{50}) was used to compare curves from each time point. For comparisons between ADP concentration–response curves in the presence of a fixed antagonist, maximum response (R_{max}) to ADP was also calculated.

Outlier identification was based on difference of fits analysis.²¹ The effect of each point on EC_{50} (ΔEC_{50}) and Hill Slope ($\Delta Hill$) was calculated by iterative removal of points and refitting the curves. ΔEC_{50} and $\Delta Hill$ values were reviewed for comparable data (all points in all concentration–response curves for one agonist at one time point) and any points falling out with two standard deviations (ΔEC_{50}) or four standard deviations ($\Delta Hill$) were considered outliers. Additionally, for concentration–response curves in the presence of inhibitors, if the final point was less than 75% of the point preceding it, it too was considered an outlier, to ensure validity of the R_{max} parameter.

Wherever possible, paired statistics were used; however, the optimal aggregometry data were unpaired. For paired comparisons of two time points, paired Student's t -tests were performed using the `scipy.stats` library.²² For paired data compared over three time points (SpO_2), a one-way repeated-measure analysis of variance (ANOVA) was conducted with a Tukey's HSD post hoc test using Prism 5.0 (GraphPad, La Jolla, California, United States). For optimal aggregometry data, comparisons between three time points were analysed using one-way ANOVA with Tukey's HSD post hoc test using R 3.4.2.²³ Whenever multiple comparisons were made, between multiple agonists or conditions, p -values reported were adjusted using a Bonferroni correction. Statistical significance was set as 0.05.

Results

Baseline characteristics of subjects are summarised in ► **Table 1**.

All but one subject completed the study, and data have been included until dropout for unpaired analyses. The sojourn at high altitude induced a marked hypoxaemia (► **Fig. 2A**), which was slightly more pronounced on day 11 than day 6. Platelet counts were also significantly elevated by

Table 1 Baseline characteristics

Characteristic	APEX 5 cohort
Sex (male/female)	9/20
Mean BMI (range)	22.3 (17.6–28.8)
Mean age (range)	20.7 (18–26)

Note: Body mass index (BMI) has units kg/m², and age is measured in years.




Fig. 2 Platelet counts and oxygen saturations. (A) Oxygen saturation was measured at baseline and on day 6 and day 11 of the expedition. Data points are represented as semi-transparent circles, with summary box plots superimposed. Data were analysed by a one-way repeated-measure analysis of variance (ANOVA) followed by a Tukey's HSD test. (B) Platelet count was measured at baseline and on day 11 of the expedition. Data points are represented as semi-transparent circles, with summary box plots superimposed. Data were analysed by paired Student's *t*-test. ****p* < 0.001, **p* < 0.05.

hypoxia (►Fig. 2B). Haemoglobin, haematocrit and white cell count data are provided in ►Supplementary Table S1 (available in the online version).

Hypoxic Platelets are Less Sensitive to TRAP-6 Amide

The effect of hypoxia on various platelet agonists was compared between baseline, day 6 and day 11. No platelet activation pathways were found to be sensitised by hypoxia; however, platelets became less sensitive to TRAP-6 amide (►Fig. 3E, ►Table 2, *p* < 0.01). This effect was only present on day 11.

Hypoxia Modulates Purinergic Signalling

In the presence of 1 μM P2Y₁ inhibitor MRS2500, both EC₅₀ and R_{max} were unchanged by hypoxia (►Fig. 4A, ►Table 2). Likewise, in the presence of 100 nM P2Y₁₂ inhibitor cangrelor, EC₅₀ remained unchanged after 11 days of hypoxia. While EC₅₀ remained unchanged at altitude, R_{max} was markedly increased on both days 6 and 11 compared with baseline (►Fig. 4B, ►Table 2, *p* < 0.01).

Hypoxia Modulates Basal VASP Phosphorylation in Platelets

Compared with baseline, after 11 days at high altitude there was a highly significant increased mf in the PBS-treated

samples (►Fig. 5A). There were no differences between baseline and day 11 in other conditions (►Fig. 5A). There was a marked reduction in the mf percentage increase induced by iloprost at high altitude (►Fig. 5B), but no change in mf reduction induced by the addition of ADP (►Fig. 5C).

Discussion

There is a growing body of evidence that hypoxia induces hyper-reactivity in platelets, a phenomenon important to both altitude physiology and sea-level pathophysiology. We found that hypoxia increases maximal aggregation to ADP in the presence of cangrelor and down-regulates platelet sensitivity to TRAP-6 amide. However, we did not demonstrate sensitisation to any of the platelet activators tested. Finally, and more strikingly, we have demonstrated modulation of basal VASP phosphorylation, a key step in the P2Y₁₂ signalling pathway.

Our results contrast with a previous study from our group which reported increased platelet sensitivity to ADP.⁶ Our previous findings were based on Multiplate analysis and discrepancies could be related to the impact of platelet count on the two different assays: Multiplate is more sensitive to platelet count than optumul aggregometry.^{24,25} This, however, would not explain why our previous findings were specific to ADP. Perhaps more likely is that differences relate to the parameter of platelet function that was tested. Multiplate is based on electrical-impedance aggregometry measured in real-time over 6 minutes, with the area under the impedance time curve as the final outcome.²⁶ This outcome is therefore dependent on both the final amplitude of the response and the rate at which this is reached. Optumul aggregometry, however, is based on one measurement at 5 minutes. Conceivably, differences in the ADP sensitivity could be time dependent, given the time sensitivity of phosphorylation²⁷ and changes in purinergic receptor expression²⁸ induced by ADP stimulation.

While we have not resolved the uncertainty surrounding ADP sensitivity, our study did reveal interesting changes in purinergic signalling. The R_{max} induced by ADP in the presence of 100 nM cangrelor was significantly higher at altitude (►Fig. 4B, ►Table 2). This could be due to increased P2Y₁ receptor activity. An increase in P2Y₁ activity may imply a difference in receptor expression or modulation of receptor activity since G-protein-coupled receptor activity can be regulated by multiple mechanisms.²⁹ Increased P2Y₁ activity would provide an interesting link to the proteomic data of Tyagi et al who described increased calpain expression and activity in rats exposed to hypoxia for 6 hours,¹² since a downstream consequence of P2Y₁ activation is increased intracellular calcium.³⁰

Well-controlled, high-altitude expeditions are excellent models of acute hypoxia³¹ and their findings are often relevant to hypoxic patients at sea level. In this regard, patients with chronic hypoxic diseases such as chronic obstructive pulmonary disease and obstructive sleep apnoea (OSA) have an increased risk of thrombotic events such as myocardial infarction and stroke.^{32,33} Furthermore, these conditions are also associated with increased platelet reactivity, which in the case of OSA is reversed on correction of hypoxia.^{34–36} However, the




Fig. 3 The effect of hypoxia on platelet activation pathways. Dose–response curves of platelet aggregation in response to (A) arachidonic acid (AA), (B) adenosine diphosphate (ADP), (C) collagen, (D) epinephrine, (E) thrombin receptor-activating peptide (TRAP)-6 amide and (F) U46619. Data are mean percentage aggregation ± standard error of the mean (SEM), and best fit curves optimised to these mean values. EC₅₀s were compared by one-way analysis of variance (ANOVA) followed by Tukey’s HSD post hoc tests where appropriate. *p*-Values reported were adjusted to account for family-wise error rate using Bonferroni corrections. ***p* < 0.01 vs. baseline.

Table 2 Summary results of optimul aggregometry data

EC ₅₀ values			
	Baseline	Day 6	Day 11
AA	-3.695 (0.053)	-3.696 (0.039)	-3.715 (0.064)
ADP	-5.990 (0.096)	-6.037 (0.122)	-5.912 (0.078)
Collagen	-6.776 (0.111)	-7.009 (0.107)	-6.853 (0.088)
Epinephrine	-6.339 (0.194)	-6.805 (0.144)	-6.584 (0.140)
TRAP-6 amide	-6.314 (0.105)	-6.221 (0.064)	-5.759 (0.084) ^a
U46619	-6.979 (0.122)	-6.652 (0.099)	-6.545 (0.122)
ADP + MRS2500	-5.280 (0.082)	-5.222 (0.063)	-5.244 (0.064)
ADP + Cangrelor	-5.186 (0.062)	-5.318 (0.068)	-5.335 (0.079)
R _{max} values			
ADP + MRS2500	95.230 (2.835)	100.413 (2.159)	96.638 (2.788)
ADP + Cangrelor	80.005 (4.177)	97.927 (2.557) ^a	97.186 (2.398) ^a

Abbreviations: AA, arachidonic acid; ADP, adenosine diphosphate; PRP, platelet-rich plasma; TRAP, thrombin receptor-activating peptide. Note: Dose–response curves of platelet aggregation were performed in response to AA, ADP, collagen, epinephrine, TRAP-6 amide, and U46619. Aggregometry was also performed on PRP incubated for 30 minutes with 1 μM MRS2500 (P2Y₁ inhibitor) or 100 nM cangrelor (P2Y₁₂ inhibitor) with ADP as the agonist. For experiments with inhibitors, the maximal response (R_{max}) was also calculated. EC₅₀ and R_{max} values were compared by one-way analysis of variance (ANOVA) followed by Tukey’s HSD post hoc tests where appropriate. *p*-Values reported were adjusted to account for family-wise error rate using Bonferroni corrections.

^a*p* < 0.01 vs. baseline.




Fig. 4 The effect of hypoxia on adenosine diphosphate (ADP)-induced platelet aggregation in the presence of fixed doses of inhibitors. Platelet-rich plasma (PRP) was incubated for 30 minutes with (A) 1 μ M MRS2500 (P2Y₁ inhibitor) or (B) 100 nM cangrelor (P2Y₁₂ inhibitor). Data are mean percentage aggregation \pm standard error of the mean (SEM), and best fit curves optimised to these mean values. EC₅₀ and R_{max} values were compared by one-way analysis of variance (ANOVA) followed by Tukey's HSD post hoc tests where appropriate. *p*-Values reported were adjusted to account for family-wise error rate using Bonferroni corrections. ***p* < 0.01 vs. baseline.




Fig. 5 The effect of hypoxia on platelet vasodilator-stimulated phosphoprotein (VASP) phosphorylation. Platelet-rich plasma (PRP) was incubated for 6 minutes with phosphate-buffered saline (PBS), iloprost or adenosine diphosphate (ADP) + Iloprost before addition of VASPFix. Flow cytometry was used to identify the fluorescein isothiocyanate (FITC) median fluorescence (MF), reflecting the degree of VASP phosphorylation. (A) Raw FITC MF for each condition at each time point. Data are mean FITC MF \pm standard error of the mean (SEM). Data were compared by paired *t*-tests with *p*-values adjusted by Bonferroni correction. (B) Percentage increase in FITC MF induced by addition of iloprost. (C) Percentage decrease induced by ADP addition. (B, C) Individual data points are represented by semi-translucent circles with box plots superimposed. Data were compared by paired *t*-tests. ***p* < 0.01, ****p* < 0.001.

clinical relevance of isolated, increased P2Y₁ activity is unclear since overall sensitivity to ADP was unaltered (**Fig. 3B**, **Table 2**). Nonetheless, increased P2Y₁ receptor activity has been postulated to contribute to P2Y₁₂ inhibitor resistance,^{37,38} although evidence for this is lacking and to our knowledge no

study has investigated the impact of hypoxia on P2Y₁₂ inhibitor resistance.

An alternative explanation for the change in R_{max} with cangrelor is an increase in P2Y₁₂ expression at altitude. Interestingly, a recent whole blood microarray study showed a twofold increase in P2Y₁₂ expression in well acclimatised sojourners at high altitude.³⁹ These findings were not validated in isolated platelets or by quantitative polymerase chain reaction. With either an increase in P2Y₁ or P2Y₁₂ expression, a change in sensitivity to ADP might be expected in the absence of inhibitors. However, as we detected no change in sensitivity to ADP, we propose that our finding of increased basal VASP phosphorylation is a compensatory mechanism that is also activated by hypoxia. Together with a lack of change in maximum phosphorylation induced by iloprost (**Fig. 5B**) and a reduction in the percentage increase in phosphorylation induced by iloprost (**Fig. 5C**), this suggests that the basal ratio of VASP:VASP-P is altered by hypoxia. Although we did not observe reduced P2Y₁₂ activity in response to ADP (**Fig. 5A, C**), we only studied a single high concentration of ADP (5 μ M), a concentration which induces near-maximal aggregation (**Fig. 3B**). It may be that lower concentrations of ADP are unable to overcome increased basal VASP phosphorylation. If this is the case, it could represent a compensatory response to increased P2Y₁ receptor activity or P2Y₁₂ expression, the explanations proposed for the reduction in cangrelor efficacy. VASP-P is regulated by several mediators that are altered in hypoxic conditions. For example, exhaled nitric oxide (NO) levels increased over 48 hours in healthy subjects exposed acutely to 4,559 m,⁴⁰ while raised circulating cyclic guanosine monophosphate (cGMP) levels and NO metabolites (nitrite and nitrate) were reported following gradual ascent to 5,200 m.⁴¹ NO inhibits platelet aggregation, at least in part via cGMP-mediated VASP phosphorylation as platelet adhesion to injured vessel walls could not be inhibited by NO in VASP-deficient mice.⁴² Interestingly, recent work demonstrated that nitrite led to VASP phosphorylation in isolated platelets in the presence of deoxygenated red blood cells.⁴³ Thus, either increased NO synthase or nitrite reductase activity could explain our finding of increased basal VASP phosphorylation at altitude.⁴⁴ **Fig. 6** summarises the proposed changes to purinergic signalling induced by hypoxia; however, this model will require further investigation to confirm its validity.

Kiouptsi et al examined the effects of brief exposure (30 minutes) of washed human platelets to extreme and moderate hypoxia (1 and 8% oxygen, respectively) on ex vivo expression of activated $\alpha_{IIb}\beta_3$ (a key integrin in platelet aggregation⁴⁵) and aggregation. They reported that platelets exposed to extreme hypoxia had reduced expression of activated $\alpha_{IIb}\beta_3$ in response to ADP stimulation.⁴⁶ Our proposed model of hypoxia-induced alterations to purinergic signalling may offer a mechanistic explanation to these findings since an increase in VASP phosphorylation would attenuate P2Y₁₂ activity and reduce ADP-induced activation of $\alpha_{IIb}\beta_3$.⁴⁷⁻⁴⁹ Kiouptsi et al also demonstrated that brief, extreme hypoxic exposure (1% oxygen, 30 minutes) reduced aggregation of hypoxic washed platelets in response to TRAP-6. However, sensitivity of PRP to TRAP-6 amide was unchanged.⁴⁶ Our data did not show any




Fig. 6 Proposed model of hypoxia-induced changes to purinergic signalling. Our findings are in red, while those of other studies examining the impact of hypoxia on platelet signalling are in blue. Our data suggest that hypoxia up-regulates the basal level of vasodilator-stimulated phosphoprotein (VASP) phosphorylation, a key determinant of platelet aggregation downstream of the P2Y₁₂ receptor. Other work demonstrating increased nitric oxide (NO) levels at altitude provides a possible mechanism for this finding.⁴¹ Phosphorylated-VASP (VASP-P) inhibits the expression of active α_{IIb}β₃ expression, which links to Kiouptsi et al's finding that hypoxic platelets have lower expression of active α_{IIb}β₃ in response to adenosine diphosphate (ADP).⁴⁶ Increased P2Y₁ pathway activity would be consistent with Tyagi et al's finding that calpains are up-regulated by hypoxia.¹² Since we found no change in overall sensitivity to ADP, it may be that the changes produced by hypoxia in these two pathways counteract one another. AC, adenylate cyclase; ADP, adenosine diphosphate; cAMP, cyclic adenosine monophosphate; cGMP, cyclic guanine monophosphate; GC, guanylate cyclase; PI3K, phosphoinositide 3-kinase.

difference in TRAP-6 sensitivity on day 6 (► **Fig. 3E**), but reduced sensitivity was observed after a longer duration of hypoxia. It could be that hypoxia-induced changes in sensitivity to TRAP-6 amide evolve over time, and only after 11 days was the effect on platelets pronounced enough to be seen in PRP. Interestingly, transcription of the receptor for TRAP-6 amide, protease-activated receptor 1, has also previously been shown to be down-regulated in hypoxic cancer cells,⁵⁰ but this has not been investigated in platelets and would be an interesting topic for further research.

This study has several limitations. First, it is difficult to predict the clinical relevance of subtle changes to signalling when overall sensitivity to the agonist is not altered. Additionally, our study only examined the purinergic pathways in detail, and it is possible that other subtle differences exist in other signalling pathways. Finally, our study had a slight gender imbalance, and sex has been shown to affect platelet aggregation and purinergic signalling,⁵¹ implying our results may be more relevant to women than men.

Detailed investigation of purinergic signalling provided evidence of increased VASP phosphorylation and impaired inhibition of aggregation by a P2Y₁₂ antagonist. We did not, however, find evidence of increased sensitivity to any platelet agonists despite the substantial evidence of hypoxia-induced

platelet hyper-reactivity in the literature. However, the observed changes in aggregation in response to an ADP antagonist are of potential therapeutic significance to high-altitude sojourners and hypoxic sea-level patients prescribed platelet inhibitors and warrant further investigation.

What is known about this topic?

- Exposure to hypobaric hypoxia is thought to induce a thrombogenic phenotype, with increased risk of stroke in those residing at high altitude.
- The platelet proteome is altered by hypoxia and previous work has suggested platelet sensitivity to ADP is increased at altitude.

What does this paper add?

- Using optimal aggregometry, we show no significant changes in the dose–response of several platelet receptor agonists.
- Hypoxia suppressed the extent to which cangrelor inhibited platelet aggregation to ADP, a finding with potential implications for hypoxic patients taking platelet inhibitors.

Authors' Contributions

G.G.P., M.V.C., T.D.W. and A.A.R.T. conceived the study and designed the experiments; G.G.P., J.M.Y., J.A.W., C.J.G., R.C. D., E.W.L., G.S.T., M.V.C. and S.R.W. performed experiments and assisted with logistics; G.G.P., M.V.C., T.D.W., J. K.B. and A.A.R.T. analysed and interpreted the data; G.G.P. and A.A.R.T. drafted the manuscript with contributions from all authors.

Funding

This study received funding from the Wilderness Medical Society Charles S. Houston Grant, supported by the Academy of Wilderness Medicine. It also received funding from British Heart Foundation, (Grant/Award Number: 'FS/18/13/3328'; 'PG/15/47/31591'), Wilderness Medicine Society (Grant/Award Number: 'Charles S. Houston grant') and Wellcome Trust, (Grant/Award Number: '103258/Z/13/Z,A').

Conflict of Interest

The authors declare no competing financial interests. A.A. R.T. reports non-financial support from Actelion UK Ltd, outside the submitted work.

Acknowledgments

We thank the APEX 5 expedition members for their participation. We thank the following companies for loan of equipment and technical support: Platelet Solutions Ltd., BMG LABTECH Ltd., Hettich Centrifuges, QInstruments, Genlab and Philip Harris Ltd. These companies had no role

in the study design or interpretation of the results. We thank Dr. Zorka Castillo (Universidad Mayor de San Andrés, La Paz, Bolivia), Dr. Martin Waterfall (Flow Cytometry Facility, University of Edinburgh) and Dr. Craig Nash (University of Michigan) for technical support. J.K.B. acknowledges funding support from a Wellcome-Beit Intermediate Clinical Fellowship (103258/Z/13/Z/A), BBSRC Institute Strategic Program Grant and the UK Intensive Care Foundation. A.A.R.T. and M.C. acknowledge funding support from the British Heart Foundation (Intermediate Clinical Fellowship to A.A.R.T., FS/18/13/3328; Project Grant supporting M.C., PG/15/47/31591). S.R.W. acknowledges support from a Wellcome Trust Senior Clinical Fellowship (209220). G.G.P. acknowledges funding support from a British Society of Haemostasis and Thrombosis travel grant and the Principal's Go Abroad fund (University of Edinburgh). The expedition received funding from the following charitable organisations: The JABBS Foundation, Scottish Mountaineering Trust, Gilchrist Educational Trust and the Mount Everest Foundation.

References

- Bärtsch P. How thrombogenic is hypoxia? *JAMA* 2006;295(19):2297–2299
- Anand AC, Jha SK, Saha A, Sharma V, Adya CM. Thrombosis as a complication of extended stay at high altitude. *Natl Med J India* 2001;14(04):197–201
- Jha SK, Anand AC, Sharma V, Kumar N, Adya CM. Stroke at high altitude: Indian experience. *High Alt Med Biol* 2002;3(01):21–27
- Schaber M, Leichtfried V, Fries D, et al. Influence of acute normobaric hypoxia on hemostasis in volunteers with and without acute mountain sickness. *BioMed Res Int* 2015;2015:593938
- Toff WD, Jones CI, Ford I, et al. Effect of hypobaric hypoxia, simulating conditions during long-haul air travel, on coagulation, fibrinolysis, platelet function, and endothelial activation. *JAMA* 2006;295(19):2251–2261
- Rocke AS, Paterson GG, Barber MT, et al. Thromboelastometry and platelet function during acclimatization to high altitude. *Thromb Haemost* 2018;118(01):63–71
- Pichler Hefti J, Risch L, Hefti U, et al. Changes of coagulation parameters during high altitude expedition. *Swiss Med Wkly* 2010;140(7-8):111–117
- Ninivaggi M, de Laat M, Lancé MMD, et al. Hypoxia induces a prothrombotic state independently of the physical activity. *PLoS One* 2015;10:e0141797
- Nieswandt B, Pleines I, Bender M. Platelet adhesion and activation mechanisms in arterial thrombosis and ischaemic stroke. *J Thromb Haemost* 2011;9(Suppl 1):92–104
- Jennings LK. Mechanisms of platelet activation: need for new strategies to protect against platelet-mediated atherothrombosis. *Thromb Haemost* 2009;102(02):248–257
- Lehmann T, Mairbäurl H, Pleisch B, Maggiorini M, Bärtsch P, Reinhart WH. Platelet count and function at high altitude and in high-altitude pulmonary edema. *J Appl Physiol* (1985) 2006;100(02):690–694
- Tyagi T, Ahmad S, Gupta N, et al. Altered expression of platelet proteins and calpain activity mediate hypoxia-induced prothrombotic phenotype. *Blood* 2014;123(08):1250–1260
- Fox JE, Taylor RG, Taffarel M, Boyles JK, Goll DE. Evidence that activation of platelet calpain is induced as a consequence of binding of adhesive ligand to the integrin, glycoprotein IIb-IIIa. *J Cell Biol* 1993;120(06):1501–1507
- Jacobson KA, Deflorian F, Mishra S, Costanzi S. Pharmacology of the platelet purinergic receptors. *Purinergic Signal* 2011;7(03):305–324
- Keyes LE, Mather L, Duke C, et al. Older age, chronic medical conditions and polypharmacy in Himalayan trekkers in Nepal: an epidemiologic survey and case series. *J Travel Med* 2016;23(06):1–6
- Chan MV, Warner TD. Standardised optical multichannel (optimum) platelet aggregometry using high-speed shaking and fixed time point readings. *Platelets* 2012;23(05):404–408
- Butt E, Abel K, Krieger M, et al. cAMP- and cGMP-dependent protein kinase phosphorylation sites of the focal adhesion vasodilator-stimulated phosphoprotein (VASP) in vitro and in intact human platelets. *J Biol Chem* 1994;269(20):14509–14517
- Hollopeter G, Jantzen H-M, Vincent D, et al. Identification of the platelet ADP receptor targeted by antithrombotic drugs. *Nature* 2001;409(6817):202–207
- Schwarz UR, Geiger J, Walter U, Eigenthaler M. Flow cytometry analysis of intracellular VASP phosphorylation for the assessment of activating and inhibitory signal transduction pathways in human platelets—definition and detection of ticlopidine/clopidogrel effects. *Thromb Haemost* 1999;82(03):1145–1152
- Glenn JR, Dovlatova N, White AE, Dhillon K, Heptinstall S, Fox SC. 'VASPFix' for measurement of VASP phosphorylation in platelets and for monitoring effects of P2Y12 antagonists. *Thromb Haemost* 2014;111(03):539–548
- Belsley DA, Kuh E, Welsch RE. Detecting Influential Observations and Outliers. In: *Regression Diagnostics: Identifying Influential Data and Sources of Collinearity*. Hoboken: John Wiley & Sons, Inc; 1980:6–84
- Jones E, Oliphant T, Peterson P, et al. *SciPy: open source scientific tools for Python* [Internet]. 2001. Available at: <http://www.scipy.org/>. Accessed October 16, 2019
- R Core Team. *R: a language and environment for statistical computing*. R Foundation for Statistical Computing Vienna, Austria [Internet]. 2019. Available at: <http://www.r-project.org/>. Accessed October 16, 2019
- Würtz M, Hvas A-M, Kristensen SD, Grove EL. Platelet aggregation is dependent on platelet count in patients with coronary artery disease. *Thromb Res* 2012;129(01):56–61
- Chan MV, Armstrong PCJ, Papalia F, Kirkby NS, Warner TD. Optical multichannel (optimum) platelet aggregometry in 96-well plates as an additional method of platelet reactivity testing. *Platelets* 2011;22(07):485–494
- Würtz M, Hvas A-M, Christensen KH, Rubak P, Kristensen SD, Grove EL. Rapid evaluation of platelet function using the Multiplate® Analyzer. *Platelets* 2014;25(08):628–633
- Beck F, Geiger J, Gambaryan S, et al. Temporal quantitative phosphoproteomics of ADP stimulation reveals novel central nodes in platelet activation and inhibition. *Blood* 2017;129(02):e1–e12
- Nisar S, Kelly E, Cullen PJ, Mundell SJ. Regulation of P2Y1 receptor traffic by sorting Nexin 1 is retromer independent. *Traffic* 2010;11(04):508–519
- Magalhaes AC, Dunn H, Ferguson SS. Regulation of GPCR activity, trafficking and localization by GPCR-interacting proteins. *Br J Pharmacol* 2012;165(06):1717–1736
- Jin J, Daniel JL, Kunapuli SP. Molecular basis for ADP-induced platelet activation. II. The P2Y1 receptor mediates ADP-induced intracellular calcium mobilization and shape change in platelets. *J Biol Chem* 1998;273(04):2030–2034
- Grocott M, Montgomery H, Vercueil A. High-altitude physiology and pathophysiology: implications and relevance for intensive care medicine. *Crit Care* 2007;11(01):203
- Feary JR, Rodrigues LC, Smith CJ, Hubbard RB, Gibson JE. Prevalence of major comorbidities in subjects with COPD and incidence of myocardial infarction and stroke: a comprehensive analysis using data from primary care. *Thorax* 2010;65(11):956–962
- Marin JM, Carrizo SJ, Vicente E, Agustí AG. Long-term cardiovascular outcomes in men with obstructive sleep apnoea-hypopnoea with or without treatment with continuous positive airway pressure: an observational study. *Lancet* 2005;365(9464):1046–1053

- 34 Sanner BM, Konermann M, Tepel M, Groetz J, Mummenhoff C, Zidek W. Platelet function in patients with obstructive sleep apnoea syndrome. *Eur Respir J* 2000;16(04):648–652
- 35 Maclay JD, McAllister DA, Johnston S, et al. Increased platelet activation in patients with stable and acute exacerbation of COPD. *Thorax* 2011;66(09):769–774
- 36 Bokinsky G, Miller M, Ault K, Husband P, Mitchell J. Spontaneous platelet activation and aggregation during obstructive sleep apnea and its response to therapy with nasal continuous positive airway pressure. A preliminary investigation. *Chest* 1995;108(03):625–630
- 37 Qureshi Z, Hobson AR. Clopidogrel “resistance”: where are we now? *Cardiovasc Ther* 2013;31(01):3–11
- 38 Topçuoğlu MA, Arsava EM, Ay H. Antiplatelet resistance in stroke. *Expert Rev Neurother* 2011;11(02):251–263
- 39 Jha PK, Sahu A, Prabhakar A, et al. Genome-wide expression analysis suggests hypoxia-triggered hyper-coagulation leading to venous thrombosis at high altitude. *Thromb Haemost* 2018; 118(07):1279–1295
- 40 Duplain H, Sartori C, Lepori M, et al. Exhaled nitric oxide in high-altitude pulmonary edema: role in the regulation of pulmonary vascular tone and evidence for a role against inflammation. *Am J Respir Crit Care Med* 2000;162(01):221–224
- 41 Levett DZ, Fernandez BO, Riley HL, et al; Caudwell Extreme Everest Research Group. The role of nitrogen oxides in human adaptation to hypoxia. *Sci Rep* 2011;1:109
- 42 Massberg S, Grüner S, Konrad I, et al. Enhanced in vivo platelet adhesion in vasodilator-stimulated phosphoprotein (VASP)-deficient mice. *Blood* 2004;103(01):136–142
- 43 Srihirun S, Piknova B, Sibmooh N, et al. Phosphorylated vasodilator-stimulated phosphoprotein (P-VASPSer239) in platelets is increased by nitrite and partially deoxygenated erythrocytes. *PLoS One* 2018;13:e0193747
- 44 Webb AJ, Milsom AB, Rathod KS, et al. Mechanisms underlying erythrocyte and endothelial nitrite reduction to nitric oxide in hypoxia: role for xanthine oxidoreductase and endothelial nitric oxide synthase. *Circ Res* 2008;103(09):957–964
- 45 Versteeg HH, Heemskerk JWM, Levi M, Reitsma PH. New fundamentals in hemostasis. *Physiol Rev* 2013;93(01):327–358
- 46 Kiouptsi K, Gambaryan S, Walter E, Walter U, Jurk K, Reinhardt C. Hypoxia impairs agonist-induced integrin $\alpha_{IIb}\beta_3$ activation and platelet aggregation. *Sci Rep* 2017;7(01):7621
- 47 Kauffenstein G, Bergmeier W, Eckly A, et al. The P2Y₁₂ receptor induces platelet aggregation through weak activation of the $\alpha_{IIb}\beta_3$ integrin—a phosphoinositide 3-kinase-dependent mechanism. *FEBS Lett* 2001;505(02):281–290
- 48 Kamae T, Shiraga M, Kashiwagi H, et al. Critical role of ADP interaction with P2Y₁₂ receptor in the maintenance of alpha (IIb)beta3 activation: association with Rap1B activation. *J Thromb Haemost* 2006;4(06):1379–1387
- 49 Horstrup K, Jablonka B, Hönig-Liedl P, Just M, Kochsiek K, Walter U. Phosphorylation of focal adhesion vasodilator-stimulated phosphoprotein at Ser157 in intact human platelets correlates with fibrinogen receptor inhibition. *Eur J Biochem* 1994;225(01): 21–27
- 50 Caradec J, Sirab N, Keumeugni C, et al. ‘Desperate house genes’: the dramatic example of hypoxia. *Br J Cancer* 2010;102(06): 1037–1043
- 51 Becker DM, Segal J, Vaidya D, et al. Sex differences in platelet reactivity and response to low-dose aspirin therapy. *JAMA* 2006; 295(12):1420–1427