

This is a repository copy of *Robertkochia solimangrovi* sp. nov., isolated from mangrove soil, and emended description of the genus *Robertkochia*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/155804/>

Version: Accepted Version

Article:

Quan Lam, Ming, Vodovnik, Masa, Zorec, Masa et al. (9 more authors) (2019)
Robertkochia solimangrovi sp. nov., isolated from mangrove soil, and emended description of the genus *Robertkochia*. International journal of systematic and evolutionary microbiology. ISSN 1466-5034

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

1 ***Robertkochia solimangrovi* sp. nov., isolated from**
2 **mangrove soil, and emended description of the genus**
3 ***Robertkochia***

4
5 Ming Quan Lam¹, Maša Vodovnik², Maša Zorec², Sye Jinn Chen¹, Kian Mau Goh¹, Adibah
6 Yahya¹, Madihah Md Salleh¹, Zaharah Ibrahim¹, Lili Tokiman³, Simon J. McQueen-Mason⁴,
7 Neil C. Bruce^{4*} and Chun Shiong Chong^{1*}

8
9 ¹ Department of Biosciences, Faculty of Science, Universiti Teknologi Malaysia, 81310
10 Skudai, Johor, Malaysia

11 ² Biotechnical Faculty, University of Ljubljana, Groblje 3, 1230 Domzale, Slovenija

12 ³ Johor National Parks Corporation, Kota Iskandar, 79575 Iskandar Puteri, Johor, Malaysia

13 ⁴ Centre for Novel Agricultural Products, Department of Biology, University of York,
14 Wentworth Way, York, YO10 5DD, United Kingdom

15
16 ***Correspondence:** Chun Shiong Chong, cschong@utm.my; Neil C. Bruce,
17 neil.bruce@york.ac.uk

18
19 **Keywords:** *Robertkochia solimangrovi*; polyphasic taxonomy; *Flavobacteriaceae*; mangrove

20
21 The full length 16S rRNA gene of strain CL23^T has been deposited at
22 EMBL/DDBJ/GenBank with accession number [MK258111](#).

23 The whole genome shotgun project of strain CL23^T and *R. marina* CC-AMO-30D^T are
24 available at EMBL/DDBJ/GenBank under accession [QKWN00000000](#) and [QXMP00000000](#)
25 respectively.

26
27
28
29

30 ABSTRACT

31 To date, there is sparse information for the genus *Robertkochia* with *Robertkochia marina*
32 CC-AMO-30D^T as the only described member. We report here a new species isolated from
33 mangrove soil of Malaysia Tanjung Piai National Park and perform polyphasic
34 characterization to determine its taxonomy position. Strain CL23^T is a Gram-negative,
35 yellow-pigmented, strictly aerobic, catalase-positive and oxidase-positive bacterium. The
36 optimal growth conditions were determined to be at pH 7.0, 30–37°C and 1–2% (w/v) NaCl.
37 The major respiratory quinone was menaquinone-6 (MK-6) and the highly abundant polar
38 lipids were four unidentified lipids, a phosphatidylethanolamine and two unidentified
39 aminolipids. The 16S rRNA similarity between CL23^T and *R. marina* CC-AMO-30D^T is
40 96.67%. Strain CL23^T and *R. marina* CC-AMO-30D^T are clustered together and were
41 distinguished from taxa of closely related genera in 16S rRNA phylogenetic analysis.
42 Genome sequencing revealed the strain CL23^T has a genome size of 4.4 Mbp and a G+C
43 content of 40.72 mol%. Overall genome indexes (OGRIs) including digital DNA-DNA
44 hybridization (dDDH) value and average nucleotide identity (ANI) are 17.70% and
45 approximately 70%, below the cut-off 70% and 95% respectively, indicated that strain CL23^T
46 is a distinct species to that of *R. marina* CC-AMO-30D^T. Collectively, based on phenotypic,
47 chemotaxonomic, phylogenetic and genomic evidence presented, strain CL23^T is proposed as
48 a new species with the name *Robertkochia solimangrovi* sp. nov. (=KCTC 72252^T =LMG
49 31418^T). An emended description of the genus *Robertkochia* is also proposed.

50

51

52

53

54

55

56

57

58

59

60

61

62

63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96

Flavobacteriaceae is one of the widely spread bacterial families composed of 158 genera at the time of writing [1]. The genus *Robertkochia* was introduced by Hameed et al. in 2014 [2] as one of the new genera in the family *Flavobacteriaceae*. Until now, the genus consisted of a single species *Robertkochia marina* CC-AMO-30D^T, which was isolated from surface seawater at Taichung harbour, Taiwan [2]. The species was described as Gram negative, strictly aerobic, orange-pigmented and with iso-C_{15:0}, iso-C_{15:1} G and iso-C_{17:0} 3-OH as predominant fatty acids. The report for *Robertkochia* is scarce as the previous study only focused on taxonomic assignment with one species reported so far [2]. Furthermore, the genome of this genus and prospective application have not been studied or reported.

Robertkochia and many other members of the *Flavobacteriaceae* are halophilic or halotolerant bacteria that reside in diverse saline environments such as seawater, mangrove forest and marine sediment [3-5]. Mangroves are inter-tidal wetlands that connect terrestrial and marine ecosystems [6]. Due to periodic tidal flats, drastic changes in salinity and nutrient availability of the mangrove environment make it a unique ecosystem [7]. Free living and symbiotic bacteria in such environment were found to play essential roles in maintaining mangrove ecosystem such as recycling of organic matter and biotransformation of minerals [8-10]. It was estimated that less than 5% of species in mangrove environment have been described so far [11]. Therefore, it could be considered as one of the interesting areas to be explored. In the present study, strain CL23^T was isolated from soil obtained from mangrove forest located at Tanjung Piai National Park, Johor, Malaysia. This strain was characterized using polyphasic approach (phenotypic, chemotaxonomic and genomic aspects) following the recommended guidelines [12, 13] and new criteria for classification [14] to elucidate its taxonomy position. The results indicated that strain CL23^T represents a new species within *Robertkochia* genus, with the name *Robertkochia solimangrovi* sp. nov. is proposed.

97
98

99 ISOLATION AND HOME HABITAT

100 Soil from the mangrove forest was sampled at Tanjung Piai National Park (GPS location:
101 1°16'06.0" N, 103°30'31.2" E) in September 2017 with permit (CJB F No. 734342) granted
102 by Johor National Parks Corporation. The soil samples were serially diluted with sterile
103 distilled water (10^{-1} to 10^{-8}). A 0.1 ml of diluted sample was spread onto marine agar 2216
104 (MA; BD Difco) and incubated at 30–35°C for 1 to 14 days. A yellow-pigmented strain
105 designated as CL23^T was isolated from MA and re-streaked twice to obtain a pure culture.
106 The strain was maintained in marine broth 2216 (MB; BD Difco) with 20 % (v/v) glycerol at
107 –80°C. Strain CL23^T was deposited at Korean Collection for Type Cultures (KCTC) and
108 Belgian Co-ordinated Collections of Micro-organisms (BCCM) under accession of KCTC
109 72252^T and LMG 31418^T, respectively. For comparative polyphasic taxonomy
110 characterization, *R. marina* CC-AMO-30D^T (=JCM 18552^T) was obtained from Japan
111 Collection of Microorganisms (JCM). Both strains were routinely cultured on MA and in MB
112 at 30°C for 48 h, unless specified otherwise.

113
114
115

116 16S rRNA PHYLOGENY

117 Genomic DNA was extracted using DNeasy Blood and Tissue kit (Qiagen) and was purified
118 by DNA Clean and ConcentratorTM-25 (Zymo Research) following manual instructions. The
119 16S rRNA gene of strain CL23^T was amplified by PCR using universal primers: 27F (5'-
120 AGAGTTTGATCMTGGCTCAG-3') and 1525R (5'-AAGGAGGTGWTCARCC-3') [15].
121 The 16S rRNA gene was sequenced at Apical Scientific Pte. Ltd., Seri Kembangan, Malaysia.
122 After the sequencing, the raw sequences were trimmed, and the sequences were aligned using
123 ClustalW. The nearly full-length 16S rRNA gene was searched against EzBioCloud database
124 for identification. The amplified 16S rRNA gene of strain CL23^T was also cross-checked
125 with the genome data to ensure the acquisition of full-length gene (1522 bp). The 16S rRNA
126 gene of strain CL23^T (MK258111) shared highest similarity (96.67%) with *R. marina* CC-
127 AMO-30D^T (JX235674), which is below the accepted threshold of 98.7% for species
128 delineation [14]. The 16S rRNA gene similarity was less than 94% between strain CL23^T and

129 other members of closely related genera: *Joostella marina* En5^T (93.82%), *Joostella*
130 *atrarenae* M1-2^T (93.82%), *Zhouia spongiae* HN-Y44^T (93.75%) and *Pustulibacterium*
131 *marinum* E403^T (93.35%).

132

133 Phylogenetic trees of 16S rRNA were built following the Neighbor-joining (NJ) [16] and
134 Maximum Likelihood (ML) [17] algorithms using MEGA 7.0 software [18] based on 1000
135 bootstrap replications [19] and Kimura-2 parameter. Following the 16S rRNA phylogenetic
136 analysis (Fig. 1), strain CL23^T and *R. marina* CC-AMO-30D^T formed a clade in NJ and ML
137 trees, confirming the placement of strain CL23^T within *Robertkochia* genus. The high
138 bootstrap value at the node separating the branch of strain CL23^T and *R. marina* CC-AMO-
139 30D^T in 16S rRNA phylogenetic tree supported that these two strains are distinct between
140 each other.

141

142

143

144 PHENOTYPIC AND CHEMOTAXONOMIC CHARACTERIZATION

145 Colony morphology was observed on MA at 30°C after 48 h of incubation. The Gram
146 staining was performed according to the protocol as described previously [20]. The malachite
147 green staining was used to assess the presence of endospore in 7-day old cultures [21]. The
148 Gram stain reaction and endospore formation were examined under light microscope (Nikon
149 ECLIPSE E200). Cell morphology was examined under scanning electron microscope (SEM;
150 JEOL JSM-IT300LV). The bacterial motility was investigated by hanging-drop approach
151 [22]. The presence of flexirubin-type pigment was determined by flooding the cells with 20 %
152 (w/v) KOH [12].

153

154 Catalase activity was detected by effervescence using 3 % (v/v) H₂O₂ while oxidase activity
155 was determined by oxidation of tetramethyl-*p*-phenylenediamine. Hydrolysis of starch, casein,
156 L-tyrosine, hypoxanthine, xanthine, Tween 20, Tween 40, Tween 60, Tween 80,
157 carboxymethyl-cellulose (CMC) and xylan were tested according to Smibert and Krieg [21].
158 Bile esculin hydrolysis was investigated using the method of Facklam and Moody [23]. Other
159 biochemical characteristics were revealed by API 20 E and API 20 NE kits (BioMérieux,
160 France). Carbohydrate utilization and enzyme activity profile of both strains were

161 investigated by API 50 CHB and API ZYM kits (BioMérieux, France), respectively. All API
162 kits were carried out by following the manufacturer's instructions with slight modification in
163 which inoculation was supplemented up to 2 % (w/v) NaCl.

164

165 Growth under anaerobic condition was tested by incubating the bacteria on MA for 14 days at
166 30°C using AnaeroGen (Oxoid) in an anaerobic jar (Mitsubishi Gas Chemical). Growth was
167 tested on the following media: Reasoner's 2A agar (R2A; HiMedia), Nutrient agar (NA;
168 Merck), Tryptic soy agar (TSA; Merck), Luria-Bertani agar (LBA; Conda) and Muller Hinton
169 agar (MHA; Sigma) supplemented with 2 % (w/v) NaCl at 30°C for 7 days. The temperature
170 range (4, 9, 15, 20, 25, 30, 37, 40, 42, 45 and 50°C) and the optimum temperature for growth
171 were determined using MB at pH 7. The pH range (in intervals of 1.0 pH unit) and optimum
172 pH for growth were investigated using MB at 30°C. The pH was adjusted with the following
173 buffer systems: 50 mM citrate phosphate (pH 4–5), 50 mM sodium phosphate (pH 6–8) and
174 50 mM glycine–NaOH (pH 9–10) [24]. The pH was verified after autoclaving. To test NaCl
175 tolerance and optimal concentration, the bacteria were grown in a medium containing yeast
176 extract (1.0 g l⁻¹), peptone (5.0 g l⁻¹), MgCl₂ (5.0 g l⁻¹), MgSO₄·7H₂O (2.0 g l⁻¹), CaCl₂ (0.5 g
177 l⁻¹), KCl (1.0 g l⁻¹) and NaCl (0, 0.5, 1–11 %, w/v) [10].

178

179 Antibiotic susceptibility of bacteria against 21 antibiotics was tested using the disk diffusion
180 method on MA at 30°C for 48 h [25]. The antibiotics disks (Oxoid) used were: ampicillin (10
181 µg), bacitracin (10 IU), carbenicillin (100 µg), chloramphenicol (100 µg), clindamycin (2 µg),
182 doxycycline (30 µg), erythromycin (60 µg), gentamicin (10 µg), kanamycin (50 µg),
183 lincomycin (2 µg), minocycline (30 µg), neomycin (30 µg), novobiocin (5 µg), oleandomycin
184 (15 µg), oxacillin (1 µg), penicillin G (10 IU), piperacillin (100 µg), polymyxin B (300 IU),
185 rifampicin (5 µg), streptomycin (10 µg) and tetracycline (30 µg).

186

187 Strain CL23^T was determined as a Gram negative, rod-shaped, non-spore forming, oxidase
188 positive and catalase positive bacterium with motile ability by gliding. The colony was in a
189 circular form with 0.5–1.0 mm diameter, smooth surface, convex elevation, entire margin and
190 has translucent property on MA after 48 h incubation. Under SEM, cells of strain CL23^T were
191 0.2–0.4 µm in width and 2.3–3.2 µm in length. The notable distinctive features to
192 differentiate strain CL23^T and *R. marina* CC-AMO-30D^T are shown in Table 1. In terms of
193 morphology, strain CL23^T is yellow pigmented while *R. marina* CC-AMO-30D^T was found
194 to be orange pigmented. Strain CL23^T grew well in 15–42°C, pH 5–9 and 0–9 % (w/v) NaCl,

195 and in general strain CL23^T demonstrated a broader growth range compared to *R. marina*
196 CC-AMO-30D^T (Table 1). The optimal growth conditions of strain CL23^T were observed at
197 30–37°C, pH 7 and 1–2 % (w/v) NaCl. Strain CL23^T was also able to produce acetoin, β-
198 galactosidase and weakly positive toward amygdaline according to API 20 E but not for *R.*
199 *marina* CC-AMO-30D^T. Based on API ZYM, strain CL23^T was able to produce α-
200 galactosidase, β-galactosidase and α-mannosidase, which were absent in *R. marina* CC-
201 AMO-30D^T. Both strains were further distinguished by the hydrolysis capability of gelatin,
202 Tween 20, Tween 40, Tween 60, and exhibiting resistance towards ampicillin, penicillin G,
203 piperacillin and bacitracin (Table 1).

204

205 For the chemotaxonomic analysis, cellular fatty acids were extracted following the protocol
206 of Microbial Identification System (MIDI, version 6.1) [26]. Biomass of strain CL23^T and its
207 reference strain *R. marina* CC-AMO-30D^T were harvested from MA after 48 h of incubation
208 at 30°C. The cells were saponified with methanolic base, then the resulting sodium salts of
209 fatty acids were methylated. In the final step, methyl esters were transferred to the organic
210 phase and washed. Fatty acid methyl esters were analyzed on an Agilent 6890 equipped with
211 Ultra-2 capillary column and subsequently identified in the RTSBA6 library. As exhibited in
212 Table 2, the predominant cellular fatty acid of strain CL23^T and *R. marina* CC-AMO-30D^T
213 were found to be iso-C_{15:0}, iso-C_{15:1} G and iso-C_{17:0} 3-OH (> 10%). Nonetheless, some fatty
214 acid patterns and abundance of strain CL23^T varied when compared to *R. marina* CC-AMO-
215 30D^T, such as summed features 3 (3.64%) and 9 (5.24%) were constituted in strain CL23^T but
216 none for *R. marina* CC-AMO-30D^T. On top of that, the amount of iso-C_{16:0}, anteiso-C_{15:0} and
217 iso-C_{16:0} 3-OH of strain CL23^T are remarkably lower than *R. marina* CC-AMO-30D^T (Table
218 2).

219

220 The polar lipids and respiratory quinone analyses of strain CL23^T were performed by Dr.
221 Brian Tindall at the Identification Service, DSMZ, Braunschweig, Germany. In brief, the
222 respiratory quinones were extracted by solvent methanol: hexane (2:1 v/v), separated by TLC
223 and High Performance Liquid Chromatography (HPLC) following the standard method by
224 Tindall [27]. The polar lipids were extracted using chloroform: methanol solvent and
225 separated by two-dimensional silica gel thin layer chromatography (TLC) [28]. Total lipid
226 material was identified using molybdato-phosphoric acid and specific functional groups were
227 determined using spray reagents specific for defined functional groups.

228

229 The major respiratory quinone of strain CL23^T was identified to be menaquinone-6 (MK-6),
230 which matched to *R. marina* [2] and other members in *Flavobacteriaceae* family [12]. In
231 terms of polar lipids, strain CL23^T has four unidentified lipids (L1, L2, L3 and L4), a
232 phosphatidylethanolamine (PE) and two unidentified aminolipids (AL1 and AL2) as major
233 polar lipids (Fig. S1). Additionally, three unidentified glycolipids (GL1, GL2 and GL3) and
234 an unknown lipid (L5) were observed in minor amounts. The unidentified lipids (L1–L3) and
235 glycolipids (GL1–GL3) were not detected in *R. marina* CC-AMO-30D^T [2]. Moreover, an
236 unidentified phospholipid (PL) was contained in *R. marina* CC-AMO-30D^T in which this
237 lipid was not found in strain CL23^T [2].

238

239

240

241 **GENOMIC CHARACTERIZATION**

242 The genome of reference strain *R. marina* CC-AMO-30D^T was not available at the time of
243 study, therefore, both the genomes of strain CL23^T (NCBI accession: QKWN000000000) and
244 *R. marina* CC-AMO-30D^T (NCBI accession: QXMP000000000) were sequenced in this study.
245 Whole genome sequencing of strain CL23^T was accomplished on an Illumina HiSeq 2500
246 platform (2 × 150 bp). The raw reads were filtered, and the quality data was *de novo*
247 assembled using SOAPdenovo 2.04 [29]. The resulting genome was annotated using NCBI
248 Prokaryotic Genome Annotation Pipeline (PGAP) [30].

249

250 The assembled genome of strain CL23^T consists of 23 contigs with 322× depth of sequencing
251 coverage (average), made up the size of genome with 4,407,290 bp in length and a GC
252 content of 40.72 mol%. The genome size of strain CL23^T is significantly larger than *R.*
253 *marina* CC-AMO-30D^T (3,571,649 bp). The GC content of strain CL23^T is slightly lower
254 than *R. marina* CC-AMO-30D^T (43.67 mol%). Based on PGAP annotation, a total of 3669
255 protein coding genes was found in genome of strain CL23^T. The genes responsible for
256 phosphatase activity were found in the genome of strain CL23^T and *R. marina* CC-AMO-
257 30D^T with a total of 12 and 7 phosphatases were encoded respectively (Table S1). This
258 correlated to API ZYM results in which both strains were positive to acidic and alkali
259 phosphatases. Notably, the number of phosphatases annotated is higher in strain CL23^T as
260 compared to *R. marina* CC-AMO-30D^T. On the other hand, strain CL23^T consists of a series

261 of genes for assimilatory sulfate reduction into sulfite (sulfate adenylyltransferase subunit
262 CysN and CysD, adenylylsulfate kinase and phosphoadenylylsulfate reductase) and then
263 sulfite reduction into sulfide (FAD-binding oxidoreductase and LLM class flavin-dependent
264 oxidoreductase) (Table S1). Nevertheless, the genes responsible for reduction of sulfite to
265 sulfide are absent in *R. marina* CC-AMO-30D^T (Table S1). Furthermore, strain CL23^T also
266 encodes a set of genes for reduction of nitrate to ammonia (*NirBD* and *NrfAH*) in which
267 *NirBD* genes were not found in genome of *R. marina* CC-AMO-30D^T (Table S1). These
268 genes suggest that strain CL23^T participates in nutrient recycling in mangrove environments.

269

270 Multilocus sequence analysis (MLSA) was conducted on five housekeeping genes of strain
271 CL23^T, *R. marina* CC-AMO-30D^T and related genera, which the sequences were retrieved
272 from genome data. The sequences of housekeeping genes were aligned individually and then
273 concatenated in the following order: *rpoB*–*gyrB*–*recA*–*mutL*–*atpD*. The phylogenetic tree of
274 concatenated housekeeping genes was constructed using MEGA 7.0 similarly as described in
275 section “16S rRNA phylogeny”. In this tree (Fig. 2), strain CL23^T and *R. marina* CC-AMO-
276 30D^T are clustered together but well distinguished from each other with high level of support
277 (>90% bootstrap value). Likewise, the phylogenetic tree based on whole genome sequences
278 that built using REALPHY 1.12 [31] also supported that both strain CL23^T and *R. marina*
279 CC-AMO-30D^T are grouped in the same clade (Fig. S2).

280

281 To further underpin the classification of strain CL23^T as a new species, the overall genome
282 related indexes (OGRIs) were determined. Average nucleotide identity based on BLAST
283 (ANIb) was calculated using JSpeciesWS [32]. ANI based on USEARCH (OrthoANIu) was
284 determined by ChunLab's online ANI calculator [33]. The digital DNA-DNA hybridization
285 (dDDH) value was calculated by Genome-to-Genome Distance Calculator [34].

286

287 The ANIb and OrthoANIu values between strain CL23^T and *R. marina* CC-AMO-30D^T are
288 69.35% and 70.47% respectively. These ANI values are below the recommended 95–96% for
289 species delineation [35]. Similarly, the dDDH value between two strains was found to be
290 17.70%, lower than 70%, the cut off for species boundary [34]. Combining the interpretation
291 of ANI and dDDH values, the result revealed the identity of strain CL23^T as a distinct species
292 within the same genus as *R. marina* CC-AMO-30D^T.

293

294 Based on polyphasic taxonomy characterization including phenotypic, chemotaxonomic,
295 phylogenetic and genomic aspects, the results clearly indicated that strain CL23^T (=KCTC
296 72252^T =LMG 31418^T) represents a new species within the genus *Robertkochia*, for which
297 the name *Robertkochia solimangrovi* sp. nov. is proposed.

298

299

300

301 **DESCRIPTION OF *ROBERTKOCHIA SOLIMANGROVI* SP. NOV.**

302

303 *Robertkochia solimangrovi* sp. nov. (so.li.man.gro'vi. L. neut. n. *solum* soil; N.L. neut. n.
304 *mangrovum* a mangrove; N.L. gen. n. *solimangrovi* of soil of a mangrove, pertaining to
305 where the type strain was isolated.)

306 The cells are Gram-negative, rod shape, approximate 0.2–0.4 µm in width and 2.3–3.2 µm in
307 length with motile ability by gliding. Colony is yellow-pigmented, in circular form with 0.5–
308 1.0 mm diameter, smooth surface, convex elevation, entire margin and has translucent
309 property after 48 hours incubation at 30°C on MA. Flexirubin-type pigment is absent. Cells
310 are positive for oxidase and catalase. Growth occurs at 15–42 °C (optimum, 30–37°C), pH
311 5–9 (optimum, pH 7) and in the presence of 0–9 % (w/v) NaCl (optimum, 1–2 % (w/v) NaCl).
312 Grows well on MA, however, no growth is observed on R2A, NA, LBA, TSA and MHA
313 media. No growth is observed on MA in anaerobic condition. The predominant fatty acids
314 are iso-C_{15:0}, iso-C_{15:1} G and iso-C_{17:0} 3-OH. The major isoprenoid quinone is menaquinone-6
315 (MK-6). The major polar lipids are four unidentified lipids, a phosphatidylethanolamine and
316 two unidentified aminolipids. Xylan, esculin, Tween 20, 40 and 60 are hydrolyzed. L-
317 Tyrosine is weakly hydrolyzed. Casein, starch, CMC, Tween 80, xanthine and hypoxanthine
318 are not hydrolyzed. In the API 20 E strip, positive for ONP-β-D-galactopyranoside and
319 acetoin production; weakly positive for fermentation/oxidation of amygdaline; negative for
320 arginine dihydrolase, lysine decarboxylase, ornithine decarboxylase, tryptophane deaminase,
321 urease and gelatinase, production of H₂S and indole, utilization of citrate,
322 fermentation/oxidation of D-glucose, D-mannitol, inositol, D-sorbitol, L-rhamnose, D-
323 saccharose, D-melibiose and L-arabinose. In the API 20 NE strip, positive for hydrolysis of
324 pNP-β-D-galactopyranoside and esculin ferric citrate; negative for nitrate reduction, indole

325 production, arginine dihydrolase, gelatinase and urease, fermentation of D-glucose and
326 assimilation of D-glucose, D-arabinose, D-mannose, D-mannitol, N-acetyl-glucosamine, D-
327 maltose, potassium gluconate, capric acid, adipic acid, malic acid and phenylacetic acid. In
328 the API 50 CHB strip, acid is produced from D-galactose, D-glucose, D-mannose, esculin
329 ferric citrate, D-cellobiose, D-maltose, D-lactose, D-melibiose, D-saccharose, D-trehalose, D-
330 melezitose, D-raffinose, amidon, glycogen and gentibiose; acid is weakly produced from
331 methyl- α -D-glucopyranoside, arbutin, salicin and D-turanose; acid is not produced from
332 glycerol, erythritol, D-arabinose, L-arabinose, D-ribose, D-xylose, L-xylose, D-adonitol,
333 methyl- β -D-xylopyranoside, D-fructose, L-sorbose, L-rhamnose, dulcitol, inositol, D-mannitol,
334 D-sorbitol, methyl- α -D-mannopyranoside, N-acetyl-glucosamine, amygdalin, inulin, xylitol,
335 D-lyxose, D-tagatose, D-fucose, L-fucose, D-arabitol, L-arabitol, potassium gluconate,
336 potassium 2-ketogluconate and potassium 5-ketogluconate. In the API ZYM strip, alkali
337 phosphatase, esterase (C4), esterase lipase (C8), leucine arylamidase, valine arylamidase,
338 cystine arylamidase, trypsin, chymotrypsin, acid phosphatase, naphthol-AS-BI-
339 phosphohydrolase, α -galactosidase, β -galactosidase, α -glucosidase, β -glucosidase, N-acetyl-
340 β -glucosaminidase and α -mannosidase are present; weak positive reaction for α -fucosidase
341 and negative results for lipase (C14) and α -fucosidase. Cells are susceptible to carbenicillin,
342 clindamycin, doxycycline, lincomycin, minocycline, novobiocin, oleandomycin, rifampicin
343 and tetracycline, but not to ampicillin, bacitracin, chloramphenicol, erythromycin, gentamicin,
344 kanamycin, neomycin, oxacillin, penicillin G, piperacillin, polymyxin B and streptomycin.

345

346 The type strain is CL23^T (=KCTC 72252^T =LMG 31418^T), isolated from soil of mangrove
347 collected from Tanjung Piai National Park, Johor, Malaysia.

348

349 The EMBL/DDBJ/GenBank accession for 16S rRNA gene of strain CL23^T is [MK258111](#).
350 Genome metrics are as follows: genome size, 4,407,290 bp; number of contigs, 23; G+C
351 content, 40.72 mol%. The Whole Genome Shotgun project of strain CL23^T is available at
352 EMBL/DDBJ/GenBank under accession [QKWN00000000](#). The version described in this
353 paper is QKWN01000000.

354

355 **EMENDED DESCRIPTION OF GENUS *ROBERTKOCHIA***

356

357 The characteristics of the genus *Robertkochia* are described according to Hameed et al. 2014
358 [2] with following amendments and additional information. Oxidase is either positive or
359 negative and catalase is positive. The DNA G+C content of the type strain of type species is
360 43.67 mol% based on genome data. The Whole Genome Shotgun project of type strain of
361 type species is available at EMBL/DDBJ/GenBank under accession [QXMP00000000](#). The
362 version described in this paper is QXMP01000000.

363

364

365 **AUTHOR STATEMENTS**

366

367 *Funding information*

368 This work was financially co-sponsored by the Ministry of Education Malaysia and
369 Biotechnology and Biological Sciences Research Council (BBSRC) United Kingdom under
370 program of United Kingdom-Southeast Asia Newton Ungku Omar Fund (UK-SEA-NUOF)
371 with project number 4B297 and BB/P027717/1, respectively. The work was also supported
372 by GUP (Tier 1) grant provided by Universiti Teknologi Malaysia under project number
373 20H43 which granted to Chun Shiong Chong. Slovenian Research Agency funded this work
374 (cellular fatty acids analysis) via P4-0097 (C), which granted to Maša Vodovnik and Maša
375 Zorec. Neil C. Bruce, Simon J. McQueen-Mason and Chun Shiong Chong are grateful for
376 BBSRC International Partnering Award (BB/P025501/1).

377

378 *Acknowledgements*

379 The authors would like to acknowledge Johor National Parks Corporation for sampling
380 permit (CJB F No. 734342) at Tanjung Piai National Park, Johor, Malaysia. Ming Quan Lam
381 is grateful to Khazanah Watan Postgraduate (PhD) scholarship (scholar ID: 40852) from
382 Yayasan Khazanah. Ming Quan Lam appreciates Dr. Suganthi Thevarajoo, Dr. Chitra
383 Selvaratnam and Mr. Jia Chun Lim for technical guidance and fruitful discussion. Sye Jinn
384 Chen acknowledges Zamalah scholarship (PhD) from Universiti Teknologi Malaysia.

385

386

387 *Ethical statement*

388 No human and animal experiments are involved.

389

390 *Conflicts of interest*

391 All authors declared that there are no conflicts of interest.

392

393

394 **ABBREVIATIONS**

395 ANI, average nucleotide identity, dDDH, digital DNA-DNA hybridization; MA, marine agar;
396 MB, marine broth; MK, menaquinone; ML, maximum likelihood; NJ, neighbor-joining;
397 OGRI, overall genome related index; PE, phosphatidylethanolamine.

398

399

400 **REFERENCES**

401

- 402 1. **Parte AC.** LPSN – List of Prokaryotic names with Standing in Nomenclature
403 (bacterio.net), 20 years on. *Int J Syst Evol Microbiol* 2018;68(6):1825-9.
- 404 2. **Hameed A, Shahina M, Lin S-Y, Lai W-A, Liu Y-C, et al.** *Robertkochia marina*
405 gen. nov., sp. nov., of the family *Flavobacteriaceae*, isolated from surface seawater, and
406 emended descriptions of the genera *Joostella* and *Galbibacter*. *Int J Syst Evol Microbiol*
407 2014;64(2):533-9.
- 408 3. **Thevarajoo S, Selvaratnam C, Goh KM, Hong KW, Chan XY, et al.** *Vitellibacter*
409 *aquimaris* sp. nov., a marine bacterium isolated from seawater. *Int J Syst Evol Microbiol*
410 2016;66(9):3662-8.
- 411 4. **Li Y, Bai S, Yang C, Lai Q, Zhang H, et al.** *Mangrovimonas yunxiaonensis* gen.
412 nov., sp. nov., isolated from mangrove sediment. *Int J Syst Evol Microbiol* 2013;63(6):2043-8.
- 413 5. **Wang B, Sun F, Du Y, Liu X, Li G, et al.** *Meridianimaribacter flavus* gen. nov., sp.
414 nov., a member of the family *Flavobacteriaceae* isolated from marine sediment of the South
415 China Sea. *Int J Syst Evol Microbiol* 2010;60(1):121-7.
- 416 6. **Alongi DM.** Mangrove forests. In: *Blue Carbon: Coastal Sequestration for Climate*
417 *Change Mitigation*. Cham: Springer; 2018. p. 23-36.
- 418 7. **Lin X, Hetharua B, Lin L, Xu H, Zheng T, et al.** Mangrove sediment microbiome:
419 adaptive microbial assemblages and their routed biogeochemical processes in Yunxiao
420 Mangrove National Nature Reserve, China. *Microb Ecol* 2019;78(1):57-69.

- 421 8. **Castro RA, Dourado MN, Almeida JRd, Lacava PT, Nave A, et al.** Mangrove
422 endophyte promotes reforestation tree (*Acacia polyphylla*) growth. *Braz J Microbiol*
423 2018;49(1):59-66.
- 424 9. **Kathiresan K.** Salt-tolerant microbes in mangroves: ecological role and
425 bioprospecting potential. In: Dagar JC, Yadav RK, Sharma PC, (editors). *Research*
426 *Developments in Saline Agriculture*. Singapore: Springer Singapore; 2019. p. 237-55.
- 427 10. **Lam MQ, Oates NC, Thevarajoo S, Tokiman L, Goh KM, et al.** Genomic analysis
428 of a lignocellulose degrading strain from the underexplored genus *Meridianimaribacter*.
429 *Genomics* 2019. doi: [10.1016/j.ygeno.2019.06.011](https://doi.org/10.1016/j.ygeno.2019.06.011) (in press)
- 430 11. **Thatoi H, Behera BC, Mishra RR, Dutta SK.** Biodiversity and biotechnological
431 potential of microorganisms from mangrove ecosystems: a review. *Ann Microbiol*
432 2013;63(1):1-19.
- 433 12. **Bernardet J-F, Nakagawa Y, Holmes B.** Proposed minimal standards for describing
434 new taxa of the family *Flavobacteriaceae* and emended description of the family. *Int J Syst*
435 *Evol Microbiol* 2002;52(3):1049-70.
- 436 13. **Tindall BJ, Rosselló-Móra R, Busse H-J, Ludwig W, Kämpfer P.** Notes on the
437 characterization of prokaryote strains for taxonomic purposes. *Int J Syst Evol Microbiol*
438 2010;60(1):249-66.
- 439 14. **Chun J, Oren A, Ventosa A, Christensen H, Arahal DR, et al.** Proposed minimal
440 standards for the use of genome data for the taxonomy of prokaryotes. *Int J Syst Evol*
441 *Microbiol* 2018;68(1):461-6.
- 442 15. **Claus D.** A standardized Gram staining procedure. *World J Microbiol Biotechnol*
443 1992;8(4):451-2.
- 444 16. **Smibert RM, Krieg NR.** Phenotypic characterization. In: Gerhardt P, Murray RGE,
445 Wood WA, Krieg NR. *Methods for General and Molecular Bacteriology*. Washington, DC:
446 American Society for Microbiology; 1994. p. 607-54.
- 447 17. **Tittsler RP, Sandholzer LA.** The use of semi-solid agar for the detection of bacterial
448 motility. *J Bacteriol* 1936;31(6):575.
- 449 18. **Facklam RR, Moody MD.** Presumptive identification of group D *Streptococci*: the
450 bile-esculin test. *Appl Microbiol* 1970;20(2):245-50.
- 451 19. **Lam MQ, Nik Mut NN, Thevarajoo S, Chen SJ, Selvaratnam C, et al.**
452 Characterization of detergent compatible protease from halophilic *Virgibacillus* sp. CD6. 3
453 *Biotech* 2018;8(2):104.

- 454 20. **Jorgensen JH, Turnidge JD.** Susceptibility test methods: dilution and disk diffusion
455 methods. In: James HJ, Michael AP, Karen CC, Guido F, Marie LL Sandra SR, David WW
456 (editors). *Manual of Clinical Microbiology*, 11th ed. Washington, DC: American Society of
457 Microbiology; 2015. p. 1253-73.
- 458 21. **Sasser M.** *Identification of bacteria by gas chromatography of cellular fatty acids.*
459 *USFCC Newsl* 1990;20:16.
- 460 22. **Tindall BJ.** Lipid composition of *Halobacterium lacusprofundi*. *FEMS Microbiol*
461 *Lett* 1990;66(1-3):199-202.
- 462 23. **Tindall BJ, Sikorski J, Smibert RA, Krieg NR.** Phenotypic characterization and the
463 principles of comparative systematics. In: Reddy CA, Beveridge TJ, Breznak JA, Maxluf G,
464 Schmidt TM, Snyder LR (editors). *Methods for General and Molecular Microbiology*, 3rd ed.
465 Washington, DC: American Society of Microbiology; 2007. p. 330-93.
- 466 24. **Lane DJ.** 16S/23S rRNA sequencing. In: Stackebrandt E and Goodfellow M (editors).
467 *Nucleic Acid Techniques in Bacterial Systematics*. Chichester, United Kingdom: Wiley; 1991.
468 p. 125-75.
- 469 25. **Saitou N, Nei M.** The neighbor-joining method: a new method for reconstructing
470 phylogenetic trees. *Mol Biol Evol* 1987;4(4):406-25.
- 471 26. **Felsenstein J.** Evolutionary trees from DNA sequences: a maximum likelihood
472 approach. *J Mol Evol* 1981;17(6):368-76.
- 473 27. **Kumar S, Stecher G, Tamura K.** MEGA7: Molecular evolutionary genetics analysis
474 version 7.0 for bigger datasets. *Mol Biol Evol* 2016;33(7):1870-4.
- 475 28. **Felsenstein J.** Confidence limits on phylogenies: an approach using the bootstrap.
476 *Evolution* 1985:783-91.
- 477 29. **Li R, Li Y, Kristiansen K, Wang J.** SOAP: short oligonucleotide alignment program.
478 *Bioinformatics* 2008;24(5):713-4.
- 479 30. **Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, et al.** NCBI
480 prokaryotic genome annotation pipeline. *Nucleic Acids Res* 2016;44(14):6614-24.
- 481 31. **Bertels F, Silander OK, Pachkov M, Rainey PB, van Nimwegen E.** Automated
482 Reconstruction of Whole-Genome Phylogenies from Short-Sequence Reads. *Mol Biol Evol*
483 2014;31(5):1077-88.
- 484 32. **Richter M, Rosselló-Móra R, Oliver Glöckner F, Peplies J.** JSpeciesWS: a web
485 server for prokaryotic species circumscription based on pairwise genome comparison.
486 *Bioinformatics* 2016;32(6):929-31.

- 487 33. **Yoon S-H, Ha S-m, Lim J, Kwon S, Chun J.** A large-scale evaluation of algorithms
488 to calculate average nucleotide identity. *Antonie van Leeuwenhoek* 2017;110(10):1281-6.
- 489 34. **Meier-Kolthoff JP, Auch AF, Klenk H-P, Göker M.** Genome sequence-based
490 species delimitation with confidence intervals and improved distance functions. *BMC*
491 *Bioinformatics* 2013;14(1):60.
- 492 35. **Richter M, Rosselló-Móra R.** Shifting the genomic gold standard for the prokaryotic
493 species definition. *Proc Natl Acad Sci* 2009;106(45):19126-31.

494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520

521

522 **FIGURES AND TABLES**

523

524 **Table 1.** Differential phenotypic characteristics of strain CL23^T and *Robertkochia marina*
 525 CC-AMO-30D^T.

526 Strains: 1, strain CL23^T; 2, *R. marina* CC-AMO-30D^T. All data were obtained from this study.
 527 +, Positive reaction; –, negative reaction; w, weakly positive reaction. All strains were
 528 positive for catalase; hydrolysis of xylan and aesculin; production of acid from D-glucose,
 529 esculin ferric citrate, D-cellobiose, D-maltose, D-saccharose, D-trehalose, D-melezitose,
 530 amidon and glycogen in API 50 CHB strips; and activity of alkali phosphatase, esterase (C4),
 531 esterase lipase (C8), leucine arylamidase, valine arylamidase, cystine arylamidase, trypsin,
 532 chymotrypsin, acid phosphatase, naphthol-AS-BI-phosphohydrolase, α-glucosidase, β-
 533 glucosidase and N-acetyl-β-glucosaminidase. Both strains were negative for flexirubin-type
 534 pigment; growth under anaerobic condition; growth on R2A, NA, LBA, TSA and MHA
 535 media; hydrolysis of casein, starch, CMC, Tween 80, xanthine and hypoxanthine; nitrate
 536 reduction; indole and H₂S production; urease; acid production from glycerol, erythritol, D-
 537 arabinose, L-arabinose, D-ribose, D-xylose, L-xylose, D-adonitol, methyl-β-D-xylopyranoside,
 538 D-fructose, L-sorbose, L-rhamnose, dulcitol, inositol, D-mannitol, D-sorbitol, methyl-α-D-
 539 mannopyranoside, N-acetyl-glucosamine, amygdalin, inulin, xylitol, D-lyxose, D-tagatose, D-
 540 fucose, L-fucose, D-arabitol, L-arabitol, potassium gluconate and potassium 2-ketogluconate
 541 and potassium 5-ketogluconate in API 50 CHB strips; and activity of lipase (C14) and β-
 542 glucuronidase (API ZYM).

Characteristics	1	2
Colony pigmentation	Yellow	Orange
Oxidase	+	–
Growth parameters		
pH range	5–9	6–7
Temperature range (°C)	15–42	20–40
Temperature optimum (°C)	30–37	30
NaCl range (% w/v)	0–9	0.5–4
NaCl optimum (% w/v)	1–2	2
Hydrolysis of		
Tween 20	+	w
Tween 40	+	–
Tween 60	+	w
Tyrosine	w	–

Gelatin	–	+
Production of Acetoin	+	–
Oxidation of Amygdaline	w	–
Utilization of D-Galactose	+	–
D-Mannose	+	–
Arbutin	w	–
Salicin	w	–
D-Lactose	+	–
D-Melibiose	+	–
D-Raffinose	+	–
Gentiobiose	+	–
D-Turanose	w	–
Enzyme activity (API ZYM)		
α-Galactosidase	+	–
β-Galactosidase	+	–
α-Mannosidase	+	w
α-Fucosidase	w	–
Antibiotic susceptibility (per disc)		
Ampicillin (10 µg)	–	+
Penicillin G (10 IU)	–	+
Piperacillin (100 µg)	–	+
Bacitracin (45 µg)	–	+

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558 **Table 2.** Cellular fatty acid profiles (%) of strain CL23^T and *Robertkochia marina* CC-AMO-
559 30D^T.

560 Strains: 1, strain CL23^T; 2, *R. marina* CC-AMO-30D^T. All data presented in the table are
561 from this study. TR, trace ($\leq 0.5\%$); –, not detected. Major components ($> 10\%$) are
562 highlighted in bold.

Fatty acid	1	2
Branched saturated		
iso-C _{13:0}	TR	2.4
iso-C _{14:0}	–	2.4
iso-C _{15:0}	21.8	19.9
iso-C _{16:0}	3.4	6.1
anteiso-C _{15:0}	2.3	5.8
Unsaturated		
C _{15:1} ω5c	0.7	–
C _{17:1} ω6c	1.7	–
C _{17:1} ω8c	0.8	–
Branched unsaturated		
iso-C _{15:1} G	10.8	23.3
iso-C _{16:1} G	–	1.6
iso-C _{16:1} H	1.0	–
anteiso-C _{15:1} A	TR	2.8
Hydroxy		
C _{15:0} 2-OH	0.9	1.5
C _{15:0} 3-OH	2.0	0.6
C _{16:0} 3-OH	1.4	TR
C _{17:0} 3-OH	1.1	TR
iso-C _{16:0} 3-OH	2.6	6.5
iso-C _{17:0} 3-OH	29.5	15.5
Summed features *		
3 †	3.6	–
9 #	5.2	–

563 * Summed features are groups of two or three fatty acids that cannot be separated by GLC
564 with the MIDI system.

565 † Summed feature 3 consisted of iso-C_{15:0} 2-OH, C_{16:1}ω6c and/or C_{16:1}ω7c and annotated here as
566 iso-C_{15:0} 2-OH based on the equivalent chain length (ECL).

567 # Summed feature 9 consisted of iso-C_{17:1}ω9c and/or C_{16:0} 10-methyl.

568


569

570

571

572 **Figure legends**

573



574

575 **Fig. 1.** Neighbor joining 16S rRNA phylogenetic tree manifesting the relationship of strain
 576 CL23^T with closely related taxa of family *Flavobacteriaceae*. Corresponding Genbank
 577 accession numbers are indicated in parentheses. Bootstrap values $\geq 70\%$ based on 1000
 578 resampled datasets are depicted as percentages at nodes. Bootstrap value from left to right for
 579 NJ and ML calculated with same sequence set. Filled circles indicate that corresponding
 580 nodes were also recovered in dendrograms generated using ML algorithm. The sequence of
 581 *Cytophaga hutchinsonii* ATCC 33406^T was used as outgroup. Bar, 0.05 substitutions per
 582 nucleotide position.

583


584

585

586

587

588



589
 590 **Fig. 2.** Neighbor joining phylogenetic tree based on the concatenated sequences of five
 591 housekeeping genes: *rpoB*–*gyrB*–*recA*–*mutL*–*atpD*, indicating the position of strain CL23^T.
 592 Bootstrap values $\geq 70\%$ based on 1000 resampled datasets are depicted as percentages at
 593 nodes; value $< 70\%$ is indicated by a dash. Bootstrap value from left to right for NJ and ML
 594 calculated with same sequence set. Filled circles indicate that corresponding nodes were also
 595 recovered in dendrograms generated using ML algorithm. The sequence of *Cytophaga*
 596 *hutchinsonii* ATCC 33406^T was used as outgroup. Bar, 0.05 substitutions per nucleotide
 597 position.

598

599

Supplementary Materials

Robertkochia solimangrovi* sp. nov., isolated from mangrove soil, and emended description of the genus *Robertkochia

Ming Quan Lam¹, Maša Vodovnik², Maša Zorec², Sye Jinn Chen¹, Kian Mau Goh¹, Adibah Yahya¹, Madihah Md Salleh¹, Zaharah Ibrahim¹, Lili Tokiman³, Simon J. McQueen-Mason⁴, Neil C. Bruce^{4*} and Chun Shiong Chong^{1*}

¹ Department of Biosciences, Faculty of Science, Universiti Teknologi Malaysia, 81310 Skudai, Johor, Malaysia

² Biotechnical Faculty, University of Ljubljana, Groblje 3, 1230 Domzale, Slovenija

³ Johor National Parks Corporation, Kota Iskandar, 79575 Iskandar Puteri, Johor, Malaysia

⁴ Centre for Novel Agricultural Products, Department of Biology, University of York, Wentworth Way, York, YO10 5DD, United Kingdom

***Correspondence:** Chun Shiong Chong, cschong@utm.my; Neil C. Bruce, neil.bruce@york.ac.uk

Supplementary figures




Fig. S1. Polar lipids profile of strain CL23^T. Unidentified lipids; L1–L5, phosphatidylethanolamine; PE, unidentified aminolipids; AL1–AL2, unidentified glycolipids; GL1–GL3.




Fig. S2. Neighbor joining phylogenomic tree manifesting the relationship of strain CL23^T with closely related taxa of family *Flavobacteriaceae*. Corresponding Genbank accession numbers are indicated in parentheses. Bootstrap values $\geq 50\%$ based on 1000 resampled datasets are depicted as percentages at nodes. Bootstrap value from left to right for NJ and ML calculated with same sequence set. Filled circles indicate that corresponding nodes were also recovered in dendrograms generated using ML algorithm. Bar, 0.005 substitutions per nucleotide position.

Supplementary tables

Table S1. List of potential genes for phosphatases, sulfur reduction and nitrate reduction encoded in the genome of strain CL23^T and *R. marina* CC-AMO-30D^T.

Category	Bacterial strain	NCBI Annotation	Accession		
Phosphatases	CL23 ^T	alkaline phosphatase family protein	TRZ44267		
		alkaline phosphatase	TRZ44500		
		alkaline phosphatase	TRZ44343		
		pyrophosphatase	TRZ44378		
		sodium-translocating pyrophosphatase	TRZ44400		
		alkaline phosphatase family protein	TRZ43533		
		alkaline phosphatase family protein	TRZ43596		
		alkaline phosphatase	TRZ42861		
		HAD family phosphatase	TRZ42760		
		alkaline phosphatase family protein	TRZ42969		
		alkaline phosphatase family protein	TRZ41972		
		HAD family phosphatase	TRZ41063		
		<i>R. marina</i> CC-AMO-30D ^T	alkaline phosphatase family protein	TRZ46762	
			alkaline phosphatase family protein	TRZ45488	
	alkaline phosphatase family protein		TRZ45685		
	sodium-translocating pyrophosphatase		TRZ44743		
	HAD family phosphatase		TRZ42656		
	pyrophosphatase		TRZ41149		
	Sulfur reduction	CL23 ^T	HAD family phosphatase	TRZ40862	
			sulfate adenylyltransferase subunit CysN	TRZ46029	
sulfate adenylyltransferase subunit CysD			TRZ46030		
adenylyl-sulfate kinase			TRZ46031		
phosphoadenylylsulfate reductase			TRZ44200		
phosphoadenylylsulfate reductase			TRZ42776		
FAD-binding oxidoreductase			TRZ41175		
LLM class flavin-dependent oxidoreductase			TRZ41182		
<i>R. marina</i> CC-AMO-30D ^T		sulfate adenylyltransferase subunit CysN	TRZ40960		
		sulfate adenylyltransferase subunit CysD	TRZ40970		
		adenylyl-sulfate kinase	TRZ40959		
		phosphoadenylylsulfate reductase	TRZ46694		
		Nitrate reduction	CL23 ^T	nitrite reductase (<i>NirBD</i>)	TRZ44395
				nitrite reductase (NAD(P)H) (<i>NirBD</i>)	TRZ42280

	nitrite reductase (NAD(P)H) small subunit (<i>NirBD</i>)	TRZ42281
	NAD(P)H-nitrite reductase (<i>NirBD</i>)	TRZ42287
	ammonia-forming cytochrome c nitrite reductase (<i>NrfAH</i>)	TRZ42033
	cytochrome c nitrite reductase small subunit (<i>NrfAH</i>)	TRZ42034
<i>R. marina</i> CC- AMO-30D ^T	ammonia-forming cytochrome c nitrite reductase (<i>NrfAH</i>)	TRZ44150
	cytochrome c nitrite reductase small subunit (<i>NrfAH</i>)	TRZ44178