



This is a repository copy of *Dental RECUR randomized trial to prevent caries recurrence in children.*

White Rose Research Online URL for this paper:  
<https://eprints.whiterose.ac.uk/155908/>

Version: Published Version

---

**Article:**

Pine, C.M., Adair, P.M., Burnside, G. et al. (12 more authors) (2020) Dental RECUR randomized trial to prevent caries recurrence in children. *Journal of Dental Research*, 99 (2). pp. 168-174. ISSN 0022-0345

<https://doi.org/10.1177/0022034519886808>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:  
<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# Dental RECUR Randomized Trial to Prevent Caries Recurrence in Children

Journal of Dental Research  
2020, Vol. 99(2) 168–174  
© International & American Associations  
for Dental Research 2019



Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/0022034519886808  
journals.sagepub.com/home/jdr

C.M. Pine<sup>1,2</sup>, P.M. Adair<sup>3</sup>, G. Burnside<sup>4</sup>, L. Brennan<sup>5</sup>, L. Sutton<sup>4</sup>, R.T. Edwards<sup>6</sup>,  
V. Ezeofor<sup>6</sup>, S. Albadri<sup>7</sup>, M.M. Curnow<sup>8</sup>, C. Deery<sup>9</sup>, M.T. Hosey<sup>10</sup> , J. Willis-Lake<sup>11</sup>,  
J. Lynn<sup>12</sup>, J. Parry<sup>13,14</sup>, and F.S.L. Wong<sup>2</sup>

## Abstract

The purpose of this study was to determine the efficacy of a dental nurse–delivered intervention—the Dental RECUR Brief Negotiated Interview for Oral Health (DR-BNI)—in reducing the recurrence of dental caries in children who have a primary tooth extracted. It was based on a 2-arm multicenter randomized controlled trial with blinded outcome assessment. Participants were 5- to 7-y-old children ( $n = 241$ ) scheduled to have primary teeth extracted in 12 UK centers. Test intervention parents ( $n = 119$ ) received DR-BNI led by trained dental nurses. DR-BNI is a 30-min structured conversation informed by motivational interviewing with a forward focus to prevent future caries. Preventive goals are agreed, and a review appointment is made with child's general dental practitioner, who is advised to treat the child as being at high caries risk. The control intervention ( $n = 122$ ) was a parent-nurse conversation about child's future tooth eruption, with advice given to visit a general dental practitioner as usual. At baseline, the DR-BNI group's mean dmft was 6.8, and the control group's was 6.3. A median of 5 teeth were extracted, mainly under general anesthesia. Final dental assessments were conducted by a single examiner visiting 189 schools 2 y after intervention; 193 (80%) of 241 children were examined. In the control group, 62% developed new caries in teeth that were caries-free or unerupted at baseline, as compared with 44% in the test group, a significant reduction ( $P = 0.021$ ). The odds of new caries experience occurring were reduced by 51% in the DR-BNI group as compared with control. There was a 29% decrease in the relative risk of new caries experience in the DR-BNI group as compared with control. This single low-cost, low-intensity intervention was successful in significantly reducing the risk of recurrence of dental caries in children. This trial has implications for changing pediatric dental practice internationally. Training in and implementation of a motivational interviewing–informed brief intervention provides opportunities for dental nurses to facilitate behavior change improving the oral health of children at high caries risk (ISRCTN 24958829).

**Keywords:** caries detection/diagnosis/prevention, caries treatment, child dentistry, clinical studies/trials, health services research, motivational interviewing

<sup>1</sup>Research and Innovation, Salford Royal NHS Foundation Trust, Northern Care Alliance NHS Group, Summerfield House, Salford, UK

<sup>2</sup>Barts and the London Schools of Medicine and Dentistry, Institute of Dentistry, Queen Mary University of London, London, UK

<sup>3</sup>Centre for Improving Health-Related Quality of Life, School of Psychology, Queen's University, Belfast, UK

<sup>4</sup>Department of Biostatistics, University of Liverpool, Liverpool, UK

<sup>5</sup>Health Education North West, Regatta Place, Liverpool, UK

<sup>6</sup>Centre for Health Economics and Medicines Evaluation, Bangor University, Bangor, UK

<sup>7</sup>Paediatric Dentistry, School of Dentistry, University of Liverpool, Liverpool, UK

<sup>8</sup>Public Dental Service, Broxden Dental Centre, NHS Tayside, Perth, UK

<sup>9</sup>School of Clinical Dentistry, University of Sheffield, Sheffield, UK

<sup>10</sup>Paediatric Dentistry, Centre of Oral, Clinical and Translational Science, Faculty of Dentistry, Oral and Craniofacial Sciences, Kings College London, London, UK

<sup>11</sup>Kent Community Health NHS Foundation Trust, Maidstone, UK

<sup>12</sup>Community Dental Service, Arches Health and Care Centre, Belfast Health and Social Care Trust, Belfast, UK

<sup>13</sup>Special Care Dental Service, Sussex Community NHS Foundation Trust, Brighton, UK

<sup>14</sup>Paediatric Dentistry, University College Cork, Cork, Ireland

A supplemental appendix to this article is available online.

## Corresponding Author:

C.M. Pine, Barts and the London Schools of Medicine and Dentistry, Institute of Dentistry, Queen Mary University of London, Turner St, Whitechapel, London E1 2AD, UK.

Email: c.m.pine@qmul.ac.uk

## Introduction

Dental extractions are the highest cause of planned admission to hospitals for children <11 y of age in England and Scotland (Royal College of Surgeons 2015; Information Services Division 2016). Surgery cannot prevent future decay because the underlying etiologic factors of high sugar intake, irregular toothbrushing with fluoride toothpaste, and symptomatic dental attendance (Public Health England 2017) are unchanged. A review of children having their first permanent molars extracted found that 40% had previous extractions of primary teeth (Albadri et al. 2007). Parents whose children develop new decay postextraction may have struggled to accept health advice or felt unable to change previous unhealthy behaviors (Amin and Harrison 2007).

In other habitual behaviors, such as smoking, motivational interviewing (MI) approaches have moved people from inaction to action (Prochaska et al. 2008). MI has been used in successful interventions influencing parents to adopt and maintain preventive child oral health behaviors (Weinstein et al. 2004; Weinstein et al. 2006; Freudenthal and Bowen 2010). A brief intervention used MI within a structured framework and was taught to practitioners in a short training program (Emmons and Rollnick 2001). The 30-min intervention was delivered in a medical setting and changed negative attitudes, beliefs, and behaviors; to date, no dental studies have been conducted.

We developed a psychosocial intervention—the Dental RECUR Brief Negotiated Interview for Oral Health (DR-BNI)—to be delivered to parents of children who have had a dental extraction of primary teeth (Pine et al. 2015). DR-BNI can be delivered by dental nurses (assistants) and is designed to develop shared understanding with parents through communication about adopting healthier behaviors to reduce recurrence of caries in their children. Our aim was to test the efficacy of a dental nurse–delivered intervention, the DR-BNI, in reducing the recurrence of dental caries in children who had a primary tooth extracted 2 y previously.

## Methods

### Study Design

This was a 2-arm multicenter randomized controlled trial with blinded outcome assessment.

### Primary Outcome

The primary outcome was dental caries experienced in the 2 y postintervention on any tooth in either dentition at the dental level of involvement, which had been caries-free at baseline.

### Sample Size

The primary outcome variable is binary, taking the value 1 where a child had caries experience after 2 y on any tooth in either primary or permanent dentition, which was caries-free

(or unerupted) at baseline; 0, otherwise. From a previous clinical trial of 5- to 7-y-olds who had extractions (Curnow et al. 2010), 87% developed new carious teeth 2 y later. Setting the minimum clinically significant difference to 20% (67% in the test group), with 80% power and a significance level of 0.05, gave a minimum sample size of 78 per group. Allowing 30% with incomplete final assessments gave a final required sample size of 112 per group.

### Governance

Research ethics and UK NHS approvals were obtained. Participants were recruited in 12 UK centers, including dental hospital clinics and secondary care centers, which provided extraction services. Principal investigators were pediatric dentists heading the centers. All staff received training in good clinical practice, trial protocol, and diagnostic criteria for baseline caries assessments. Each site had investigator site files and participant case report forms (protocol details in Pine et al. 2015).


### Recruitment

Inclusion criteria were written consent from parents/legal guardians of patients aged 5 to 7 y who were scheduled to have at least 1 primary tooth extracted for dental caries under general anesthesia, inhalation sedation, or local anesthesia. Exclusion criteria were having all first permanent molar teeth extracted, participation in another trial or in the previous 3 months, severe disability, or no parental consent given. Randomization and intervention delivery occurred after enrollment and up to 6 wk postextraction (Appendix 1; Fig. 1). Randomization was stratified by site, and participants were randomized with sequentially numbered sealed envelopes.

### Test Intervention

DR-BNI is a “talking” intervention, a 30-min therapeutic conversation between dental nurse (assistant) and parent/caregiver, structured in 6 segments (Build Rapport, Ask about Pros and Cons, Feedback, Readiness to Change, Action Plan, Dental Appointment and Thanks). The intervention, developed by a clinical and health psychologist (P.M.A.), is informed by MI techniques. The focus is forward-looking about keeping the newly erupting dentition healthy. DR-BNI seeks to increase parental self-efficacy for 3 child oral health–related behaviors: twice-daily toothbrushing with fluoride toothpaste; controlling free sugars intake, especially at bedtime; and attending a dentist regularly for preventive care rather than symptomatically.

Dental nurses attended 1 d of training by P.M.A. in DR-BNI. Training followed MI principles combined with health behavior change techniques (Miller and Moyers 2006) for promoting oral health. The aim was to explore opportunities with parents that might lead to change in past behaviors rather than telling them what to do. Nurses were advised to try to agree on 1 or 2 goals with parents, if appropriate, using the behaviors described in a modified dental contemplation ladder (Coolidge



**Figure 1.** CONSORT flow diagram. DR-BNI, Dental RECUR Brief Negotiated Interview for Oral Health; GDP, general dental practitioner.

et al. 2011; Appendix 1). Nurses were trained in change talk, developing a change plan and consolidating commitment. After training, nurses practiced in their clinics (Appendix 1).

The agreed-on goals were tailored for each family, committing to a specific dental health-related behavior for their child, such as changing from sugar-containing drinks to sugar-free or brushing their child's teeth at bedtime with fluoridated toothpaste. At intervention end, the nurse assisted parents to make a recall appointment with their child's general dental practitioner (GDP) 3 to 4 mo after the intervention, and in addition, the nurse sent a text reminder. Parents left the clinics with a copy of their agreed-on goals and the child's review dental appointment (Appendix 2).

### Placebo Control Intervention

This was developed by C.M.P. with the same structure as the DR-BNI, but the delivery mode was educational, giving information on tooth eruption between the ages of 6 and 14 y. The conversation was structured around concepts of growing up,

shedding teeth, and growing new ones, with illustrations. There was no discussion on caries prevention, and parents were advised to attend their child's dental practice as usual. All families in both groups received the same leaflet on dental development to take home.

### Intervention Delivery

Most parents received the intervention at a routine appointment (assessment, pre-extraction, or extraction), and where this was not possible, an appointment was made between enrollment and 6 wk postoperatively. With parents' permission, an audio recording was made of the intervention conversation (Appendix 1).

### Contacts with Dentists

Letters to GDPs of DR-BNI participants noted the agreed preventive goals, and dentists were sent a booklet containing advice on preventive care advised for children at high caries risk, including 3-mo recalls (Scottish Intercollegiate Guidelines Network 2014; Public Health England 2017). GDP booklets contained case report forms to be completed and returned about child dental care in the first year. At the end of the second year, we contacted GDPs about appointments attended and dental care provided. Similarly, we contacted control group participants' GDPs at 1 and 2 y ( $\pm 3$  mo) postenrollment. GDPs received

payments from research funds to contribute to costs of completing case report forms.

### Measures

At the intervention appointment, parents completed the Oral Health Behaviours Questionnaire incorporating measures of parental self-efficacy for child toothbrushing and dietary sugar intake (Adair et al. 2004). A general parental self-efficacy scale (Prochaska et al. 2008) and contemplation ladder measuring readiness to change behavior (Coolidge et al. 2011) and modified for oral health (Appendix 1) were also completed. Prior to extractions, pediatric dentists at the centers undertook dental examinations noting teeth with caries into dentine and teeth to be extracted.

A single examiner (C.M.P.) undertook the final clinical assessments in participants' primary schools or homes 2 y after the intervention,  $\pm 3$  mo. Children were examined supine with a single-use plane mouth mirror, and teeth were illuminated by Daray light of 2,000 lux. Upper anterior buccal plaque was

recorded as an indicator of oral cleanliness. Teeth were examined for untreated caries into dentine, restorations, and fissure sealants (Pitts et al. 1997). Cotton wool rolls were used to dry teeth, and probes were available to remove debris and to check the integrity of restorations and the presence of sealants (Appendix 1). All dental assessments were blind to group assignment.

### Statistical Analysis

Analysis of the primary outcome variable used logistic regression, adjusted for the stratification variable center and baseline dmft. Unadjusted relative risk estimates were calculated. The primary outcome was analyzed with the full analysis set by using the intention-to-treat principle as far as possible, with participants excluded only where outcome data were unavailable. Per-protocol analysis was conducted to test the robustness of the main results to departures from intention to treat. Sensitivity analyses were conducted with multiple imputation to investigate robustness of analysis to missing primary outcome data.

### Results

The first patient was randomized in April 2015; more centers entered, and the last patient was randomized in November 2016. Ten centers were in England, 1 in Scotland, and 1 in Northern Ireland. A total of 119 children were randomized to the DR-BNI group and 122 to the placebo control (Fig. 1). Of these 241 children, 235 (98%) received the interventions.

Baseline characteristics (Table 1) were similar. Children were, on average, 6 y old. There were similar numbers of boys and girls; a third of mothers completed education at secondary school or earlier. Over half of parents reported that their children had sweets every day or most days, with over a third having sugary drinks frequently. Children had high levels of deciduous caries experience, with a mean dmft of 6.8 in the DR-BNI group and 6.3 in the control group. At recruitment, children had a median 5 teeth extracted, almost all under general anesthesia. Not all first permanent molars were erupted, with a mean DMFT of 0.1 in the DR-BNI group and 0.0 in the control group.

Intervention compliance was >95% for both groups, with 96% of parents agreeing on preventive goals, such as to reduce specific sugar behaviors and/or improve toothbrushing frequency (Appendix 2).

Two years after intervention, final dental examinations were undertaken across the United Kingdom by a single examiner (C.M.P.) visiting 189 schools and 2 children at home. Of 241 children, 193 (80%) were examined. For 2 children, baseline assessments had not been completed; therefore, 191 of 241 (79%) were analyzed, comprising 88 (74%) in the DR-BNI group and 103 (84%) in the control group.

Table 2 shows that 62% of children in the control group developed new caries in teeth that were previously caries-free or unerupted, as compared with 44% of children in the test

**Table 1.** Baseline Parameters.

	Children, Mean $\pm$ SD (Range) or n (%)	
	DR-BNI (n = 119)	Control (n = 122)
Age, y	6.3 $\pm$ 0.8 (4.8 to 8.0)	6.4 $\pm$ 0.8 (4.9 to 8.0)
Sex		
Female	63 (53)	61 (50)
Male	56 (47)	61 (50)
Mother's education		
No formal education/primary school	6 (5)	6 (5)
Secondary school	34 (29)	30 (25)
Further education (college)	34 (29)	46 (38)
Higher education (university)	34 (29)	27 (22)
Missing	11 (9)	13 (11)
Sweets consumption <sup>a</sup>		
Every day/most days	64 (54)	68 (56)
Once a week/occasionally/never	46 (39)	50 (41)
Missing	9 (8)	4 (3)
Sugary drinks consumption <sup>a</sup>		
Every day/most days	43 (36)	57 (47)
Once a week/occasionally/never	66 (56)	61 (50)
Missing	10 (8)	4 (3)
Toothbrushing <sup>a</sup>		
Twice/three times a day	94 (79)	99 (81)
Not every day/once a day	19 (16)	20 (16)
Missing	6 (5)	3 (3)
dmft	6.8 $\pm$ 3.4 (1.0 to 15.0)	6.5 $\pm$ 3.0 (1.0 to 14.0)
No. of teeth extracted <sup>b</sup>	5.5 $\pm$ 3.3 (1.0 to 15.0)	5.2 $\pm$ 2.9 (1.0 to 14.0)
DMFT <sup>c</sup>	0.1 $\pm$ 0.4	0.0 $\pm$ 0.2

DR-BNI, Dental RECUR Brief Negotiated Interview for Oral Health.

<sup>a</sup>Oral Health Behaviours Questionnaire.

<sup>b</sup>Median for each group: 5.0.

<sup>c</sup>Median for each group: 0.0.

group, which was a significant reduction ( $P = 0.021$ ). Figure 2 gives the distribution of new caries in teeth, providing a measure of severity. The odds of new caries experience occurring were reduced by 51% in the DR-BNI group as compared with control. There was a 29% decrease in the relative risk of new caries experience in the DR-BNI group as compared with control. Similar significant differences were found in 2 sensitivity analyses: 1 with the per-protocol data set and 1 with multiple imputation to replace missing outcome data.

To explore whether the differences arose from a single center, perhaps due to a particularly effective nurse intervention, the direction of differences in proportions for all centers was analyzed (Table 3). Sufficient numbers were available in 9 centers, and in 8 of the 9, the direction was the same, showing consistency in benefit to the DR-BNI group.

Results from the ninth center, L, were in the opposite direction. Families were almost entirely of Bangladeshi heritage, with very high levels of childhood caries (Public Health England 2018). Although 1 of the 2 dental nurses delivering


**Table 2.** Primary Outcome: Proportion of Children with Dental Caries in Previously Caries-free or Unerupted Teeth 2 y after Intervention.

DR-BNI	Control	Difference in Proportions (95% CI)	Adjusted Odds Ratio (95% CI)
0.44 (n = 88)	0.62 (n = 103)	0.18 (0.04 to 0.32), P = 0.014	0.49 (0.26 to 0.90), P = 0.021 (n = 191)
		Per protocol	
0.46 (n = 81)	0.63 (n = 94)	0.17 (0.02 to 0.32), P = 0.024	0.52 (0.27 to 0.99), P = 0.046 (n = 175)
		Multiple imputation <sup>a</sup>	
—	—	—	0.44 (0.24 to 0.81), P = 0.008 (n = 241)

The odds of new caries experience occurring were reduced by 51% in the DR-BNI group as compared with control. Relative risk = 0.71 (95% CI, 0.54 to 0.94), P = 0.014. There was a 29% decrease in the risk of new caries experience in the DR-BNI group as compared with the control. Significance at P < .05.

DR-BNI, Dental RECUR Brief Negotiated Interview for Oral Health.

<sup>a</sup>1,000 imputations.



**Figure 2.** Proportion of children by group who developed new caries and number of teeth affected. DR-BNI, Dental RECUR Brief Negotiated Interview for Oral Health.

the interventions was bilingual in English and Sylheti, it is likely that additional interventions may be needed to facilitate changes in cultural norms in this community.

Over 2 y, around 60% of children returned to the same dental practice that had referred them for extractions (Table 4). There was a nonsignificant trend for DR-BNI children to return sooner, 3 to 4 mo after extractions. At the practices, similar proportions of children were given fluoride varnish applications (around 80%) and had fissure sealants placed (around 30%). The difference between the groups was in the proportion of children who had fillings placed—namely, 22% in DR-BNI group as compared with 40% in control. This mirrored the higher caries experience found in the independent final dental examinations.

## Discussion

This trial tested the efficacy of a brief negotiated interview, which changed how dental teams traditionally talk to patients. Instead of direct advice on what not to do, nurses had a structured conversation supporting families to decide how they could improve their children's dental future. The intervention

was theory driven and targeted to children at the highest risk for developing new caries. The decision to extract multiple teeth is a “teachable moment” for many families, when they may be more receptive to considering making things better in the future (Papies 2016).

Formal training for nurses took 1 d, with posttraining practice in their clinics to develop conversational skills. During training, we challenged some nurses' criticism of parents' behaviors that had led to so much caries. We focused on empathic communication to support development of healthier routines for families.

Recruitment took >18 mo, as participants were a hard-to-reach group, intermittently engaging with dental care (Huntington et al. 2017). Some families were very disadvantaged and known to social services, as families came from many countries with diverse cultures. The

importance of family environment (Mattila et al. 2000) and social determinants of health underpin barriers to healthy behaviors (World Health Organization 2008). Taking a non-judgmental approach in the DR-BNI led families to engage in considering changes that they identified as possible to undertake for their children in their day-to-day life.

Nearly two-thirds of children returned to their referring dental practice, and higher numbers did not return in the DR-BNI group. As similar proportions of children in both groups had fluoride varnish and fissure sealants, this does not explain the reduced caries levels in the DR-BNI group. Therefore, it appears that it was the nurses' intervention with parents addressing underlying etiologic factors, potentially reinforced in dental practices, that was critical to achieving a significant benefit for children in the DR-BNI group.

Undertaking final dental assessments in schools was a major endeavor, as children attended 189 schools across the United Kingdom. Nevertheless, it was worthwhile, given that 80% of children were examined, far more than if parents had been asked to bring children to clinics. Critically, this comprehensive data collection allowed demonstration of the consistency in the direction of benefit across 8 of the 9 centers. This

**Table 3.** Primary Outcome by Center: Proportion of Children with Dental Caries in Previously Caries-free or Unerupted Teeth 2 y after Intervention.

Center	Children, <sup>a</sup> <i>n</i>	DR-BNI ( <i>n</i> = 88)	Control ( <i>n</i> = 103)	Difference in Proportion of Children with New Caries
A	8	0.40	0.67	-0.27
B	31	0.33	0.56	-0.23
C	9	0.40	1.00	-0.60
D	24	0.44	0.60	-0.16
E	18	0.29	0.45	-0.16
F	24	0.55	0.77	-0.22
G	21	0.10	0.45	-0.35
H	23	0.36	0.67	-0.31
I	2	—	0.50	—
J	2	1.00	—	—
K	5	1.00	1.00	0
L	24	0.82	0.62	+0.20

Of 12 centers, 8 showed a reduction in the proportions of children with new caries in the DR-BNI group as compared with control; 3 centers' numbers were too low to compare; and for 1 center, L, the difference was in the other direction.

DR-BNI, Dental RECUR Brief Negotiated Interview for Oral Health.

<sup>a</sup>Total, *N* = 191.

**Table 4.** Dental Attendance during the 2 y Postintervention and Dental Treatment Provided.

	Children, <i>n</i> (%)	
	DR-BNI ( <i>n</i> = 119)	Control ( <i>n</i> = 122)
Attended dental practice		
At least once	72 (61)	78 (64)
Did not attend	28 (24)	30 (25)
Missing data	19 (16)	14 (12)
Dental treatment for those who attended at least once		
Fluoride varnish	61 (85)	61 (78)
Fissure sealants	21 (29)	25 (32)
At least one restoration	16 (22)	31 (40)

DR-BNI, Dental RECUR Brief Negotiated Interview for Oral Health.

supports the conclusion that the effect was not dependent on a single outperforming center or individual nurse but demonstrated that positive outcomes were general and attainable.

## Conclusions and Implications for Clinical Practice

In a wide range of children at high caries risk, this single low-cost, low-intensity intervention was successful in significantly reducing the risk of new caries experience.

This trial has implications for changing pediatric dental practice internationally. Training in, and implementation of, an MI-informed brief intervention provides opportunities for dental nurses to go beyond clinical prevention to facilitate behavior change and to support oral health improvements for children at high caries risk. The lead research team has been invited by

Health Education England (North West) to develop the DR-BNI into a training program for dental nurses in the NHS.

## Author Contributions

C.M. Pine, P.M. Adair, G. Burnside, contributed to conception, design, data analysis, and interpretation, drafted and critically revised the manuscript; L. Brennan, contributed to data acquisition, drafted and critically revised the manuscript; L. Sutton, contributed to data analysis and interpretation, drafted and critically revised the manuscript; R.T. Edwards, contributed to design and data analysis, drafted and critically revised the manuscript; V. Ezeofor, contributed to data analysis, critically revised the manuscript; S. Albadri, M.M. Curnow, C. Deery, M.T. Hosey, J. Willis-Lake, J. Lynn, J. Parry, F.S.L. Wong, contributed to data acquisition, critically revised the manuscript. All authors gave final approval and agree to be accountable for all aspects of the work.

## Acknowledgments

Research ethics and governance approval via the IRAS system from the Greater Manchester Central NRES Committee (13/NW/0466) and the Salford Royal Foundation Trust R&D Department, the trial sponsor. Thanks are due to Beverley Greenhalgh, sponsor representative, and Stephen Brown, database manager. Further details are provided in Appendix 3. Thanks are due to all those who assisted in recruitment and program delivery of this clinical trial—notably, to the families, children, GPs, and dental nurses and to Geoffrey Pine, who supported the logistics for visits to many of the 189 schools. Dental nurses delivering the interventions at each center are thanked and recognized: Julie Bonar, Denise Champion, Rizwana Dakri, Francesca Davies, Beverley Gayle, Claire Hulme, Megan Lane, Emma Mahoney, Charlotte Marlow, Nicola McDowell, Alison McNealy, Louise Morris, Bethany Powe, Sutinder Saroi, Aimee Shaw, Eleanor Taylor, Stacey Teece; Kerry Whitmore, and Vanessa Wilson. Appendix 3 gives details of their NHS Trust sites. This article reports independent research funded by the UK National Institute for Health Research under the Research for Patient Benefit Programme (grant PB-PG-0610-22310). The views expressed in this publication are those of the authors and not necessarily those of the National Institute for Health Research or the Department of Health and Social Care. The grant was awarded to Salford Royal NHS Foundation Trust with C.M. Pine as chief investigator. NOCLOR NHS Research Support Services provided recruitment costs to the Barts Health NHS Trust site. The authors declare no potential conflicts of interest with respect to the authorship and/or publication of this article.

## ORCID iD

M.T. Hosey  <https://orcid.org/0000-0003-1178-4106>

## References

- Adair PM, Pine CM, Burnside G, Nicoll AD, Gillett A, Anwar S, Broukal Z, Chestnutt IG, Declerck D, Ping FX, et al. 2004. Familial and cultural perceptions and beliefs of oral hygiene and dietary practices among ethnically and socio-economically diverse groups. *Community Dent Health*. 21(1 Suppl):102–111.

- Albadri S, Zaitoun H, McDonnell S, Davidson L. 2007. Extraction of first permanent molar teeth: results from three dental hospitals. *Br Dent J*. 203(7):E14.
- Amin MS, Harrison RL. 2007. A conceptual model of parental behaviour change following a child's dental general anaesthesia procedure. *Pediatr Dent*. 29(4):278–286.
- Coolidge T, Skaret E, Heima M, Johnson EK, Hillstead MB, Farjo N. 2011. Thinking about going to the dentist: a contemplation ladder to assess dentally-avoidant individuals' readiness to go to a dentist. *BMC Oral Health*. 11(4):1–12.
- Curnow M, Pine C, Burnside G, Nicholson J, Chesters R, Huntington E. 2010. A randomised controlled trial of the efficacy of supervised toothbrushing in high-caries-risk children. *Caries Res*. 36(4):294–300.
- Emmons KM, Rollnick S. 2001. Motivational interviewing in health care settings: opportunities and limitations. *Am J Prev Med*. 20 (1):68–74.
- Freudenthal JJ, Bowen DM. 2010. Motivational interviewing to decrease parental risk-related behaviors for early childhood caries. *J Dent Hyg*. 84(1):29–34.
- Huntington C, Newton JT, Donaldson N, Lioffi C, Reynolds PA, Alharatani R, Hosey MT. 2017. Lessons learned on recruitment and retention in hard-to-reach families in a phase III randomised controlled trial of preparatory information for children undergoing general anaesthesia. *BMC Oral Health*. 17(1):122.
- Information Services Division. 2016. Childhood admissions summary data, 2011/12–2015/16. Edinburgh (Scotland): Information Services Division.
- Mattila M-L, Rautava P, Sillanpää M, Paunio P. 2000. Caries in five-year-old children and associations with family-related factors. *J Dent Res*. 79(3):875–881.
- Miller WR, Moyers TB. 2006. Eight stages in learning motivational interviewing. *J Teach Addict*. 5:3–17.
- Papies EK. 2016. Health goal priming as a situated intervention tool: how to benefit from non-conscious motivational routes to health behaviour. *Health Psychol Rev*. 10(4):408–424.
- Pine CM, Adair P, Burnside G, Robinson L, Edwards RT, Albadri S, Curnow M, Ghahreman M, Henderson M, Malies C, et al. 2015. A new primary dental care service compared with standard care for child and family to reduce the re-occurrence of childhood dental caries (Dental RECUR): study protocol for a randomised controlled trial. *Trials*. 16:505–514.
- Pitts NB, Evans DJ, Pine CM. 1997. British Association for the Study of Community Dentistry (BASCD) diagnostic criteria for caries prevalence surveys—1996/97. *Community Dent Health*. 14 Suppl 1:6–9.
- Prochaska JJ, Spring B, Nigg CR. 2008. Multiple health behavior change research: an introduction and overview. *Prev Med*. 46(3):181–188.
- Public Health England. 2017. Delivering better oral health: an evidence based toolkit for prevention. Third ed. London (UK): Public Health England.
- Public Health England. 2018. National Dental Epidemiology Programme for England: oral health survey of five-year-old children. A report on the inequalities found in prevalence and severity of dental decay. London (UK): Public Health England. PHE 2018081.
- Royal College of Surgeons. 2015. The state of children's oral health in England. London (UK): Royal College of Surgeons.
- Scottish Intercollegiate Guidelines Network. 2014. Dental interventions to prevent caries in children. Edinburgh (Scotland): Scottish Intercollegiate Guidelines Network.
- Weinstein P, Harrison R, Benton T. 2004. Motivating mothers to prevent caries in their young children: one-year findings. *J Am Dent Assoc*. 135(6):731–738.
- Weinstein P, Harrison R, Benton T. 2006. Motivating mothers to prevent caries: confirming the beneficial effect of counselling. *J Am Dent Assoc*. 137(6):789–793.
- World Health Organization. 2008. Closing the gap in a generation: health equity through action on the social determinants of health. Final report of the Commission on Social Determinants of Health. Geneva (Switzerland): World Health Organization.