



This is a repository copy of *Porphyromonas gingivalis* lipopolysaccharide rapidly activates trigeminal sensory neurons and may contribute to pulpal pain.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/157154/>

Version: Published Version

Article:

Kaewpitak, A., Bauer, C.S., Seward, E.P. orcid.org/0000-0001-5311-1343 et al. (2 more authors) (2020) *Porphyromonas gingivalis* lipopolysaccharide rapidly activates trigeminal sensory neurons and may contribute to pulpal pain. *International Endodontic Journal*, 53 (6). pp. 846-858. ISSN 0143-2885

<https://doi.org/10.1111/iej.13282>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:
<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

***Porphyromonas gingivalis* lipopolysaccharide rapidly activates trigeminal sensory neurons and may contribute to pulpal pain**

A. Kaewpitak^{1,†}, C. S. Bauer^{2,‡}, E. P. Seward², F. M. Boissonade¹  & C. W. I. Douglas¹

¹School of Clinical Dentistry; and ²Department of Biomedical Science, University of Sheffield, Sheffield, UK

Abstract

Kaewpitak A, Bauer CS, Seward EP, Boissonade FM, Douglas CWI. *Porphyromonas gingivalis* lipopolysaccharide rapidly activates trigeminal sensory neurons and may contribute to pulpal pain. *International Endodontic Journal*, 53, 846–858, 2020.

Aim To determine whether *Porphyromonas gingivalis* lipopolysaccharide (LPS) can directly activate trigeminal neurons, to identify which receptors are involved and to establish whether activation leads to secretion of the neuropeptide calcitonin gene-related peptide (CGRP) and/or the translocation of NF- κ B.

Methodology Mouse trigeminal ganglion (TG) cells were cultured *in vitro* for 2 days. The effect of *P. gingivalis* LPS (20 μ g mL⁻¹) on calcium signalling was assessed (by calcium imaging using Cal-520 AM) in comparison with the transient receptor potential channel A1 (TRPA1) agonist cinnamaldehyde (CA; 100 μ mol L⁻¹), the TRP channel V1 (TRPV1) agonist capsaicin (CAP; 1 μ mol L⁻¹) and high potassium (60 mmol L⁻¹ KCl). TG cultures were pre-treated with either 1 μ mol L⁻¹ CLI-095 to block Toll-like receptor 4 (TLR4) signalling or with 3 μ mol L⁻¹ HC-

030031 to block TRPA1 signalling. CGRP release was determined using ELISA, and nuclear translocation of NF- κ B was investigated using immunocytochemistry. Data were analysed by one-way analysis of variance, followed by Bonferroni's *post hoc* test as appropriate.

Results *Porphyromonas gingivalis* LPS directly exerted a rapid excitatory response on sensory neurons and non-neuronal cells ($P < 0.001$ to $P < 0.05$). The effects on neurons appear to be mediated via TLR4- and TRPA1-dependent pathways. The responses were accompanied by an increased release of CGRP ($P < 0.001$) and by NF- κ B nuclear translocation ($P < 0.01$).

Conclusions *Porphyromonas gingivalis* LPS directly activated trigeminal sensory neurons (via TLR4 and TRPA1 receptors) and non-neuronal cells, resulting in CGRP release and NF- κ B nuclear translocation. This indicates that *P. gingivalis* can directly influence activity in trigeminal sensory neurons and this may contribute to acute and chronic inflammatory pain.

Keywords: mouse, pain, symptomatic pulpitis, TLR4, trigeminal, TRPA1.

Received 11 November 2018; accepted 12 February 2020

[†]Faculty of Dentistry, Prince of Songkla University, Hat Yai, Thailand

[‡]Sheffield Institute for Translational Neuroscience, University of Sheffield, Sheffield, UK

Correspondence: Fiona M. Boissonade, School of Clinical Dentistry, University of Sheffield, Claremont Crescent, Sheffield S10 2TA, UK (e-mail: f.boissonade@sheffield.ac.uk).

[The copyright line for this article was changed on 8 April 2020 after original online publication.]

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Introduction

Infection of the dental pulp initiates a variety of pathophysiological changes including the generation of pain. Whilst immune mechanisms are known to indirectly influence nociception, bacteria or their products have also been reported to activate sensory neurons directly. A range of oral bacteria are known to infect dental pulps, many of which are Gram-negative, and amongst which is *Porphyromonas gingivalis* (Rôças et al. 2002).

A major component of Gram-negative bacterial surfaces is lipopolysaccharide (LPS), and previous investigators have shown that LPS might directly activate sensory neurons (Diogenes et al. 2011, Tse et al. 2014). *Escherichia coli* LPS has been shown to increase neuronal excitability of dorsal root ganglion (DRG) and trigeminal ganglion (TG) neurons (Meseguer et al. 2014). However, *P. gingivalis* LPS differs from that of *E. coli* and it is not known whether this and other nonenterobacterial LPS that would be found in pulpal infection have the same effects and, if so, which neuronal receptors are involved.

There are two categories of cell surface receptors that are thought to be key participants in a rapid nociceptor response to LPS and the initiation of neurogenic inflammation: Toll-like receptors (TLRs) and transient receptor potential channels (TRPs). TLRs are pattern-recognition receptors highly conserved throughout evolution (Santoni et al. 2015), and it is known that they can be activated by some forms of LPS (Diogenes et al. 2011, Meseguer et al. 2014, Tse et al. 2014). Of the 11 TLRs identified in humans so far (Qi et al. 2011), TLR4 specifically recognizes LPS of Gram-negative bacteria (Coats et al. 2005) and previous workers have shown that TLR4 is expressed by many cell types within the dental pulp, including odontoblasts (Jiang et al. 2006), fibroblasts (Hirao et al. 2009) and nerve fibres (Wadachi & Hargreaves 2006). TRPs are a family of cation channels that respond to a range of thermal and chemical stimuli including inflammatory stimuli (Geppetti et al. 2008, Veldhuis & Bunnett 2013, Santoni et al. 2015). *E. coli* LPS is capable of sensitizing TRPV1 via a mechanism involving TLR4 (Diogenes et al. 2011). Also, the TRPA1 receptor can be activated by the anionic lipid A of LPS (Meseguer et al. 2014). Although TRPV1 and TLR4 co-expression has been demonstrated (Diogenes et al. 2011), a complete picture of the crosstalk with other receptors and the actual pathways that these receptors feed into has not been elucidated. Therefore, the aim of this study was to determine whether *P. gingivalis* LPS could induce

pulpal pain by directly activating trigeminal neurons and lead to secretion of the neuropeptide calcitonin gene-related peptide (CGRP) and the translocation of NF- κ B.

Materials and methods

Animals

Healthy, 3- to 9-month-old C57BL/6 wild-type adult mice (30–35 g) were used. All animals were maintained on a 12-h light/dark cycle in a temperature-controlled environment and given food *ad libitum*. All animal procedures were conducted under the Animal (Scientific Procedures) Act 1986, and approved by the UK Home Office.

Mouse trigeminal ganglia (TG) primary cultures

Mice were sacrificed by an overdose of isoflurane anaesthetic and cervical dislocation, and the TG were removed immediately. TG were placed in ice-cold, calcium- and magnesium-free Hanks' balanced salt solution (HBSS; Invitrogen, Paisley, UK), and then quickly washed twice with HBSS supplemented with 1% penicillin/streptomycin (PS) (Gibco, Paisley, UK) at 37 °C in 5% CO₂. Ganglia were then treated with 1.25% w/v type IV collagenase (Sigma, Gillingham, UK) once for 60 min and then again for 45 min, followed by 1.25% w/v trypsin (Sigma) for 15 min at 37 °C. After digestion, the TG were homogenized mechanically by pipetting up and down, and centrifuged at 1500 *g* for 3 min, then re-suspended in supplemented medium (DMEM/F12 media, 1% w/v BSA, 1% N2 supplement, 1% PS; Bottenstein & Sato 1979). The mixtures were transferred to poly-l-lysine and laminin-coated coverslips that were placed at the bottom of 24-well tissue culture trays and cultured for 2–3 days with media change once after 24 h. All experiments on cultured cells were performed on day 2–3.

Intracellular calcium measurements

Cells were loaded for 60 min at 37 °C with 5 μ mol L⁻¹ Cal-520 AM (Abcam, Cambridge, UK) in DMEM/F12 medium after which coverslips were transferred into a chamber (Warner Instruments RC-25F; Harvard Apparatus Ltd, Edenbridge, UK) mounted on the stage of a Zeiss Axiovert S100TV microscope (Zeiss, Jena, Germany) and continuously perfused with buffer (142 mmol L⁻¹ NaCl,

5 mmol L⁻¹ NaHCO₃, 10 mmol L⁻¹ HEPES, 16 mmol L⁻¹ glucose, 2 mmol L⁻¹ KCl, 2 mmol L⁻¹ CaCl₂, 1 mmol L⁻¹ MgCl₂, and 0.1% BSA, pH 7.3) at a flow rate of 5 mL min⁻¹. Cells were illuminated at 488 nm with a monochromator (TILL Photonics, Kaufbeuren, Germany), and viewed using a band-pass filter (510–540 nm) and 40× objective (NA1.3; Zeiss). Images were acquired at 6-second intervals using a Photometrics Cascade 512B CCD camera (Photometrics UK Ltd, Marlow, UK) and time-lapse sequences recorded using WinFluor[®] software (V3.8.7; Strathclyde Imaging Software, University of Strathclyde, Glasgow, UK). Following background subtraction, fluorescence values (measured for every image from regions of interest [ROIs] placed over the soma of cells) were exported and plotted using GraphPad Prism 6 (GraphPad Software, San Diego, CA, USA). Cells were classified as responsive to a particular stimulus when fluorescence, measured in an ROI, increased by more than five standard deviations over the baseline fluorescence. Baseline fluorescence was measured as the average of the last 10 frames before application of tested products or agonists. Data are presented as the changes in intensity of fluorescence between the baseline and peak of the activated curve. This calculation was used to account for variations in the starting fluorescence of individual cells.

The effect of *P. gingivalis* LPS (20 µg mL⁻¹) on calcium signalling was assessed in comparison with the TRPA1 agonist cinnamaldehyde (CA; 100 µmol L⁻¹), the TRPV1 agonist capsaicin (CAP; 1 µmol L⁻¹) and high potassium (60 mmol L⁻¹ KCl). To determine whether signalling was occurring via TLR4 or the TRPA1 receptor, we used specific blocking agents applied to the TG cultures prior to exposure to LPS. TG cultures were pre-treated with either 1 µmol L⁻¹ CLI-095 (InvivoGen, Toulouse, France) for 3 h to block TLR4 or for 1 min with 3 µmol L⁻¹ HC-030031 (Tocris Bioscience, Bristol, UK) to block TRPA1 signalling (Kunkler *et al.* 2011, Meseguer *et al.* 2014, Tse *et al.* 2014), and the resultant calcium flux of cells was recorded as described above.

Calcitonin gene-related peptide (CGRP) release assay

Experiments were performed at 37 °C in modified HBSS (Invitrogen) (10.9 mmol L⁻¹ HEPES, 4.2 mmol L⁻¹ NaHCO₃, 10 mmol L⁻¹ dextrose, and 0.1% BSA in 1 × HBSS). After two initial washes in HBSS, a 15 min baseline sample was collected. The cells were

then exposed to either HBSS alone or HBSS containing *P. gingivalis* LPS (20 µg mL⁻¹) for 15 min. Where indicated, inhibitors of TLR4, TRPA1 and TRPV1 were administered at 1 µmol L⁻¹ CLI-095 for 3 h, 3 µmol L⁻¹ HC-030031 for 2 min or 10 µmol L⁻¹ capsazepine for 2 min prior to the exposure to LPS, as described by previous investigators (Eid *et al.* 2008, Henrich & Buckler 2009, Tse *et al.* 2014). The supernatants were collected and assayed by a competitive enzyme-linked immunosorbent assay (ELISA) for CGRP. Briefly, the ELISA was performed by coating microtitre plate wells with 1 : 1000 anti-CGRP mouse monoclonal capture antibody (Sigma) in 0.05 mol L⁻¹ carbonate buffer (pH 9.5) overnight at 4 °C to immobilize the capture antibody to the wells. Plates were then washed three times with 0.05% PBS-Tween and blocked with 5% BSA in 0.05% PBS-Tween (pH 7.4) for 3 h at room temperature. After washing, 50 µL of standard CGRP or samples was added to the plate, together with 50 µL of 2 ng mL⁻¹ biotin-conjugated CGRP (Biorbyt Ltd, Cambridge, UK), and incubated for 1 h at 37 °C. After washing six times, the plates were incubated with 1:400 streptavidin–horseradish peroxidase conjugate (strep-HRP) for 45 min. Tetramethylbenzidine/hydrogen peroxide substrate solution (TMB; Sigma) was then added until colour developed (10–20 min), and the reaction was stopped by addition of 2 mol L⁻¹ HCl. Resulting colour was read using a spectrophotometer (Infinite 200 PRO; Tecan, Reading, UK) at 450 nm. Unknown CGRP concentrations were calculated by interpolation from a standard curve.

NF-κB translocation

Following application of LPS or PBS for 15 min, cells were washed twice with PBS and then fixed in 4% paraformaldehyde (PFA) for 40 min. To block non-specific antibody binding, coverslips were incubated in PBS containing 0.5% Triton X-100 (PBST) and 10% normal donkey serum (NDS) for 1 h at room temperature. The coverslips were then incubated in mouse anti-β-tubulin III monoclonal antibody (1 : 1000, Covance, Harrogate, UK), and rabbit anti-NF-κB polyclonal antibody (1 : 250, ThermoFisher Scientific, Paisley, UK) diluted in PBST containing 1% NDS, at room temperature for 1 h. Cultures were washed twice in PBST for 15 min and incubated with Cy3-conjugated donkey anti-rabbit, and FITC-conjugated donkey anti-mouse secondary antibodies (1 : 400, Jackson ImmunoResearch, West Grove, PA, USA) diluted in PBST containing 1% NDS, at room

temperature for 1 h. Finally, the cultures were washed twice in PBS for 15 min and slides were mounted with Vectashield containing DAPI (Vector Laboratories, Burlingame, CA, USA) and visualized by fluorescent microscopy. Immunohistochemistry controls for β -tubulin III and NF- κ B were performed by omitting the primary antibodies.

NF- κ B nuclear translocation was determined by immunohistochemistry and co-localizing with DAPI and β -tubulin III, markers of DNA and neurons, respectively. The translocation of NF- κ B could be determined by analysing the proportion of NF- κ B fluorescent label in the nuclear and cytoplasmic compartments. Positive nuclear translocations were quantified by measuring the relative fluorescence intensity of NF- κ B staining at each intracellular location. DAPI nuclear staining was used to confirm the location of the nuclei. The translocation of NF- κ B was counted as positive when the nuc/cyt ratio was >1 . Neurons and non-neuronal cells positive for NF- κ B nuclear translocation were quantified.

Statistical analysis

Data were analysed by one-way ANOVA, and individual groups were compared by Bonferroni's *post hoc* test using GraphPad Prism 6 (GraphPad Software, San Diego, CA, USA). The statistical significance was set at $P < 0.05$.

Results

Porphyromonas gingivalis LPS can rapidly activate trigeminal ganglion neurons and non-neuronal cells

The responses of trigeminal neurons to *P. gingivalis* LPS, cinnamaldehyde (CA), capsaicin (CAP) and high potassium were assessed using the fluorescent calcium indicator, Cal-520 AM (Fig. 1). A total of 90 neurons were assessed from three to four coverslips per experiment, and responses were examined from three independent experiments. *P. gingivalis* LPS was found to directly activate approximately 27% of the trigeminal neurons present (24/90). Our study also demonstrated that approximately 13% and 33% of mouse TG neurons responded to CA and CAP, respectively. Application of high potassium was performed to assess neuron viability and to help distinguish neuronal from non-neuronal cells (Fig. 1g,h).

Neuronal cells show a very clear rapid rising and transient response to KCl resulting from depolarization and action potential-evoked calcium signals, the

amplitude and duration of calcium transients being determined by the contribution of different ion channels contributing to the action potential (e.g. Mohammed *et al.* 2017); in contrast, non-neuronal cells do not fire action potentials and where calcium signals are observed, these are notably slower and are likely related to transporters. As shown, neuronal cells also show clear rapid responses to cinnamaldehyde or capsaicin, whereas non-neuronal cells show no or only very weak responses to these compounds. As well as differences in the calcium signalling, the morphology of cells can also be used as a discriminator, as indicated in Fig. 1a. Note the non-neuronal cells are much smaller than neurons and lack the extensive thin branching neurites, characteristic of neurons.

Approximately 40% of the non-neuronal cells present, such as satellite glial cells and Schwann cells, also responded to *P. gingivalis* LPS (representative data are shown in Fig. 1). This suggests that neuronal supporting cells can form part of the responsive population to *P. gingivalis* LPS in the peripheral nervous system (see Fig. 1h).

Calcium influx activated by LPS can be abolished by TRPA1 antagonist and modified by TLR4 antagonist

Whether *P. gingivalis* LPS triggered calcium influx via TLRs or the TRPs was then investigated. For this, TLR4 and TRPA1 were functionally blocked by pre-treating TG neurons with either CLI-095 or HC-030031, respectively. Responses were examined from three independent experiments. A total of 72 neurons in the presence of HC-030031 and 63 neurons in the presence of CLI-095 were assessed from three to four coverslips per experiment. The TRPA1 antagonist ($3 \mu\text{mol L}^{-1}$ HC-030031) completely abolished the neuronal response to *P. gingivalis* LPS (0/72), whereas pre-treatment of neurons with a specific TLR-4 signalling inhibitor ($1 \mu\text{mol L}^{-1}$ CLI-095) reduced but did not completely inhibit the response. The percentage of neurons responding to LPS in the presence of CLI-095 was reduced to less than 5%, suggesting a contributory role for the TLR4 receptor. Representative, individual traces are shown in Fig. 2.

Neuropeptide (CGRP) release by *P. gingivalis* LPS can be prevented by TLR4, TRPA1 and TRPV1 antagonists

Lipopolysaccharide (LPS) is thought to be a mediator of neurogenic inflammation (Hou *et al.* 2003,




Figure 1 Analysis of intracellular calcium in primary trigeminal cell culture. (a) Phase-contrast and (b–f) time-lapse calcium imaging in trigeminal ganglion (TG) neurons (white arrow) and non-neuronal cells (white arrowhead) loaded with Cal-520 AM, during application of *Porphyromonas gingivalis* lipopolysaccharide (LPS), cinnamaldehyde (CA), capsaicin (CAP) and high potassium chloride (KCl). The high potassium application is to assess neuron viability and to help distinguish neuronal from non-neuronal cells. (g,h) Graphs show traces of the transient change in calcium in individual cells over time after application of LPS and the agonists in neurons (g), and non-neuronal cells (h); data plotted are in arbitrary fluorescent units.

Diogenes *et al.* 2011, Ferraz *et al.* 2011) initiated by neuropeptide release. Whether *P. gingivalis* LPS could mediate neuropeptide release from the TG neurons *in vitro* was therefore determined.

TG cultures were either stimulated with vehicle, *P. gingivalis* LPS (20 μg mL⁻¹) or capsaicin (1 μmol L⁻¹) for 15 min at 37 °C and the resulting levels of CGRP released into the culture medium determined by ELISA. *P. gingivalis* LPS significantly

increased CGRP release from TG neurons (Fig. 3, $P < 0.001$) as did treatment with the positive control, 1 μmol L⁻¹ capsaicin (Fig. 3).

Since the calcium influx in response to LPS could be abolished by the TRPA1 blocker, and partially removed by TLR4 signalling inhibition, whether TLR4, TRPA1 and TRPV1 inhibitors could prevent CGRP secretion in response to LPS was explored. CGRP release experiments were again performed by




Figure 2 The effects of *Porphyromonas gingivalis* lipopolysaccharide (LPS) after TRPA1 and TLR4 blocking. Representative samples of responses to ultrapure *P. gingivalis* LPS in trigeminal ganglion (TG) neurons following pre-incubation with (a) no block, (b) 1 $\mu\text{mol L}^{-1}$ CLI-095 [TLR4 block] and (c) 3 $\mu\text{mol L}^{-1}$ HC-030031 [TRPA1 block]; data plotted are in arbitrary fluorescent units. Note that panel a is identical to the image in Fig. 1g and is included here for ease of comparison with panels b and c. (d) shows the percentage of LPS-responding neurons in three groups; no block, TLR4 block and TRPA1 block.

exposing TG cultures to LPS following pre-treatment with 3 $\mu\text{mol L}^{-1}$ HC-030031, 1 $\mu\text{mol L}^{-1}$ CLI-095 or 1 $\mu\text{mol L}^{-1}$ capsazepine. CGRP release from TG cultures in the presence of *P. gingivalis* LPS (20 $\mu\text{g mL}^{-1}$) was inhibited by all the blocking agents to the level obtained in vehicle-only stimulated samples (Fig. 4, $P < 0.001$).

Porphyromonas gingivalis LPS induces NF- κ B nuclear translocation in TG neurons, which can be modified by TLR4 antagonist

Porphyromonas gingivalis LPS is known to activate nuclear translocation of NF- κ B in a range of cell types (Hashimoto *et al.* 2004, Herath *et al.* 2013).

Consequently, whether LPS induces NF- κ B nuclear translocation in neurons and/or the support cells present in the TG cultures was determined.

Nuclear translocation of NF- κ B was considered positive when the nuclear/cytoplasm ratio (nuc/cyt ratio) of NF- κ B in neurons was >1 , calculated using the methods of Hunot *et al.* (1997) and Kuiken *et al.* (2012). Immunofluorescence revealed that *P. gingivalis* LPS stimulation resulted in positive nuclear translocation of NF- κ B in TG neurons and non-neuronal cells over a period of 30 min (Fig. 5). The percentage of neuronal cells positive for NF- κ B translocation was significantly greater in LPS-treated cultures compared with control cultures (62% vs. 18%, respectively; $P < 0.01$). The percentage of NF- κ B translocation-




Figure 3 Calcitonin gene-related peptide (CGRP) levels in lipopolysaccharide (LPS)- and capsaicin-stimulated trigeminal ganglion (TG) neurons. The secreted CGRP response to *Porphyromonas gingivalis* LPS or capsaicin was determined by the incubation of TG cells for 15 min with ultrapure *P. gingivalis* LPS ($20 \mu\text{g mL}^{-1}$) or capsaicin ($1 \mu\text{mol L}^{-1}$) at 37°C in 5% CO_2 . Measurement of CGRP levels was performed using enzyme-linked immunosorbent assay. Data are the mean \pm SD from three independent experiments using cells cultured from six ganglia collected from three different animals; each independent experiment assessed responses from four coverslips. Error bars are \pm SD; *** $P < 0.001$.

positive non-neuronal cells was also significantly greater following LPS treatment (67% vs. 37% in stimulated and control cultures, respectively; $P < 0.05$).

Further work showed that both neuronal and non-neuronal nuclear translocation of NF- κ B in response to LPS could be inhibited by blocking TLR4 signalling ($1 \mu\text{mol L}^{-1}$ CLI-095), but not by blocking TRPA1 ($3 \mu\text{mol L}^{-1}$ HC-030031) or TRPV1 signalling ($10 \mu\text{mol L}^{-1}$ capsazepine). The percentage of neurons showing nuclear translocation of NF- κ B in the presence of $1 \mu\text{mol L}^{-1}$ CLI-095 inhibitor along with LPS decreased significantly from 65% to approximately 30%, and in non-neuronal cells it decreased from 65% to approximately 20% (Fig. 6; $P < 0.05$).

Discussion

This study reports for the first time that *P. gingivalis* LPS directly activates TLR4 and TRPA1 receptors in trigeminal sensory neurons and non-neuronal support cells. The study also shows that this activation leads to release of the neuropeptide calcitonin gene-related peptide (CGRP) and activation of the NF- κ B signalling pathway.

The rationale for the study comes from evidence that some bacterial pathogens can directly stimulate a




Figure 4 Calcitonin gene-related peptide (CGRP) levels in trigeminal ganglion (TG) neuron cultures in response to lipopolysaccharide (LPS) in the presence of specific agonist inhibitors. The secreted CGRP response to *Porphyromonas gingivalis* LPS or LPS in the presence of TLR4, TRPA1 and TRPV1 blockers was determined by incubation of TG cells for 15 min with ultrapure *P. gingivalis* LPS ($20 \mu\text{g mL}^{-1}$), or $20 \mu\text{g mL}^{-1}$ ultrapure *P. gingivalis* LPS with $1 \mu\text{mol L}^{-1}$ CLI-095 [TLR4 block], $3 \mu\text{mol L}^{-1}$ HC-030031 [TRPA1 block] or $1 \mu\text{mol L}^{-1}$ capsazepine [TRPV1 block] at 37°C in 5% CO_2 . Measurement of CGRP levels was performed using enzyme-linked immunosorbent assay. Data presented are the means from two independent experiments using cells cultured from four ganglia collected from two different animals; each independent experiment assessed responses from four coverslips. Error bars are \pm SD; *** $P < 0.001$.

nociceptor neuronal response (Chiu *et al.* 2013). There is relatively little information, however, about the neuronal response to oral Gram-negative bacteria and since *P. gingivalis* is commonly reported in association with symptomatic pulpitis (Rôças *et al.* 2002, Jacinto *et al.* 2006), whether *P. gingivalis* LPS could directly activate sensory neurons was investigated. For this, an *in vitro* culture model of TG neurons was developed. It was not possible to obtain a culture of neurons free from support cells (glial and Schwann cells) because they are essential for survival of the neurons *in vitro*. However, it was possible to differentiate the responses of neurons from those of non-neuronal cells to various agonists by a combination of calcium imaging and immunofluorescence.

There are a number of receptors that could be involved in the detection and transduction of dental pain following pulpal infection. Some of these involve responses to mediators released as part of the inflammatory response (Igwe 2003, Gonzalez-Rey *et al.* 2010), but there are two categories of cellular receptors that could potentially interact with Gram-




Figure 5 Photomicrographs of p65 NF- κ B nuclear translocation in *Porphyromonas gingivalis* LPS-treated TG neurons and non-neuronal cells. The TG cultures were (a) left untreated or (b) stimulated with *P. gingivalis* LPS for 30 min. Cells were permeabilized with 0.1% Triton X-100 and subsequently stained with primary antibodies against anti-p65 NF- κ B and appropriate secondary antibodies. The β -tubulin III and p65 NF- κ B control samples showed no positive labelling (c and d, respectively). The β -tubulin III is shown in green (a1,b1) and p65 NF- κ B appears in red (a2,b2), and nuclear staining with DAPI appears blue (a3, b3). The merged images (a4,b4) show the combined p65 NF- κ B, β -tubulin III and DAPI staining. Scale bars = 70 μ m. Localization of p65 NF- κ B is indicated in neurons by yellow arrows and non-neuronal cells by white arrows. The translocation of NF- κ B was counted as positive when nuc/cyt ratio was >1. Quantitative analysis of neuronal nuclear translocation stimulated by *P. gingivalis* LPS is shown in figure (e). Neurons and non-neuronal cells positive for NF- κ B nuclear translocation were quantified. Data are the mean \pm SD from three independent experiments. Error bars are \pm SD; * P < 0.05 and ** P < 0.01.

negative bacterial LPS. First, TLRs are thought to be primary key participants in inflammatory responses; secondly, TRP receptors have emerged as having critically important roles in peripheral sensitization (Rang *et al.* 1991, Chen *et al.* 2011, Chung *et al.* 2011, Tóth *et al.* 2011). TLRs have been reported to be present on trigeminal and DRG neurons (Wadachi & Hargreaves 2006, Ochoa-Cortes *et al.* 2010, Qi *et al.* 2011) and in the present study ultrapure *P. gingivalis* LPS directly activated more than 20% of TG neurons in the cultures, with a resultant increase in intracellular calcium. This is the first time that *P. gingivalis*

LPS has been shown to directly activate TG neurons *in vitro*. Also, the neuronal response could be partially blocked by the specific TLR4 inhibitor, CLI-095, with the proportion of responding cells reducing from around 20% to only 5%.

LPS of Gram-negative oral anaerobic bacteria is known to elicit different host responses from those observed with the classic enterobacterial endotoxin (Darveau & Hancock 1983, Dixon & Darveau 2005, Herath *et al.* 2013). Indeed, *P. gingivalis* LPS has been considered to have relatively low biological activity because of its low phosphate content, altered length




Figure 6 Percentages of p65 NF-κB nuclear translocation in (a) *Porphyromonas gingivalis* LPS-treated TG neurons and (b) non-neuronal cells in the presence of TLR4, TRPA1 and TRPV1 signalling inhibitors. Positive nuclear translocations were quantified by measuring the fluorescence intensity at each intracellular location. The translocation of NF-κB was counted as positive when nuc/cyt ratio was >1. Neurons and non-neuronal cells positive for NF-κB nuclear translocation were quantified. Data are the mean from two independent experiments. Error bars are ±SD; * $P < 0.05$.

and positions of fatty acids in the lipid A moiety, compared with *E. coli* LPS (Takada & Kotani 1989). Also, Ogawa & Uchida (1996) have described differential cytokine production by mononuclear cells in response to *P. gingivalis* LPS. Notwithstanding these differences, *P. gingivalis* LPS was found to induce inflammatory responses in LPS non-responder mice (TLR4 knockout mice) (Kirikae *et al.* 1999), which suggests a different molecular recognition pathway is involved from that for *E. coli* LPS.

The other category of potential receptors is the transient receptor potential channels (TRPs). Numerous studies have shown that TRP channels are a broad group of ion channels located on nociceptors; TRPA1 and TRPV1 are thought to be involved in responses to bacterial infection (Chung *et al.* 2011, Gibbs *et al.* 2011, Huang *et al.* 2012). TRPV1 is expressed in small and medium diameter nerve fibres throughout the human tooth pulp, which comprise about 20%–35% of the TG neurons (Chung *et al.* 2011, Gibbs *et al.* 2011). The data in the present study support this, with 33% of TG neurons functionally responding to the TRPV1 agonist, capsaicin. As well as TRPV1, the receptor TRPA1 could also act as a sensor since it is known to respond to a broad spectrum of endogenous compounds and irritants. However, there has been a report that TRPA1 expression in the trigeminal ganglion is distributed in only a small population of TG neurons (6%–10%; Huang *et al.* 2012). The present findings concur with this as it was noted the proportion of neurons in the cultures responding to the TRPA1 agonist CA was less than 15%.

The present study demonstrates that *P. gingivalis* LPS activates trigeminal neurons through TRPA1-dependent pathways, since the response was completely blocked by the TRPA1 inhibitor, HC-030031. This is consistent with the finding of Meseguer *et al.* (2014) that *E. coli* LPS exerted a fast excitatory action on TG neurons via the TRPA1 receptor. However, the neuronal activation by *P. gingivalis* LPS could occur via both TRPA1 and TLR4. This difference could be explained by the different properties of *P. gingivalis* LPS and *E. coli* LPS (Coats *et al.* 2005, Herath *et al.* 2013). It is also possible that activation of TLR4 sensitizes TRPA1. As indicated by the present and previous studies (Calvo-Rodríguez *et al.* 2017), TLR4 activation of neurons can increase levels of intracellular calcium; this rise in calcium could then act on TRPA1 to create a positive reinforcement of calcium signalling. Previous studies have reported that a rise in calcium can sensitize TRPA1 (Zurborg *et al.* 2007).

Since TRPA1 has been shown to control vascularity in neurogenic inflammation (Kunkler *et al.* 2011), levels of CGRP release in response to *P. gingivalis* LPS were measured. The present study demonstrated that a relatively high concentration of *P. gingivalis* LPS ($20 \mu\text{g mL}^{-1}$) could enhance neuronal activation and produce a significant increase in CGRP release ($P < 0.001$). In contrast, Ferraz *et al.* (2011) reported that a low concentration ($2 \mu\text{g mL}^{-1}$) of *P. gingivalis* LPS failed to induce CGRP release. However, since only supraphysiological levels of LPS (here $20 \mu\text{g mL}^{-1}$) triggered a neuronal response, it may indicate a different processing mechanism from that of the classical LPS receptor complex of TLR4 with




Figure 7 Proposed model for LPS stimulation of TG neurons. LPS can activate TG neurons via TRPA1 and TLR4; this activation can be blocked by their specific antagonists HC-030031 and CLI-095, respectively. The neuronal activation can result in both rapid responses (e.g. CGRP release) and delayed responses (e.g. NF- κ B nuclear translocation).

MD2 and CD14 co-factors. With supraphysiological LPS concentrations, LPS signalling could occur via CD14-independent mechanisms (Ulevitch & Tobias 1995) and so the mechanisms revealed in this study may not completely represent the canonical pathway of TLR4-LPS interaction. Moreover, the effect of LPS on the release of CGRP could be reversed by not only TRPA1 or TLR4 signalling blockers, but also by a TRPV1 blocker. This finding may support the notion that *P. gingivalis* LPS somehow induces a TRPV1-dependent release of CGRP, which would be consistent with the findings of Diogenes *et al.* (2011). These workers showed that TLR4 stimulation could enhance TRPV1 activation, leading to the release of CGRP. Even though there has been very limited evidence to show how LPS modulates peripheral sensitization, it has been thought that Ca^{2+} permeation through TRPV1 leads to exocytosis, which underlies CGRP and substance P release (Henrich & Buckler 2009) and which can account for a rapid neuronal response. However, it is also possible that CGRP release may also be influenced by the triggering of PKC/PLC pathways (Vellani *et al.* 2010), as a latent consequence. Taken together, the present data cannot rule out the possibility that LPS sensitivity of TG neurons requires TRPV1-mediated release of CGRP.

In order to have more prolonged effects on neurons, LPS would be expected to result in activation of the NF- κ B pathway because this is the major transcription factor responsible for modulating gene expression in response to LPS (Arias-Salvatierra *et al.*

2011). Also, both TRPA1 and TLR4 mediate the production of inflammatory mediators through the NF- κ B pathway and this pathway is thought to be related to voltage-gated channel over-expression (e.g. Ca^{2+} and Na^{+} channels) and possibly inducing allodynia, thermal hyperplasia or chronic pain (Huang *et al.* 2006). Thus, the observed secretion of CGRP may relate to involvement of both these receptors (TRPA1 and TLR4), though CGRP secretion itself appears to involve the TRPV1-dependent pathway. These data lead us to propose a possible model for LPS stimulation of TG cells as shown in Fig. 7.

Several previous reports have suggested that satellite glial cells could be a possible therapeutic target for modulation of pain (Watkins & Maier 2002, Takeda *et al.* 2009). This is because neuronal excitability could be developed and maintained through the sensory neuron along with the surrounding satellite glial cells acting as a functional unit (Takeda *et al.* 2009, Villa *et al.* 2010, Kushnir *et al.* 2011, Poulsen *et al.* 2014, Costa & Moreira Neto 2015, Hanani 2015). In the present study, TG neuronal supporting cells were directly activated by *P. gingivalis* LPS (e.g. using calcium influx and NF- κ B nuclear translocation measurements). This is consistent with the finding of Yoon *et al.* (2012) who reported that *E. coli* LPS could increase expression of satellite cell markers in DRG. Taken together, the activation of neuronal supporting cells, such as satellite glial cells and/or Schwann cells, may be a crucial element in generation and maintenance of orofacial

pain; however, studies are currently being carried out to further characterize their contribution.

Conclusion

This study sheds light on the mechanisms involved in the pain pathways that can be directly activated by bacterial infection of the pulp. Contact between oral Gram-negative bacteria, such as *P. gingivalis*, and TG neuronal and non-neuronal cells can involve their direct activation through a combination of TRPA1, TLR4 and TRPV1 receptors. This results in NF- κ B nuclear translocation and release of the neuropeptide CGRP.

Acknowledgements

This study was supported by a PhD Studentship to A Kaewpitak from the Royal Thai Government, to whom we are grateful.

Conflict of interest

The authors have stated explicitly that there are no conflicts of interest in connection with this article.

References

- Arias-Salvatierra D, Silbergeld EK, Acosta-Saavedra LC, Calderon-Aranda ES (2011) Role of nitric oxide produced by iNOS through NF- κ B pathway in migration of cerebellar granule neurons induced by Lipopolysaccharide. *Cellular Signalling* **23**, 425–35.
- Bottenstein JE, Sato GH (1979) Growth of a rat neuroblastoma cell line in serum-free supplemented medium. *Proceedings of the National Academy of Sciences of the United States of America* **76**, 514–7.
- Calvo-Rodríguez M, de la Fuente C, García-Durillo M, García-Rodríguez C, Villalobos C, Núñez L (2017) Aging and amyloid β oligomers enhance TLR4 expression, LPS-induced Ca²⁺ responses, and neuron cell death in cultured rat hippocampal neurons. *Journal of Neuroinflammation* **14**, 24.
- Chen Y, Yang C, Wang ZJ (2011) Proteinase-activated receptor 2 sensitizes transient receptor potential vanilloid 1, transient receptor potential vanilloid 4, and transient receptor potential ankyrin 1 in paclitaxel-induced neuropathic pain. *Neuroscience* **193**, 440–51.
- Chiu IM, Heesters BA, Ghasemlou N et al. (2013) Bacteria activate sensory neurons that modulate pain and inflammation. *Nature* **501**, 52–7.
- Chung M-K, Lee J, Duraes G, Ro JY (2011) Lipopolysaccharide-induced pulpitis up-regulates TRPV1 in trigeminal ganglia. *Journal of Dental Research* **90**, 1103–7.
- Coats SR, Pham TT, Bainbridge BW, Reife RA, Darveau RP (2005) MD-2 mediates the ability of tetra-acylated and penta-acylated lipopolysaccharides to antagonize *Escherichia coli* lipopolysaccharide at the TLR4 signaling complex. *Journal of Immunology* **175**, 4490–8.
- Costa FAL, Moreira Neto FL (2015) Satellite glial cells in sensory ganglia: its role in pain. *Revista Brasileira de Anestesiologia* **65**, 73–81.
- Darveau RP, Hancock RE (1983) Procedure for isolation of bacterial lipopolysaccharides from both smooth and rough *Pseudomonas aeruginosa* and *Salmonella typhimurium* strains. *Journal of Bacteriology* **155**, 831–8.
- Diogenes A, Ferraz CCR, Akopian AN, Henry MA, Hargreaves KM (2011) LPS sensitizes TRPV1 via activation of TLR4 in trigeminal sensory neurons. *Journal of Dental Research* **90**, 759–64.
- Dixon DR, Darveau RP (2005) Lipopolysaccharide heterogeneity: innate host responses to bacterial modification of lipid A structure. *Journal of Dental Research* **84**, 584–95.
- Eid SR, Crown ED, Moore EL et al. (2008) HC-030031, a TRPA1 selective antagonist, attenuates inflammatory- and neuropathy-induced mechanical hypersensitivity. *Molecular Pain* **4**, 48.
- Ferraz CCR, Henry MA, Hargreaves KM, Diogenes A (2011) Lipopolysaccharide from *Porphyromonas gingivalis* sensitizes capsaicin-sensitive nociceptors. *Journal of Endodontics* **37**, 45–8.
- Geppetti P, Nassini R, Materazzi S, Benemei S (2008) The concept of neurogenic inflammation. *BJU International* **101**(Suppl 3), 2–6.
- Gibbs JL, Melnyk JL, Basbaum AI (2011) Differential TRPV1 and TRPV2 channel expression in dental pulp. *Journal of Dental Research* **90**, 765–70.
- Gonzalez-Rey E, Ganea D, Delgado M (2010) Neuropeptides: keeping the balance between pathogen immunity and immune tolerance. *Current Opinion in Pharmacology* **10**, 473–81.
- Hanani M (2015) Role of satellite glial cells in gastrointestinal pain. *Frontiers in Cellular Neuroscience* **9**, 412.
- Hashimoto M, Asai Y, Ogawa T (2004) Separation and structural analysis of lipoprotein in a lipopolysaccharide preparation from *Porphyromonas gingivalis*. *International Immunology* **16**, 1431–7.
- Henrich M, Buckler KJ (2009) Acid-evoked Ca²⁺ signalling in rat sensory neurones: effects of anoxia and aglycaemia. *Pflügers Archiv: European Journal of Physiology* **459**, 159–81.
- Herath TDK, Darveau RP, Seneviratne CJ, Wang C-Y, Wang Y, Jin L (2013) Tetra- and penta-acylated lipid A structures of *Porphyromonas gingivalis* LPS differentially activate TLR4-mediated NF- κ B signal transduction cascade and immuno-inflammatory response in human gingival fibroblasts. *PLoS ONE* **8**, e58496.
- Hirao K, Yumoto H, Takahashi K, Mukai K, Nakanishi T, Matsuo T (2009) Roles of TLR2, TLR4, NOD2, and NOD1 in pulp fibroblasts. *Journal of Dental Research* **88**, 762–7.

- Hou L, Li W, Wang X (2003) Mechanism of interleukin-1 β -induced calcitonin gene-related peptide production from dorsal root ganglion neurons of neonatal rats. *Journal of Neuroscience Research* **73**, 188–97.
- Huang J, Zhang X, McNaughton PA (2006) Inflammatory pain: the cellular basis of heat hyperalgesia. *Current Neuropharmacology* **4**, 197–206.
- Huang D, Li S, Dhaka A, Story GM, Cao Y-Q (2012) Expression of the transient receptor potential channels TRPV1, TRPA1 and TRPM8 in mouse trigeminal primary afferent neurons innervating the dura. *Molecular Pain* **8**, 66.
- Hunot S, Brugg B, Ricard D et al. (1997) Nuclear translocation of NF- κ B is increased in dopaminergic neurons of patients with Parkinson disease. *Proceedings of the National Academy of Sciences of the United States of America* **94**, 7531–6.
- Igwe OJ (2003) c-Src kinase activation regulates prepro-tachykinin gene expression and substance P secretion in rat sensory ganglia. *European Journal of Neuroscience* **18**, 1719–30.
- Jacinto RC, Gomes BPFA, Shah HN, Ferraz CC, Zaia AA, Souza-Filho FJ (2006) Incidence and antimicrobial susceptibility of *Porphyromonas gingivalis* isolated from mixed endodontic infections. *International Endodontic Journal* **39**, 62–70.
- Jiang H-W, Zhang W, Ren B-P, Zeng J-F, Ling J-Q (2006) Expression of toll like receptor 4 in normal human odontoblasts and dental pulp tissue. *Journal of Endodontics* **32**, 747–51.
- Kirikae T, Nitta T, Kirikae F et al. (1999) Lipopolysaccharides (LPS) of oral black-pigmented bacteria induce tumor necrosis factor production by LPS-refractory C3H/HeJ macrophages in a way different from that of *Salmonella* LPS. *Infection and Immunity* **67**, 1736–42.
- Kuiken HJ, Egan DA, Laman H, Bernards R, Beijersbergen RL, Dirac AM (2012) Identification of F-box only protein 7 as a negative regulator of NF- κ B signalling. *Journal of Cellular and Molecular Medicine* **16**, 2140–9.
- Kunkler PE, Ballard CJ, Oxford GS, Hurley JH (2011) TRPA1 receptors mediate environmental irritant-induced meningeal vasodilatation. *Pain* **152**, 38–44.
- Kushnir R, Cherkas PS, Hanani M (2011) Peripheral inflammation upregulates P2X receptor expression in satellite glial cells of mouse trigeminal ganglia: a calcium imaging study. *Neuropharmacology* **61**, 739–46.
- Meseguer V, Alpizar YA, Luis E et al. (2014) TRPA1 channels mediate acute neurogenic inflammation and pain produced by bacterial endotoxins. *Nature Communications* **5**, 3125.
- Mohammed ZA, Doran C, Grundy D, Nassar MA (2017) Veratridine produces distinct calcium response profiles in mouse dorsal root ganglia neurons. *Scientific Reports* **7**, 45221.
- Ochoa-Cortes F, Ramos-Lomas T, Miranda-Morales M et al. (2010) Bacterial cell products signal to mouse colonic nociceptive dorsal root ganglia neurons. *American Journal of Physiology: Gastrointestinal and Liver Physiology* **299**, G723–32.
- Ogawa T, Uchida H (1996) Differential induction of IL-1 beta and IL-6 production by the nontoxic lipid A from *Porphyromonas gingivalis* in comparison with synthetic *Escherichia coli* lipid A in human peripheral blood mononuclear cells. *FEMS Immunology and Medical Microbiology* **14**, 1–13.
- Poulsen JN, Larsen F, Duroux M, Gazerani P (2014) Primary culture of trigeminal satellite glial cells: a cell-based platform to study morphology and function of peripheral glia. *International Journal of Physiology, Pathophysiology and Pharmacology* **6**, 1–12.
- Qi J, Buzas K, Fan H et al. (2011) Painful pathways induced by TLR stimulation of dorsal root ganglion neurons. *Journal of Immunology* **186**, 6417–26.
- Rang HP, Bevan S, Dray A (1991) Chemical activation of nociceptive peripheral neurones. *British Medical Bulletin* **47**, 534–48.
- Rôças IN, Siqueira JF, Andrade AFB, de Uzeda M (2002) Identification of selected putative oral pathogens in primary root canal infections associated with symptoms. *Anaerobe* **8**, 200–8.
- Santoni G, Cardinali C, Morelli MB, Santoni M, Nabissi M, Amantini C (2015) Danger- and pathogen-associated molecular patterns recognition by pattern-recognition receptors and ion channels of the transient receptor potential family triggers the inflammasome activation in immune cells and sensory neurons. *Journal of Neuroinflammation* **12**, 21.
- Takada H, Kotani S (1989) Structural requirements of lipid A for endotoxicity and other biological activities. *Critical Reviews in Microbiology* **16**, 477–523.
- Takeda M, Takahashi M, Matsumoto S (2009) Contribution of the activation of satellite glia in sensory ganglia to pathological pain. *Neuroscience and Biobehavioral Reviews* **33**, 784–92.
- Tóth DM, Szőke E, Bölcskei K et al. (2011) Nociception, neurogenic inflammation and thermoregulation in TRPV1 knockdown transgenic mice. *Cellular and Molecular Life Sciences* **68**, 2589–601.
- Tse K-H, Chow KBS, Leung WK, Wong YH, Wise H (2014) Lipopolysaccharide differentially modulates expression of cytokines and cyclooxygenases in dorsal root ganglion cells via Toll-like receptor-4 dependent pathways. *Neuroscience* **267**, 241–51.
- Ulevitch RJ, Tobias PS (1995) Receptor-dependent mechanisms of cell stimulation by bacterial endotoxin. *Annual Review of Immunology* **13**, 437–57.
- Veldhuis NA, Bunnett NW (2013) Proteolytic regulation of TRP channels: implications for pain and neurogenic inflammation. *Proceedings of the Australian Physiological Society* **44**, 101–3.
- Vellani V, Kinsey AM, Prandini M et al. (2010) Protease activated receptors 1 and 4 sensitize TRPV1 in nociceptive neurones. *Molecular Pain* **6**, 61.

- Villa G, Fumagalli M, Verderio C, Abbracchio MP, Ceruti S (2010) Expression and contribution of satellite glial cells purinoceptors to pain transmission in sensory ganglia: an update. *Neuron Glia Biology* **6**, 31–42.
- Wadachi R, Hargreaves KM (2006) Trigeminal nociceptors express TLR-4 and CD14: a mechanism for pain due to infection. *Journal of Dental Research* **85**, 49–53.
- Watkins LR, Maier SF (2002) Beyond neurons: evidence that immune and glial cells contribute to pathological pain states. *Physiological Reviews* **82**, 981–1011.
- Yoon S-Y, Patel D, Dougherty PM (2012) Minocycline blocks lipopolysaccharide induced hyperalgesia by suppression of microglia but not astrocytes. *Neuroscience* **221**, 214–24.
- Zurborg S, Yurgionas B, Jira JA, Caspani O, Heppenstall PA (2007) Direct activation of the ion channel TRPA1 by Ca^{2+} . *Nature Neuroscience* **10**, 277–9.