



This is a repository copy of *Histological characterization of interneurons in Alzheimer's disease reveals a loss of somatostatin interneurons in the temporal cortex.*

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/159357/>

Version: Accepted Version

Article:

Waller, R. orcid.org/0000-0001-5815-8829, Mandeya, M., Viney, E. et al. (2 more authors) (2020) Histological characterization of interneurons in Alzheimer's disease reveals a loss of somatostatin interneurons in the temporal cortex. *Neuropathology*, 40 (4). pp. 336-346. ISSN 0919-6544

<https://doi.org/10.1111/neup.12649>

This is the peer reviewed version of the following article: Waller, R., Mandeya, M., Viney, E., Simpson, J.E. and Wharton, S.B. (2020), Histological characterization of interneurons in Alzheimer's disease reveals a loss of somatostatin interneurons in the temporal cortex. *Neuropathology*, which has been published in final form at <https://doi.org/10.1111/neup.12649>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Histological characterization of interneurons in Alzheimer's Disease reveals a loss of somatostatin interneurons in the temporal cortex

Rachel Waller ^a, Memory Mandeya ^a, Edward Viney ^a, Julie E. Simpson ^a, Stephen. B. Wharton ^a

^a Sheffield Institute for Translational Neuroscience, The University of Sheffield, Sheffield, United Kingdom

Corresponding author:

Dr Rachel Waller
Sheffield Institute for Translational Neuroscience
University of Sheffield
385a Glossop Road
Sheffield
S10 2HQ
Tel: +44 (0)114 222 2272
Email : r.waller@sheffield.ac.uk

Running title: Interneurons in Alzheimer's Disease

Abstract

Neuronal dysfunction and synaptic loss are major hallmarks of Alzheimer's Disease (AD) which correlate with symptom severity. Impairment of the GABAergic inhibitory interneurons, which form around 20% of the total neuronal network, may be an early event contributing to neuronal circuit dysfunction in neurodegenerative diseases. This study examined the expression of two of the main classes of inhibitory interneuron, parvalbumin (PV) and somatostatin (SST) interneurons in the temporal cortex and hippocampus of AD and control cases, using immunohistochemistry. We report significant regional variation in the number of PV and SST interneurons with a higher number identified per mm² in the temporal cortex compared to the hippocampus. Fewer SST⁺ interneurons, but not PV⁺ interneurons, were identified per mm² in the temporal cortex of AD cases compared to control subjects. Our results support regional neuroanatomical effects on selective interneuron classes in AD, and suggest that impairment of the interneuronal circuit may contribute to neuronal dysfunction and cognitive decline in AD.

Keywords

Alzheimer disease, Interneurons, Pathology, Humans, Somatostatin

1 Introduction

2 Neuronal dysfunction and synaptic loss are key features of Alzheimer's Disease (AD) which
3 correlate with symptom severity ¹. Pyramidal neurons have been the main focus of study in
4 relation to AD pathology ², with cholinesterase inhibitors commonly used for symptomatic
5 relief in AD. However, interneurons are also essential to the normal functioning of the
6 neuronal network. Forming around 20% of the total neuronal network ³, most interneurons
7 are γ -aminobutyric acid (GABA)ergic and inhibitory, also acting as a source of neuropeptides
8 that help to regulate cortical function ⁴.

9 There are three main classes of inhibitory interneuron with calcium-binding proteins and
10 neuropeptides used to distinguish between them. The different types of interneuron provide
11 GABA, a major inhibitory neurotransmitter, to different subcellular places with the most
12 common class of cortical interneurons, being the calcium-binding protein parvalbumin (PV)
13 expressing interneurons. These cells comprise of the chandelier cells which innervate the
14 axon initial segment, basket cells that target the soma of excitatory cells, and trans-laminar
15 interneurons ⁵. These interneurons are responsible for their fast-spiking activity and precise
16 inhibitory control of pyramidal cell output ⁶. Somatostatin (SST) expressing interneurons, the
17 Martinotti and non-Martinotti cells, fire more randomly and gradually to stimulus targeting
18 dendrites of excitatory neurons ⁷. SST⁺ cells release GABA that acts on a wide array of
19 GABA receptors (GABAR), alongside fast synaptic GABA_AR, the slower more persistent
20 metabotropic GABA_BRs become activated on pyramidal cells silencing connections between
21 these excitatory cells ⁸. Furthermore it has been proposed that the GABA released from SST
22 cells affects the connectivity between excitatory neurons, regardless of whether these
23 excitatory neurons are synaptically connected with SST neurons ⁹.

24 The third class of interneurons express the serotonin receptor 5HT_{3a}R as well as expressing
25 vasoactive intestinal peptide (VIP) and are distinguishable for synapsing PV⁺ and SST⁺
26 interneurons ¹⁰. Typical VIP expressing interneurons are bipolar displaying continuous
27 adapting firing properties ¹¹.

28 Cognitive symptoms associated with AD are the result of altered neural networks including
29 abnormal oscillatory rhythmic activity and network hypersynchrony that can occur many
30 years prior to clinical symptom onset ¹²⁻¹⁴. Studies have shown the involvement of
31 interneurons in regulating these neural circuits and networks to be altered in AD, suggesting
32 that interneuron dysfunction could play a role in neuronal network failure and cognitive
33 dysfunction in AD ¹⁵. The Tg8CRND8 mouse model of AD exhibits spatial memory deficits
34 and altered anxiety alongside a reduced number of interneurons. Treatment with α -

35 melanocyte stimulating hormone (α -MSH) improved spatial memory and prevented the loss
36 of SST⁺ interneurons, demonstrating its role in restoring GABAergic inhibition to improve
37 cognition in this model ¹⁶. Similarly, in the APP/PS1 mouse model of AD SST⁺ interneurons
38 decrease with aging compared to control mice, while PV⁺ interneurons were increased
39 suggesting a diverse interneuron population vulnerability that may be related to amyloid beta
40 (A β) pathology in AD ¹⁷.

41 Previous work has shown that the enzymes responsible for degrading A β are expressed in
42 interneurons ¹⁸. The presence of endothelin-converting enzyme (ECE-2) and neprilysin in
43 interneurons suggests the possibility that A β may have a role in the regulation of inhibition,
44 acting as a neuropeptide important for interneuronal function ¹⁸. Additionally, γ -aminobutyric
45 acid (GABA) receptors are altered in AD ¹⁹ with a varied pattern of GABA_B receptor R1
46 protein (GBR1) expression throughout the hippocampus of AD patients. An increased GBR1
47 expression was identified in the CA4 and CA3/2 areas, yet was rapidly reduced in the CA1
48 region with advanced AD pathology and progression of neurofibrillary tangles (NFT) prior to
49 neuronal cell death ¹⁹. These GABA receptors have been shown to be involved in the
50 inhibitory neurotransmission system that could contribute to neuronal resistance seen in the
51 initial stages of disease pathology. Thus changes in the balance between inhibitory and
52 excitatory neurotransmission are likely to contribute to AD development. The learning and
53 memory deficits that correlated with the age-dependent decline in SST⁺ interneurons in the
54 apoE4-KI mouse model of AD were rescued following the restoration of GABA signaling
55 using pentobarbital, a GABA_A receptor enhancer ⁵. Another mutated amyloid precursor
56 protein (APP) familial mouse model of AD exhibited memory deficits and a reduction in
57 GABA related proteins and GABAergic interneurons as early as 4 months ²⁰. With APP
58 known to play a role in GABAergic synaptic formations, by administering diazepam and
59 correcting the APP function an improvement in memory and also reduced A β accumulation
60 was seen. Clearly, the GABAergic deficiency caused memory deficits and contributed to A β
61 accumulation in this model ²⁰.

62 We have investigated changes in two of the main subclasses of inhibitory interneurons, the
63 PV- and SST- expressing interneuron in temporal structures of AD and age-matched control
64 brain using an immunohistochemistry approach. We hypothesized that inhibitory neurons are
65 lost in areas of the brain typically affected by pathology in AD.

66 **Materials and Methods**

67 **Human central nervous system tissue**

68 All formalin fixed paraffin embedded (FFPE) lateral temporal cortex (Brodmann areas 21/22)
69 and hippocampus tissue was obtained from the Sheffield Brain Tissue Bank (SBTB). The
70 SBTB gave full ethical approval for the use of tissue in this study (ref. 08/MRE00/103+5) as a
71 Research Tissue Bank approved by the Scotland Research Ethics Committee. A summary of
72 the cohort used in the study is provided in Table 1.

73 **Immunohistochemistry**

74 Immunohistochemistry (IHC) was performed using a standard avidin-biotin complex-horse
75 radish peroxidase (ABC-HRP) method, and visualized with diaminobenzidine (DAB) (Vector
76 Laboratories, UK). Briefly, deparaffinised 5µm sections were rehydrated and endogenous
77 peroxidase quenched by blocking the sections in 3% H₂O₂/methanol for 20 minutes at room
78 temperature (RT). Following antigen retrieval (Table 2) sections were blocked in 1.5% normal
79 serum for 30 minutes at RT. Sections were incubated in the relevant optimal antibody dilution,
80 washed thoroughly in tris buffered saline (TBS, pH7.5) and incubated in 0.5% biotinylated
81 secondary antibody for 30 minutes at RT. After thorough washing in TBS the sections were
82 incubated in ABC-HRP for 30 minutes at RT before a final wash in TBS and incubation in the
83 substrate DAB for 5 minutes at RT, sections were dehydrated, cleared in xylene and mounted
84 for image analysis. Negative controls (omission of the primary antibody and isotype controls)
85 were incubated in each run. Additional dual labelling was completed where SST stained
86 sections (using a standard ABC-HRP method as earlier) were incubated with the avidin-biotin
87 blocking kit (Vector Laboratories, UK), and incubated overnight at 4°C with anti-PV, followed
88 by the alkaline-phosphatase-conjugated ABC (Vectastain Elite kit, Vector Laboratories, UK),
89 developed with alkaline phosphatase substrate (Vector Laboratories, UK; red) and
90 counterstained with haematoxylin. Negative controls consisted of sections incubated in the
91 absence of primary antibody.

92

93 **Quantification of antibody staining and statistical analysis**

94 Immunostained sections were imaged using either a Nikon Eclipse 80i microscope (Nikon UK,
95 Kingston upon Thames) or digitally scanned under a 40x objective lens using a Nanozoomer
96 XR (Hamamatsu, Photonics Ltd., Hertfordshire, UK). Scanned sections were stored as
97 NanoZoomer Digital Pathology Image (.ndpi) files, viewed and exported using NDP.View 2.

98 Quantification of PV-specific immunoreactivity within the temporal cortex was performed by
99 capturing non-overlapping bright-field microscope images at x20 magnification in three
100 contiguous belt transects covering the total cortical thickness (Supp Fig 1). Within the
101 hippocampus, five non-contiguous images along the pyramidal layer were exported from the

102 scanned sections at x20 magnification in the CA1 region, and three images in the subiculum.
103 Non-overlapping images were exported from the scanned sections at x20 magnification
104 across the thickest region of the entorhinal cortex in two contiguous belt transects (Supp Fig
105 2). PV immunopositive cells were manually counted in each region as well as the total
106 percentage PV area immunoreactivity analysed in Analysis^D, (Olympus Biosystems,
107 Watford, UK).

108 SST⁺ cells were more infrequent than PV, and therefore quantification was performed solely
109 by manual cell counts rather than using computer-aided image analysis. Within the temporal
110 cortex, SST immunopositive cells were counted in the area of highest expression in six non-
111 overlapping images exported from the scanned sections at x20 magnification in a contiguous
112 belt. Within the CA1 and enthorinal cortex the areas of highest expression within the
113 hippocampus, SST immunopositive cells were counted in six random images exported from
114 the scanned sections at x20 magnification in both the CA1 and entorhinal cortex regions.

115 All statistical analyses were performed using IBM SPSS Statistic v24. For variation between
116 neuroanatomical regions, Friedman's Two-Way Analysis of Variance was used to compare
117 immunoreactivity in the four brain regions (temporal cortex, hippocampus CA1, subiculum and
118 entorhinal cortex) across the full cohort. Post hoc differences were assessed by Wilcoxon
119 Signed Rank Test. Statistical comparisons of quantitative data between the control and AD
120 cases was performed using Mann-Whitney U Tests.

121

122 **Results**

123 **Somatostatin (SST) interneurons were reduced in the temporal cortex of AD patients**

124 SST immunoreactivity was discretely associated with the cytoplasm of neuronal cell bodies
125 and immediate extending dendrites (Figures 1a&b), therefore for quantification total cell
126 counts were used (Table 3). In addition, immunoreactivity was located predominantly at the
127 grey matter/white matter border (Figures 1c&d). There was significant neuroanatomical
128 regional variation in the number of SST⁺ interneurons ($F=18.96$ 2df $p<0.001$) with an
129 increased number in the temporal cortex compared to CA1 ($p=0.001$) and entorhinal cortex
130 ($p=0.006$). There was a slight increase in the number of SST⁺ cells in the entorhinal cortex
131 compared to CA1 regions ($p=0.025$) (Figure 2a). Overall, there was a reduction (of
132 approximately 30%) in the number of SST⁺ interneurons in the temporal cortex of AD patients
133 compared to control cases (Mann-Whitney U test $p=0.040$), although this did not achieve
134 statistical significance if corrected for multiple testing using the Bonferroni method ($p=0.102$).
135 No significant differences in the number of SST⁺ interneurons were detected between the

136 entorhinal cortex or CA1 regions of the hippocampus of AD patients and control cases (Mann-
137 Whitney U test $p=0.382$, $p=0.673$ respectively (Figure 2b).

138 **Parvalbumin (PV) immunoreactivity of neuronal cell bodies was more pronounced in a**
139 **band like pattern of the outer layers of the cortex**

140 PV immunoreactivity was detected in the cytoplasm of neuronal cell bodies and immediate
141 extending dendrites (Figures 3a&b). For quantification both the total percentage area of PV
142 immunoreactivity and total cell count were used (Table 3). Immunoreactivity appeared higher
143 in the more outer cortical layers (I-IV) of the temporal and entorhinal cortex in a band like
144 pattern (Figure 3c). Within the hippocampus the staining pattern varied greatly with PV
145 immunoreactivity of cell bodies in CA1 being more sparsely distributed than in the temporal
146 cortex (Figure 3d), while in the subiculum the immunoreactivity appeared in clusters (Figure
147 3e). One of the AD cases showed very limited cytoplasmic cell body staining with
148 immunoreactivity restricted to the surface of neurons in a beaded string like manner,
149 suggesting a synaptic bouton labelling pattern (Figure 3f).

150 There was a significant difference in PV immunoreactive area ($F= 37.87$ 3df $p<0.001$) and
151 cell count ($F= 32.50$ 3df $p<0.001$) across all four brain regions investigated. The temporal
152 cortex had the highest total PV immunoreactivity and cell count per mm^2 compared to the
153 other three areas of the hippocampus (CA1, subiculum, entorhinal cortex) ($p<0.001$). Within
154 the hippocampus PV immunoreactivity was significantly higher in the subiculum compared to
155 CA1 ($p<0.001$) and entorhinal cortex ($p=0.001$) (Figure 4a). There was no significant
156 difference in PV immunoreactivity between the CA1 and entorhinal cortex ($p=0.485$) (Figure
157 4a).

158 For cell count, the temporal cortex had significantly more PV immunopositive neurons per mm^2
159 compared to the CA1 and entorhinal cortex ($p<0.001$) (Figure 4b). In contrast, there was no
160 significant difference in the number of PV positive neurons per mm^2 in the subiculum
161 compared to the temporal cortex ($p=0.372$), likely reflecting extensive case to case variation
162 in the immunoreactive profile (Figure 4b), Within the hippocampus the subiculum contained
163 significantly more PV immunoreactive cells per mm^2 than CA1 ($p<0.001$) and entorhinal
164 regions ($p=0.004$), with an increase number of PV positive cells per mm^2 in the entorhinal
165 regions compared to CA1 ($p=0.044$) (Figure 4b).

166 **Parvalbumin (PV) immunoreactivity was not altered in AD and did not colocalise with**
167 **somatostatin (SST) immunoreactivity**

168 There were no significant differences in total PV immunoreactive area or cell count per mm²
169 between AD and control cases regardless of the brain region investigated (Supp Fig 3).
170 Additionally no colocalisation of PV and SST immunoreactivity was present (Supp Fig 4).

171

172 **Discussion**

173 This study suggests a trend to a decrease in SST⁺ interneurons per mm² in the temporal
174 cortex of AD cases whilst, in contrast, no significant difference in PV⁺ interneurons per mm²
175 was identified in AD cases compared to control subjects, in several brain area investigated.
176 Overall there are significant differences in the number of PV⁺ and SST⁺ interneurons per
177 mm² across neocortical and hippocampal subregions within the temporal lobe, with more of
178 both types of interneuron identified in the temporal cortex compared to hippocampal
179 subfields.

180 A major role of interneurons is to influence neuronal circuits by modulating the action of
181 excitatory neurons. SST and PV expressing cells are subsets of GABAergic interneurons ⁹,
182 each providing GABAergic input to specific subcellular domains at defined rates and times ⁶.
183 The unique entity of each interneuron is shown by the lack of colocalisation between PV and
184 SST expressing interneuron which is supported by other work identifying the non-
185 overlapping groups of interneurons ²¹⁻²⁴. Alterations in the gene expression, neural activity
186 and anatomy of SST⁺ interneurons have been identified in a number of psychiatric and
187 neurological disease including schizophrenia, seizure disorders and epilepsy ⁹. In particular
188 SST depletion in the cortex and hippocampus of AD patients has been connected to memory
189 and learning impairment ^{25, 26}. Similarly, SST interneuron decline has been identified in a
190 number of AD animal models strongly correlating with memory and learning impairments ¹⁶,
191 ²⁷⁻²⁹. By restoring GABA receptor signalling with pentobarbitol following GABAergic
192 interneuronal loss, memory and learning deficits were rescued in the apoE4-KI mice ⁵. Also,
193 the use of the neuroprotective peptide α -MSH attenuated GABAergic interneuron loss and
194 improved cognition in the TgCRND8 mouse model of AD ¹⁶.

195 Repetitive activity in pyramidal neurons can drive SST⁺ interneurons into providing feedback
196 inhibition ^{30, 31}. Consequently, a decrease in SST⁺ interneurons in AD temporal cortex as seen
197 in the current study, could act as a protective measure, enabling the continuing excitation of
198 surviving pyramidal neurons to compensate for their progressive loss seen in the disease.

199 Sub-regions of the temporal cortex have important roles in coordinating hippocampal
200 functions, therefore the reduction in SST⁺ interneurons within the temporal cortex of AD
201 patients could suggest a disruption in the overall cortico-hippocampal network and loss of

202 inhibition downstream causing over excitation of pyramidal neurons in the hippocampus ³².
203 Ultimately this could lead to associated negative effects such as an increase in oxidative
204 stress, DNA damage and dysregulation of intracellular calcium that could contribute to
205 neuronal death associated with AD ³³.

206 Previous work has shown several hundred somatostatin labelled neurons in layer II/III of the
207 temporal cortex ³⁴ however, this work investigated three epilepsy cases aged 25-30 yrs, thus
208 differing in both age and pathology of the subjects in the current study. SST decrease in the
209 brain with increasing age ³⁵; this is further heightened in AD ³⁶⁻³⁹. Remaining SST⁺ neuronal
210 processes in AD are located in close proximity to neuritic plaques in the cingulate, frontal,
211 temporal cortex ⁴⁰ and hippocampus ⁴¹. An early reduction in SST⁺ interneurons in the olfactory
212 cortex of an A β PP/PS1 double transgenic mouse model of AD ³⁷ has since been shown in
213 human AD post mortem tissue where SST also colocalised with amyloid-beta (A β) in the
214 olfactory cortex ³⁸. In contrast SST⁺ interneurons rarely colocalised with tau protein ³⁸. In AD
215 the accumulation of A β has been suggested to be caused by the impaired clearance of the
216 protein ⁴². It may be speculated that a loss of SST⁺ interneurons in AD, as seen in this study
217 and others, may lead to a loss in A β degrading enzymes, including endothelin-converting
218 enzyme ECE-2 and therefore reduced A β metabolism and clearance ¹⁸ resulting in A β
219 accumulation and induced cell death ⁴³. Through mass spectrometry studies the most
220 pronounced peptide to bind to A β was the cyclic neuroendocrine peptide somatostatin-14
221 (SST-14) ⁴⁴ highlighting the likely importance of the role of SST interaction with A β surrounding
222 AD pathology.

223 There is conflicting literature as to the significance of PV⁺ interneurons in AD. Human post
224 mortem brain studies have shown a loss of PV⁺ interneurons in areas known to be affected
225 early in the disease, including the entorhinal cortex and hippocampus ⁴⁵⁻⁴⁷ as well as in the
226 hippocampus of patients with dementia with Lewy bodies ⁴⁸. However, in contrast, other
227 human studies showed little variation in PV⁺ interneurons in AD subjects, similar to our
228 current findings, suggesting PV⁺ interneurons are resistant to degeneration in AD ⁴⁹⁻⁵¹.

229 However, despite no changes, PV-expressing synaptic boutons were identified surrounding
230 pyramidal neurons in the hippocampus of some AD patients. This could suggest that as
231 neurons are lost in the hippocampus, an area involved in early AD pathology, an increasing
232 lack of stimulus onto GABAergic interneurons occurs. Ultimately this lack of innervation
233 could lead to PV translocating to axonal terminals to maintain calcium homeostasis and
234 synaptic inhibition in the remaining pyramidal neurons, as the dysregulation of intracellular
235 calcium homeostasis due to synaptic impairment has been previously identified as an
236 initiating factor in AD ⁵². However, a further detailed quantification of these synaptic boutons

237 is required in a larger sample size to confirm this. Therefore the overall loss of PV⁺
238 interneurons may be delayed and not seen until later in the disease as suggested in an
239 A β PP/PS1 double transgenic mouse model of AD³⁷ where the differential vulnerability
240 among interneuron populations was possibly related to A β pathology¹⁷. Parvalbumin
241 immunoreactivity in the subiculum was localised to the parvopyramdial clusters in the current
242 study which has previously shown to be areas immunopositive for depositions of amyloid-Bri,
243 an amyloidogenic fragment associated with a stop codon mutation in the *BRI* gene⁵³.
244 However, no difference in parvalbumin immunoreactivity in the parvopyramdial clusters was
245 identified between controls and AD subjects in this study possibly reflecting the small sample
246 size.

247 Alternatively, neuronal loss in AD and resulting loss of excitable input may cause interneuron
248 dysfunction, rather than degeneration, which could explain the translocation of PV to synaptic
249 boutons^{54, 55}. This translocation of PV could cause detrimental changes in the protein's
250 function causing calcium dysregulation and impaired interneuron inhibition, increasing
251 pyramidal neuron excitability and ultimately neuronal loss associated with the disease.
252 Restoring the function of PV⁺ interneurons has been shown to restore inhibitory synaptic
253 transmission, network activity and cognitive deficits in human amyloid precursor protein
254 (hAPP) transgenic mice⁵⁶.

255 Development of effective therapeutics for treating AD will come about through a better
256 understanding of the mechanisms underlying neuronal dysfunction and loss in the disease.
257 Our current findings, the first neuropathological study investigating PV and SST interneuron
258 distribution throughout the temporal cortex and hippocampus of human AD patients compared
259 to control subjects suggest interneuron changes in AD may be selective to specific interneuron
260 populations and anatomical location. However, these conclusions are based on investigations
261 carried out on only a small number of cases and two subclasses of interneuron. In order to
262 better understand the significance and of interneurons in AD, a much larger study examining
263 more cases and investigating the various interneuron populations across different anatomical
264 regions in the human AD brain is warranted.

265 **Acknowledgements**

266 We would like to thank the donors and their relatives for the gracious gift of tissues for
267 research. We wish to thank the staff of the Sheffield Brain Tissue Bank for their help. We
268 acknowledge the support of the National Institute for Health Research (NIHR) Sheffield
269 Biomedical Research Centre (Translational Neuroscience). RW, and work in SBW's laboratory
270 is supported by the Alzheimer's Society [242(AS-PG-14-015), 386(AS-PG-17-007)].

271

272 **Figure legends**

273 **Figure 1 Somatostatin immunoreactivity.** Somatostatin immunoreactivity was associated
274 with the cytoplasm of interneuronal cell bodies (**a**, black arrows) and immediate dendritic
275 processes (**b**, blue arrows) throughout the temporal cortex and hippocampus.
276 Immunoreactivity was primarily located at the grey matter / white matter border (**c**, low
277 magnification, **d** high magnification). *Scale bar a* 100µm, **b** 100µm, **c** 500µm, **d** 100µm.

278 **Figure 2 Quantification of somatostatin immunoreactivity.** There was significantly higher
279 number of somatostatin immunoreactive interneurons per mm² in the temporal cortex
280 compared to CA1 and entorhinal cortex across all cases (AD and controls) investigated in
281 this study (**a**). There was a reduction in the number of somatostatin interneurons per mm² in
282 the temporal cortex of AD patients compared to control cases (**b**). No differences were seen
283 in the number of somatostatin interneurons per mm² between AD and control cases in the
284 other areas investigated (CA1 and entorhinal cortex) (**b**). *P=<0.05, **P=<0.01 ***P=<0.001.

285 **Figure 3 Parvalbumin immunoreactivity.** Temporal and entorhinal cortex showed a
286 regular pattern of parvalbumin positive interneurons with cytoplasmic (black arrow) and
287 dendritic immunoreactivity (blue arrow) (**a** lower magnification, **b** higher magnification). This
288 pattern of staining appeared in a band like pattern concentrating in the outer layers (I-IV) of
289 the cortex (**c**). Within the CA1 region the immunoreactivity was more dispersed with
290 cytoplasmic (black arrow) and dendritic immunoreactivity (blue arrow) (**d**) and
291 immunoreactivity in the subiculum appeared in clusters (**e** black arrows). A bead-string like
292 pattern of immunoreactivity was present lining neuronal cell bodies of areas of AD
293 hippocampus (circled) (**f**). *Scale bar a* 250µm, **b** 100µm, **c** 500µm, **d** 100µm, **e** 500µm, **f**
294 50µm.

295 **Figure 4 Quantification of parvalbumin immunoreactivity.** Parvalbumin positive cells
296 were counted and expressed as the total number of positive cells per mm² (**a**). There overall
297 total percentage parvalbumin immunoreactivity was calculated per total area examined (**b**).
298 *P=<0.05, **P=<0.01 ***P=<0.001.

299 **Supplementary Figure 1. An illustration of transect belt sampling in the temporal**
300 **cortex.** Non-overlapping images were taken across the cortical layers at X20 magnification
301 in three contiguous belts. The number of images in each belt varied according to the
302 thickness of the cortex.

303 **Supplementary Figure 2. An illustration of plot sampling and transect belt sampling in**
304 **the hippocampus.** Five randomly distributed images were taken from the CA1 region (blue)

305 and three randomly distributed images from the subiculum (red). Non-overlapping images
306 were taken across the entorhinal cortex in two contiguous belts (green).

307 **Supplementary Figure 3.** Boxplots showing total parvalbumin percentage area in the
308 temporal cortex (**a**), subiculum (**b**), CA1 (**c**), and entorhinal cortex (**d**) and total cell count in
309 the temporal cortex (**e**), subiculum (**f**), CA1 (**g**), and entorhinal cortex (**h**) in Alzheimer's
310 disease compared to control cases.

311 **Supplementary Figure 4. Somatostatin and parvalbumin interneurons are distinct**
312 **cells.** (a) Dual immunoreactivity of somatostatin interneurons (brown cells, blue arrows)
313 showed no colocalisation with parvalbumin interneurons (red cells, black arrows) lower
314 magnification. (b) Parvalbumin⁺ interneurons, higher magnification, (c,d) somatostatin⁺
315 interneuron, higher magnification.

ID	Median age (years) (min-max)	Sex (F/M)	PMD (hrs)*	Cause of death (based on clinical information)
Control (9 hippocampus blocks, 9 temporal cortex blocks)	74 (59-84)	(5/4)	47 (5-75)	<ol style="list-style-type: none"> 1. No neurological disease. Brain age-related changes. 2. Atypical pneumonia. Brain no abnormality. 3. Sudden death, history of epilepsy. Brain no abnormality 4. Guillan-Barre Syndrome, metastatic carcinosarcoma. Brain lacunar infarct. 5. Subarachnoid haemorrhage. No neurodegenerative pathology. 6. Pneumonia, carcinoma of the bladder. Brain - basal Ganglia calcification. 7. Cardiac failure, liver failure, chronic hepatitis C. Brain age-related changes. 8. Hepatocellular carcinoma, cirrhosis. Brain no significant abnormality. 9. Renal failure. Brain - no significant abnormality.
AD (9 hippocampus blocks, 9 temporal cortex blocks)	75 (59-93)	(5/4)	62 (24-96)	<p>Alzheimer's Disease</p> <p>Braak Stage 5 (n=4), Braak Stage 6 (n=5).</p> <p>Neuritic plaque staging according to CERAD, moderate neuritic plaques (n=3), severe neuritic plaques (n=6).</p>

317 **Table 1.** Age, sex, post mortem delay (PMD) and cause of death of SBTB brain donors.

318 *Information not available for 10 individuals (6 controls, 4 AD)

319 Key: SBTB, Sheffield Brain Tissue Bank; F, female; M, male

Antibody	Specificity	Isotype	Dilution (time, Temp)	Antigen retrieval method	Supplier
Parvalbumin	Ca ²⁺ binding protein	Rabbit IgG	1:500 (1hr, RT)	MW, 10 mins, TSC	Abcam (ab11427)
Somatostatin	neuropeptide	Rabbit IgG	1:100 (1hr, RT)	MW, 10 mins, EDTA	Abcam (ab108456)

Table 2. Antibody source and specificity

Key: RT, room temperature; MW, microwave; TSC, trisodium citrate buffer pH 6.5; EDTA, Ethylenediamine Tetra-acetic Acid pH 8.0.

	Parvalbumin				Somatostatin	
	Number of positive cells per mm ²		Total percentage immunoreactivity		Number of positive cells per mm ²	
	Mean (SD)	Median (IQR)	Mean (SD)	Median (IQR)	Mean (SD)	Median (IQR)
Temporal cortex						
Control	29.68 (6.85)	28.04 (11.92)	8.87 (4.54)	7.26 (8.42)	2.07 (0.63)	2.05 (0.82)
AD	26.41 (9.19)	21.84 (8.71)	9.30 (3.35)	8.18 (2.35)	1.49 (0.51)	1.43 (0.41)
CA1						
Control	4.58 (5.40)	3.77 (4.72)	0.15 (0.20)	0.05 (0.19)	0.49 (0.60)	0.41 (1.23)
AD	3.63 (4.69)	1.89 (7.55)	0.18 (0.17)	0.14 (0.28)	0.82 (0.37)	0.82 (0.62)
Subiculum						
Control	21.78 (14.53)	17.30 (25.15)	1.80 (2.45)	0.63 (2.01)		
AD	25.15 (17.10)	15.72 (34.60)	0.76 (0.68)	0.44 (1.07)		
Entorhinal cortex						
Control	8.87 (3.53)	9.14 (4.32)	0.14 (0.08)	0.13 (0.14)	1.14 (0.56)	1.23 (1.23)
AD	9.63 (8.00)	15.09 (14.99)	0.23 (0.35)	0.12 (0.24)	1.05 (0.83)	0.82 (1.03)

Table 3. Quantification of Parvalbumin and Somatostatin staining

Key: SD, standard deviation; IQR, inter-quartile range

References

1. Huang Y, Mucke L. Alzheimer mechanisms and therapeutic strategies. *Cell*. 2012; **148**: 1204-1222.
2. Niikura T, Tajima H, Kita Y. Neuronal cell death in Alzheimer's disease and a neuroprotective factor, humanin. *Curr Neuropharmacol*. 2006; **4**: 139-147.
3. Vargova G, Vogels T, Kostecka Z, Hromadka T. Inhibitory interneurons in Alzheimer's disease. *Bratisl Lek Listy*. 2018; **119**: 205-209.
4. Rudy B, Fishell G, Lee S, Hjerling-Leffler J. Three groups of interneurons account for nearly 100% of neocortical GABAergic neurons. *Dev Neurobiol*. 2011; **71**: 45-61.
5. Tong LM, Yoon SY, Andrews-Zwilling Y, Yang A, Lin V, Lei H, et al. Enhancing GABA Signaling during Middle Adulthood Prevents Age-Dependent GABAergic Interneuron Decline and Learning and Memory Deficits in ApoE4 Mice. *J Neurosci*. 2016; **36**: 2316-2322.
6. Lapray D, Lasztocki B, Lagler M, Viney TJ, Katona L, Valenti O, et al. Behavior-dependent specialization of identified hippocampal interneurons. *Nat Neurosci*. 2012; **15**: 1265-1271.
7. Tan Z, Hu H, Huang ZJ, Agmon A. Robust but delayed thalamocortical activation of dendritic-targeting inhibitory interneurons. *Proc Natl Acad Sci U S A*. 2008; **105**: 2187-2192.
8. Urban-Ciecko J, Fanselow EE, Barth AL. Neocortical somatostatin neurons reversibly silence excitatory transmission via GABA_B receptors. *Curr Biol*. 2015; **25**: 722-731.
9. Urban-Ciecko J, Barth AL. Somatostatin-expressing neurons in cortical networks. *Nat Rev Neurosci*. 2016; **17**: 401-409.
10. Jiang X, Shen S, Cadwell CR, Berens P, Sinz F, Ecker AS, et al. Principles of connectivity among morphologically defined cell types in adult neocortex. *Science*. 2015; **350**: aac9462.
11. Lim L, Mi D, Llorca A, Marin O. Development and Functional Diversification of Cortical Interneurons. *Neuron*. 2018; **100**: 294-313.
12. Anticevic A, Cole MW, Murray JD, Corlett PR, Wang XJ, Krystal JH. The role of default network deactivation in cognition and disease. *Trends Cogn Sci*. 2012; **16**: 584-592.
13. Sperling RA, Laviolette PS, O'Keefe K, O'Brien J, Rentz DM, Pihlajamaki M, et al. Amyloid deposition is associated with impaired default network function in older persons without dementia. *Neuron*. 2009; **63**: 178-188.
14. Zott B, Busche MA, Sperling RA, Konnerth A. What Happens with the Circuit in Alzheimer's Disease in Mice and Humans? *Annu Rev Neurosci*. 2018; **41**: 277-297.
15. Palop JJ, Mucke L. Network abnormalities and interneuron dysfunction in Alzheimer disease. *Nat Rev Neurosci*. 2016; **17**: 777-792.

16. Ma K, McLaurin J. alpha-Melanocyte stimulating hormone prevents GABAergic neuronal loss and improves cognitive function in Alzheimer's disease. *J Neurosci.* 2014; **34**: 6736-6745.
17. De la Rosa-Prieto C, Saiz-Sanchez D, Ubeda-Banon I, Flores-Cuadrado A, Martinez-Marcos A. Neurogenesis, Neurodegeneration, Interneuron Vulnerability, and Amyloid-beta in the Olfactory Bulb of APP/PS1 Mouse Model of Alzheimer's Disease. *Front Neurosci.* 2016; **10**: 227.
18. Pacheco-Quinto J, Eckman CB, Eckman EA. Major amyloid-beta-degrading enzymes, endothelin-converting enzyme-2 and neprilysin, are expressed by distinct populations of GABAergic interneurons in hippocampus and neocortex. *Neurobiol Aging.* 2016; **48**: 83-92.
19. Iwakiri M, Mizukami K, Ikonovic MD, Ishikawa M, Hidaka S, Abrahamson EE, et al. Changes in hippocampal GABABR1 subunit expression in Alzheimer's patients: association with Braak staging. *Acta Neuropathol.* 2005; **109**: 467-474.
20. Umeda T, Kimura T, Yoshida K, Takao K, Fujita Y, Matsuyama S, et al. Mutation-induced loss of APP function causes GABAergic depletion in recessive familial Alzheimer's disease: analysis of Osaka mutation-knockin mice. *Acta Neuropathol Commun.* 2017; **5**: 59.
21. Lee S, Hjerling-Leffler J, Zaghera E, Fishell G, Rudy B. The largest group of superficial neocortical GABAergic interneurons expresses ionotropic serotonin receptors. *J Neurosci.* 2010; **30**: 16796-16808.
22. Gonchar Y, Burkhalter A. Three distinct families of GABAergic neurons in rat visual cortex. *Cereb Cortex.* 1997; **7**: 347-358.
23. Kawaguchi Y, Kubota Y. GABAergic cell subtypes and their synaptic connections in rat frontal cortex. *Cereb Cortex.* 1997; **7**: 476-486.
24. Petilla Interneuron Nomenclature G, Ascoli GA, Alonso-Nanclares L, Anderson SA, Barrionuevo G, Benavides-Piccione R, et al. Petilla terminology: nomenclature of features of GABAergic interneurons of the cerebral cortex. *Nat Rev Neurosci.* 2008; **9**: 557-568.
25. Cervia D, Bagnoli P. An update on somatostatin receptor signaling in native systems and new insights on their pathophysiology. *Pharmacol Ther.* 2007; **116**: 322-341.
26. Davis KL, Mohs RC, Marin DB, Purohit DP, Perl DP, Lantz M, et al. Neuropeptide abnormalities in patients with early Alzheimer disease. *Arch Gen Psychiatry.* 1999; **56**: 981-987.
27. Andrews-Zwilling Y, Bien-Ly N, Xu Q, Li G, Bernardo A, Yoon SY, et al. Apolipoprotein E4 causes age- and Tau-dependent impairment of GABAergic interneurons, leading to learning and memory deficits in mice. *J Neurosci.* 2010; **30**: 13707-13717.
28. Leung L, Andrews-Zwilling Y, Yoon SY, Jain S, Ring K, Dai J, et al. Apolipoprotein E4 causes age- and sex-dependent impairments of hilar GABAergic interneurons and learning and memory deficits in mice. *PLoS One.* 2012; **7**: e53569.

29. Levenega J, Krishnamurthy P, Rajamohamedsait H, Wong H, Franke TF, Cain P, et al. Tau pathology induces loss of GABAergic interneurons leading to altered synaptic plasticity and behavioral impairments. *Acta Neuropathol Commun.* 2013; **1**: 34.
30. Kapfer C, Glickfeld LL, Atallah BV, Scanziani M. Supralinear increase of recurrent inhibition during sparse activity in the somatosensory cortex. *Nat Neurosci.* 2007; **10**: 743-753.
31. Silberberg G, Markram H. Disynaptic inhibition between neocortical pyramidal cells mediated by Martinotti cells. *Neuron.* 2007; **53**: 735-746.
32. Ueno H, Suemitsu S, Murakami S, Kitamura N, Wani K, Okamoto M, et al. Postnatal development of GABAergic interneurons and perineuronal nets in mouse temporal cortex subregions. *Int J Dev Neurosci.* 2017; **63**: 27-37.
33. Cotman CW, Su JH. Mechanisms of neuronal death in Alzheimer's disease. *Brain Pathol.* 1996; **6**: 493-506.
34. Gonzalez-Albo MC, Elston GN, DeFelipe J. The human temporal cortex: characterization of neurons expressing nitric oxide synthase, neuropeptides and calcium-binding proteins, and their glutamate receptor subunit profiles. *Cereb Cortex.* 2001; **11**: 1170-1181.
35. Lu T, Pan Y, Kao SY, Li C, Kohane I, Chan J, et al. Gene regulation and DNA damage in the ageing human brain. *Nature.* 2004; **429**: 883-891.
36. Gahete MD, Rubio A, Duran-Prado M, Avila J, Luque RM, Castano JP. Expression of Somatostatin, cortistatin, and their receptors, as well as dopamine receptors, but not of neprilysin, are reduced in the temporal lobe of Alzheimer's disease patients. *J Alzheimers Dis.* 2010; **20**: 465-475.
37. Saiz-Sanchez D, Ubeda-Banon I, De la Rosa-Prieto C, Martinez-Marcos A. Differential expression of interneuron populations and correlation with amyloid-beta deposition in the olfactory cortex of an AbetaPP/PS1 transgenic mouse model of Alzheimer's disease. *J Alzheimers Dis.* 2012; **31**: 113-129.
38. Saiz-Sanchez D, De la Rosa-Prieto C, Ubeda-Banon I, Martinez-Marcos A. Interneurons, tau and amyloid-beta in the piriform cortex in Alzheimer's disease. *Brain Struct Funct.* 2015; **220**: 2011-2025.
39. Davies P, Katzman R, Terry RD. Reduced somatostatin-like immunoreactivity in cerebral cortex from cases of Alzheimer disease and Alzheimer senile dementia. *Nature.* 1980; **288**: 279-280.
40. Morrison JH, Rogers J, Scherr S, Benoit R, Bloom FE. Somatostatin immunoreactivity in neuritic plaques of Alzheimer's patients. *Nature.* 1985; **314**: 90-92.
41. Armstrong DM, LeRoy S, Shields D, Terry RD. Somatostatin-like immunoreactivity within neuritic plaques. *Brain Res.* 1985; **338**: 71-79.

42. Weller RO, Subash M, Preston SD, Mazanti I, Carare RO. Perivascular drainage of amyloid-beta peptides from the brain and its failure in cerebral amyloid angiopathy and Alzheimer's disease. *Brain Pathol.* 2008; **18**: 253-266.
43. Solarski M, Wang H, Wille H, Schmitt-Ulms G. Somatostatin in Alzheimer's disease: A new Role for an Old Player. *Prion.* 2018; **12**: 1-8.
44. Wang H, Muiznieks LD, Ghosh P, Williams D, Solarski M, Fang A, et al. Somatostatin binds to the human amyloid beta peptide and favors the formation of distinct oligomers. *Elife.* 2017; **6**.
45. Brady DR, Mufson EJ. Parvalbumin-immunoreactive neurons in the hippocampal formation of Alzheimer's diseased brain. *Neuroscience.* 1997; **80**: 1113-1125.
46. Solodkin A, Veldhuizen SD, Van Hoesen GW. Contingent vulnerability of entorhinal parvalbumin-containing neurons in Alzheimer's disease. *J Neurosci.* 1996; **16**: 3311-3321.
47. Mikkonen M, Alafuzoff I, Tapiola T, Soininen H, Miettinen R. Subfield- and layer-specific changes in parvalbumin, calretinin and calbindin-D28K immunoreactivity in the entorhinal cortex in Alzheimer's disease. *Neuroscience.* 1999; **92**: 515-532.
48. Bernstein HG, Johnson M, Perry RH, LeBeau FE, Dobrowolny H, Bogerts B, et al. Partial loss of parvalbumin-containing hippocampal interneurons in dementia with Lewy bodies. *Neuropathology.* 2011; **31**: 1-10.
49. Hof PR, Cox K, Young WG, Celio MR, Rogers J, Morrison JH. Parvalbumin-immunoreactive neurons in the neocortex are resistant to degeneration in Alzheimer's disease. *J Neuropathol Exp Neurol.* 1991; **50**: 451-462.
50. Ferrer I, Soriano E, Tunon T, Fonseca M, Guionnet N. Parvalbumin immunoreactive neurons in normal human temporal neocortex and in patients with Alzheimer's disease. *J Neurol Sci.* 1991; **106**: 135-141.
51. Leuba G, Kraftsik R, Saini K. Quantitative distribution of parvalbumin, calretinin, and calbindin D-28k immunoreactive neurons in the visual cortex of normal and Alzheimer cases. *Exp Neurol.* 1998; **152**: 278-291.
52. Bezprozvanny I, Mattson MP. Neuronal calcium mishandling and the pathogenesis of Alzheimer's disease. *Trends Neurosci.* 2008; **31**: 454-463.
53. Holton JL, Ghiso J, Lashley T, Rostagno A, Guerin CJ, Gibb G, et al. Regional distribution of amyloid-B β deposition and its association with neurofibrillary degeneration in familial British dementia. *Am J Pathol.* 2001; **158**: 515-526.
54. Khundakar AA, Hanson PS, Erskine D, Lax NZ, Roscamp J, Karyka E, et al. Analysis of primary visual cortex in dementia with Lewy bodies indicates GABAergic involvement associated with recurrent complex visual hallucinations. *Acta Neuropathol Commun.* 2016; **4**: 66.

55. Lunghi C, Emir UE, Morrone MC, Bridge H. Short-term monocular deprivation alters GABA in the adult human visual cortex. *Curr Biol.* 2015; **25**: 1496-1501.
56. Verret L, Mann EO, Hang GB, Barth AM, Cobos I, Ho K, et al. Inhibitory interneuron deficit links altered network activity and cognitive dysfunction in Alzheimer model. *Cell.* 2012; **149**: 708-721.