



UNIVERSITY OF LEEDS

This is a repository copy of *The multiple mechanisms of spatially discordant alternans in the heart*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/159506/>

Version: Accepted Version

Article:

Colman, MA orcid.org/0000-0003-2817-8508 (2020) The multiple mechanisms of spatially discordant alternans in the heart. *Biophysical Journal*, 118 (10). pp. 2336-2338. ISSN 0006-3495

<https://doi.org/10.1016/j.bpj.2020.03.027>

© 2020, Elsevier. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

The multiple mechanisms of spatially discordant alternans in the heart

Michael A. Colman, University of Leeds, UK

Cardiac alternans, referring to beat-to-beat alternations to the action potential (AP) duration (APD) of cardiomyocytes (Fig. 1), are associated with contractile dysfunction and arrhythmogenesis in multiple disease conditions. In tissue, these cellular phenomena can manifest as spatially concordant alternans (SCA), wherein all regions alternate with the same phase, or spatially discordant alternans (SDA), wherein different regions alternate with opposite (period-2; Fig. 1) or otherwise offset (higher period) phase. Whereas SCA can directly affect cardiac output (mechanical force can alternate strong/weak associated with long/short APD), SDA present also the possibility for transition to potentially fatal arrhythmias (1).

Previous theoretical studies (1,2) have demonstrated the critical role of conduction velocity restitution (CVR) in the development of SDA (Fig. 1), mechanisms which are supported by experimental studies (3); this paradigm, which requires rapid pacing and/or slow recovery of the fast I_{Na} current (I_{Na}), is now well established and has received detailed non-linear analysis (2,4). However, other experimental studies, such as those associated with heart failure, Brugada syndrome, and long QT syndromes, observe SDA emerging at normal or slow pacing rates and/or in the absence of slow I_{Na} recovery (5–7); these observations cannot be explained by mechanisms dependent on CVR. This is far from a trivial discrepancy: the efficacy of pharmacological treatment strategies is strongly linked to the governing arrhythmia mechanisms and it is therefore vital to establish if the dynamics of SDA in the absence of CVR are similar to or distinct from those in its presence.

To explore these pertinent gaps in the understanding of SDA mechanisms, Huang *et al.* in this issue (8) build on two decades' work from Zhilin Qu and colleagues to theoretically study the occurrence and dynamics of SDA emerging at slow pacing rates, or otherwise in the absence of CVR. These investigations reveal distinct dynamics compared to CVR-dependent SDA, highlighting the clinical importance of developing a greater understanding of SDA under all conditions.

In single cells, rapid pacing in combination with certain conditions (such as slow recovery of the calcium current or I_{Na}) can induce bifurcations of the APD restitution curve dependent on refractory properties – alternans. Incomplete recovery leads to a smaller current amplitude on the subsequent beat which, for depolarising currents, results in a reduced AP upstroke velocity and/or plateau phase, and ultimately shorter APD; this incomplete current activation permits full recovery for the next beat, and the cycle repeats. These refractory properties are also present in the intracellular calcium handling system: rapid pacing can induce beat-to-beat alternations to the magnitude of intracellular calcium release and consequently the whole-cell calcium transient, which may induce APD alternans through interactions with calcium-sensitive currents such as that carried by the sodium-calcium exchanger (9,10).

At slow pacing rates and in the absence of slow ion current recovery, however, other mechanisms are necessary to induce alternans, such as abnormal dynamics of the transient outward current (6) or alternating emergence of early-after-depolarisations (EADs; Fig. 1). These mechanisms may be the most relevant for clinically observed alternans in the absence of rapid pacing, yet it is unclear how the manifestation in tissue may differ for alternans resulting from these disparate single-cell mechanisms.

The dynamics of alternans in tissue can be sensitive to heterogeneous initial conditions, which alone can be sufficient to initially induce transient SDA. Heterogeneous initial conditions can

be imposed directly in simulation studies, but it is less clear how these conditions translate to the clinic. In this context, heterogeneous initial conditions may refer to repolarisation heterogeneity, either caused by underlying AP heterogeneity or spatially complex previous excitations in electrically homogeneous tissue, or the presence of abnormal focal excitations responsible for premature ventricular complexes (PVCs). Thus, AP heterogeneity, prior patterns of excitation, and prior PVCs all present the possibility to induce perturbations which can impact the long-term dynamics of SDA. Huang *et al.* in this issue study the impact of all of these factors in combination with multiple conditions of CVR disengagement on the long-term dynamics and stability of nodal points (in 1D) or lines (in 2D) which form at the boundary between different tissue regions in SDA (Fig. 1).

When CVR was engaged, multiple conditions (simple rapid pacing in fully homogeneous tissue; directly imposed heterogeneous initial conditions; PVCs; AP heterogeneity) all led to stable SDA (Fig. 1). Moreover, the final state, referring to the number and location of nodal points or lines, was independent of the specific initial condition heterogeneity and largely independent of AP heterogeneity; nodal lines form perpendicular to AP propagation for multiple underlying heterogeneity structures. Thus, initial and underlying heterogeneities failed to influence the long-term dynamics emerging from rapid-pacing induced alternans; CVR-induced SDA is the only stable solution.

The outcome is more variable in the absence of CVR, with long-term dynamics dependent on the method by which CVR was disengaged and the specific conditions of the simulation. In all cases without CVR, rapid pacing in isolation was insufficient for SDA; perturbations due to heterogeneous initial conditions or PVCs were necessary to induce a transient or sustained period of SDA. In conditions where CVR was disengaged during rapid pacing by increasing the I_{Na} recovery rate, heterogeneous initial conditions- or PVC-induced SDA was unstable, with nodes disappearing to form SCA; in the presence of AP propagation, CVR is necessary for rapid-pacing induced alternans to maintain stable SDA.

A second method to disengage CVR which the authors implemented was to apply rapid-pacing stimuli to the whole-tissue simultaneously, i.e. global pacing; no conduction throughout the tissue occurs and CVR cannot be engaged. Under these non-conduction conditions, weakly heterogeneous initial conditions behaved similarly to the rapid I_{Na} recovery condition, with SDA nodes disappearing and SCA forming (Fig. 1). However, in the presence of strongly heterogeneous initial conditions and/or underlying electrical heterogeneity, stable SDA could be obtained. The emergent SDA pattern was dependent on the specific initial condition heterogeneity and/or underlying tissue heterogeneity; SCA and heterogeneity-induced SDA are both stable solutions.

Finally, CVR was disengaged by slow pacing, with alternans induced via I_{to} or EAD-dependent mechanisms (Fig. 1). In combination with global pacing, similar features emerged to rapid global pacing, with both SCA and SDA being stable solutions in homogeneous and heterogeneous tissue. However, in the presence of conduction and at slow pacing rates, SDA induced in homogeneous tissue was not stable and always resulted in SCA; underlying electrical heterogeneity was necessary to induce stable SDA. This final condition may be the most relevant for the clinical presentation of SDA observed at normal or slow pacing rates.

In summary, the stability of SDA and dependence on prior tissue state and perturbations are all affected by underlying CVR conditions and cellular alternan mechanism. These observations may be critical for successful management of clinical conditions associated with

T-wave alternans, and mechanistic analysis such as presented here is essential to direct future research into novel treatment strategies.




Fig. 1. Different mechanisms of SCA and SDA. The upper panel illustrates single-cell alternans emerging during rapid pacing or alternating EADs. The middle panel summarises the different condition-dependent pathways to the final state of SCA or SDA. *Het. IC* refers to conditions with heterogeneous initial conditions; *PVC* refers to conditions which included an initial applied focal excitation; *Het. AP* refers to conditions with underlying electrophysiological heterogeneity; *Unstable SDA* refers to a transient period of SDA which eventually forms SCA. The lower panel, taken from Huang *et al.* in this issue, illustrates SCA and SDA in a 1D strand of tissue.

References

1. Qu Z, Garfinkel A, Chen PS, Weiss JN. Mechanisms of discordant alternans and induction of reentry in simulated cardiac tissue. *Circulation*. 2000 Oct 3;102(14):1664–70.
2. Echebarria B, Karma A. Instability and spatiotemporal dynamics of alternans in paced cardiac tissue. *Phys Rev Lett*. 2002 May 20;88(20):208101.
3. Gizzi A, Cherry EM, Gilmour RF, Luther S, Filippi S, Fenton FH. Effects of pacing site and stimulation history on alternans dynamics and the development of complex spatiotemporal patterns in cardiac tissue. *Front Physiol*. 2013;4:71.
4. Watanabe MA, Fenton FH, Evans SJ, Hastings HM, Karma A. Mechanisms for discordant alternans. *J Cardiovasc Electrophysiol*. 2001 Feb;12(2):196–206.
5. Narayan SM. T-wave alternans and the susceptibility to ventricular arrhythmias. *J Am Coll Cardiol*. 2006 Jan 17;47(2):269–81.
6. Qu Z, Xie Y, Garfinkel A, Weiss JN. T-wave alternans and arrhythmogenesis in cardiac diseases. *Front Physiol*. 2010;1:154.
7. Holley CL, Cooper JA. Macrovolt T-wave alternans and polymorphic ventricular tachycardia. *Circulation*. 2009 Aug 4;120(5):445–6.
8. Huang C, Song Z, Landaw J, Qu Z. Spatially Discordant Repolarization Alternans in the Absence of Conduction Velocity Restitution. *Biophys J*. 2020 In press. DOI: 10.1016/j.bpj.2020.02.008
9. Clusin WT. Mechanisms of calcium transient and action potential alternans in cardiac cells and tissues. *Am J Physiol Heart Circ Physiol*. 2008 Jan;294(1):H1–10.
10. Restrepo JG, Weiss JN, Karma A. Calsequestrin-mediated mechanism for cellular calcium transient alternans. *Biophys J*. 2008 Oct;95(8):3767–89.