



UNIVERSITY OF LEEDS

This is a repository copy of *Ultrasound erosions in the feet best predict progression to inflammatory arthritis in anti-CCP positive at-risk individuals without clinical synovitis*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/160005/>

Version: Supplemental Material

Article:

Di Matteo, A, Mankia, K, Duquenne, L et al. (5 more authors) (2020) Ultrasound erosions in the feet best predict progression to inflammatory arthritis in anti-CCP positive at-risk individuals without clinical synovitis. *Annals of the Rheumatic Diseases*, 79 (7). pp. 901-907. ISSN 0003-4967

<https://doi.org/10.1136/annrheumdis-2020-217215>

© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ. This is an author produced version of a paper published in *Annals of the Rheumatic Diseases*. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Supplementary Table 1. Prevalence and distribution of x-rays bone erosions at baseline			
Bone erosions	MCP2 joints	MCP5 joints	MTP5 joints
Absence	775 (99.9%)	775 (99.9%)	772 (99.5%)
Presence	1 (0.1%)	1 (0.1%)	4 (0.5%)
<p>Baseline hands and feet x-rays were available for 388 CCP+ at-risk individuals. A total of 6 bone erosions, in 4 out of 388 (1.0%) CCP+ at risk individuals, were identified on x-rays. Two out of 6 bone erosions were also detected on US. These two bone erosions (one grade 1 and one grade 2) were both detected at MTP5 joint level, in absence of concomitant synovitis.</p>			