

This is a repository copy of *Exact nonadiabatic part of the Kohn-Sham potential and its fluidic approximation*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/160118/>

Version: Published Version

Article:

Entwistle, M. T. and Godby, R. W. orcid.org/0000-0002-1012-4176 (2020) Exact nonadiabatic part of the Kohn-Sham potential and its fluidic approximation. PHYSICAL REVIEW MATERIALS. 035002. ISSN 2475-9953

<https://doi.org/10.1103/PhysRevMaterials.4.035002>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Exact nonadiabatic part of the Kohn-Sham potential and its fluidic approximation

M. T. Entwistle  and R. W. Godby

Department of Physics, University of York, and European Theoretical Spectroscopy Facility, Heslington, York YO10 5DD, United Kingdom



(Received 15 November 2019; accepted 25 February 2020; published 16 March 2020)

We present a simple geometrical “fluidic” approximation to the nonadiabatic part of the Kohn-Sham potential, v_{KS} , of time-dependent density-functional theory (DFT). This part of v_{KS} is often crucial, but most practical functionals utilize an adiabatic approach based on ground-state DFT, limiting their accuracy in many situations. For a variety of model systems, we calculate the exact time-dependent electron density and find that the fluidic approximation corrects a large part of the error arising from the “exact adiabatic” approach, even when the system is evolving far from adiabatically.

DOI: [10.1103/PhysRevMaterials.4.035002](https://doi.org/10.1103/PhysRevMaterials.4.035002)

I. INTRODUCTION

Time-dependent Kohn-Sham (KS) density-functional theory [1–3] (TDDFT) is in principle an exact and efficient theory of the dynamics of systems of interacting electrons. In practical applications, while performing well in some cases, its validity is often restricted by the limitations of available approximate functionals for electron exchange and correlation (xc). Typically, an adiabatic approximation to the xc potential is used, in which the instantaneous electron density is implicitly assumed to be in its ground state, thereby neglecting all “memory effects.” While these ground-state approximations have steadily improved [3–14], by definition they cannot approach the exact TDDFT potential: It is necessary to address the nonadiabatic contributions for TDDFT to be capable of predictive accuracy in relation to a multitude of applications to diverse fields such as the determination of electronic excitation energies, including those of a charge-transfer nature [15], electron dynamics [16] including nonperturbative charge transfer dynamics [17], time-resolved spectroscopy [18], and electron scattering [19].

In this paper, to clearly distinguish between adiabatic and nonadiabatic contributions, we consider the purest application of the concept of the adiabatic functional to the *complete* KS potential, v_{KS} : at each instant, the DFT KS potential whose *ground-state* density is equal to the exact time-dependent density. The remainder of the exact v_{KS} constitutes the unambiguously nonadiabatic part, to which we also propose an approximation.

We work in the Runge-Gross formalism [1] of TDDFT, in which the exact xc potential, v_{xc} , at time t [20] depends on the density at all points in space and all nonfuture times. It has been argued [21–24] that the exact nonadiabatic functional often requires strong nonlocal temporal and spatial dependence on the density. A number of properties of the exact functional, such as the harmonic potential theorem (HPT) [21] and zero-force theorem (ZFT) [22], have been used to identify limitations of previous approximate TDDFT functionals. Adiabatic functionals trivially satisfy many of these exact conditions through their complete lack of memory dependence, yet prove inadequate in many applications [15–19, 25–35]. The

development of nonadiabatic functionals that continue to satisfy these exact properties is nontrivial. For example, it was shown that modifying the adiabatic local density approximation by introducing time nonlocality, such as in the Gross-Kohn [36] approximation, is inappropriate [21, 22].

The best-known approximate nonadiabatic functional is that developed by Vignale and Kohn [24, 37, 38] (VK). This was constructed by studying the responses to slowly varying perturbations of the homogeneous electron gas, and they found a time-dependent xc vector potential as a functional of the local current and charge densities j and n , thereby implicitly obtaining a scalar potential which depends nonlocally on the density. While the VK formalism has proved promising [39–49], not least through it obeying the HPT and ZFT, its validity is limited [50–54], owing to the constraints under which it was derived.

II. CALCULATIONS

Our calculations employ the iDEA code [55] which solves the many-electron Schrödinger equation exactly for small, one-dimensional prototype systems of spinless electrons [56, 57]. This gives us access to the exact electron density $n(x, t)$. We then determine the exact $v_{\text{KS}}(x, t)$ through reverse engineering [58]. We also obtain the *exact adiabatic* KS potential [26, 34, 59] v_{KS}^{A} by applying *ground-state* reverse engineering to the instantaneous density at each time [60]. The *exact nonadiabatic* component Δv_{KS} is then $v_{\text{KS}} - v_{\text{KS}}^{\text{A}}$.

A. Fluidic approximation

In developing an approximation to Δv_{KS} , it is helpful to consider the situation in different inertial frames, related through a Galilean transformation, as noted by Tokatly *et al.* [31, 61–64]. While v_{KS}^{A} requires zero correction in any inertial frame when the density is fully static in one of these frames, in the more general case the nonadiabatic corrections to v_{KS}^{A} may be expected to be at their smallest in the local, instantaneous rest frame of the density, defined by a transformation velocity of the local velocity field $u(x, t) = j(x, t)/n(x, t)$. In particular, the effects of acceleration ($\dot{u} \neq 0$) and dispersion ($\partial_x u \neq$

0) have the least effect in a frame where u itself is zero [65]. Conveniently, introducing a vector potential $A = -u(x, t)$ in the original frame of reference is (apart from an unimportant temporal phase factor) equivalent to a Galilean transformation to the local instantaneous rest frame [61,62,66]. As described above, the nonadiabatic correction should be minimal in the latter frame, and here we adopt the simple assumption that it is zero. We term this the *fluidic* approximation. The resulting nonadiabatic correction in the original frame is therefore

$$\Delta v_{\text{KS}}(x, t > 0) = - \int_{-\infty}^x \frac{\partial}{\partial t} u(x', t > 0) dx', \quad (1)$$

where we have gauge transformed A into a scalar potential. It is evident that the density dependence of this Δv_{KS} is nonlocal in both space and time [24].

B. System 1

As a first test of the fluidic approximation, we consider two interacting electrons in a potential well, which takes the form of an inverted Gaussian function. Initially, in the ground state, a uniform electric field, $-\varepsilon x$, is applied at $t = 0$, driving the electrons to the right and inducing a current [Fig. 1(a)]. The sudden application of the perturbation means that we are well outside of the adiabatic limit, and this can be seen by solving the time-dependent KS equations with the exact adiabatic KS potential, $v_{\text{KS}}(t) = v_{\text{KS}}^{\text{A}}(t)$. By plotting the change in the electron density from the ground state, δn , we find $v_{\text{KS}}^{\text{A}}(t)$ on its own to be wholly inadequate ($\approx 13\%$ error in n [67] at $t = 8$ a.u.), while adding the fluidic approximation substantially reduces this error to less than 1% [Fig. 1(b)].

To understand these results, we analyze the nonadiabatic correction to the KS potential in both its scalar and its vector forms. We find very good agreement between the exact ΔA_{KS} and that obtained using the fluidic approximation $-u(x, t)$ [Fig. 2(a)]. The velocity field u [the negative of the fluidic curve in Fig. 2(a)] quickly becomes strongly nonuniform in both space and time as the electrons explore excited states—far removed from a universal rest frame. Similarly close agreement between the exact and fluidic Δv_{KS} [Fig. 2(b)] is evident when the nonadiabatic correction is cast into its scalar form through Eq. (1).

C. Systems 2A, 2B, 2C

We now consider a set of systems of interacting electrons in atomlike external potentials which decay much more slowly at large x , $v_{\text{ext}} = -a/(|x| + a)$ with $a = 20$, thereby increasing correlation. At time $t = 0$, a static sinusoidal perturbation of the form $\varepsilon \cos(0.75x)$ is applied, where ε is 0.02 for system 2A (two electrons), 0.02 for system 2B (three electrons), and 0.1 for system 2C (three electrons).

In system 2A, the sudden perturbation at $t = 0$ acts to push the two electrons apart. This results in a velocity field that is varying in both space and time, as in system 1; in this case, even the sign of u is not the same for all x , which takes us even further away from a universal rest frame. Correspondingly, we find the exact adiabatic potential to be insufficient ($\approx 5\%$ error in n at $t = 5$ a.u.), while adding the fluidic approximation reduces this error to $\approx 1\%$. System 2B




FIG. 1. System 1: Two interacting electrons in a Gaussian potential well, with a uniform electric field applied at $t = 0$, driving the electrons to the right and inducing a current. (a) The ground-state external potential (dashed purple) and exact ground-state electron density (dashed blue), along with the perturbed external potential (solid purple) and exact time-dependent electron density at $t = 8$ a.u. (solid blue). (b) The change in the exact electron density [$\delta n(x, t) = n(x, t) - n(x, 0)$] at $t = 8$ a.u. (short-dashed green), along with that obtained when using the exact v_{KS}^{A} (solid blue), and when adding the exact v_{KS}^{A} with the fluidic approximation $\Delta A_{\text{KS}} = -u$ (dashed red). The exact adiabatic potential is clearly inadequate, but its error is substantially reduced by the fluidic approximation.

contains three interacting electrons in the same v_{ext} as system 2A. The additional electron results in a ground-state density that is much less spatially uniform. We run the simulation for 5 a.u. of time and find similar results: v_{KS}^{A} produces an error in n of $\approx 5\%$, and the fluidic approximation reduces this to $\approx 1\%$.

As mentioned above, the fluidic approximation assumes that a system remains close to its ground state in the local instantaneous rest frame. To stretch this approximation severely, in system 2C the perturbing potential is much stronger, resulting in a much larger response of the density [Fig. 3(a)]. The fluidic approximation still succeeds in reducing the error in the density, from $\approx 25\%$ where only the exact adiabatic potential is used, to $\approx 6\%$ at $t = 5$ a.u. [Fig. 3(b)]. At later times, the dynamic (time-dependent) xc effects become very significant. To confirm this, we replace the xc component of the exact time-dependent v_{KS} with the fixed ground-state v_{xc} , thereby suppressing the dynamic part, and find this potential to be wholly inadequate ($\approx 62\%$ error in n at $t = 18$ a.u.). Here, the exact adiabatic KS potential is better ($\approx 17\%$ error), while adding the fluidic approximation improves it further ($\approx 15\%$ error) [Fig. 3(c)].




FIG. 2. The nonadiabatic correction to the KS potentials for system 1. (a) The exact ΔA_{KS} (short-dashed green) and that obtained using the fluidic approximation $\Delta A_{KS} = -u$ (dashed red), at $t = 8$ a.u. (b) The corresponding exact (short-dashed green) and fluidic (dashed red) Δv_{KS} in its scalar form. The fluidic approximation performs very well, even though the velocity field is non-uniform in both space and time. (The exact adiabatic approximation, of course, amounts to setting $\Delta A_{KS} = \Delta v_{KS} = 0$)

D. Exact conditions

A number of properties of the exact xc functional are known, and these are often used to identify the limitations of approximate functionals. We now explore whether the fluidic approximation satisfies these exact conditions.

We begin with the *one-electron limit*, where the exact xc functional, when applied to a one-electron system, reduces to the negative of the Hartree potential v_H , thereby canceling the spurious self-interaction. This means that v_{KS} is described exactly by a known functional [16,26,34], which has been termed [68] the single orbital approximation—itsself capable of capturing features such as steps in the KS potential [16,69]—whose nonadiabatic part is

$$\Delta v_{KS}(x, t) = - \int_{-\infty}^x \frac{\partial}{\partial t} u(x', t) dx' - \frac{1}{2} u^2(x, t). \quad (2)$$

We note that the first term is the fluidic approximation [Eq. (1)]. We have studied systems of one electron in the external potentials from systems 1, 2A, and 2C, and confirm that the full Eq. (2) yields the exact v_{KS} ; here, the effect on the density of including the $-u^2/2$ term ranges from $<0.1\%$ (potential 2A) to 14% (potential 2C), so the fluidic approximation alone is already satisfactory. Indeed, in our two- and three-electron systems, the effect of adding the additional term to the fluidic approximation is small and typically slightly deleterious.




FIG. 3. System 2C: Three interacting electrons in an atomiclike potential, with a static sinusoidal perturbation applied at $t = 0$, pushing the electrons apart. (a) The ground-state external potential (dashed purple) and exact ground-state electron density (dashed blue), along with the perturbed external potential (solid purple) and exact time-dependent electron density at $t = 5$ a.u. (short-dashed blue) and $t = 18$ a.u. (solid blue). (b) The change in the exact electron density at $t = 5$ a.u. (short-dashed green), along with that obtained when using the exact v_{KS}^A (solid blue), and when adding the exact v_{KS}^A with the fluidic approximation (dashed red). Even though the density is strongly disrupted, the fluidic approximation remains successful. (c) The same as (b) but at $t = 18$ a.u., where the dynamic xc contribution is very significant, evident by the completely inadequate result obtained with the fixed v_{xc} (short-dashed gray) (see main text). Here, the exact v_{KS}^A is better, but adding the fluidic approximation improves it further.

The ZFT [22] follows from Newton's third law and requires the net force exerted on the system by v_H and v_{xc} to vanish. At the level of the KS potential, $\int n(x, t) \partial_x \Delta v_{KS}(x, t) dx = \int n(x, t) \partial_x v_{ext}(x, t) dx$, since the exact v_{KS}^A satisfies the theorem in its own right. In the fluidic approximation for system 1 [70], the left- and right-hand sides of this equation are within 11% of one another so the theorem appears to be approximately obeyed.




FIG. 4. System 3: Two interacting electrons in a tunneling system. Inset: The exact total electron number on the left-hand side ($x < 0$) (short-dashed green); also the exact adiabatic (solid blue) and fluidic approximation (dashed red).

The HPT [21] shows that in a system of interacting electrons in a harmonic potential, subject to a uniform electric field at $t = 0$, the density rigidly moves in the manner of the underlying classical harmonic oscillator. We have shown that the fluidic approximation adds exactly the nonadiabatic correction required [71] by the HPT. We have also confirmed this numerically for two interacting electrons in a harmonic potential.

A constraint that can be challenging for nonadiabatic functionals is the *memory condition* [72], which notes that $v_{xc}(t)$ and hence $v_{KS}(t)$ must be independent of which previous instant in the evolution of the system is to be used to designate the “initial state.” This is violated by the VK functional [34]. Equation (1) demonstrates that the fluidic approximation satisfies this memory condition by virtue of its dependence only on the instantaneous rate of change of u , and not its full history.

E. System 3

As a challenging test of the fluidic approximation, we finally consider two interacting electrons in a tunneling system.

Initially, v_{ext} is a symmetric double-well potential, with one electron localized in each well. At $t = 0$, the left-hand well is raised and the right-hand well lowered, initiating tunneling through the barrier [Fig. 4]. A tunneling electron has an imaginary momentum, meaning that the (real) velocity field is of less physical significance. Correspondingly, the fluidic approximation recovers less of the adiabatic density error, but nevertheless reduces it from $\approx 8\%$ to $\approx 4\%$, at $t = 15$ a.u. Accordingly, the tunneling rate from the left-hand side to the right-hand side is initially improved, but this is not the case at later times [inset of Fig. 4].

III. CONCLUSIONS

In summary, we have calculated the *exact* adiabatic and nonadiabatic parts of the KS potential, v_{KS}^A and Δv_{KS} , for a variety of model systems. Δv_{KS} is precisely defined by our procedure, and represents the part of the time-dependent KS potential that is *intrinsically unobtainable from a ground-state functional*. Our key finding is that a simple geometrical approximation to this nonadiabatic KS potential—making use of a Galilean transformation to the local instantaneous rest frame—recovers most of the density error attributable to the exact adiabatic approach: typically 80–95% in the ballistic systems studied. Studies of additional systems should further illuminate this decomposition of the KS potential of TDDFT in highly nonadiabatic situations, with the fluidic approximation providing a solid foundation for a hierarchy of approximations to Δv_{KS} .

Data created during this research is available from the York Research Database [73].

ACKNOWLEDGMENTS

We thank Jack Wetherell and Nick Woods for recent developments in the iDEA code, and Matt Hodgson and Carsten Ullrich for helpful comments.

-
- [1] E. Runge and E. K. U. Gross, *Phys. Rev. Lett.* **52**, 997 (1984).
 - [2] R. van Leeuwen, *Phys. Rev. Lett.* **82**, 3863 (1999).
 - [3] W. Kohn and L. J. Sham, *Phys. Rev.* **140**, A1133 (1965).
 - [4] J. P. Perdew and A. Zunger, *Phys. Rev. B* **23**, 5048 (1981).
 - [5] J. P. Perdew and Y. Wang, *Phys. Rev. B* **45**, 13244 (1992).
 - [6] J. P. Perdew, *Phys. Rev. Lett.* **55**, 1665 (1985).
 - [7] K. Burke, J. P. Perdew, and Y. Wang, in *Derivation of a Generalized Gradient Approximation: The pw91 Density Functional*, edited by J. F. Dobson, G. Vignale, and M. P. Das (Plenum, NY, 1997), p. 81.
 - [8] J. P. Perdew, J. A. Chevary, S. H. Vosko, K. A. Jackson, M. R. Pederson, D. J. Singh, and C. Fiolhais, *Phys. Rev. B* **46**, 6671 (1992).
 - [9] J. P. Perdew, K. Burke, and M. Ernzerhof, *Phys. Rev. Lett.* **77**, 3865 (1996).
 - [10] J. P. Perdew, M. Ernzerhof, and K. Burke, *J. Chem. Phys.* **105**, 9982 (1996).
 - [11] E. Proynov, S. Sirois, and D. Salahub, *Int. J. Quantum Chem.* **64**, 427 (1997).
 - [12] T. Van Voorhis and G. E. Scuseria, *J. Chem. Phys.* **109**, 400 (1998).
 - [13] J. P. Perdew, S. Kurth, A. Zupan, and P. Blaha, *Phys. Rev. Lett.* **82**, 2544 (1999).
 - [14] J. Sun, A. Ruzsinszky, and J. P. Perdew, *Phys. Rev. Lett.* **115**, 036402 (2015).
 - [15] N. T. Maitra, *J. Phys.: Condens. Matter* **29**, 423001 (2017).
 - [16] P. Elliott, J. I. Fuks, A. Rubio, and N. T. Maitra, *Phys. Rev. Lett.* **109**, 266404 (2012).
 - [17] J. I. Fuks, *European Phys. J. B* **89**, 236 (2016).

- [18] J. I. Fuks, K. Luo, E. D. Sandoval, and N. T. Maitra, *Phys. Rev. Lett.* **114**, 183002 (2015).
- [19] Y. Suzuki, L. Lacombe, K. Watanabe, and N. T. Maitra, *Phys. Rev. Lett.* **119**, 263401 (2017).
- [20] The real system of interacting electrons is mapped onto an auxiliary system of noninteracting electrons moving in the effective potential v_{KS} .
- [21] J. F. Dobson, *Phys. Rev. Lett.* **73**, 2244 (1994).
- [22] G. Vignale, *Phys. Rev. Lett.* **74**, 3233 (1995).
- [23] G. Vignale, *Phys. Lett. A* **209**, 206 (1995).
- [24] G. Vignale and W. Kohn, *Phys. Rev. Lett.* **77**, 2037 (1996).
- [25] C. Jamorski, M. E. Casida, and D. R. Salahub, *J. Chem. Phys.* **104**, 5134 (1996).
- [26] P. Hessler, N. T. Maitra, and K. Burke, *J. Chem. Phys.* **117**, 72 (2002).
- [27] A. Dreuw, J. L. Weisman, and M. Head-Gordon, *J. Chem. Phys.* **119**, 2943 (2003).
- [28] N. T. Maitra, F. Zhang, R. J. Cave, and K. Burke, *J. Chem. Phys.* **120**, 5932 (2004).
- [29] J. Neugebauer, E. J. Baerends, and M. Nooijen, *J. Chem. Phys.* **121**, 6155 (2004).
- [30] R. J. Cave, F. Zhang, N. T. Maitra, and K. Burke, *Chem. Phys. Lett.* **389**, 39 (2004).
- [31] C. A. Ullrich and I. V. Tokatly, *Phys. Rev. B* **73**, 235102 (2006).
- [32] K. J. H. Giesbertz, E. J. Baerends, and O. V. Gritsenko, *Phys. Rev. Lett.* **101**, 033004 (2008).
- [33] J. I. Fuks, A. Rubio, and N. T. Maitra, *Phys. Rev. A* **83**, 042501 (2011).
- [34] N. T. Maitra, *J. Chem. Phys.* **144**, 220901 (2016).
- [35] N. Singh, P. Elliott, T. Nautiyal, J. K. Dewhurst, and S. Sharma, *Phys. Rev. B* **99**, 035151 (2019).
- [36] E. K. U. Gross and W. Kohn, *Phys. Rev. Lett.* **55**, 2850 (1985).
- [37] J. Dobson, G. Vignale, and M. Das, *Electronic Density Functional Theory: Recent Progress and New Directions* (Springer US, New York, US, 2013).
- [38] G. Vignale, C. A. Ullrich, and S. Conti, *Phys. Rev. Lett.* **79**, 4878 (1997).
- [39] M. van Faassen, P. L. de Boeij, R. van Leeuwen, J. A. Berger, and J. G. Snijders, *Phys. Rev. Lett.* **88**, 186401 (2002).
- [40] M. van Faassen, P. L. de Boeij, R. van Leeuwen, J. A. Berger, and J. G. Snijders, *J. Chem. Phys.* **118**, 1044 (2003).
- [41] P. L. de Boeij, F. Kootstra, J. A. Berger, R. van Leeuwen, and J. G. Snijders, *J. Chem. Phys.* **115**, 1995 (2001).
- [42] C. A. Ullrich and G. Vignale, *Phys. Rev. B* **58**, 7141 (1998).
- [43] C. A. Ullrich and G. Vignale, *Phys. Rev. Lett.* **87**, 037402 (2001).
- [44] C. A. Ullrich and G. Vignale, *Phys. Rev. B* **65**, 245102 (2002).
- [45] J. Berger, P. Romaniello, R. van Leeuwen, and P. Boeij, *Phys. Rev. B* **74**, 245117 (2006).
- [46] R. D'Agosta and G. Vignale, *Phys. Rev. Lett.* **96**, 016405 (2006).
- [47] V. U. Nazarov, J. M. Pitarke, Y. Takada, G. Vignale, and Y.-C. Chang, *Phys. Rev. B* **76**, 205103 (2007).
- [48] I. D'Amico and C. A. Ullrich, *Phys. Rev. B* **74**, 121303(R) (2006).
- [49] N. Sai, M. Zwolak, G. Vignale, and M. Di Ventura, *Phys. Rev. Lett.* **94**, 186810 (2005).
- [50] J. A. Berger, P. L. de Boeij, and R. van Leeuwen, *Phys. Rev. B* **75**, 035116 (2007).
- [51] M. van Faassen and P. L. de Boeij, *J. Chem. Phys.* **121**, 10707 (2004).
- [52] M. van Faassen and P. L. de Boeij, *J. Chem. Phys.* **120**, 8353 (2004).
- [53] C. A. Ullrich and K. Burke, *J. Chem. Phys.* **121**, 28 (2004).
- [54] J. A. Berger, P. L. de Boeij, and R. van Leeuwen, *Phys. Rev. B* **71**, 155104 (2005).
- [55] M. J. P. Hodgson, J. D. Ramsden, J. B. J. Chapman, P. Lillystone, and R. W. Godby, *Phys. Rev. B* **88**, 241102(R) (2013).
- [56] The use of spinless electrons gives access to richer correlation for a given number of electrons. The electrons interact via the appropriately softened Coulomb repulsion [65] $u(x, x') = (|x - x'| + 1)^{-1}$. We use Hartree atomic units: $m_e = \hbar = e = 4\pi\epsilon_0 = 1$.
- [57] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevMaterials.4.035002> for the parameters of the model systems, and details of the convergence.
- [58] J. D. Ramsden and R. W. Godby, *Phys. Rev. Lett.* **109**, 036402 (2012).
- [59] M. Thiele, E. K. U. Gross, and S. Kümmel, *Phys. Rev. Lett.* **100**, 153004 (2008).
- [60] Our graphs show the various adiabatic and non-adiabatic KS potentials, etc., evaluated on the *exact* time-dependent density, so any errors in the potentials or densities are entirely attributable to errors in the functionals, not the input to the functionals.
- [61] I. V. Tokatly, *Phys. Rev. B* **71**, 165104 (2005).
- [62] I. V. Tokatly, *Phys. Rev. B* **71**, 165105 (2005).
- [63] I. Tokatly, in *Time-Dependent Density Functional Theory*, edited by M. A. L. Marques, C. A. Ullrich, F. Nogueira, A. Rubio, K. Burke, and E. Gross (Springer-Verlag, Berlin, 2006), Chap. 8, pp. 123–136.
- [64] C. Ullrich, *Time-Dependent Density-Functional Theory: Concepts and Applications*, Oxford Graduate Texts (OUP, Oxford, 2012), pp. 465–476.
- [65] The rate of change of kinetic energy is proportional to ui (as in classical mechanics), and so is smallest when u is zero. Also, if the density is moving with velocity u it will more rapidly encounter a region in which a larger nonadiabatic correction is required.
- [66] The stated A causes the wave function in the original frame to become the wave function in the instantaneous rest frame multiplied by $\exp(-iu^2t/2)$.
- [67] The integrated absolute error, $\int dx |n_1(x, t) - n_2(x, t)|$, expressed as a percentage of the total number of electrons.
- [68] M. J. P. Hodgson, J. D. Ramsden, T. R. Durrant, and R. W. Godby, *Phys. Rev. B* **90**, 241107(R) (2014).
- [69] M. J. P. Hodgson, *Electrons in Model Nanostructures*, Ph.D. thesis, University of York, 2016.
- [70] Systems 2A, 2B, and 2C satisfy the theorem owing to their symmetry and so do not form a useful test.
- [71] Apart from an unimportant time-dependent constant.
- [72] N. T. Maitra, K. Burke, and C. Woodward, *Phys. Rev. Lett.* **89**, 023002 (2002).
- [73] M. T. Entwistle and R. W. Godby, Data related to “Exact non-adiabatic part of the Kohn-Sham potential and its fluidic approximation,” <http://dx.doi.org/10.15124/8570b943-498b-4690-8044-b2208d318ef0> (2020).