



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/161528/>

Version: Accepted Version

Article:

Webb, T.L., du Plessis, H., Christian, H. et al. (2020) Understanding obesity among companion dogs: New measures of owner's beliefs and behaviour and associations with body condition scores. *Preventive Veterinary Medicine*, 180. 105029. ISSN: 0167-5877

<https://doi.org/10.1016/j.prevetmed.2020.105029>

Article available under the terms of the CC-BY-NC-ND licence
(<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Journal Pre-proof

Understanding Obesity among Companion Dogs: New Measures of Owner's Beliefs and Behaviour and Associations with Body Condition Scores

Thomas L. Webb, Hugues du Plessis, Hayley Christian, Eleanor Raffan, Vanessa Rohlf, Gavin A. White



PII: S0167-5877(20)30101-X

DOI: <https://doi.org/10.1016/j.prevetmed.2020.105029>

Reference: PREVET 105029

To appear in: *Preventive Veterinary Medicine*

Received Date: 3 February 2020

Revised Date: 5 May 2020

Accepted Date: 12 May 2020

Please cite this article as: Webb TL, Plessis Hd, Christian H, Raffan E, Rohlf V, White GA, Understanding Obesity among Companion Dogs: New Measures of Owner's Beliefs and Behaviour and Associations with Body Condition Scores, *Preventive Veterinary Medicine* (2020), doi: <https://doi.org/10.1016/j.prevetmed.2020.105029>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

Running head: OBESITY AMONG COMPANION DOGS

Understanding Obesity among Companion Dogs: New Measures of Owner's Beliefs and Behaviour and Associations with Body Condition Scores

Thomas L. Webb, The University of Sheffield, UK Hugues du Plessis, Purina Studios,
Barcelona

Hayley Christian, Telethon Kids Institute, University of Western Australia, Australia

Eleanor Raffan, Wellcome Trust-MRC Institute of Metabolic Science-Metabolic Research
Laboratories, University of Cambridge, Addenbrooke's Hospital, Cambridge, UK

Vanessa Rohlf, La Trobe University, Australia

Gavin A. White, University of Nottingham, Sutton Bonington Campus, UK

For resubmission to *Preventive Veterinary Medicine*

Direct correspondence to: Thomas L. Webb, Department of Psychology, The University of
Sheffield, Cathedral Court, 1 Vicar Lane, Sheffield, S1 2LT. t.webb@sheffield.ac.uk

Abstract

This research aimed to improve our understanding of how owners' beliefs and behaviour are associated with obesity in companion dogs. To do this, we employed new theoretical frameworks and integrated previously reported measures to curate a collection of brief, user-friendly self-report measures to assess owner factors. The reliability and validity of these was examined in two phases of empirical research, each with a cross-sectional questionnaire design that also examined the validity of assessing body condition score (BCS) from photographs submitted by owners. Phase 1 ($n = 47$ dog owners from France) found that the brief owner-report measures correlated with the long-form measures (all correlations except one exceeded $r = 0.70$). BCS as coded from photographs were highly correlated with a vet's assessment of the same dogs ($r = 0.67$). Phase 2 ($n = 3,339$ dog owners from France, Germany, the UK, Italy, and Russia) investigated which measures are associated with obesity among companion dogs. Perceptions of the dog's vulnerability to the threat of obesity, perceived weight status, perceived costs associated with ownership, normative beliefs about feeding, social support from friends, and being in the precontemplation stage of change predicted BCS alongside demographic factors (e.g., dog's age, neutered status). Taken together, the findings provide a method for assessing a wide range of factors that may be associated with obesity among companion dogs and point to potential targets for interventions designed to reduce obesity.

Word count: 232 words (max. 400)

Keywords: Obesity, Overweight, Dog, Owner perceptions

Understanding Obesity among Companion Dogs: New Measures of Owner's Beliefs and Behaviour and Associations with Body Condition Scores

Risk factors for obesity among companion dogs are multifactorial and include factors pertaining to the dog (e.g., genetics, breed, neutered status, age, sex, and responsiveness to food). However, owners typically control food intake and can moderate energy expenditure among companion animals (Association for Pet Obesity Prevention, 2014). This occurs via owners' feeding and exercise practices (Kienzle et al., 1998; Bland et al., 2009; Courcier et al., 2010), which are influenced by their knowledge and the presence of barriers to implementing good practice (Cutt et al., 2008; Endenburg et al., 2018; Webb et al., 2018), and their behavioural, normative, and control beliefs with respect to feeding and exercise (Cutt et al., 2008; Rohlf et al., 2010; 2012; Endenburg et al., 2018). Other factors that have been shown to be associated with obesity include owner perceptions and motivations for treat giving (White et al., 2016; Morelli et al., 2020), and the strength of the bond between dog and owner (Rohlf et al., 2012; Westgarth et al., 2014; 2016). Taken together then, it is clear that factors related to both owner and dog are associated with obesity in companion dogs.

While significant progress has been made in understanding how owners' beliefs and behaviours are associated with obesity among companion animals, research in this area lags behind understanding of behavioural factors in other domains, such as with respect to obesity in people. For example, although research has drawn on social cognitive frameworks like the Theory of Planned Behaviour (Ajzen, 1991) to identify the factors that are likely to be associated with obesity among companion animals (Rohlf et al., 2010; Cairns-Taylor & Fordyce, 2016), many theoretical frameworks that have proved useful in other areas have yet to be applied in this context and, often, research on factors associated with obesity among companion animals is conducted without the guidance of theory (e.g., Perry et al., 2020). While using theory is not a panacea for behaviour change (for a review of issues, see

Prestwich et al., 2015), there are clear advantages to so doing (e.g., theory can suggest potential predictors and explain how and why they are associated with outcomes) and so we suggest that it might be valuable for researchers and practitioners tasked with the challenge of reducing obesity among companion animals to consider theoretical frameworks that have proved useful in other areas. Specifically, we propose that five additional theoretical frameworks might provide useful insights.

Protection Motivation Theory (Rogers, 1983) suggests that two beliefs determine how motivated people are to protect themselves (or in this case their dog). The first belief is their appraisal of the threat, which comprises beliefs about how vulnerable their dog is to becoming overweight or to the negative consequences of obesity (termed *threat vulnerability*) and how severe they deem the threat (termed *threat severity*). In support of the role of severity, Muñoz-Prieto et al. (2018) found that dog owners who did not consider obesity to be a disease were more likely to have obese dogs. The second belief is owner's appraisal of their ability to cope with the threat, which includes beliefs about the efficacy of a potential response (e.g., whether altering feeding or exercise regimes would reduce the risk of obesity, termed *response efficacy*) and their personal ability to enact the required response (e.g., whether they would be able to alter feeding or exercise, termed *self-efficacy*).

Theoretical models that consider how people move through a series of stages of behaviour change are also worthy of consideration. For example, the **Transtheoretical – or 'stages of change' – model** (Prochaska & DiClemente, 1984) suggests that people progress through a series of five stages of behaviour change: *Precontemplation* (where they have not thought about the issue), *contemplation* (where they are intending to take action at some point in the (not immediate) future), *preparation* (where they are intending to take action and have started making plans as to how they will do so), *action* (where they have started to take action), and finally, *maintenance* (where they have taken action, achieved the desired

outcomes, and are now working to prevent relapse). As such, the model may provide a useful summary of the extent to which owners have begun to think about their dog's weight or started taking action and help to target interventions toward those for whom they are likely to be most appropriate and effective (Norcross et al., 2011).

The third framework that may offer useful insights is **Control Theory** (Carver & Scheier, 1982). Control Theory views goal directed behaviour as involving three main processes: goal setting, goal monitoring, and goal operating. The processes involved in setting goals are well accounted for by social cognition models such as the Theory of Planned Behaviour and Protection Motivation Theory, as described above. However, having set a goal (e.g., to feed a dog a certain amount of food a day, or to reduce overall weight), Control Theory suggests that the person needs to monitor the relation between the goal and the current state. In the context of striving to feed an appropriate amount of food to a dog, *monitoring* may involve keeping track of both the amount of food that is provided to the animal (at mealtime and as treats), and/or the weight of the animal.

Finally, volitional – or post-intentional – factors, may help to understand when and how motivation is translated into action. Evidence suggests that people often struggle to translate their motivation (as reflected by behavioural intentions) into action (Sheeran & Webb, 2016). For example, many dog owners are motivated to feed and exercise their dog appropriately, yet struggle to do so, possibly because they fail to appropriately restrict food or succumb to begging (Webb, 2015). Theoretical models like the **Health Action Process Approach** (Schwarzer, 2008) and the **Rubicon Model of Action Phases** (Heckhausen & Gollwitzer, 1987), therefore suggest volitional – or post-intentional – factors, which can help to understand when and how motivation is translated into action. Two key factors in this regard are *action planning* (planning when, where, and how to act) and *coping planning*

(identifying barriers that might derail intended actions and generating plans to manage or overcome them) (Zhang et al., 2019).

The Present Research

The present research sought to understand how owners' beliefs and behaviour are associated with obesity in companion dogs by developing brief measures of beliefs, behaviours, and risk factors, and exploring which factors are associated with canine obesity. Data collection was undertaken in two phases. In the first phase of the research, dog owners completed an online questionnaire containing brief and longer measures of each of the factors of interest, and we investigated whether brief measures can capture the same information as longer measures. In the second phase, a larger sample of owners across five countries completed the brief measures and a subsample completed them a second time so that test-retest reliability could be assessed. In both phases of the research, owners also uploaded photographs of their dog from which trained coders assessed body condition score (BCS; Laflamme, 1997).

Material and Methods

The methods were reviewed and approved by the Research Ethics Committee in the Department of Psychology at The University of Sheffield, UK (Application #022521).

Phase 1: Participants and measures. Power analysis suggested that 42 participants would provide 95% power to detect large-sized (i.e., $r = 0.50$) bivariate relationships between brief and full measures. Therefore, in November 2018, CEN Nutrition Animale recruited 47 dog owners living in France by sending an email to members of their panel. Figure 1 provides a visual representation of the recruitment process. To be eligible to take part in the study, participants needed to be over 18 years old and own (at least one) dog over the age of 1 and be primarily responsible for its care. Participants with dogs that had been diagnosed with a chronic or terminal illness were excluded.

Supplementary Material 1 shows the factors that were measured in Phase 1 mapped onto the theoretical frameworks from which they are derived, and the associated brief and complete (in italics) measures of each factor. Most measures were taken as is or slightly adapted (for country-specific differences) from established questionnaires (Cutt et al., 2008; Rohlf et al. 2010; Richards et al., 2013; Raffan et al., 2015), and recent studies on factors associated with human and canine obesity (Muñoz-Prieto et al., 2018). In addition, owners were asked to self-assess the body condition of their dog using the 5-point BCS scale, where 1 = severely underweight, 2 = thin, 3 = ideal weight, 4 = overweight, and 5 = obese. Participants were also asked to upload two photographs of their dog (one from above and one from the side), alongside a reference object for scale. Five veterinary students at the National Veterinary School of Toulouse were trained to use the submitted photographs to rate the dog's BCS. Finally, owners were asked to bring their dog to a vet who provided an additional assessment of body condition and weight.

Phase 2: Participants and measures. Power analysis suggested that 2,384 participants would provide 95% power to detect small-sized relationships (i.e., $f^2 = .02$) between 62 potential predictors and body condition scores. Therefore, between February and April 2019, IPSOS MORI recruited 3,339 dog owners across five countries; France ($n = 599$), Germany ($n = 626$), the UK ($n = 714$), Russia ($n = 687$) and Italy ($n = 713$) by emailing pre-identified dog owners an invitation to take part in the research. Figure 2 provides a visual representation of the recruitment process. The eligibility criteria were the same as in Phase 1, except that only people aged between 18 and 65 were approached. A subsample of the dog owners living in France (18%; $n = 107$) completed the questionnaire a second time two weeks later to assess test-retest reliability.

Phase 2 used the brief measures developed in Phase 1 (see Supplementary Material 2 for full details of the factors that were measured in Phase 2 mapped onto the theoretical

frameworks from which they are derived, along with the associated measures) and asked owners to upload two pictures of their dog so that coders could assess BCS. $n = 1645$ of the participants (49%) provided two photographs in accordance with the specified guidelines. $n = 870$ participants (26%) submitted photographs that were not suitable (e.g., a photo of their dog jumping into a swimming pool), 241 (7%) of the participants submitted the same photo twice, and a further 583 (17%) of participants submitted photographs, but not of their dog (e.g., stock photographs from Google images). BCS was coded from these photographs by the same trained coders as in phase 1. There was good agreement between the coders (average weighted kappa = 0.62, $SD = 0.10$, range 0.49 to 0.74), according to Altman's (1991) guidelines for interpreting kappa.

The breed of the dogs (where specified by the owners) was coded into those prone to obesity (Cairn Terriers, Basset hounds, Cavalier King Charles Spaniels, Beagles, Cocker Spaniels, Dachshunds, Dalmatians, Labrador Retrievers, Golden Retrievers, Shetland Sheepdogs, Rottweilers) versus those not prone (other breeds, including mixed breed dogs), on the basis of Hand et al. (2000), Lund et al. (2006) and Delaney (2010).

Approach to Analyses. To evaluate whether the brief self-report measures of owner factors provided an adequate measure of the respective constructs and thus could be used in Phase 2 of the research, correlations were examined between the brief and complete measures of each factor. These are reported in the final column of Supplementary Material 1. A correlation of $r = 0.70$ or above was taken to indicate a substantial (and therefore acceptable) correlation between the brief and complete measures.

In order to assess the test-retest reliability of the brief measures in Phase 2 of the research, the intra-class correlation co-efficient (ICC) was computed between the two administrations of the questionnaire. Values between 0.40 and 0.60 indicate moderate

agreement, 0.61 to 0.80 indicate good agreement, values > 0.80 indicate excellent agreement (see Landis & Koch, 1977).

A three-step approach was used to identify which factors were associated with coders' ratings of BCS in Phase 2. Firstly, factors were identified that correlated with BCS at a level that would not occur by chance in more than 10% of the sample (i.e., $p < .10$). Second, a series of hierarchical multiple regressions were conducted to identify the factors within each domain (i.e., beliefs about overweight and obesity, dog-owner bond, feeding, and exercise) that were correlated with BCS scores over and above demographic factors (age of owner, age, gender, and size of dog; along with neutered status), which were entered in Step 1. Finally, the factors that were significantly associated with BCS (i.e., $p < .05$) within each domain were entered in a single hierarchical multiple regression, again controlling for demographic factors.

Results

Phase 1

Dog and owner characteristics. Owners were typically female (23% male), with an average age of 44 years ($SD = 13$, range: 26 to 70). Dogs were, on average 6.5 years old ($SD = 3.2$, range: 1 to 13); 51% were male and 51% were neutered.

Measures of body condition. Table 1 shows the correlation between the three measures of BCS that were employed in Phase 1 (owner's ratings, the average rating of five coders from the pictures that owners submitted, and ratings by the vets when the owners brought their dog to the clinic). The three measures were correlated; coders' ratings of BCS from the photographs that owners submitted were highly correlated with the vet's assessment of BCS ($r = 0.67$) and more so than with the owners' self-ratings ($r = 0.58$), suggesting that coding BCS from photographs is a valid way to assess BCS; and likely more accurate than owners' self-ratings.

Validating brief measures of owner factors. All of the brief measures were correlated at $r = 0.70$ or above with the complete measures, with the exception of the measure of normative beliefs with respect to exercise ($r = 0.39$).

Phase 2

Dog and owner characteristics. Owners were typically female (37% male), with an average age of 44 years ($SD = 12$, range: 18 to 66). Dogs were, on average 5.9 years old ($SD = 3.4$, range: 1 to 19); 57% were male and 46% were neutered. Supplementary Material 3 shows this information by country of recruitment.

Test-retest reliability of the measures. All of the measures showed at least moderate test-retest reliability and most indicated good agreement (see Table 2).

Owner and coder-rated assessments of body condition. Figure 3 shows the BCS ratings provided by the owners against the ratings of BCS made by the coders on the basis of the photographs that the owners submitted. The two ratings were correlated ($r = 0.41$); although owners' typically rated their dog's BCS lower than the coders ($M = 3.13$ vs. 3.54, $SD = 0.50$ vs. 0.62, for owner and coder ratings, respectively, $t(1644) = 26.92$, $p < .001$) and $N = 539$ owners (33% of the sample) rated the BCS of their dog as 'normal' (i.e., BCS = 3), when the coders rated the dog as overweight (BCS = 4 or 5). We therefore used coders' ratings of BCS in our analyses to identify the factors associated with BCS.

Identifying the factors associated with BCS. Table 3 shows the factors that correlated with BCS at a level that would not occur by chance in more than 10% of the sample (i.e., $p < .10$).¹ Table 4 shows the findings of a series of hierarchical multiple regressions to identify the factors within each domain (i.e., beliefs about overweight and obesity, dog-owner bond, feeding, and exercise) that were correlated with BCS scores over and above demographic factors (age of owner, age, gender, and size of dog; along with

¹ The full correlations by country of recruitment are shown in Supplementary Material 3.

neutered status), which were entered in Step 1. Finally, Table 5 shows the findings of a regression of (vet-rated) BCS on factors that were significant predictors of BCS in the within-domain models.²

The overall model was statistically significant, $F(11, 1644) = 35.88, p < .001$, and explained 19% of the variance in BCS. In the final step, the factors that explained significant variance included (in order of the amount of variance explained): Threat vulnerability ($beta = 0.20$: Owners who believed that their dog easily puts on weight were more likely to have an overweight dog), the age of dog ($beta = 0.14$: Older dogs were more likely to be overweight), weight status ($beta = -0.12$: Owners who thought that their dog was fit were less likely to have an overweight dog), perceived costs ($beta = -0.20$: Owners who associate dog ownership with more costs were less likely to have an overweight dog), normative beliefs about feeding ($beta = 0.10$: Owners who think that others believe they feed their dog too much were more likely to have an overweight dog), social support from friends ($beta = -0.07$: Owners whose friends support them to exercise their dog were less likely to have an overweight dog), stage of change: Precontemplation ($beta = -0.06$: Owners who think about their dogs weight were more likely to have an overweight dog), and neutered status ($beta = 0.05$: Neutered dogs were more likely to be overweight).

Discussion

The present research drew on a range of theoretical frameworks to identify and measure factors that are potentially associated with obesity among companion dogs. Having established that brief self-report measures of each factor were appropriate (Phase 1), a large sample of dog owners across five countries completed the measures and submitted

² For parsimony, we focus on the findings of the final hierarchical multiple regression in this report. However, the raw data can be accessed on the Open Science Framework (<https://osf.io/6tz9a/>) and further details of the other analyses are available by contact with the corresponding author.

photographs of their dogs, from which trained coders assessed their body condition (Phase 2). The findings indicated that, in addition to age, sex and neuter status, six owner-related factors were associated with BCS. Specifically, the owners of overweight or obese dogs were *more* likely to (i) think about their dog's weight (i.e., have moved past the precontemplation stage of change), (ii) believe that their dog is unfit, (iii) that it is vulnerable to gaining weight, and (iv) that others think their dog is fed too much. They were *less* likely to (v) believe that dog ownership is costly and (vi) have social support from friends for exercising their dog.

Some of these findings support those of previous research. For example, previous work has also found that normative beliefs are associated with owners' behaviour toward their dogs – e.g., Rohlf et al. (2012) found that, if other people important to the respondent were supportive of their performing beneficial behaviours (including the provision of a nutritionally balanced diet), then those behaviours were more likely to be carried out. In the present research, norms were framed in terms of whether owners' belief that others are critical of their actions (e.g., believe that they feed their dog too much, as in Rohlf et al., 2010), rather than whether they support positive behaviours (e.g., providing a nutritionally balanced diet). However, the association found between norms and outcomes like weight reinforces the idea that owners are likely to consider others beliefs when deciding how to behave with respect to their dog.

In agreement with the current research, a number of studies have also shown that social support can be an important influence on dog walking. However, the majority of these studies focus on the support provided by the *dog* for walking (e.g., Kushner et al., 2006; Cutt et al., 2008; Johnson & Meadows, 2010; Higgins et al., 2013). The present research found that support from *friends* for dog walking (but not support from family or the dog) was associated with better outcomes for the dog. Such findings build on evidence that social support promotes physical activity among people (e.g., Wing & Jeffery, 1999, found that

participants who joined a weight loss programme with three friends, family members, or co-workers lost more weight than those who joined alone) and provide a clear basis for interventions designed to provide social support (at least from friends, with respect to exercise) in an effort to help people to manage the weight of their dogs. For example, Richards et al. (2016) sent emails to dog owners encouraging them to walk their dog with friends and family in an effort to increase self-efficacy for overcoming barriers (e.g., it being dark or having family commitments) and Schneider et al. (2015) developed an online social network intended to increase walking in dog owners.

Novel findings from the current study extend our current understanding of obesity in companion dogs. For instance, we found that the owners of healthy weight dogs associate more costs with owning a dog, than the owners of overweight dogs. The perceived costs of dog ownership were measured using items from the Monash Dog Owner Relationship Scale (MDORS; Dwyer et al., 2006) that reflect the extent to which ownership interferes with life etc, rather than financial costs. For example, participants were asked to what extent they feel that looking after their dog is a chore, find it hard to look after their dog, or find it annoying that they sometimes have to change their plans because of their dog. It seems possible therefore that people who have an overweight dog associate less costs with ownership simply because they don't do as much – i.e., they don't exercise their dog much, think about what to feed them etc. In other words, providing appropriate care for a dog and helping it to maintain a healthy weight is likely more onerous and time consuming than not doing so. Therefore, interventions designed to promote healthy weight might consider managing owner's expectations with respect to the effort involved in owning a dog.

The present findings also indicate that people with overweight dogs are more likely to think about their dog's weight. In other words, they are more likely than the owners of healthy weight dogs to believe that their dog has a weight problem and is vulnerable to

gaining weight. On the one hand, these findings are intelligible in that they suggest that owners notice when their dog is overweight and have at least thought about trying to do something about it; however, they stand in contrast to evidence that people with overweight dogs are often not aware that their dog is overweight (Eastland-Jones et al., 2014) and our finding that many owners assess their dogs BCS as lower than that given by an objective, independent observer. In other words, although these findings support the idea that owners of overweight dogs typically underestimate the BCS of their dogs (White et al., 2011; Eastland-Jones et al., 2014), there was also evidence that the owners of overweight dogs were more likely to respond to measures in a way that acknowledges that their dogs are overweight. The implication is that, although the owners of overweight dogs typically underestimate their dogs' obesity problem using an objective measure, they typically do recognise that the dog is overweight to some extent and so may be willing to engage with weight loss programs. Indeed, Raffan et al. (2015) found that the owners of overweight dogs were more likely to try to restrict their dog's food intake – although, evidently, they were not successful in doing so, suggesting that additional support may be needed to translate willingness into effective action.

The present study did not find any association between factors reflecting the strength of bond / attachment between dog and owner and BCS scores. However, this is not the first study to find no association; a study of risk factors for canine obesity in Denmark (Bjørnvad et al., 2019) found that attachment (as measured by the Lexington Attachment to Pets Scale, Johnson et al., 1992) was not associated with BCS. One explanation is that a strong human-animal bond can have both positive effects on behaviours that influence weight (e.g., lead owners to want to provide high quality diet and exercise regime/training etc., Westgarth et al., 2016) and negative effects (e.g., lead owners to give unsuitable feedstuffs / treats, not to walk the dog when it is raining etc.) These opposing influences may cancel each other out,

suggesting that future research might try to distinguish between owners whose strong attachment leads to healthy versus unhealthy behaviours.

Finally, we did not find any association between feeding treats and obesity as has been reported by Kienzle et al. (1998) and Perry et al. (2020). However, other studies have reported similar findings. For example, German et al. (2011) did not find an association between feeding treats and the success of weight management, and Muñoz-Prieto et al. (2018) found that dogs were more prone to be overweight/obese if they did *not* receive treats. Taken together, these findings suggest that it is important to identify when feeding treats is associated with increased weight and when it is not. It may be that the term ‘treats’ is too generic, as Heuberger and Wakshlag (2011) found that feeding treats high in crude fibre actually reduced a dog’s risk for obesity. This may suggest that the nutritional/energy content of the treats, and other related behaviours (such as reducing the main ration to compensate for treat feeding) are important; and also point to the importance of manufacturer providing sufficient nutritional information on treats to consumers; something which is currently variable (Morelli et al., 2018; 2020). Future research should also ask owners what kind of treats they feed and whether they adjust the main meal to account for the additional calories provided by treats, in addition to how often they provide treats order to better understand the relationship between treats and weight.

Theoretical Frameworks for Understanding Factors Associated with Obesity

The additional theoretical frameworks that were considered in the present research (namely, Protection Motivation Theory, the Transtheoretical or ‘stages of change’ model, Control Theory, the Health Action Process Approach and the Rubicon Model of Action Phases), alongside more traditional theoretical frameworks (e.g., the Theory of Planned Behaviour, Social Cognitive Theory) identified some useful additional factors that may be important for understanding obesity among companion dogs. For example, as discussed

above, threat appraisals from Protection Motivation Theory were associated with body condition scores, such that owners of overweight dogs tended to believe that their dog was more vulnerable to the threat of obesity. Similarly, identifying owners 'stages of change' with respect to tackling obesity proved useful, in the sense that having an overweight dog likely prompted owners to think about taking action (i.e., move beyond the precontemplation stage of action).

There was less clear evidence that monitoring behaviours (e.g., activity levels, food consumed) or outcomes (e.g., weight) was associated with BCS. The bivariate correlations suggest that monitoring weight was associated with BCS, but monitoring weight was not predictive alongside other variables in the multivariate regressions. Furthermore, monitoring diet and exercise was predictive of BCS in the within-domain regressions; however, so doing was associated with higher BCS, suggesting that owners may monitor these behaviours in an effort to tackle an existing weight problem. Similarly, the within-domain regressions provided some evidence that people with overweight dogs were more likely to have made plans specifying how to deal with barriers to appropriate feeding. However, for the most part, volitional factors like monitoring and planning were not associated with BCS, supporting assertions that interventions may need to focus on providing information and helping owners to set appropriate goals before encouraging them to think about how to translate these goals into action (Webb et al., 2018).

Assessing BCS from Photos

It is worth briefly reflecting on the finding that it was possible for trained coders to assess dogs' BCS from photographs. That is, Phase 1 of the present research found a strong correlation ($r = 0.67$) between BCS assessed from photographs submitted by owners (termed vBCS by Gant et al., 2016) and assessments of BCS made by a vet. This supports Gant et al. who suggested that it is possible to indirectly estimate body condition from photographs; a

conclusion that is strengthened by their finding that age, sex, breed, coat length, and coat colour did not significantly affect vBCS. However, one caveat to this conclusion is that only around half of the owners submitted two photographs in accordance with the instructions (namely, from above and from the side). Therefore, it may be more appropriate to say that it was possible for trained coders to assess dogs' BCS if owners provide appropriate photographs, but that clear instructions and guidance are likely needed for owners to do so.

Limitations and Future Directions

The correlational, cross-sectional design of the research means that it is not possible to use the data to distinguish between factors that precede obesity (and thus might cause overweight) and consequences. Indeed, the direction of the relations between some of the factors and BCS scores suggests that some of the factors are likely to be consequences, rather than predictors, of obesity. For example, the negative association between precontemplation and BCS scores suggests that having an overweight dog leads people to think about their dog's weight (and thus not be in the precontemplation stage of change), rather than the converse (i.e., that not thinking about the dog's weight predicts healthy weight). While such relationships are interesting in the sense that they help to understand how people with overweight dogs think about their situation and help to identify those who need to change (i.e., the challenge of diagnosis), these factors are unlikely to explain why dogs become overweight in the first place. In contrast, factors such as a lack of social support from friends for exercising seem more likely to represent predictors of obesity and, thus, targets for intervention. We therefore hope that the brief measures developed in the present research provide the tools needed for longitudinal and / or experimental studies that can examine the relationship between owners' beliefs and behaviour and obesity over time.

Declarations of interest: This research was funded by Nestlé Purina as part of their commitment to reducing the risk of pet obesity through collaborative prevention programs.

Acknowledgements: The authors are grateful to Laurence Colliard for helping to train the students who coded Body Condition Scores from the photographs submitted by owners, to the VETEVEN Association for their collaboration in this process, and to Xuemei Si for computing the reliability of this coding. Hayley Christian's contribution was supported by a Future Leader Fellowship from the Australian National Heart Foundation (#100794).

References

- Ajzen, I., 1991. The theory of planned behavior. *Organ. Behav. Hum. Decis. Process.* 50, 179-211. [https://doi.org/10.1016/0749-5978\(91\)90020-T](https://doi.org/10.1016/0749-5978(91)90020-T)
- Altman, D.G., 1991. *Practical statistics for medical research*. London: Chapman and Hall.
- Association for Pet Obesity Prevention. Obesity facts and risks. June 11, 2014. <http://www.petobesityprevention.org/pet-obesity-fact-risks/>
- Bjørnvad, C.R., Gloor, S., Johansen, S.S., Sandøe, P., Lund, T.B., 2019. Neutering increases the risk of obesity in male dogs but not in bitches - A cross-sectional study of dog- and owner-related risk factors for obesity in Danish companion dogs. *Prev. Vet. Med.* 170, <https://doi.org/10.1016/j.prevetmed.2019.104730>
- Bland, I.M., Guthrie-Jones, A., Taylor, R.D., Hill, J., 2009. Dog obesity: Owner attitudes and behaviour. *Prev. Vet. Med.* 92, 333–340. <https://doi.org/10.1016/j.prevetmed.2009.08.016>
- Cairns-Haylor, T., Fordyce, P., 2017. Mapping discussion of canine obesity between veterinary surgeons and dog owners: a provisional study. *Vet. Rec.* 180, 149. <https://doi.org/10.1136/vr.103878>

- Carver, C.S., Scheier, M.F., 1982. Control theory: a useful conceptual framework for personality-social, clinical, and health psychology. *Psychol. Bull.* 92, 111–135. <https://doi.org/10.1037/0033-2909.92.1.111>.
- Courcier, E.A., Thomson, R.M., Mellor, D.J., Yam, P.S., 2010. An epidemiological study of environmental factors associated with canine obesity. *J. Small Anim. Pract.* 51, 362-367. <https://doi.org/10.1111/j.1748-5827.2010.00933.x>
- Cutt, H.E., Giles-Corti, B., Knuiaman, M.W., Pikora, T.J., 2008. Physical activity behavior of dog owners: Development and reliability of the Dogs and Physical Activity (DAPA) Tool. *J. Phys. Act. Health.* 5, Supp 1, S73-S89. <https://doi.org/10.1123/jpah.5.s1.s73>
- Delaney, S.J., 2010. Overweightness / Obesity: Nestlé Purina PetCare Handbook of Canine and Feline Clinical Nutrition (pp. 32-35). Nestlé Purina Petcare Company.
- Dwyer, F., Bennett, P.C., & Coleman, G. J., 2006. Development of the Monash Dog Owner Relationship Scale (MDORS). *Anthrozoös*, 19, 243-256, <https://doi.org/10.2752/089279306785415592>
- Eastland-Jones, R.C., German, A.J., Holden, S.L., Biourge, V., Pickavance, L.C., 2014. Owner misperception of canine body condition persists despite use of a body condition score chart. *J. Nutr. Sci.* 3, e45. <https://doi.org/10.1017/jns.2014.25>
- Endenburg, N., Soontararak, S., Charoensuk, C., van Lith, H.A., 2018. Quality of life and owner attitude to dog overweight and obesity in Thailand and the Netherlands. *BMC Vet. Res.* 14, 221. <https://doi.org/10.1186/s12917-018-1531-z>
- Gant, P., Holden, S. L., Biourge, V., German, A.J., 2016. Can you estimate body composition in dogs from photographs? *BMC Vet. Res.* 12, 18. <https://doi.org/10.1186/s12917-016-0642-7>

- German, A.J., Holden, S.L., Gernon, L.J., Morris, P.J., Biourge, V., 2011. Do feeding practices of obese dogs, before weight loss, affect the success of weight management? *Br. J. Nutr.* 106, S97–S100. <https://doi.org/10.1017/S0007114511000596>
- Hand, M.S., Thatcher, C.D., Remillard, R.L., Roudebush, P. & Novotny, B.J., 2000. *Small animal clinical nutrition* (4th Ed). Topeka, KS: Mark Morris Institute.
- Heckhausen, H., Gollwitzer, P.M., 1987. Thought contents and cognitive functioning in motivational versus volitional states of mind. *Motiv. Emot.* 11, 101-120. <https://doi.org/10.1007/BF00992338>
- Heuberger, R., Wakshlag, J., 2011. The relationship of feeding patterns and obesity in dogs. *J. Anim. Physiol. Anim. Nutr.* 95, 98-105. <https://doi.org/10.1111/j.1439-0396.2010.01024.x>
- Higgins, J.W., Temple, V., Murray, H., Kumm, E., Rhodes, R., 2013. Walking sole mates: Dogs motivating, enabling and supporting guardians' physical activity. *Anthrozoos*, 26, 237-252. <https://doi.org/10.2752/175303713X13636846944286>
- Johnson, T.P., Garrity, T.F., Stallones, L., 1992. Psychometric evaluation of the Lexington Attachment to Pets Scale (Laps). *Anthrozoos*, 5, 160–175. <https://doi.org/10.2752/089279392787011395>
- Johnson, R.A., Meadows, R.L., 2010. Dog-walking: Motivation for adherence to a walking program. *Clin. Nurs. Res.* 19, 387-402. <https://doi.org/10.1177/1054773810373122>
- Kienzle, E., Bergler, R., Mandernach, A., 1998. A comparison of the feeding behavior and the human-animal relationship in owners of normal and obese dogs. *J. Nutr.* 128, 2779S–2782S. <https://doi.org/10.1093/jn/128.12.2779S>
- Kushner, R.F., Blatner, D.J., Jewell, D.E., Rudloff, K., 2006. The PPET Study: People and pets exercising together. *Obesity*. 14, 1762-1770. <https://doi.org/10.1038/oby.2006.203>

- Laflamme, D., 1997. Development and validation of a body condition score system for dogs. *Canine Pract.* 22, 10-15.
- Landis, J.R., Koch, G.G., 1977. The measurement of observer agreement for categorical data. *Biometrics.* 33, 159-74.
- Lund, E.M., Armstrong, P.J., Kirk C.A. Klausner J. S., 2006. Prevalence and risk factors for obesity in adult dogs from private US veterinary practices. *Int. J. Appl. Res. Vet. Med.* 4, 177-186.
- Morelli, G., Fusi, E., Tenti, S., Serva, L., Marchesini, G., Diez, M., Ricci, R., 2018. Study of ingredients and nutrient composition of commercially available treats for dogs. *Vet. Rec.* 182, 351. <https://doi.org/10.1136/vr.104489>
- Morelli, G., Marchesini, G., Contiero, B., Fusi, E., Diez, M., Ricci, R., 2020. A survey of dog owners' attitudes toward treats. *J. Appl. Anim. Welf. Sci.* 23, 1-9. <https://doi.org/10.1080/10888705.2019.1579095>
- Muñoz-Prieto, A., et al., 2018. European dog owner perceptions of obesity and factors associated with human and canine obesity. *Sci. Rep.* 8, 13353. <https://doi.org/10.1038/s41598-018-31532-0>
- Norcross, J.C., Krebs, P.M., Prochaska, J.O., 2011. Stages of change. *J. Clin. Psychol.* 67, 143-154. <https://doi.org/10.1002/jclp.20758>
- Perry, L.M., Shmalberg, J., Tanprasertsuk, J., Massey, D., Honaker, R.W., Jha, A.R., 2020. Risk factors associated with canine overweightness and obesity in an owner-reported survey. *bioRxiv preprint*, retrieved from <https://www.biorxiv.org/content/10.1101/2020.01.06.896399v1> 16th January 2020. <https://doi.org/10.1101/2020.01.06.896399>

- Prestwich, A., Webb, T.L., Conner, M., 2015. Using theory to develop and test interventions to promote changes in health behaviour: evidence, issues, and recommendations. *Curr. Opin. Psychol.* 5, 1-5. <https://doi.org/10.1016/j.copsyc.2015.02.011>
- Prochaska, J.O., DiClemente, C.C., 1984. *The transtheoretical approach: Crossing the traditional boundaries of change*. Homewood, IL: Irwin.
- Raffan, E., Smith, S.P., O'Rahilly, S., & Wardle, J., 2015. Development, factor structure and application of the Dog Obesity Risk and Appetite (DORA) questionnaire. *PeerJ*, 3, e1278. <https://doi.org/10.7717/peerj.1278>
- Richards, E.A., McDonough, M.H., Edwards, N.E., Lyle, R.M., Troped, P.J., 2013. Development and psychometric testing of the Dogs and WalkinG Survey (DAWGS). *Res. Q. Exerc. Sport.* 84, 492-502. <https://doi.org/10.1080/02701367.2013.839935>
- Richards, E.A., Ogata, N., Cheng, C., 2016. Evaluation of the Dogs, Physical Activity, and Walking (Dogs PAW) Intervention: A randomized controlled trial. *Nurs. Res.* 65, 191-201. <https://doi.org/10.1097/NNR.0000000000000155>
- Rogers, R.W., 1983. Cognitive and physiological processes in fear appeals and attitude change: A revised theory of protection motivation. In J. T. Cacioppo & R. E. Petty (Eds.), *Social psychophysiology: A sourcebook*. London: Guilford Press.
- Rohlf, V., Toukhsati, S., Coleman, G.I., Bennett, P.C., 2010. Dog obesity: Can dog caregivers' (owners') feeding and exercise intentions and behaviors be predicted from attitudes? *J. Appl. Anim. Welf. Sci.* 13, 213-236. <https://doi.org/10.1080/10888705.2010.483871>
- Rohlf, V., Bennett, P.C., Toukhsati, S., Coleman, G., 2012. Beliefs underlying dog owners' health care behaviors: Results from a large, self-selected, internet sample. *Anthrozoös*, 25, 171-185. <https://doi.org/10.2752/175303712X13316289505341>

- Schneider, K.L., Murphy, D., Ferrara, C., Oleski, J., Panza, E., Savage, C., Gada, K., Bozzella, B., Olendzki, E., Kern, D., Lemon, S.C., 2015. An online social network to increase walking in dog owners: A randomized trial. *Med. Sci. Sports. Exerc.* 47, 631–639. <https://doi.org/10.1249/MSS.0000000000000441>.
- Schwarzer, R., 2008. Modeling health behavior change: How to predict and modify the adoption and maintenance of health behaviors. *Appl. Psychol.* 57, 1–29. <https://doi.org/10.1111/j.1464-0597.2007.00325.x>
- Sheeran, P., Webb, T.L., 2016. The intention-behavior gap. *Soc. Personal. Psychol. Compass.* 10, 503-518. <https://doi.org/10.1111/spc3.12265>
- Webb, T.L., 2015. Why pet owners overfeed: A self-regulation perspective. Proceedings of the Nestlé Purina Companion Animal Nutrition Summit on “Pet Obesity: The Future of Weight Management.
- Webb, T.L., Krasuska, M., Toth, Z., du Plessis, H.R., Colliard, L., 2018. Using research on self-regulation to understand and tackle the challenges that owners face helping their (overweight) dogs to lose weight. *Prev. Vet. Med.* 159, 227-231. <https://doi.org/10.1016/j.prevetmed.2018.08.017>
- Westgarth, C., Christley, R.M., Christian, H.E., 2014. How might we increase physical activity through dog walking? A comprehensive review of dog walking correlates. *Int. J. Behav. Nutr. Phys. Act.* 11, 83. <https://doi.org/10.1186/1479-5868-11-83>
- Westgarth, C., Knuiman, M., Christian, H.E., 2016. Understanding how dogs encourage and motivate walking: cross-sectional findings from RESIDE. *BMC Public Health*, 16. <https://doi.org/10.1186/s12889-016-3660-2>
- White, G.A., Ward, L, Pink, C., Craigon, J., Millar, K.M., 2016. “Who’s been a good dog?” – Owner perceptions and motivations for treat giving. *Prev. Vet. Med.* 132, 14-19. <https://doi.org/10.1016/j.prevetmed.2016.08.002>

- White, G.A., Hobson-West, P., Cobb, K., Craigon, J., Hammond, R., Millar, K.M., 2011. Canine obesity: Is there a difference between veterinarian and owner perception? *J. Small Anim. Pract.* 52, 622-626. <https://doi.org/10.1111/j.1748-5827.2011.01138.x>
- Wing, R.R., Jeffery, R.W., 1999. Benefits of recruiting participants with friends an increasing social support of weight loss and maintenance. *J. Consult. Clin. Psychol.* 67, 132–138. <https://doi.org/10.1037/0022-006X.67.1.132>
- Zhang, C.-Q., Zhang, R., Schwarzer, R., Hagger, M.S., 2019. A meta-analysis of the Health Action Process Approach. *Health Psychol.* 38, 623-637. <https://doi.org/10.1037/hea0000728>


Figure 1*Recruitment Process (Phase 1)*

Figure 2

Recruitment Process (Phase 2)





Figure 3*Owner Versus Coder-Rated BCS (Phase 2)*

Table 1*Descriptive Statistics and Correlations between the Measures of BCS (Phase 1)*

	<i>Mean (SD)</i>	Owner-rated	Vet-rated	Coder-rated
Owner-rated	3.47 (0.55)	1.00		
Vet-rated	3.60 (0.54)	.58***	1.00	
Coder-rated	3.51 (0.51)	.57***	.67***	1.00

Note. *** $p < .001$

Table 2*Descriptive Statistics, Internal Reliability (Alpha) and Test-Retest Reliability (ICC) of**Measures in Phase 2*

Factor	<i>M</i> (or median)	<i>SD</i>	Alpha	ICC
Dog and owner characteristics				
Age of owner	43.65	11.75	Single item	n/a
Gender of owner	Female		Single item	n/a
Age of dog	5.89	3.41	Single item	n/a
Gender of dog	Male		Single item	n/a
Breed (not prone vs. prone)	No		Single item	n/a
Size of dog (height)	Reaches knee		Single item	n/a
Neutered status	No		Single item	n/a
Coder-rated BCS	3.54	0.62	kappa = 0.62	n/a
Owner-rated BCS	3.14	0.53	Single item	0.88
Beliefs about obesity / overweight				
Knowledge of dog's weight	Yes		Single item	0.55
Stage of change: Precontemplation	2.36	1.17	Single item	0.65
Stage of change: Contemplation	3.32	1.18	Single item	0.59
Stage of change: Preparation	3.60	1.06	Single item	0.61
Stage of change: Action	3.80	0.99	Single item	0.55
Stage of change: Maintenance	3.50	1.25	Single item	0.65
Threat appraisal: Severity	4.50	0.64	0.73	0.62
Threat appraisal: Vulnerability	2.60	1.08	0.77	0.86
Coping appraisal: Response efficacy	4.46	0.61	0.74	0.66
Coping appraisal: Self-efficacy	4.16	0.70	0.67	0.62
Self-monitoring (of outcomes)	4.12	0.86	Single item	0.61
Weight status	4.16	0.79	0.67	0.88
Dog-owner bond				
Time spent with dog	3.40	1.39	Single item	0.65
Play games with dog	4.70	0.70	Single item	0.41
Take dog to visit people	2.85	1.31	Single item	0.56
Perceived emotional closeness	4.40	0.67	0.68	0.86
Perceived costs	1.94	0.95	0.56	0.84
Feeding				
What kind of food do you typically buy?	Kibble		n/a	Categorical
How many times a day do you feed your dog	1.96	0.73	n/a	0.90
How much do you feed your dog each day? (cups)	1.5 to 2		n/a	Categorical
Knowledge	3.96	0.83	0.55	0.59
Importance of appropriate feeding	4.37	0.64	0.66	0.71
Behavioural beliefs - feed to please	2.69	0.96	0.48	0.67
Normative beliefs	1.88	0.93	0.88	0.80

PBC / self-efficacy	4.05	0.83	0.83	0.75
Intention	4.09	0.87	0.76	0.61
Food monitoring	3.55	0.95	0.64	0.70
Action planning	4.01	0.92	0.78	0.58
Coping planning	3.02	0.99	0.58	0.68
Goal operating (restrictions on human food)	2.73	1.13	0.75	0.68
Responsiveness to food	3.30	0.96	0.64	0.73
Fussiness	3.21	1.06	0.54	0.65
Interest in food	3.47	0.94	0.59	0.72
Barriers	1.87	0.81	0.86	0.74
Beliefs about treats	3.38	0.74	0.71	0.79
Knowledge about treats	3.73	0.96	n/a	0.51
Follow treat guidelines	3.48	1.09	n/a	0.69
Exercise				
Knowledge	3.51	1.09	0.80	0.66
Behavioural beliefs: Value of exercise	4.19	0.81	0.61	0.45
Behavioural beliefs: Dog centred	3.94	0.71	0.74	0.73
Outcome expectations (owner)	4.26	0.71	0.74	0.70
Outcome expectations (dog)	4.39	0.65	0.72	0.72
Normative beliefs	2.21	1.01	0.83	0.77
PBC / self-efficacy	3.78	0.96	0.85	0.79
Intentions	3.93	1.03	0.87	0.78
Goal monitoring	2.80	1.14	0.79	0.64
Action planning	3.34	1.09	0.82	0.67
Coping planning	3.13	1.11	Single item	0.39
Behaviour	3.61	0.95	0.62	0.79
Barriers	1.97	0.79	0.91	0.73
Facilitators	3.95	1.00	0.78	0.65
Social support (dog)	3.92	0.90	0.73	0.79
Social support (family)	3.17	1.00	0.51	0.71
Social support (friends)	2.52	1.00	0.53	0.59

Note. Alpha = Cronbach's alpha. ICC = Intraclass Correlation Co-efficient between the two administrations of the questionnaire (i.e., test-retest reliability). Values between 0.40 and 0.60 indicate moderate agreement, 0.61 to 0.80 indicates good agreement, values >.80 indicate excellent agreement (see Landis & Koch, 1977).

Table 3*Factors Correlated with (Coder-rated) BCS ($p < .10$), Phase 2*

Factor	<i>r</i>
Dog and owner characteristics	
Age of owner	0.06
Age of dog	-0.06
Gender of dog	-0.08
Size of dog (height)	0.22
Breed (not prone vs. prone)	0.12
Neutered status	0.14
Beliefs about obesity / overweight	
Knowledge of dog's weight	-0.06
Stage of change: Precontemplation	-0.11
Stage of change: Contemplation	0.10
Stage of change: Action	-0.05
Stage of change: Maintenance	0.06
Threat vulnerability	0.35
Coping self-efficacy	-0.14
Self-monitoring weight	-0.05
Thoughts / feelings about dog's weight	-0.33
Dog-owner bond	
Play games with dog	-0.04
Take dog to visit people	-0.05
Perceived costs	-0.07
Feeding	
Knowledge (feeding)	-0.12
Knowledge (treats)	-0.10
Restrictions on human food	0.09
Interest in food	0.09
Importance of appropriate feeding	-0.04
Normative beliefs	0.26
PBC / self-efficacy	-0.08
Coping planning	0.07
Barriers	0.20
Exercise	
Knowledge	-0.06
Exercise behaviour	-0.12
Monitoring activity levels	-0.04
Normative beliefs	0.12
PBC / self-efficacy	-0.07
Intentions	-0.05
Barriers	0.04
Facilitators	-0.10
Social support (dog)	-0.06
Social support (friends)	-0.06

Table 4

Hierarchical Regressions of (vet-rated) BCS on Factors Reflecting (i) Beliefs about Outcomes (Regression 1), Human-Animal Bond (Regression 2), Feeding (Regression 3), and Exercise (Regression 4), Controlling for Demographics

	Step 1	Step 2
Age of owner	0.03	
Gender of owner	-0.01	
Size of dog	-0.03	
Gender of dog	-0.05*	
Age of dog	0.20***	
Neutered status	0.10***	
<i>F</i>	23.02***	
<i>R</i> ²	0.07	
Regression 1: Beliefs about outcomes		
Knowledge of dog's weight		-0.04
Stage of change: Precontemplation		-0.07*
Stage of change: Contemplation		-0.03
Stage of change: Action		-0.06*
Stage of change: Maintenance		0.02
Threat vulnerability		0.24***
Self-efficacy		0.03
Self-monitoring		0.01
Weight status		-0.16***
<i>F</i> change		26.47***
<i>R</i> ² change		0.12
Regression 2: Human-animal bond		
Play games		0.00
Visit people		-0.04
Perceived costs		-0.06*
<i>F</i> change		2.51†
<i>R</i> ² change		0.00
Regression 3: Feeding		
Knowledge (feeding)		-0.03
Knowledge (treats)		-0.03
Restrictions on human food		0.01
Importance of appropriate feeding		0.01
PBC / self-efficacy		0.03
Coping planning		0.06*
Barriers		0.17***

<i>F</i> change	9.84***
<i>R</i> ² change	0.04

Regression 4: Exercise

Knowledge	-0.01
Behaviour	-0.05
Monitoring	0.06*
PBC / self-efficacy	-0.07
Intention	0.07
Barriers	0.00
Facilitators	-0.02
Social support (dog)	-0.00
Social support (friends)	-0.03
<i>F</i> change	2.29***
<i>R</i> ² change	0.01

Note. Values alongside factor names represent beta values. *F* and *R*² change refer to additional variance explained by inclusion of predictors in Step 2, over and above those included in Step 1.

Table 5

Regression of (vet-rated) BCS on Factors that were Significant Predictors of BCS in the Within-Domain Models, Controlling for (Significant) Demographics

	Step 1	Step 2
Gender of dog	-0.06**	-0.02
Age of dog	0.20***	0.14***
Neutered status	0.10***	0.05*
Threat vulnerability	-	0.20***
Weight status	-	-0.12***
Perceived costs	-	-0.10***
Normative beliefs (feeding)	-	0.10**
Social support (friends)	-	-0.07**
Stage: Precontemplation	-	-0.06*
Stage of change: Action	-	-0.05
Normative beliefs (exercise)	-	-0.00
<i>F</i>		37.50***
Adj. <i>R</i> ²		0.06
<i>F</i> change		33.07***
<i>R</i> ² change		0.13

Note. Values alongside factor names represent beta values. *F* and *R*² change refer to additional variance explained by inclusion of predictors in Step 2, over and above those included in Step 1.