



UNIVERSITY OF LEEDS

This is a repository copy of *International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: General Introduction*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/161685/>

Version: Accepted Version

Article:

Levin, L, Day, P orcid.org/0000-0001-9711-9638, Hicks, L et al. (4 more authors) (2020) International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: General Introduction. *Dental Traumatology*, 36 (4). pp. 309-313. ISSN 1600-4469

<https://doi.org/10.1111/edt.12574>

This article is protected by copyright. All rights reserved. This is the peer reviewed version of the following article: Levin, L., Day, P., Hicks, L., O'Connell, A., Fouad, A.F., Bourguignon, C. and Abbott, P.V. (2020), International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: General Introduction. *Dent Traumatol*. Accepted Author Manuscript. , which has been published in final form at <https://doi.org/10.1111/edt.12574>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>



International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: General Introduction.

Liran Levin¹, Peter Day², Lamar Hicks³, Anne O'Connell⁴, Ashraf F. Fouad⁵, Cecilia Bourguignon⁶, Paul V. Abbott⁷.

¹Faculty of Medicine and Dentistry, University of Alberta, Canada.

²School of Dentistry at the University of Leeds and Community Dental Service Bradford District Care NHS Trust.

³Division of Endodontics, University of Maryland School of Dentistry, UMB, Baltimore, Maryland, USA.

⁴Paediatric Dentistry, Dublin Dental University Hospital, Trinity College Dublin, The University of Dublin, Ireland.

⁵Adams School of Dentistry, University of North Carolina, Chapel Hill, NC, USA.

⁶Cecilia Bourguignon, Private Practice, Paris, France

⁷UWA Dental School, University of Western Australia.

Keywords: Trauma, avulsion, tooth fracture, prevention, luxation

Short title: IADT guidelines for management of traumatic dental injuries

Correspondence and reprint requests to:

Prof. Liran Levin – Chair of the IADT Guidelines Committee

University of Alberta, Faculty of Medicine & Dentistry

5-468 Edmonton Clinic Health Academy

11405 - 87 Avenue NW, 5th Floor

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/EDT.12574](https://doi.org/10.1111/EDT.12574)

This article is protected by copyright. All rights reserved

Edmonton AB T6G 1C9

E-mail: liran@ualberta.ca

Acknowledgement and Competing Interest

The authors declare there are no competing interest for the above manuscript. No funding was received for the presented work.

Ethical Statement – No ethic approval was required for this paper.

PROFESSOR LIRAN LEVIN (Orcid ID : 0000-0002-8123-7936)

PROFESSOR ASHRAF F. FOUAD (Orcid ID : 0000-0001-6368-1665)

PROFESSOR PAUL VINCENT ABBOTT (Orcid ID : 0000-0001-5727-4211)

Article type : Comprehensive Review

International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: General Introduction.

Abstract:

Traumatic dental injuries (TDIs) occur most frequently in children and young adults. Older adults also suffer TDIs but at significantly lower rates than individuals in the younger cohorts. Luxation injuries are the most common TDIs in the primary dentition, whereas crown fractures are more commonly reported for the permanent teeth. Proper diagnosis, treatment planning and follow-up are very important to assure a favorable outcome. These updates of the International Association of Dental Traumatology's (IADT) Guidelines include a comprehensive review of the current dental literature using EMBASE, MEDLINE, PUBMED, Scopus and Cochrane Databases for Systematic Reviews searches from 1996-2019 and a search of the journal Dental Traumatology from 2000-2019.

The goal of these guidelines is to provide information for the immediate or urgent care of TDIs. It is understood that some follow-up treatment may require secondary and tertiary interventions involving dental and medical specialists with experience in dental trauma.

As with previous guidelines, the current working group included experienced investigators and clinicians from various dental specialties and general practice. The current revision represents the best evidence

based on the available literature and expert opinions. In cases where the published data were not conclusive, recommendations were based on the consensus opinions of the working group. They were then reviewed and approved by the members of the IADT Board of Directors.

It is understood that guidelines are to be applied using careful evaluation of the specific clinical circumstances, the clinician's judgment and the patient's characteristics, including the probability of compliance, finances, and a clear understanding of the immediate and long-term outcomes of the various treatment options versus non-treatment. The IADT does not, and cannot, guarantee favorable outcomes from adherence to the Guidelines. However, the IADT believes that their application can maximize the probability of favorable outcomes.

Introduction

Traumatic dental injuries (TDIs) occur frequently in children and young adults, comprising 5% of all injuries. Twenty-five percent of all school children experience dental trauma and 33% of adults have experienced trauma to the permanent dentition, with the majority of the injuries occurring before age 19. Luxation injuries are the most common TDIs in the primary dentition, whereas crown fractures are more commonly reported for the permanent teeth. Proper diagnosis, treatment planning and follow-up are important to assure a favorable outcome.

These updates of the International Association of Dental Traumatology's (IADT) Guidelines include a review of the current dental literature using EMBASE, MEDLINE, PUBMED and Scopes searches from 1996-2019 and a search of the journal *Dental Traumatology* from 2000-2019.

The goal of these guidelines is to provide information for the immediate and urgent care of TDIs. It is understood that some of the subsequent treatment may require secondary and tertiary interventions involving specialists with experience in dental trauma.

The IADT published its first set of guidelines in 2001, and updated them in 2007. A further update was published in *Dental Traumatology* in 2012. As with previous guidelines, the current working group included experienced investigators and clinicians from various dental specialties and general practice. The current revision represents the best evidence based on the available literature and expert professional judgment. In cases where the data were not conclusive, recommendations were based on the consensus opinion of the working group, then reviewed and approved by the members of the IADT Board of Directors.

It is understood that guidelines are to be applied with evaluation of the specific clinical circumstances, clinicians' judgment and patients' characteristics, including but not limited to the probability of compliance, finances, and an understanding of the immediate and long-term outcomes of treatment options versus non-treatment. The IADT does not, and cannot, guarantee favorable outcomes from adherence to the Guidelines, but the IADT believes that their application can maximize the chances of a favorable outcome.

These Guidelines offer recommendations for the diagnosis and treatment of specific TDIs. However, they provide neither the comprehensive nor the detailed information found in textbooks, the scientific literature or the Dental Trauma Guide (DTG). The DTG can be accessed at <http://www.dentaltraumaguide.org>. In addition, the IADT website, <http://www.iadt-dentaltrauma.org> provides connection to the journal *Dental Traumatology* and other dental trauma information.

GENERAL RECOMMENDATIONS

Special considerations for trauma to primary teeth

A young child is often difficult to examine and treat due to lack of cooperation and because of fear. This situation is distressing for both the child and the parents. It is important to keep in mind that there is a close relationship between the root apex of the injured primary tooth and the underlying permanent tooth germ. Tooth malformation, impacted teeth, and eruption disturbances in the developing permanent dentition are some of the consequences that can occur following severe injuries to primary teeth and/or alveolar bone. A child's maturity and ability to cope with the emergency situation, the time for shedding of the injured tooth, and the occlusion are all important factors that influence treatment. Multiple traumatic episodes are also common in children and this may affect the outcomes following trauma to a tooth.

Immature versus Mature Permanent Teeth

Every effort should be made to preserve the pulp in the immature permanent tooth to ensure continued root development. A large majority of TDIs occur in children and teenagers where loss of a tooth has lifetime consequences. The immature permanent tooth has considerable capacity for healing after traumatic pulp exposure, luxation injury or root fracture.

Avulsion of Permanent Teeth

The prognosis for avulsed permanent teeth is heavily dependent on the actions taken at the place of accident. Promotion of public awareness of first-aid treatment for the avulsed tooth is strongly encouraged. Treatment choices and prognosis for the avulsed tooth are largely dependent on the viability of the periodontal ligament (PDL), and the maturity of the root. *See the IADT's specific Guidelines for managing avulsed teeth.*

Patient/Parent Instructions

Patient compliance with follow-up visits and home care contributes to better healing following a TDI. Both the patient and the parents of a young patient should be advised regarding care of the injured tooth or teeth for optimal healing, preventing further injury, employing meticulous oral hygiene and rinsing with an antibacterial agent such as alcohol-free chlorhexidine gluconate 0.12% for 1-2 weeks.

Alternatively, with a young child, it is desirable to apply the chlorhexidine to the affected area with a cotton swab.

Summary tables for follow-up, splinting duration and core outcomes

To help summarise activities for the follow-up appointment and splinting regimes, Tables 1, 2 and 3 have been produced. These are presented for different injuries in the primary and permanent dentition. The core outcome variables, explained in the next paragraph, are also included.

Core Outcome Set:

When the worldwide trauma literature is reviewed, it is dominated by one centre in Copenhagen⁶⁷. The lifetime work of Dr Andreasen and his research group is remarkable in both its longevity and the prolific publication of their results. One of the key fundamentals of scientific research is replication, where the results found in one centre with one group of patients are also consistently seen across other patient groups. It is essential that the results from other centres are published even when they confirm the findings from earlier studies. By increasing the number of studies available for clinicians and researchers to analyse, the ability to compare, contrast and combine studies as appropriate is enhanced.

The IADT recently developed a core outcome set (COS) for Traumatic Dental Injuries (TDI) in children and adults.¹ This is one of the first COS developed in dentistry and follows a robust consensus methodology and is underpinned by a systematic review of the outcomes used in the trauma literature.² A number of outcomes were identified as recurring throughout the different injury types. These outcomes were then included as “generic”—that is relevant to all TDI. Injury-specific outcomes were also determined as those outcomes related only to one or more particular TDI. Additionally, the study established what, how, when and by whom these outcomes should be measured. Tables 1 and 2 show the generic and injury specific outcomes to be recorded at the follow-up review appointments for the different traumatic injuries. Further information for each outcome is described in the original paper¹ and the supplementary materials that are available on the journal’s website.

References

1. Kenny KP, Day PF, Sharif MO, Parashos P, Lauridsen E, Feldens CA, et al: What are the important outcomes in traumatic dental injuries? An international approach to the development of a core outcome set. *Dent Traumatol* 2018;34:4-11.
2. Sharif MO, Tejani-Sharif A, Kenny K, Day PF: A systematic review of outcome measures used in clinical trials of treatment interventions following traumatic dental injuries. *Dent Traumatol* 2015;31:422-428.

Table 1 - Primary Dentition follow-up regimes

At these follow-up visits consider collecting the generic and injury specific outcomes as identified by the Core Outcome Set - Kenny et al. Dent Traumatol 2018.

	1W	4W	8W	3M	6M	1 Year	At 6 Years old		Generic outcomes to consider collecting as identified by the Core Outcome Set	Injury specific outcomes to consider collecting as identified by the Core Outcome Set
Enamel fracture										
Enamel/dentin fracture			*						Periodontal healing (including bone loss, gingival recession, mobility, ankylosis/resorption) Pulp healing (including infection) Pain Discoloration Tooth loss Quality of life (days off work, school, sport) Aesthetics (patient perception) Trauma related dental anxiety Number of clinic visits Impact on development of permanent successor	Quality of restoration Loss of restoration
Crown fracture	*		*			*	(Radiograph only if endodontic treatment carried out)			Quality of restoration Loss of restoration
Crown/root fracture	*		*			*	(Radiograph only if endodontic treatment carried out)			If crown restored: Quality of restoration Loss of restoration
Root fracture	*	*S	*			*				Realignment – where spontaneous repositioning undertaken
Alveolar fracture	*	*SR	*			*R		*		
Concussion	*		*							Periodontal healing (including bone loss, gingival

Subluxation	*		*						recession, mobility, ankylosis/resorption)	
Extrusion	*		*			*			Pulp healing (including infection)	Realignment – where spontaneous repositioning undertaken
Lateral luxation	*	*S	*		*	*			Pain	Realignment – where spontaneous repositioning undertaken
Intrusion	*		*		*	*	*		Discoloration	Realignment – where spontaneous repositioning undertaken
									Tooth loss	Realignment – where spontaneous repositioning undertaken
									Quality of life (days off work, school, sport)	Realignment – where spontaneous repositioning undertaken
									Aesthetics (patient perception)	Realignment – where spontaneous repositioning undertaken
									Trauma related dental anxiety	
									Number of clinic visits	
									Impact on development of permanent successor	Infra-occlusion
Avulsion	*		*				*		Pain	
									Tooth loss	
									Aesthetics	
									Quality of life	
									Trauma related dental anxiety	
									Number of clinic visits	
									Impact on development of permanent successor	

* = clinical review appointment

S = splint removal

R = radiograph advised even if no clinical signs or symptoms

Table 2 - Permanent Dentition follow-up regimes

At these follow-up visits consider collecting the generic and injury specific outcomes as identified by the Core Outcome Set - Kenny et al. Dent Traumatol 2018.

	2W	4W	6-8W	3M	4M	6M	1 Year	Yearly up to at least 5 years	Generic outcomes to consider collecting as identified by the Core Outcome Set	Injury specific outcomes to consider collecting as identified by the Core Outcome Set
Infraction	No follow up									
Enamel fracture			*R				*R		Periodontal healing (including bone loss, gingival recession, mobility, ankylosis/resorption) Pulp healing (including infection) # Pain Discoloration Tooth loss Quality of life (days off work, school, sport) Aesthetics (patient perception) Trauma related dental anxiety Number of clinic visits	Quality of restoration Loss of restoration
Enamel/dentin fracture			*R				*R			
Crown fracture			*R	*R		*R	*R			Quality of restoration Loss of restoration
Crown/root fracture			*R	*R		*R	*R	*R		
Root fracture (apical third, mid-third)		*S*R	*R		*R	*R	*R	*R		Root fracture repair
Root fracture (cervical third)		*R	*R		*S*R	*R	*R	*R		
Alveolar fracture		*S*R	*R		*R	*R	*R	*R		

Concussion		*R					*R		Periodontal healing (including bone loss, gingival recession, mobility, ankylosis/resorption) Pulp healing (including infection) # Pain Discoloration Tooth loss Quality of life (days off work, school, sport) Aesthetics (patient perception) Trauma related dental anxiety Number of clinic visits	
Subluxation	(*S) *R			*R		*R	*R			
Extrusion	*S*R	*R	*R	*R		*R	*R	*R		Infra-occlusion
Lateral luxation	*R	*S*R	*R	*R		*R	*R	*R		
Intrusion	*R	(*S) *R	*R	*R		*R	*R	*R		Infra-occlusion Realignment – where spontaneous repositioning undertaken
Avulsion (mature tooth)	*S*R	*R		*R		*R	*R	*R		
Avulsion (immature tooth)	*S*R	*R	*R	*R		*R	*R	*R		Infra-occlusion

* = clinical review appointment

S = splint removal

R = radiograph advised even if no clinical signs or symptoms

= for immature permanent teeth with necrotic and and infected pulps, consider the following additional outcomes: root length, root width and late stage crown fracture

Table 3 - Splinting durations for the permanent and primary dentitions

	2W	4W	4M
Permanent Dentition			
Subluxation	• (if splinted)		
Extrusion	*		
Lateral luxation		*	
Intrusion		*	
Avulsion	*		
Root fracture (apical third, mid-third)		*	
Root fracture (cervical third)			*
Alveolar fracture		*	
Primary Dentition			
Root fracture		• (if splinting required)	

Lateral luxation		<ul style="list-style-type: none">• (if splinting required)	
Alveolar fracture		<ul style="list-style-type: none">•	