



This is a repository copy of *Macrophage polarisation associated with atherosclerosis differentially affects their capacity to handle lipid.*

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/162021/>

Version: Accepted Version

Article:

Baidžajevas, K., Hadadi, É., Lee, B. et al. (6 more authors) (2020) Macrophage polarisation associated with atherosclerosis differentially affects their capacity to handle lipid. *Atherosclerosis*, 305. pp. 10-18. ISSN 0021-9150

<https://doi.org/10.1016/j.atherosclerosis.2020.05.003>

Article available under the terms of the CC-BY-NC-ND licence
(<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Manuscript Number: ATH-D-19-00697R2

Title: Macrophage polarisation associated with atherosclerosis differentially affects their capacity to handle lipid

Article Type: Research paper

Section/Category: Basic Research

Keywords: Inflammation, Atherosclerosis, Macrophage, Innate, Lipid

Corresponding Author: Dr. Heather Wilson,

Corresponding Author's Institution: University of Sheffield Medical School

First Author: Kajus Baidžajevs

Order of Authors: Kajus Baidžajevs; Éva Hadadi; Bernett Lee; Josephine Lum; Foo Shihui; Ian Sudbury; Endre Kiss-Tóth; Siew Cheng Wong; Heather Wilson

Abstract: Background & Aims: Lipid-rich foam cell macrophages drive atherosclerosis via several mechanisms, including inflammation, lipid uptake, lipid deposition and plaque vulnerability. The atheroma environment shapes macrophage function and phenotype; anti-inflammatory macrophages improve plaque stability while pro-inflammatory macrophages promote rupture. Current evidence suggests a variety of macrophage phenotypes occur in atherosclerotic plaques with local lipids, cytokines, oxidised phospholipids and pathogenic stimuli altering their phenotype. In this study, we addressed differential functioning of macrophage phenotypes via a systematic analysis of in vitro polarised, human monocyte-derived macrophage phenotypes, focussing on molecular events that regulate foam-cell formation.

Methods: We examined transcriptomes, protein levels and functionally determined lipid handling and foam cell formation capacity in macrophages polarised with IFN γ +LPS, IL-4, IL-10, oxPAPC and CXCL4.

Results: RNA sequencing of differentially polarised macrophages revealed distinct gene expression changes, with enrichment in atherosclerosis and lipid-associated pathways. Analysis of lipid processing activity showed IL-4 and IL-10 macrophages have higher lipid uptake and foam cell formation activities, while inflammatory and oxPAPC macrophages displayed lower foam cell formation. Inflammatory macrophages showed low lipid uptake, while higher lipid uptake in oxPAPC macrophages was matched by increased lipid efflux capacity.

Conclusions: Atherosclerosis-associated macrophage polarisation dramatically affects lipid handling capacity underpinned by major transcriptomic changes and altered protein levels in lipid-handling gene expression. This leads to phenotype-specific differences in LDL uptake, cellular cholesterol levels and cholesterol efflux, informing how the plaque environment influences atherosclerosis progression by influencing macrophage phenotypes.

Research Data Related to this Submission

Title: Data for: Macrophage polarisation associated with atherosclerosis
differentially affects their capacity to handle lipid

Repository: Mendeley Data


<https://data.mendeley.com/datasets/snwtjhx7h3/draft?a=f9f585e0-f497-4c9f-87aa-cb40823f91bc>

Highlights:


- Total RNA sequencing of differentially polarised macrophages showed enrichment in genes associated with atherosclerosis and lipid pathways
- Anti-inflammatory macrophages had higher lipid uptake and foam cell formation activities
- Inflammatory and oxPAPC macrophages displayed lower foam cell formation
- Inflammatory macrophages showed low lipid uptake, while higher lipid uptake in oxPAPC macrophages was matched by increased lipid efflux capacity

Graphical abstract



A. M^{IFN γ +LPS}: lower cholesterol uptake, storage & efflux



B. M^{IL-4/IL-10}: high cholesterol uptake, storage & efflux



C. M^{oxPAPC}: moderate cholesterol uptake/processing, high efflux



1
2
3
4
5
6
7 **Macrophage polarisation associated with atherosclerosis differentially affects**
8 **their capacity to handle lipid**
9

10
11
12 Short Title: Macrophage polarisation & lipid handling capacity
13

14 Kajus Baidžajevas*^{a,b}, Éva Hadadi*^{a,b,c}, Bernett Lee^b, Josephine Lum^b, Foo Shihui^b,
15 Ian Sudbery^d, Endre Kiss-Tóth^a, Siew Cheng Wong^{b§}, Heather L Wilson^{a§}.
16
17

18 *These authors contributed equally.
19

20 a Department of Infection, Immunity & Cardiovascular Disease, University of Sheffield, Beech Hill
21 Road, Sheffield, S10 2RX, UK.
22

23 b Singapore Immunology Network (SIgN), Agency for Science, Technology and Research (A*STAR),
24 Singapore.
25

26 c Current address: INSERM U935, Villejuif, France.
27

28 d Department of Molecular Biology and Biotechnology, University of Sheffield, Western Bank,
29 Sheffield, S10 2TN, UK.

30 §Corresponding authors:

31 Heather Wilson: Department of Infection, Immunity & Cardiovascular Disease, University of
32 Sheffield, Beech Hill Road, Sheffield, S10 2RX, UK. Tel. + 44 114 215 9560. Email:
33 H.L.Wilson@sheffield.ac.uk
34

35
36 Siew Cheng Wong: Singapore Immunology Network (SIgN), Agency for Science, Technology and
37 Research (A*STAR), 8A Biomedical Grove, Immunos, Level 4, Singapore 138648. Tel:
38 +65 6407 0030. Email: wong_siew_cheng@immunol.a-star.edu.sg
39
40

41
42 Category: Original Research Paper / Basic Research Paper.
43

44
45 Keywords: Inflammation, Atherosclerosis, Macrophage, Innate
46

47 Total word count (main text, excluding references, including Figure legends): 3,940.
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 **(ii) Abstract**

5 *Background & Aims:* Lipid-rich foam cell macrophages drive atherosclerosis via several mechanisms,
6 including inflammation, lipid uptake, lipid deposition and plaque vulnerability. The atheroma
7 environment shapes macrophage function and phenotype; anti-inflammatory macrophages improve
8 plaque stability while pro-inflammatory macrophages promote rupture. Current evidence suggests a
9 variety of macrophage phenotypes occur in atherosclerotic plaques with local lipids, cytokines,
10 oxidised phospholipids and pathogenic stimuli altering their phenotype. In this study, we addressed
11 differential functioning of macrophage phenotypes via a systematic analysis of *in vitro* polarised,
12 human monocyte-derived macrophage phenotypes, focussing on molecular events that regulate foam-
13 cell formation.
14

15 *Methods:* We examined transcriptomes, protein levels and functionally determined lipid handling and
16 foam cell formation capacity in macrophages polarised with IFN γ +LPS, IL-4, IL-10, oxPAPC and
17 CXCL4.
18

19 *Results:* RNA sequencing of differentially polarised macrophages revealed distinct gene expression
20 changes, with enrichment in atherosclerosis and lipid-associated pathways. Analysis of lipid
21 processing activity showed IL-4 and IL-10 macrophages have higher lipid uptake and foam cell
22 formation activities, while inflammatory and oxPAPC macrophages displayed lower foam cell
23 formation. Inflammatory macrophages showed low lipid uptake, while higher lipid uptake in oxPAPC
24 macrophages was matched by increased lipid efflux capacity.
25

26 *Conclusions:* Atherosclerosis-associated macrophage polarisation dramatically affects lipid handling
27 capacity underpinned by major transcriptomic changes and altered protein levels in lipid-handling gene
28 expression. This leads to phenotype-specific differences in LDL uptake, cellular cholesterol levels and
29 cholesterol efflux, informing how the plaque environment influences atherosclerosis progression by
30 influencing macrophage phenotypes.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1. Introduction

Arterial macrophages form disease-associated lipid-rich foam cells in atherosclerotic plaques [1]. Progressive atherosclerosis results in cardiovascular diseases, which are a leading cause of death, worldwide [2]. Plaque formation arises from the accumulation of lipid in the artery wall over decades; acute myocardial infarction or stroke can occur following a loss of plaque stability due to cellular changes in the artery wall [3]. Macrophages are a predominant cell type within the artery wall, whose major functions in the plaque are to regulate inflammation, clear apoptotic cells via efferocytosis, regulate plaque stability by secreting matrix metalloproteases (MMPs) or protease inhibitors and to uptake, process, accumulate and efflux lipid [4]. The capacity of macrophages to accumulate or process lipid and to regulate proteases and cytokines in their local environment is a critical determinant of plaque development, lipid core formation and plaque stability [5,6].

Macrophages regulate lipoproteins within their local environment [7]. Uptake of lipoproteins occurs via receptor-mediated endocytosis, such as scavenger receptor, pinocytosis and phagocytosis [4]. Macrophages express scavenger receptors including SR-A1, CD36 and LOX-1 for lipoprotein uptake [8], while phagocytosis mediated internalisation delivers lipids to late endolysosomes where lysosomal acid lipase (LAL) digests cholesteryl esters to produce free cholesterol [9]. In the case of lipoprotein, cellular lipases such as LAL break down the lipoprotein particles that have been taken up, to cholesterol, triglycerides and fatty acids [10–12]. Cholesteryl esters and acylglycerides can be stored in lipid droplets, resulting in foam cell formation. Alternatively, Neutral cholesterol ester hydrolase (NCEH1) can metabolise these molecules into free cholesterol [13–15]; this can be removed from the cell by efflux or converted to fatty acids (oxidised to produce ATP) and glycerol [14,15]. Sterol O-acyltransferase 1 (SOAT1) in the endoplasmic reticulum can re-esterify free cholesterol into cholesterol ester or fatty acids into acylglycerides, for subsequent storage in lipid droplets [4,16–18]. Cholesterol, produced by NCEH1, is then effluxed from the cell by transporters ABCA1, ABCG1 and SR-BI [19] or re-esterified and stored as lipid droplets [19]. The build-up of cellular cholesterol activates transcription factors RXR, LXR α , LXR β , PPAR α and PPAR γ [20]. Heterodimerisation of LXR and RXR increases protein levels of ABCA1 and ABCG1, which actively transport HDL loaded-cholesterol out of the cell [19]. The cytosolic accumulation of esterified cholesterol and acylglyceride droplets in macrophages results in the generation of foam cells, driving atherosclerotic plaque formation [9]. In addition to lipid uptake and intracellular processing, lipid and cholesterol content are influenced by the cellular capacity for cholesterol efflux. These efflux pathways in macrophages function to remove excess lipid to prevent its accumulation, which can be cytotoxic [21–23].

The local cellular environment determines macrophage function and phenotype [24]. Traditionally, macrophages have been described as classically activated M1 and alternatively activated M2 macrophages [24]. Pro-inflammatory M1 macrophages arise from exposure to bacterial LPS and interferon-gamma [25]. Cytokines such as IL-4 and IL-13 drive polarisation to M2 macrophages [26]. In arterial atherosclerotic plaques, pro-inflammatory macrophages localise to areas of increased vulnerability, such as the shoulder regions [27]. In these vulnerable areas, they promote rupture by secreting MMPs and inflammatory cytokines thus weakening the cap structure [27]. In contrast, anti-inflammatory macrophages are associated with increased plaque stability, at times with plaque regression and less severe disease [1]. The traditional classification of M1 and M2 macrophages is now widely understood to be simplistic where other “intermediate” or differently polarised phenotypes exist according to the local environment [25]. Macrophages expressing markers for both M1 and M2 phenotypes have been labelled within atherosclerotic plaques [25]. The plaque environment contains

1
2
3
4 oxidised phospholipid [28], platelet-derived CXCL4 [29] and IL-10 [26] as well as haemoglobin [30],
5 which can each influence macrophages to polarise to different functional states.
6

7
8 The accumulation of lipid within the macrophage, and propensity to form foam cells, is altered
9 according to macrophage phenotype [31]. Pro-inflammatory stimuli have been reported to alter
10 expression of some scavenger receptors [32] [33] and to reduce cholesterol efflux transporter
11 expression [34], reducing their capacity to clear lipid from the plaque. In contrast, IL-4 macrophages
12 are considered to have high lipid handling and foam cell formation capacities [35]. The effect of IL-
13 10, CXCL4 and oxidised phospholipid on human macrophage lipid handling is less well understood.
14
15

16
17 Compelling evidence shows that a variety of macrophage phenotypes occur in atherosclerotic plaques
18 dependent on their environment. However, it is noteworthy that the majority of our current
19 understanding of macrophage polarisation and its impact on atherosclerosis derives from murine
20 models of atherosclerosis and analysis of mouse bone marrow derived macrophages [36,37]. The
21 relevance to some of these findings to atherogenesis in humans is less clear. For example, whilst *Nos2*
22 and *Arg1* expression are robust markers of proinflammatory vs. alternatively activated macrophages,
23 expression of these genes does not correlate with such macrophage subsets in humans [38–40]. In
24 order to address how the variety of plaque macrophage phenotypes function differentially, we
25 undertook a systematic study to model human macrophages polarised towards factors relevant to the
26 atherosclerotic plaque environment. Using human samples, we sequenced and analysed whole
27 transcriptomes, protein levels and functionally determined lipid handling and foam cell formation
28 capacity in human macrophages polarised with IFN γ +LPS, IL-4, IL-10, oxPAPC (oxidised
29 phospholipid 1-palmitoyl-2-arachidonyl-sn-glycero-3-phosphorylcholine macrophages) and CXCL4.
30 Our data revealed transcriptionally distinct phenotypes, differential lipid uptake, processing and efflux
31 capacities and functional differences to account for low foam cell formation in differentially polarised
32 inflammatory macrophages.
33
34
35
36

37 **2. Materials and methods**

38 Detailed methods are described in supplementary information. In brief, peripheral blood mononuclear
39 cells (PBMCs) were isolated from whole blood donated by healthy adult donors and monocytes
40 purified by CD14 positive magnetic selection. Monocytes were differentiated into monocyte-derived
41 macrophages (MDMs) over 7 days in M-CSF, resulting in CD68+ macrophages expressing receptors
42 for polarising agents (Supplementary methods; Supplementary Table 1). On day 7, the media was
43 replaced for 24 h with fresh complete media containing the following polarising agents: IFN γ and LPS
44 (R515); IL-4; IL-10; oxPAPC; CXCL4; or unpolarised as controls in each experiment (Supp.Fig.1).
45 RNA was extracted from unpolarised and polarised MDM from 8 separate donors for RNA-seq
46 analysis. RNA-seq data is deposited at DOI: 10.17632/j2hmt7k9fh.1. Differential gene expression was
47 validated by qPCR, and protein by immunofluorescence and flow cytometry. Functional changes in
48 foam cell formation and lipid handling were assessed by oil-red-o lipid staining, acLDL uptake,
49 cholesterol content and efflux determination. Replicate values (n) represent separate human donors.
50
51
52
53

54 **3. Results**

55 **Polarisation causes atherosclerosis-related changes in MDM transcriptomes**

56
57 We set out to systematically extend the current knowledge of transcriptional changes associated with
58 macrophage polarisation by assessing gene expression changes specifically in *human* macrophage
59
60
61
62
63
64
65

1
2
3
4 phenotypes arising from the same donors, treated under a variety of conditions reflecting the arterial
5 plaque environment. Monocyte-derived macrophages from 8 donors were polarised with IFN γ +LPS,
6 IL-4, IL-10, oxPAPC or CXCL4 and examined by RNA-seq for global and specific pathway changes
7 in their transcriptomes that may influence atherosclerosis progression.
8

9 Principal component analysis (PCA) clearly showed M^{IFN γ +LPS} as the most transcriptionally
10 distinct phenotype with > 5,500 differentially expressed genes (DEGs) compared to unpolarised
11 macrophages and to other phenotypes (Figure 1A). M^{IFN γ +LPS} displayed unique and high upregulation
12 of common pro-inflammatory markers (Supplementary Tables 2 and 3). M^{IL-4} was the next most
13 transcriptionally distinct phenotype as evidenced by the PCA, and characterised by *IRF4* (interferon
14 regulatory factor 4) and *MRC1* (mannose receptor C-type 1) upregulation. The PCA shows that
15 M^{IFN γ +LPS} and M^{IL-4} represent the extreme ends of the *in vitro* MDM polarisation spectrum, since they
16 exhibited the highest number of differentially expressed genes amongst the polarisation conditions
17 applied. Interestingly, in the PCA M^{IL-10}, M^{oxPAPC} and M^{CXCL4} clustered more closely to unpolarised
18 macrophages. Of note, M^{IL-10}, M^{oxPAPC} and M^{CXCL4} also showed a cytokine/chemokine (and their
19 regulators) expression profile, similar to M^{IL-4}. These included a notably lower upregulation of *CXCL8*
20 (*IL-8*), *CXCL9*, *CXCL11*, *IL1B*, *IL2RA* and *SOCS3* compared to M^{IFN γ +LPS}, but no changes in *IRF1*,
21 *NFKB1*, *NFKB2*, *RELA* and *RELB* expression (Supplementary Table 4). M^{IL-10} showed upregulation
22 of *CD163* RNA, consistent with recent findings [41], while M^{oxPAPC} was clearly distinguished by
23 increased RNA for *HMOX1* (heme oxygenase 1) and *TXNRD1* (thioredoxin reductase 1).
24
25

26 CXCL4 macrophages showed a remarkably similar transcriptome profile to unpolarised
27 macrophages as indicated by their close clustering in PCA and only 207 DEGs (1.53% of transcripts
28 detected from a total of 13,531 genes) between the two conditions (Figure 1B). These results were
29 largely in agreement with transcriptomic data on human monocyte-derived macrophages with and
30 without CXCL4 polarisation from another study[42], where monocyte differentiation into unpolarised
31 macrophages and M^{CXCL4} with M-CSF or CXCL4 respectively over 6 days resulted in only 460
32 differentially expressed probes or 375 DEGs (1.77% out of 26,051 probes with signals above the
33 detection limit [42]). *S100A8* (*S100 calcium binding protein A8*) and *MMP7* (*matrix metalloproteinase*
34 *7*) have previously been reported as M^{CXCL4} markers [43]; our RNA-seq data showed their expression
35 in M^{CXCL4} was highly variable among donors and was up-regulated in most other phenotypes,
36 especially M^{IFN γ +LPS} (Supplementary Tables 2 and 3). Given this low difference between M^{CXCL4} and
37 unpolarised macrophage transcriptomes, we omitted M^{CXCL4} from further analysis (Figure 1C, 1D).
38
39

40 Each MDM phenotype displayed significant enrichment of DEGs in atherosclerosis-related
41 pathways (Supplementary Table 5). In most phenotypes, these pathways included genes clearly
42 involved in macrophage lipid handling, such as *ABCA1* (ATP binding cassette subfamily A member
43 1), *ABCG1* (ATP binding cassette subfamily G member 1), *LPL* (lipoprotein lipase) and *MSR1*
44 (macrophage scavenger receptor 1) prompting further investigation into the lipid handling capacity of
45 these phenotypes. The RNA-sequencing results were validated by qRT-PCR on parallel samples
46 (Supplementary Figure 3), confirming differences in selected atherosclerosis-related genes.
47
48
49
50

51 **Reduced foam cell formation capacity in M^{IFN γ +LPS} and M^{oxPAPC}**

52 Our transcriptome analyses of macrophage phenotypes revealed differences in lipid handling and other
53 associated pathway enrichment for differentially expressed genes (Supplementary Table 5). We
54 therefore sought to assess foam cell formation capacity of the various polarised macrophages since
55 this underlies the pathophysiology of atherosclerosis [44]. After loading with acLDL, we measured
56 foam cell formation by oil-red-O staining (measured as % control – % acLDL-loaded Area ORO >
57
58
59
60
61
62
63
64
65

1
2
3
4 Area haematoxylin cells) in $M^{IFN\gamma+LPS}$ and M^{oxPAPC} , while M^{IL-4} and M^{IL-10} showed an increase in foam
5 cell formation, similar to that of unpolarised macrophages (Figures 2A & 2B).
6
7

8 **Reduced LDL uptake in $M^{IFN\gamma+LPS}$**

9 Since $M^{IFN\gamma+LPS}$ and M^{oxPAPC} showed reduced foam cell formation compared to other phenotypes, we
10 assessed lipoprotein uptake, which impacts macrophage lipid handling and resultant foam cell
11 formation [45]. We therefore measured fluorescently-labelled acLDL uptake in the different MDM
12 phenotypes by flow cytometry. Analysis of MDMs exposed to AlexaFluor-488-conjugated acLDL
13 showed significantly lower uptake in $M^{IFN\gamma+LPS}$ and non-significant reduction in M^{oxPAPC} ($p = 0.10$). In
14 contrast, M^{IL-4} and M^{IL-10} showed intracellular acLDL fluorescence levels similar to unpolarised
15 macrophages (Figure 3A). **Using reactome.org analysis, in $M^{IFN\gamma+LPS}$ we detected a global down-**
16 **regulation of ‘Ligand binding and uptake by scavenger receptors’ pathways** (Figure 3B) and in key
17 lipid uptake genes (Figure 3C), such as modified LDL receptors *CD36* (cluster of differentiation 36)
18 and *MSR1* (macrophage scavenger receptor 1) [46,47]. This suggests that differential polarisation
19 influences a transcriptional programme that affects downstream pathways regulating LDL uptake. **In**
20 **agreement with this, a reduction in both CD36 and MSR1 scavenger receptors was detected at the cell**
21 **surface protein level in $M^{IFN\gamma+LPS}$ (Figure 3D).**
22
23
24
25

26 **Cellular free cholesterol content in polarised macrophages after LDL loading**

27 In addition to lipid uptake, macrophage foam cell formation capacity is influenced by the ability of the
28 cell to regulate its internal lipid **content**. **Altering** the balance of internal lipid processing and storage
29 may result in changes in total, free and esterified cholesterol content in macrophage phenotypes. While
30 **$IFN\gamma$ -treated macrophages** have been shown to contain different levels of free cholesterol and its ester
31 [48], the other phenotypes described in this study have not been assessed in this regard. We therefore
32 measured the abundance of total, free and esterified cholesterol in control and acLDL-loaded cells to
33 determine if intracellular cholesterol processing capacity affected differences in foam cell formation.
34
35

36 Unpolarised macrophages showed a larger change in total cholesterol (73.5 ± 19.7 ng
37 cholesterol / ng protein), higher than for all polarisation-treated macrophage phenotypes (Figure 4A).
38 $M^{IFN\gamma+LPS}$ exhibited no significant change in total cholesterol following loading ($p = 0.10$ for the
39 difference between this condition and unpolarised cells, Figure 4A). We also calculated the cholesterol
40 ester content, but did not observe statistically significant differences in cholesterol ester by phenotype,
41 when compared to unpolarised macrophages (Figure 4B). Similar to the total cholesterol changes,
42 $M^{IFN\gamma+LPS}$ showed no significant differences ($p = 0.11$) in the change of free cholesterol content (Figure
43 4C).
44
45

46 We examined the ‘LDL clearance’ pathway for DEGs according to MDM polarisation, to
47 identify transcriptional differences in the cholesterol clearance pathways. In agreement with the
48 cholesterol content determination, $M^{IFN\gamma+LPS}$ showed largely down-regulated expression of the entire
49 clearance pathway (Figure 4D). Among key genes, *SOAT1* was not differentially expressed in any
50 phenotype, whereas, expression of *NCEH1* was significantly reduced in $M^{IFN\gamma+LPS}$ (Figure 4E), in
51 keeping with the cholesterol content assessment. In addition, *LIPA* (lipase A, lysosomal acid type,
52 involved in lipoprotein particle breakdown) showed reduced expression in $M^{IFN\gamma+LPS}$ and increased in
53 M^{IL-4} , likely contributing to the differences in intracellular lipid handling between these phenotypes
54 (Figure 4E).
55
56
57

58 **Reduced cholesterol efflux capacity in $M^{IFN\gamma+LPS}$**

1
2
3
4 **Lipid and cholesterol content are influenced by the cellular capacity for cholesterol efflux.**
5 Macrophages mainly unload intracellular lipid to apolipoprotein AI (Apo-AI) via ABCA1 transporter
6 while high-density lipoprotein (HDL) is capable of accepting lipid from ABCG1, SCARB1 (scavenger
7 receptor class B member 1) [49] and other pathways. Previous reports indicated that pro-inflammatory
8 macrophages had lower cholesterol efflux capacity [34] than anti-inflammatory macrophages [50]. We
9 tested cholesterol efflux via both acceptors, 24 h after macrophage polarisation, to determine the
10 reverse cholesterol transport capacity of each phenotype.
11
12

13 Total TopFluor cholesterol efflux, involving both Apo-AI and HDL, was significantly reduced
14 in $M^{IFN\gamma+LPS}$ while M^{IL-4} did not show a significantly altered efflux compared to unpolarised
15 macrophages (Figure 5A). $M^{IFN\gamma+LPS}$ also showed a significantly lower capacity to efflux cholesterol
16 regardless of whether HDL or Apo-AI was used as cholesterol acceptor (Figure 5B and C). M^{IL-4}
17 showed no significant difference in cholesterol efflux capacity in the presence of Apo-AI compared
18 to unpolarised macrophages (Figure 5C). Following polarisation, expression of key active cholesterol
19 efflux receptors ABCA1 and ABCG1 in ‘HDL assembly’ and ‘HDL remodelling’ pathways (Figure
20 5D) showed an opposite pattern (Figure 6E) from the downstream functional data (Figure 5A–C).
21 Expression of the major passive cholesterol efflux receptor SCARB1 (functions via HDL) was
22 significantly lower in $M^{IFN\gamma+LPS}$, but also uniquely increased in M^{IL-4} (Figure 5E), suggesting that
23 expression of SCARB1 and the aforementioned ABC transporters are under different transcriptional
24 regulation for lipid handling, post-polarisation. Protein **levels** of ABCA1 and ABCG1 upon MDM
25 polarisation (Figure 5F) were similar to the cholesterol efflux capacities of each of the macrophage
26 phenotypes (Figure 5A–C). SCARB1 cell-surface protein was uniquely up-regulated in M^{IL-4} ,
27 matching our observations at the transcript level and further suggesting differences between regulation
28 of ABCA1/ABCG1 and SCARB1 expression.
29
30
31
32
33
34

35 **4. Discussion**

36
37 Macrophages polarise to a spectrum of phenotypes in response to their environment [24]. This
38 systematic study compared human macrophage lipid handling capacities at the transcriptome, protein
39 and functional levels for cells polarised towards factors relevant to the atherosclerotic plaque
40 environment. Polarisation with $IFN\gamma+LPS$, $IL-4$, $IL-10$, $oxPAPC$ and $CXCL4$ identified distinct
41 transcriptional changes, $M^{IFN\gamma+LPS}$ being the most transcriptionally distinct phenotype in agreement
42 with other related studies [51,52]. Transcriptional changes showed enrichment in atherosclerosis and
43 lipid-associated pathways. Analysis of lipid processing activity showed M^{IL-4} and M^{IL-10} to have
44 higher lipid uptake and foam cell formation activities, while inflammatory $M^{IFN\gamma+LPS}$ and M^{oxPAPC} have
45 lower foam cell activity. Functionally, there are key differences between the $M^{IFN\gamma+LPS}$ and M^{oxPAPC}
46 inflammatory macrophage phenotypes that account for their low foam cell formation: $M^{IFN\gamma+LPS}$ exhibit
47 low lipid uptake while a higher lipid uptake in M^{oxPAPC} is compensated by increased lipid efflux
48 activity.
49
50
51
52

53 Lipid handling by macrophages is clearly critical in atherogenesis. Thus, we also considered using
54 lipids and lipoprotein particles as stimulating agents alone or in combination with the polarising factors
55 we used to treat MDMs. However, our own pilot data (unpublished) and previous work has shown that
56 lipid exposure of MDMs does not lead to notable transcriptomic changes or modify the impact of the
57 polarising stimuli used in this study [52]. We therefore considered this as unlikely to provide additional
58 insight beyond the current analysis without complicating data interpretation.
59
60
61
62
63
64
65

1
2
3
4
5
6 While previous studies, and more frequently in murine models, have compared macrophage
7 polarisation in response to specific stimuli, we undertook a systematic approach to directly compare
8 the impact of IFN γ +LPS, IL-4, IL-10, oxPAPC and CXCL4 in human macrophages at the level of
9 transcriptomes, protein through to functional lipid processing analysis. The analysis using 8 individual
10 donors showed a clear distinction in transcriptional programming due to polarisation. All phenotypes
11 showed a unique expression profile compared to unpolarised controls with exception of CXCL4
12 polarisation, similar to work by Gleissner *et al* [42]. We noted that only a subset of the differentiated
13 macrophages expressed CXCR3, the receptor for CXCL4, in all donors (46.3 ± 16.3 %). This may
14 well have contributed to the minimal transcriptomic difference in this phenotype compared to
15 unpolarised macrophages. Consistent with this, *MMP7* and *S100A8*, which are considered markers for
16 the CXCL4 phenotype [43], were not upregulated consistently in all donors, again suggesting
17 incomplete polarisation.
18
19
20

21 For the IFN γ +LPS, IL-4, IL-10 and oxPAPC polarisation conditions, the largest divergence in DEGs
22 was between M^{IFN γ +LPS} and M^{IL-4} polarisation, in keeping with other findings [51]. DEG enrichment
23 in atherosclerosis and lipid associated IPA canonical pathways suggests a logical rationale for studying
24 differences in lipid handling between macrophage phenotypes.
25
26

27 The lipid handling capacities were similar for M^{IL-4}, M^{IL-10} and unpolarised macrophages where each
28 displayed efficient foam cell formation and acLDL uptake abilities alongside increased expression of
29 *MSRI*, *CD36*, *SOAT1*, *NCEH1*, *LIPA* and other key genes in these scavenger receptor and LDL
30 clearance pathways. At earlier stages of atherogenesis, M^{IL-4} and M^{IL-10} may prevent atherosclerosis
31 progression by clearing lipid from within the artery wall due to their effective lipid processing capacity
32 alongside effective efflux [4]. Our oil-red-O measurements showed high foam cell formation in these
33 macrophages which may reflect a role in promoting atherosclerosis at later stages of disease where
34 high levels of LDL and other lipids may overwhelm their capacity to process extracellular lipid[20,53].
35 The cholesterol transporter SCARB1 plays a role in both cholesterol efflux and influx[54]. At the
36 protein level, SCARB1 was most highly expressed in M^{IL-4} compared to other polarisation states while
37 ABCA1 and ABCG1 were significantly increased in both M^{IL-4} and M^{IL-10}, but not to the same extent
38 as SCARB1. It is likely that post-transcriptional regulation accounts for differences in all of these
39 cholesterol transporters since protein changes were more pronounced and not always in the same
40 direction to that of the mRNA levels. These findings suggest that SCARB1, in combination with
41 MRC1, offers the potential as an additional marker for the M^{IL-4} phenotype given its level of
42 upregulation. Increased SCARB1 in M^{IL-4} compared to M^{IFN γ +LPS} likely accounts for their higher
43 cholesterol efflux capacity.
44
45
46
47
48

49 A pro-inflammatory and oxidised atherosclerotic plaque environment is associated with later stage
50 disease, plaque instability and vulnerability to rupture [31]. Infection is a risk factor for myocardial
51 infarction[55]while LPS has been detected within human plaques[56]. M^{IFN γ +LPS} and M^{oxPAPC}
52 phenotypes can arise due to differences in sterile inflammatory stimuli where the effect of oxidised
53 phospholipids on macrophage lipid metabolism is less well understood, particularly in human
54 macrophages. Oxidised phospholipids, derived from oxidation of lipoproteins and apoptotic cell
55 membranes, induce TLR2-mediated acute inflammatory responses in macrophages [57]. Some reports
56 of M^{oxPAPC} polarised mouse macrophages have shown that the phagocytic activity of these
57 macrophages is impaired [57]. Interestingly, in this study we found that M^{oxPAPC} showed effective lipid
58
59
60
61
62
63
64
65

1
2
3
4 uptake and processing, similar to the levels observed in M^{IL-4} and M^{IL-10} macrophages. At the same
5 time, M^{oxPAPC} showed low foam cell formation, which is likely accounted for by their high capacity to
6 efflux cholesterol. M^{oxPAPC} showed lower expression of *ABCG1* and *SCARB1* compared to M^{IL-4} and
7 M^{IL-10} suggesting that efflux may be coupled to other transporters or they can process lipids more
8 efficiently due to other regulatory factors. In contrast, M^{IFN γ +LPS} showed lower acLDL uptake
9 compared to other phenotypes, accompanied by low foam cell formation.

10
11
12 Atherosclerosis associated macrophage polarisation dramatically affects the cell's capacity to handle
13 lipid. Whole transcriptome sequencing of differentially polarised macrophages revealed distinct
14 phenotypes where differences in key lipid handling gene and protein levels have a lasting effect on
15 downstream lipid handling, as observed by differences in LDL uptake, cellular cholesterol levels and
16 the capacity to efflux excess cellular cholesterol. Functional differences account for low foam cell
17 formation in differentially polarised inflammatory macrophages: M^{IFN γ +LPS} exhibit low lipid uptake
18 while increased lipid uptake in M^{oxPAPC} is matched by increased lipid efflux activity. All of these
19 processes influence the formation of lipid-laden macrophage foam cells, underlying the progression
20 of atherosclerosis and vulnerability to cardiovascular disease.

21 22 23 24 25 **Funding**

26 This work was supported by the Biomedical Research Council (BMRC), Agency for Science,
27 Technology and Research (A*STAR) Research Attachment Programme to HLW, EKT and SCW to
28 support EH and KB. BL and JL are part of the SIgN Immunomonitoring platform (supported by a
29 BMRC IAF 311006 grant and BMRC transition funds #H16/99/b0/011).

30 31 32 33 **Acknowledgements**

34 We thank Sarah Lai for data analysis support. We thank Jon Kilby for technical support in monocyte
35 isolations.

36 37 38 **Author contributions:**

39 Conception and design: HLW, SCW, EKT. Performed experiments: KB, EH, JL, FS. Experimental
40 analysis: KB, EH, BL, IS. Interpretation: KB, EH, HLW, EKT, SCW. Wrote manuscript: HLW, KB.
41 Edited manuscript: EH, BL, JL, EKT, SCW.

42 43 44 45 **Conflict of Interest.**

46 No conflicts of interest.

47 48 49 **References**

50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
- 2
- 3
- 4
- 5
- 6
- 7 [1] Tabas, I, Bornfeldt, KE, Macrophage Phenotype and Function in Different Stages
- 8 of Atherosclerosis. *Circ Res* 2016;118(4):653–67.
- 9 [2] Nascimento, BR, Brant, LC, Moraes, DN et al., Global health and cardiovascular
- 10 disease. *Heart* 2014;100(22):1743–9.
- 11 [3] Kavurma, MM, Rayner, KJ, Karunakaran, D, The walking dead: macrophage
- 12 inflammation and death in atherosclerosis. *Curr Opin Lipidol* 2017;28(2):91–8.
- 13 [4] Moore, KJ, Sheedy, FJ, Fisher, EA, Macrophages in atherosclerosis: a dynamic
- 14 balance. *Nat Rev Immunol* 2013;13(10):709–21.
- 15 [5] de Gaetano, M, Crean, D, Barry, M et al., M1- and M2-Type Macrophage
- 16 Responses Are Predictive of Adverse Outcomes in Human Atherosclerosis. *Front*
- 17 *Immunol* 2016;7:275.
- 18 [6] Jinnouchi, H, Guo, L, Sakamoto, A et al., Diversity of macrophage phenotypes and
- 19 responses in atherosclerosis. *Cell Mol Life Sci* 2019
- 20 [7] Bories, GFP, Leitinger, N, Macrophage metabolism in atherosclerosis. *FEBS Lett*
- 21 2017;591(19):3042–60.
- 22 [8] Johnston, JM, Angyal, A, Bauer, RC et al., Myeloid Tribbles 1 induces early
- 23 atherosclerosis via enhanced foam cell expansion. *Sci Adv* 2019;5(10):eaax9183.
- 24 [9] Chistiakov, DA, Melnichenko, AA, Myasoedova, VA et al., Mechanisms of foam cell
- 25 formation in atherosclerosis. *J Mol Med (Berl)* 2017;95(11):1153–65.
- 26 [10] Huang, SC, Everts, B, Ivanova, Y et al., Cell-intrinsic lysosomal lipolysis is
- 27 essential for alternative activation of macrophages. *Nat Immunol* 2014;15(9):846–
- 28 55.
- 29 [11] Sheriff, S, Du, H, Grabowski, GA, Characterization of lysosomal acid lipase by site-
- 30 directed mutagenesis and heterologous expression. *J Biol Chem*
- 31 1995;270(46):27766–72.
- 32 [12] Vargas-Alarcón, G, Posadas-Romero, C, Villarreal-Molina, T et al., Single
- 33 nucleotide polymorphisms within LIPA (Lysosomal Acid Lipase A) gene are
- 34 associated with susceptibility to premature coronary artery disease. a replication in
- 35 the genetic of atherosclerotic disease (GEA) Mexican study. *PLoS One*
- 36 2013;8(9):e74703.
- 37 [13] Igarashi, M, Osuga, J, Uozaki, H et al., The critical role of neutral cholesterol ester
- 38 hydrolase 1 in cholesterol removal from human macrophages. *Circ Res*
- 39 2010;107(11):1387–95.
- 40 [14] Zhao, B, Song, J, Chow, WN et al., Macrophage-specific transgenic expression of
- 41 cholesteryl ester hydrolase significantly reduces atherosclerosis and lesion
- 42 necrosis in Ldlr mice. *J Clin Invest* 2007;117(10):2983–92.
- 43 [15] Zhao, B, Song, J, St Clair, RW et al., Stable overexpression of human macrophage
- 44 cholesteryl ester hydrolase results in enhanced free cholesterol efflux from human
- 45 THP1 macrophages. *Am J Physiol Cell Physiol* 2007;292(1):C405–12.
- 46 [16] Brown, MS, Ho, YK, Goldstein, JL, The cholesteryl ester cycle in macrophage
- 47 foam cells. Continual hydrolysis and re-esterification of cytoplasmic cholesteryl
- 48 esters. *J Biol Chem* 1980;255(19):9344–52.
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

- 1
2
3
4 [17] Ghosh, S, Macrophage cholesterol homeostasis and metabolic diseases: critical
5 role of cholesteryl ester mobilization. *Expert Rev Cardiovasc Ther* 2011;9(3):329–
6 40.
7
8 [18] Yang, L, Yang, JB, Chen, J et al., Enhancement of human ACAT1 gene
9 expression to promote the macrophage-derived foam cell formation by
10 dexamethasone. *Cell Res* 2004;14(4):315–23.
11
12 [19] Remmerie, A, Scott, CL, Macrophages and lipid metabolism. *Cell Immunol*
13 2018;330:27–42.
14 [20] Chawla, A, Boisvert, WA, Lee, CH et al., A PPAR gamma-LXR-ABCA1 pathway in
15 macrophages is involved in cholesterol efflux and atherogenesis. *Mol Cell*
16 2001;7(1):161–71.
17
18 [21] Ghosh, J, Guha, R, Das, S et al., Liposomal cholesterol delivery activates the
19 macrophage innate immune arm to facilitate intracellular *Leishmania donovani*
20 killing. *Infect Immun* 2014;82(2):607–17.
21
22 [22] Tabas, I, Consequences of cellular cholesterol accumulation: basic concepts and
23 physiological implications. *J Clin Invest* 2002;110(7):905–11.
24
25 [23] Warner, GJ, Stoudt, G, Bamberger, M et al., Cell toxicity induced by inhibition of
26 acyl coenzyme A:cholesterol acyltransferase and accumulation of unesterified
27 cholesterol. *J Biol Chem* 1995;270(11):5772–8.
28
29 [24] Sica, A, Erreni, M, Allavena, P et al., Macrophage polarization in pathology. *Cell*
30 *Mol Life Sci* 2015;72(21):4111–26.
31
32 [25] Chistiakov, DA, Bobryshev, YV, Nikiforov, NG et al., Macrophage phenotypic
33 plasticity in atherosclerosis: The associated features and the peculiarities of the
34 expression of inflammatory genes. *Int J Cardiol* 2015;184:436–45.
35
36 [26] Gordon, S, Martinez, FO, Alternative activation of macrophages: mechanism and
37 functions. *Immunity* 2010;32(5):593–604.
38
39 [27] Stöger, JL, Gijbels, MJ, van der Velden, S et al., Distribution of macrophage
40 polarization markers in human atherosclerosis. *Atherosclerosis* 2012;225(2):461–
41 8.
42
43 [28] Que, X, Hung, MY, Yeang, C et al., Oxidized phospholipids are proinflammatory
44 and proatherogenic in hypercholesterolaemic mice. *Nature* 2018;558(7709):301–6.
45
46 [29] Gleissner, CA, Macrophage Phenotype Modulation by CXCL4 in Atherosclerosis.
47 *Front Physiol* 2012;3:1.
48
49 [30] Boyle, JJ, Johns, M, Kampfer, T et al., Activating transcription factor 1 directs
50 Mhem atheroprotective macrophages through coordinated iron handling and foam
51 cell protection. *Circ Res* 2012;110(1):20–33.
52
53 [31] Cochain, C, Zerneck, A, Macrophages in vascular inflammation and
54 atherosclerosis. *Pflugers Arch* 2017;469(3-4):485–99.
55
56 [32] Canton, J, Neculai, D, Grinstein, S, Scavenger receptors in homeostasis and
57 immunity. *Nat Rev Immunol* 2013;13(9):621–34.
58
59 [33] Hashizume, M, Mihara, M, Atherogenic effects of TNF- α and IL-6 via up-regulation
60 of scavenger receptors. *Cytokine* 2012;58(3):424–30.
61
62 [34] Panousis, CG, Zuckerman, SH, Interferon-gamma induces downregulation of
63 Tangier disease gene (ATP-binding-cassette transporter 1) in macrophage-derived
64 foam cells. *Arterioscler Thromb Vasc Biol* 2000;20(6):1565–71.
65

- 1
2
3
4 [35] van Tits, LJ, Stienstra, R, van Lent, PL et al., Oxidized LDL enhances pro-
5 inflammatory responses of alternatively activated M2 macrophages: a crucial role
6 for Krüppel-like factor 2. *Atherosclerosis* 2011;214(2):345–9.
7
8 [36] Johnston, JM, Francis, SE, Kiss-Toth, E, Experimental models of murine
9 atherosclerosis: does perception match reality. *Cardiovasc Res*
10 2018;114(14):1845–7.
11
12 [37] Zhang, X, Li, J, Luo, S et al., IgE Contributes to Atherosclerosis and Obesity by
13 Affecting Macrophage Polarization, Macrophage Protein Network, and Foam Cell
14 Formation. *Arterioscler Thromb Vasc Biol* 2020;40(3):597–610.
15
16 [38] Murray, HW, Teitelbaum, RF, L-arginine-dependent reactive nitrogen intermediates
17 and the antimicrobial effect of activated human mononuclear phagocytes. *J Infect*
18 *Dis* 1992;165(3):513–7.
19
20 [39] Schneemann, M, Schoedon, G, Species differences in macrophage NO production
21 are important. *Nat Immunol* 2002;3(2):102.
22
23 [40] Young, R, Bush, SJ, Lefevre, L et al., Species-Specific Transcriptional Regulation
24 of Genes Involved in Nitric Oxide Production and Arginine Metabolism in
25 Macrophages. *Immunohorizons* 2018;2(1):27–37.
26
27 [41] Thornton, S, Tan, R, Sproles, A et al., A Multiparameter Flow Cytometry Analysis
28 Panel to Assess CD163 mRNA and Protein in Monocyte and Macrophage
29 Populations in Hyperinflammatory Diseases. *J Immunol* 2019;202(5):1635–43.
30
31 [42] Gleissner, CA, Shaked, I, Little, KM et al., CXC chemokine ligand 4 induces a
32 unique transcriptome in monocyte-derived macrophages. *J Immunol*
33 2010;184(9):4810–8.
34
35 [43] Erbel, C, Tyka, M, Helmes, CM et al., CXCL4-induced plaque macrophages can be
36 specifically identified by co-expression of MMP7+S100A8+ in vitro and in vivo.
37 *Innate Immun* 2015;21(3):255–65.
38
39 [44] Chistiakov, DA, Bobryshev, YV, Orekhov, AN, Macrophage-mediated cholesterol
40 handling in atherosclerosis. *J Cell Mol Med* 2016;20(1):17–28.
41
42 [45] Sorci-Thomas, MG, Thomas, MJ, Microdomains, Inflammation, and
43 Atherosclerosis. *Circ Res* 2016;118(4):679–91.
44
45 [46] Kunjathoor, VV, Febbraio, M, Podrez, EA et al., Scavenger receptors class A-I/II
46 and CD36 are the principal receptors responsible for the uptake of modified low
47 density lipoprotein leading to lipid loading in macrophages. *J Biol Chem*
48 2002;277(51):49982–8.
49
50 [47] Nozaki, S, Kashiwagi, H, Yamashita, S et al., Reduced uptake of oxidized low
51 density lipoproteins in monocyte-derived macrophages from CD36-deficient
52 subjects. *J Clin Invest* 1995;96(4):1859–65.
53
54 [48] Panousis, CG, Zuckerman, SH, Regulation of cholesterol distribution in
55 macrophage-derived foam cells by interferon-gamma. *J Lipid Res* 2000;41(1):75–
56 83.
57
58 [49] Brewer, HB, High-density lipoproteins: a new potential therapeutic target for the
59 prevention of cardiovascular disease. *Arterioscler Thromb Vasc Biol*
60 2004;24(3):387–91.
61
62 [50] Han, X, Kitamoto, S, Lian, Q et al., Interleukin-10 facilitates both cholesterol uptake
63 and efflux in macrophages. *J Biol Chem* 2009;284(47):32950–8.
64
65

- 1
2
3
4 [51] Martinez, FO, Gordon, S, Locati, M et al., Transcriptional profiling of the human
5 monocyte-to-macrophage differentiation and polarization: new molecules and
6 patterns of gene expression. *J Immunol* 2006;177(10):7303–11.
7 [52] Xue, J, Schmidt, SV, Sander, J et al., Transcriptome-based network analysis
8 reveals a spectrum model of human macrophage activation. *Immunity*
9 2014;40(2):274–88.
10 [53] Nomura, M, Liu, J, Yu, ZX et al., Macrophage fatty acid oxidation inhibits
11 atherosclerosis progression. *J Mol Cell Cardiol* 2019;127:270–6.
12 [54] Ji, A, Meyer, JM, Cai, L et al., Scavenger receptor SR-BI in macrophage lipid
13 metabolism. *Atherosclerosis* 2011;217(1):106–12.
14 [55] Kwong, JC, Schwartz, KL, Campitelli, MA et al., Acute Myocardial Infarction after
15 Laboratory-Confirmed Influenza Infection. *N Engl J Med* 2018;378(4):345–53.
16 [56] Carnevale, R, Nocella, C, Petrozza, V et al., Localization of lipopolysaccharide
17 from *Escherichia Coli* into human atherosclerotic plaque. *Sci Rep* 2018;8(1):3598.
18 [57] Kadl, A, Sharma, PR, Chen, W et al., Oxidized phospholipid-induced inflammation
19 is mediated by Toll-like receptor 2. *Free Radic Biol Med* 2011;51(10):1903–9.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure Legends

Figure 1: Global analyses of human macrophage phenotype transcriptomes.

Human monocyte-derived macrophages from 8 donors were polarised with IFN γ +LPS, IL-4, IL-10, oxPAPC or CXCL4 and examined by RNA-seq for global and specific pathway changes in their transcriptomes. (A) Principal component analysis (PCA) and (B) numbers of differentially expressed genes ($|\log_2FC| > \log_2(1.5)$ and FDR < 0.05 (Benjamini and Hochberg method)) among all conditions. (C) Principal component analysis (PCA) and (D) numbers of differentially expressed genes excluding M^{CXCL4}. PC – principal component, DEG – differentially expressed gene.

Figure 2: Foam cell formation capacity by macrophage phenotypes.

(A) Representative Oil-Red-O staining of control and acLDL (25 μ g/ml for 24 h) loaded human monocyte-derived macrophages displaying presence of foam cells. Red arrow indicates positive Oil-Red-O staining where % control – % acLDL-loaded Area_{ORO} > Area_{haematoxylin}, scale bars = 100 μ m. (B) Quantification of foam cell formation capacity of human monocyte-derived macrophages, unpolarised (M^{un}), or polarised with IFN γ +LPS, IL-4, IL-10 or oxPAPC, determined using Image J polygonal selection and area measurements, where foam cells were considered positive for oil-red-O staining greater than the area of the nucleus. Δ % foam cells = control % foam cells – acLDL % foam cells; mean \pm SEM, n = 5 separate donors, matched/repeated measures one-way ANOVA with Dunnett's post-hoc test, ** $p \leq 0.01$ compared to M^{un}.

Figure 3: LDL uptake by macrophage phenotypes.

(A) Flow cytometry quantification of AlexaFluor-488-acLDL internalisation by human monocyte-derived macrophages, (M^{un}), or polarised with IFN γ +LPS, IL-4, IL-10 or oxPAPC, n = 8. (B) Differentially expressed mRNAs in the 'Binding and uptake of ligands by scavenger receptors' pathway using reactome.org analysis (annotations retrieved from reactome.org on 16 October 2018) in M^x compared to unpolarised (M^{un}) measured using RNA-seq; n = 8, thresholds for differential gene expression were $|\log_2FC| > \log_2(1.5)$ and FDR < 0.05 (Benjamini and Hochberg method). RNA-seq. (C) MSR1 and CD36 mRNA expression among MDM phenotypes relative to unpolarised (M^{un}); n = 8 donors, dotted lines = $|\log_2(1.5)|$, *** FDR ≤ 0.001 . (D) Flow cytometry quantification of MSR1 and CD36 cell surface protein levels by MDM phenotypes, n = 5. (A and D) Mean \pm SEM, matched/repeated measures one-way ANOVA with Dunnett's post-hoc test, * $p < 0.05$, *** $p \leq 0.001$, compared to M^{un}. MFI – geometric mean fluorescence intensity.

Figure 4: Internal lipid content and lipoprotein processing in macrophage phenotypes.

Colorimetric quantification of change in (A) total cholesterol, (B) cholesterol ester and (C) free cholesterol; mean \pm SEM, n = 4 donors, matched/repeated measures one-way ANOVA with Dunnett's post-hoc test, by human monocyte-derived macrophages polarised with IFN γ +LPS, IL-4, IL-10 or oxPAPC compared to unpolarised (M^{un}). (D) Differentially expressed mRNAs in the 'LDL clearance' pathway (annotations retrieved from reactome.org on 16 October 2018) of M^x compared to M^{un} measured using RNA-seq; n = 8, thresholds for differential gene expression were $|\log_2FC| > \log_2(1.5)$ and FDR < 0.05 (Benjamini and Hochberg method). (E) RNA-seq quantification of SOAT1, NCEH1, LIPA mRNA expression among MDM phenotypes compared to M^{un}, n = 8, dotted lines = $|\log_2(1.5)|$, * FDR < 0.05, ** FDR ≤ 0.01 , *** FDR ≤ 0.001 .

Figure 5: Cholesterol efflux by MDM phenotypes.

1
2
3
4 Fluorescence quantification of MDM cholesterol efflux for total cholesterol (A) via both lipid
5 acceptors, HDL (B) and Apo-AI (C) separately; mean \pm SEM, n = 4 donors, matched/repeated
6 measures one-way ANOVA with Dunnett's post-hoc test, ** $p \leq 0.01$, *** $p \leq 0.001$ by human
7 monocyte-derived macrophages polarised with IFN γ +LPS, IL-4, IL-10 or oxPAPC compared to
8 unpolarised (M^{un}). (D) Differentially expressed genes in the 'HDL assembly' and 'HDL remodelling'
9 pathways (annotations retrieved from reactome.org on 16 October 2018) in M^x compared to M^{un}
10 measured using RNA-seq; n = 8 donors, thresholds for differential gene expression were $|\log_2FC| >$
11 $\log_2(1.5)$ and FDR < 0.05 (Benjamini and Hochberg method). (E) RNA-seq quantification of *ABCA1*,
12 *ABCG1*, *SCARB1* mRNA expression among MDM phenotypes compared to M^{un}, n = 8 donors, dotted
13 lines = $|\log_2(1.5)|$, ** FDR ≤ 0.01 , *** FDR ≤ 0.001 . (F) Flow cytometry quantification of MDM
14 phenotype for ABCA1 (n = 5), ABCG1 (n = 6) and SCARB1 (n = 4 donors) cell surface protein levels;
15 mean \pm SEM, matched/repeated measures one-way ANOVA with Dunnett's post-hoc test, * $p < 0.05$,
16 *** $p \leq 0.001$, compared to M^{un}.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Response to Reviews.
Ref.: Ms. No. ATH-D-19-00697R1

Macrophage polarisation associated with atherosclerosis differentially affects their capacity to handle lipid

Atherosclerosis

Please find responses to the reviewers' comments as outlined in red text below.

“Dear Dr. Wilson,

Thank you for submitting your paper to Atherosclerosis. It has been reviewed and the referees' reports are appended below.

We are pleased to tell you that the manuscript has been found potentially acceptable for publication. However, there are some minor points, as detailed by the reviewers, that need to be addressed before we can proceed further, and we would ask you to revise your paper addressing each issue raised. **ALL CHANGES TO THE TEXT ARE TO BE SHOWN IN RED** (not as tracked changes).

When you resubmit, you should also include a point-by-point response to each critique; if you revise to a satisfactory standard, your paper will be accepted for publication.

Editors' comments:

The manuscript has been evaluated by the Editors and the two original Reviewers. While the revised version has improved, there are still mostly textual changes that need to be addressed before the paper can be accepted. “

Reviewers' comments:

Reviewer #1: This manuscript provides a wealth of data using the state-of-art methodologies. Validation data provided, including good functional assays. Informative Graphical Abstract.

Introduction.

1. I am not sure I understand the statement Macrophages regulate lipoproteins within their own environment.

We have also noted the change requested by reviewer 2, to the start of this second paragraph of the introduction. We have therefore altered the second sentence according to reviewer 2's request (see below), and kept the first sentence as it was, since reviewer 2 implied this first sentence should remain.

Similarly: The accumulation of free and esterified cholesterol droplets.....

We have changed the sentence “The cytosolic accumulation of free and esterified cholesterol droplets in macrophages results in the generation of foam cells, underlying atherosclerotic plaque formation [9].”

Now reads: “The cytosolic accumulation of esterified cholesterol and acylglyceride droplets in macrophages results in the generation of foam cells, driving atherosclerotic plaque formation”

2. Reference missing to original work for the statement: In arterial atherosclerotic plaques, pro-inflammatory macrophages.....

We have now cited Stöger et al., 2012, after this sentence.

3. I would encourage the authors to cite original work more to back up their statements rather than review articles.

We have now cited additional original articles in our manuscript, including Murray and Teitelbaum, 1992; Stöger et al., 2012; Hashizume and Mihara, 2012 to support specific statements. Analysis of our reference list shows 70% of the articles we now cite are original, 30% are review articles. We have chosen to cite review articles where they support the statements most clearly and provide a comprehensive assimilation of the literature.

4. Page 4, macrophage phenotypes occur change to develop? No references supporting your next statement. This whole paragraph would benefit from better citation of the literature.

We have now cited Johnston et al., 2018; Zhang et al., 2020, as examples in the literature to support the statement “the majority of our current understanding of macrophage polarisation and its impact on atherosclerosis derives from murine models of atherosclerosis and analysis of mouse bone marrow derived macrophages” in this section on page 4.

Results. I have been able to read further through the results but found it really hard towards the end. I think it all comes back to needing a better Introduction so that the results can be extracted more quickly.

We reconfigured the introduction and results comprehensively in our last revision to support this reviewer’s recommendations. Since the recommendations for a better introduction are not specific we consider we are this reviewer may not wish us to make further changes without such specifications. We are open to editorial guidance on this specific point.

1). Please be careful with the use of the word expression-you are measuring CD163 RNA levels. Same for rest of sentence. Also later on you refer to protein expression rather than levels.

In the sentence on page 5 “M^{IL-10} showed upregulation of *CD163*..” we have changed the wording expression to RNA. Throughout the manuscript we have italicised the gene names when referring to their RNA and kept the names in regular font when referring to protein, as per scientific convention.

We have changed any wording of “protein expression” to “protein levels” consistently in the manuscript, and have added mRNA where appropriate (to also comply with the suggestion in point 7 below).

2). Next, paragraph-the word transcription not correct here-you are looking at a transcriptome profile.

Page 5, we have changed “CXCL4 macrophages showed remarkably similar transcription to...” to “CXCL4 macrophages showed a remarkably similar transcriptome profile...”

3). Up-regulated at higher magnitudes than what? The beginning of the sentence not referenced.

Page 5, sentence “In addition, previously reported M^{CXCL4} markers S100A8 (S100 calcium binding protein A8) and MMP7 (matrix metalloproteinase 7) showed expression changes in our study that were highly variable among donors and up-

regulated at higher magnitudes in most other phenotypes, especially M^{IFN γ +LPS} (Supplementary Tables 2 and 3).”

Is now changed to:

“S100A8 (S100 calcium binding protein A8) and MMP7 (matrix metalloproteinase 7) have previously been reported as M^{CXCL4} markers [Erbel et al., 2015]; our RNA-seq data showed their expression in M^{CXCL4} was highly variable among donors and was up-regulated in most other phenotypes, especially M^{IFN γ +LPS} (Supplementary Tables 2 and 3).”

4). P5. Our transcriptome analysis of the differently polarised-otherwise making an assumption that is not proven? Also differences in RNA levels of

We have changed (page 5/6):

“Our transcriptome analyses of macrophage phenotypes revealed key differences in atherosclerosis and lipid handling genes (Supplementary Table 5).”

To

“Our transcriptome analyses of macrophage phenotypes revealed differences in lipid handling and other associated pathway enrichment for differentially expressed genes (Supplementary Table 5).”

5). Data in Fig 2A and B good-but the sentence in the text confusing to read. Fig legend A, I think I would say that red arrow indicate first-think sentence is round the wrong way.

Legend to Fig 2A is changed as suggested.

In B not clear what n=5 means.

“n = 5 separate donors” is now written in the legend to Figure 2.

I think the text-describing Figure 2A and B and Fig 3A would be better under one Section and in one Figure.

Thank you for this suggestion: we suggest that Figures 2 and 3A and associated text (main text and figure legends) should probably remain as they are to maintain the narrative as per our previous response to reviewer 1. In our last revision we altered the introduction, and ordered the results with the following narrative structure: foam cell formation which is then broken down mechanistically to determine lipid accumulation, processing and efflux. Since this was a response to reviewer 1, and since reviewer 2 accepted our revision on this basis, we suggest we do keep the figures as currently presented.

I think you have too much Introduction in the Results-I think after your first sentence in paragraph 2 of the Introduction, you could Introduce how acLDL is internalised by saying For Example.... I think most of the two paragraphs coming after the Subheading Cellular free cholesterol content in polarised macrophages on acLDL loading should be in the Introduction.

We have moved the following text from the results to the second paragraph of the introduction as advised:

“In the case of lipoprotein, cellular lipases such as LIPA break down the lipoprotein particles that have been taken up... for subsequent storage in lipid droplets.”

Please be careful, the uptake of LDL and modified LDL are quite different animals. We agree that LDL and modified LDL behave differently. We have checked the manuscript and ensured we only refer to acLDL where this is what we have measured experimentally, or “LDL” when we are making more generalised and discussion-related statements.

6). If you removed the Introduction in the Results to the Introduction,

In our last revision we altered the start of the results section considerably in response to both reviewers’ recommendations. The start of the results section now comprises a brief paragraph that serves to introduce the rationale. This rationale is included in the final section of the introduction. We consider that the narrative in the results section will be less clear if we remove the introduction to the first section of the results.

you could say more in the Subheading Cellular free-cholesterol content.....just not very informative.

We have re-worded the “cellular free-cholesterol content...” results subheading to improve informativity.

7). Fig 3C etc are you looking at mRNA or Total RNA?

We are looking at mRNA. We have corrected figure legends to make this clear in the descriptions. We have adjusted some wording in the figure legends to make the information more accurate and informative as shown in red text.

Description of Figure 3B really muddled me at first-overall down-regulation of what in what.

We have re-worded this results section (page 6), along with the legend to Fig. 3B, to make this description clearer. This was a down-regulation of a pathway identified using Ingenuity Pathway Analysis, which we now describe.

A similar expression profile-I think here you are trying to say RNA levels translated into similar reduction in protein levels.

We have re-worded the last sentence in this same paragraph (page 6, Reduced LDL uptake section) to make the description clearer.

8). I would argue that it is agreement with rather consistent with. Very vague in lipoprotein particle breakdown.

We have re-worded the last sentence in this same paragraph (page 6, Reduced LDL uptake section) to state “In agreement with...”

9). The last section can be reduced-the first two sentences perhaps in the Introduction, and then start with the third.

We have made these changes as suggested.

Reviewer 2:

The sentence "Macrophages regulate lipoproteins within their local environment [7]. Uptake of lipoproteins occurs via receptor-mediated endocytosis, pinocytosis, phagocytosis and scavenger receptor mediated mechanisms [4]" should read Macrophages regulate lipoproteins within their local environment [7]. Uptake of lipoproteins occurs via receptor-mediated endocytosis, such as scavenger receptor, pinocytosis and phagocytosis [4].

We have made these exact changes to the second paragraph of the introduction, as suggested here.

Editorial Office comments:

-Atherosclerosis applies formatting guidelines to all accepted papers, with the aim of improving their readability.

Manuscripts that do not conform to the format guidelines of the Atherosclerosis Journal will be returned to the authors for reformatting.

When revising your manuscript, please follow carefully the recommendations of our Atherosclerosis Style Guide to be downloaded from the following link (http://cdn.elsevier.com/promis_misc/Atherosclerosis_style_guide_checklist.docx).

- Make sure to apply the formatting requirements to all figures and tables where necessary (e.g. style of p values, gene and protein nomenclature).

- Make sure to use uniform lettering and sizing of your original artwork, including letters to indicate panels, throughout all figures.

- Make sure to submit high resolution versions of each figure. **We can improve these if considered necessary by the journal.**

- A graphical abstract is required at revision. **Supplied**

Statement of Originality.

The authors declare that the work described in our manuscript entitled "**Macrophage polarisation associated with atherosclerosis differentially affects their capacity to handle lipid**", for submission as a research article to *Atherosclerosis*, has not been published previously.

The article is not under consideration for publication elsewhere, publication of the article is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out. If the article is accepted, it will not be published elsewhere by the authors, including electronically in the same form, in English or in any other language, without the written consent of the copyright-holder.

Figure 1





Figure 2

A



B




Figure 3




Figure 4





Figure 5



Supplementary Material for online publication only

[Click here to download Supplementary Material for online publication only: Supp lipid handling Figs 05Feb2020pdf.pdf](#)

Supplementary Material for online publication only

[Click here to download Supplementary Material for online publication only: Supplementary tables Athero Rev 05Feb2020.pdf](#)

Supplementary Material for online publication only

[Click here to download Supplementary Material for online publication only: M&Methods Supplementary 05Feb2020.pdf](#)

Author contributions:

Kajus Baidžajevas: Conceptualisation, Methodology, Investigation, Formal analysis, Interpretation, Writing of original draft, Writing – Review & Editing. **Éva Hadadi:** Conceptualisation, Methodology, Investigation, Formal analysis, Interpretation, Writing – Review & Editing. **Bernett Lee:** Formal analysis, Writing – Review & Editing. **Josephine Lum:** Methodology & Investigation. **Foo Shihui;** Methodology & Investigation. **Ian Sudbery:** Formal analysis. **Endre Kiss-Tóth:** Supervision, Project administration, Funding acquisition, Conceptualisation, Interpretation, Writing of original draft, Writing – Review & Editing. **Siew Cheng Wong:** Supervision, Project administration, Funding acquisition, Conceptualisation, Interpretation, Writing – Review & Editing. **Heather L Wilson:** Supervision, Project administration, Funding acquisition, Conceptualisation, Interpretation, Writing of original draft, Writing – Review & Editing.