



This is a repository copy of *Paternal effects in a wild-type zebrafish implicate a role of sperm-derived small RNAs*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/163131/>

Version: Published Version

Article:

Ord, J., Heath, P.R. orcid.org/0000-0002-8385-1438, Fazeli, A. orcid.org/0000-0003-0870-9914 et al. (1 more author) (2020) Paternal effects in a wild-type zebrafish implicate a role of sperm-derived small RNAs. *Molecular Ecology*, 29 (14). pp. 2722-2735. ISSN 0962-1083

<https://doi.org/10.1111/mec.15505>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:
<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>



Paternal effects in a wild-type zebrafish implicate a role of sperm-derived small RNAs

James Ord^{1,2,3} | Paul R. Heath⁴ | Alireza Fazeli^{2,3} | Penelope J. Watt¹

¹Department of Animal and Plant Sciences, University of Sheffield, Sheffield, UK

²Academic Unit of Reproductive and Developmental Medicine, Department of Oncology and Metabolism, University of Sheffield, Sheffield, UK

³Department of Pathophysiology, Institute of Biomedicine and Translational Medicine, University of Tartu, Tartu, Estonia

⁴Sheffield Institute of Translational Neuroscience, University of Sheffield, Sheffield, UK

Correspondence

James Ord, Department of Animal and Plant Sciences, University of Sheffield, Alfred Denny Building, Western Bank, Sheffield S10 2TN, UK.
Email: jms.ord18@gmail.com

Funding information

Horizon 2020 Framework Programme, Grant/Award Number: 668989

Abstract

While the importance of maternal effects has long been appreciated, a growing body of evidence now points to the paternal environment having an important influence on offspring phenotype. Indeed, research on rodent models suggests that paternal stress leaves an imprint on the behaviour and physiology of offspring via nongenetic information carried in the spermatozoa; however, fish have been understudied with regard to these sperm-mediated effects. Here, we investigated whether the zebrafish was subjected to heritable influences of paternal stress by exposing males to stressors (conspecific-derived alarm cue, chasing and bright light) before mating and assessing the behavioural and endocrine responses of their offspring, including their behavioural response to conspecific-derived alarm cue. We found that after males are exposed to stress, their larval offspring show weakened responses to stressors. Small RNA sequencing subsequently revealed that the levels of several small noncoding RNAs, including microRNAs, PIWI-interacting RNAs and tRNA-derived small RNAs, were altered in the spermatozoa of stressed fathers, suggesting that stress-induced alterations to the spermatozoal RNA landscape may contribute to shaping offspring phenotype. The work demonstrates that paternal stress should not be overlooked as a source of phenotypic variation and that spermatozoal small RNAs may be important intergenerational messengers in fish.

KEYWORDS

alarm cues, epigenetic inheritance, paternal effects, small RNA, sperm, stress

1 | INTRODUCTION

The ancestral environment is already well-established as a factor which can shape the development of offspring phenotypes, and the existence of parental effects is now suspected to have a large bearing on the long-term responses of organisms to selective pressures (Räsänen & Kruuk, 2007). The contribution of the maternal environment has long been appreciated, for instance during gestation

(Feil & Fraga, 2012; Lupien, McEwen, Gunnar, & Heim, 2009) or via the nutritional and endocrine composition of the oocyte (Best, Kurrasch, & Vijayan, 2017; Davis & Sandman, 2010; Redfern et al., 2017). Although relatively less studied, the paternal environment too can have important influences on offspring development in species with high levels of paternal investment postfertilization (McGhee & Bell, 2014; Otero-Ferrer et al., 2020). Meanwhile, the sperm epigenome and its role in paternal nongenetic inheritance

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Molecular Ecology* published by John Wiley & Sons Ltd

currently represent a highly active area of research (Champroux, Cocquet, Henry-Berger, Drevet, & Kocer, 2018; Chen, Yan, & Duan, 2016).

Stress can be defined as a state of threatened homeostasis induced by real or perceived danger, necessitating a suite of adaptive physiological and behavioural responses (stress responses) to restore it (Barton, 2002; Godoy, Rossignoli, Delfino-Pereira, Garcia-Cairasco, & de Lima Umeoka, 2018). Neuroendocrine pathways involving glucocorticoid signalling typically (but not exclusively) underlie these responses, although the functions of glucocorticoids are far from limited to mediation of stress responses (MacDougall-Shackleton, Bonier, Romero, & Moore, 2019). The perceived threat of predation can be considered a form of stress for prey species which rely on cues to detect nearby predators, as exposure to these cues alone is often sufficient to induce behavioural and physiological responses reflective of a neuroendocrine stress response. For instance, several species of teleost fish release a chondroitin-based alarm cue from the skin upon damage (Stensmyr & Maderspacher, 2012) which, when detected, induces behavioural responses and elevated cortisol in conspecifics (Mathuru, 2016; Stephenson, 2016).

Although stress responses are inherently adaptive in the short term, overactivation of these responses over a prolonged period, known as chronic stress, can lead to a maladaptive state through overproduction of glucocorticoids (McEwen, 2004). Aside from regulating the neuroendocrine stress response, glucocorticoids are also important in the regulation of foraging behaviour, energy mobilization and use and the circadian rhythm (Rao & Androulakis, 2019; Vitousek, Taff, Hallinger, Zimmer, & Winkler, 2018). Excessive glucocorticoid secretion can disrupt these functions and shift many physiological parameters away from their normal homeostatic ranges, leading to dysregulation (Maestriperi & Hoffman, 2011). Although previously regarded as a phenomenon inherent to industrialized society, it has become clear that the chronic stress concept is applicable in the context of wild animals and their fitness. For example, chronic stress induced by elevated predation threat can impact the reproductive success of prey species through physiological disruption and can thus influence population dynamics even in the absence of direct killing (Clinchy, Sheriff, & Zanette, 2013).

Stress, particularly in the context of predation, can be a prominent source of maternal and paternal effects, as documented in a range of species with high levels of parental investment postfertilization including fish (Bell, McGhee, & Stein, 2016; Ord, Holmes, Holt, Fazeli, & Watt, 2020), reptiles (Bestion, Teyssier, Aubret, Clobert, & Cote, 2014) and mammals (Chen et al., 2018). However, evidence for effects of paternal stress mediated by changes to the spermatozoa before fertilization has been limited to studies on laboratory rodents. For instance, male mice stressed by maternal separation in early life sire offspring with reduced anxiety-like behaviours and glucose production (Gapp et al., 2014), while males exposed to chronic stress throughout spermatogenesis sire offspring with reduced corticosterone production (Rodgers, Morgan, Bronson, Revello, & Bale, 2013).

Meanwhile, evidence for sperm-mediated nongenetic inheritance in fish has been increasingly documented in recent years, albeit

not explicitly in the context of stress. For instance, rearing environment induces specific changes to salmon sperm DNA methylation which appear to be inherited by the offspring (Rodriguez Barreto et al., 2019), while sperm-mediated paternal effects on phenotype have been demonstrated in zebrafish in response to increased sperm competition (S. Zajitschek, Hotzy, Zajitschek, & Immler, 2014) and paternal social status (Zajitschek, Herbert-Read, Abbasi, Zajitschek, & Immler, 2017).

The zebrafish has long justified its place in the pantheon of vertebrate models in translational research, particularly given its short life cycle, ease of manipulation and relative genetic similarity to humans. Core components of the neuroendocrine stress response pathways are highly conserved between mammals and fish, both of which, for instance, employ cortisol as the principle glucocorticoid regulator of these pathways (Barton, 2002). Readily measurable anxiety-like behavioural patterns influenced by glucocorticoid signalling are also similar between mammals and fish, such as edge preference behaviour, termed thigmotaxis (Steenbergen, Richardson, & Champagne, 2011). Furthermore, the spermatogenic cycle in adult zebrafish is remarkably rapid, lasting only six days from the onset of meiosis (Leal et al., 2009), allowing for experiments concerning sperm-mediated effects on the offspring to be carried out in a short timeframe. Therefore, zebrafish readily lend themselves to the study of paternal stress effects.

Although several possible epigenetic mechanisms to explain sperm-mediated paternal effects have been postulated, the transmission of noncoding RNA, such as microRNA (miRNA), in spermatozoa has garnered the most experimental evidence (Chen, Yan, & Duan, 2016). MiRNAs, which are 18–24 nucleotides in length, have the capacity to alter gene expression via the incorporation into RNA-induced silencing complexes (RISC) and subsequent cleavage of target transcripts. Remarkably, phenotypic effects of paternal stress have been shown to be recapitulated by injection of either total spermatozoal RNA from stressed males (Gapp et al., 2014) or specific stress-associated spermatozoal miRNAs into normal zygotes (Rodgers, Morgan, Leu, & Bale, 2015). MiRNAs are abundant in teleost fish (Babiak, 2014) and have important functions in early zebrafish development (Wienholds et al., 2005). It is therefore possible that paternal miRNA and/or other noncoding RNAs delivered in spermatozoa could also influence developmental outcomes in fish.

Although they could also be maladaptive, changes to the RNA consignment or other components of the spermatozoal epigenome could facilitate intergenerational adaptive responses to precondition offspring if necessitated by changes to environmental conditions during the parents' lifetime (Immler, 2018). Such soft inheritance has been suggested to provide a "buffer" which protects underlying genetic variation against environmental changes which occur too quickly for adaptation to occur via conventional mutations and in doing so could even facilitate later genetic adaptation (O'Dea, Noble, Johnson, Hesselson, & Nakagawa, 2016). However, intergenerational effects are predicted to be adaptive only when the parental environment is a good predictor of offspring environment (Guillaume, Monro, & Marshall, 2016).

Here, we hypothesized that paternal stress could induce measurable influences on stress response physiology and behaviour in the offspring of zebrafish and that such alterations would be linked to alterations in the small RNA composition of the paternal sperm. We exposed males to stressors, which included alarm substance derived from the skin of conspecifics, and measured the responses to stressors in their offspring, specifically thigmotaxis (edge preference behaviour, which we took to be reflective of stimulus avoidance), and cortisol production. We further applied small RNA sequencing of paternal spermatozoa to investigate whether any intergenerational effects could be linked with differential expression of small RNAs putatively delivered to the offspring at fertilization. Although miRNAs are the most well-known class of small RNA to be implicated in paternal effects, we also investigated whether the spermatozoal composition of other small RNA classes was subjected to stress-induced alterations, specifically PIWI-interacting RNAs (piRNAs) and tRNA-derived small RNAs (tsRNAs).

2 | METHODS

2.1 | Animals and housing

Mature male London wild-type (LWT) zebrafish (aged 6–10 months) were selected from healthy stocks, reared in tanks on a recirculatory system with water heated to 26°C and kept on a 12:12-hr light/dark cycle. Animals were fed brine shrimp or flake food twice daily.

Adult males were housed in individual compartments (30 × 15 × 25 cm) of the recirculatory system for at least one week prior to the onset of the 12- to 14-day treatment period and for the entire duration thereof. Five individuals in both control and stress groups comprised one experimental batch. A total of six experimental batches were carried out. Table S1 gives the number of adult male zebrafish used in each batch of the experiment.

2.2 | Alarm substance extraction

Alarm substance was derived from mature LWT zebrafish (indiscriminate of sex), using a previously described method (Ord et al., 2020). Briefly, for every 3 ml of extract, five fish were euthanized, and 7–10 lacerations were made to the epidermis on both sides of each fish. All five fish were then placed in a single 50-ml tube with 3 ml water and gently shaken to allow the alarm substance to seep out. The water containing the extract was then eluted, incubated at 95°C for 16 hr, centrifuged at 10,000 × g for 10 min to remove debris and filtered through a microfilter.

2.3 | Paternal chronic stress

Stress treatments were administered during a 12- to 14-day treatment period. To administer stress, each male was removed from its

home tank into a separate exposure tank (17 × 11 × 12 cm) once per day and administered one of three randomly allocated stressors: 1: alarm substance (200 µl in 600 ml water) for 20 min, 2: chasing with a small net in 5 × 1-min episodes over a 20-min period or 3: exposure to bright light via a 10w LED tube light suspended above the tank for 20 min. Individuals were not visible to each other during exposures. Following alarm substance exposure, animals were moved to separate “washing” tanks to remove traces of alarm substance before returning them to their home tanks. Males exposed to stressors were classed as paternal chronic stress (PCS) animals. Control animals were handled in an identical fashion but did not receive stress treatments.

2.4 | Mate pairing, embryo collection and rearing

Two days after the final stress treatment, male zebrafish were paired with mature naïve females randomly selected from healthy stock. Dishes of marbles (12 cm diameter × 5 cm height) were placed in the compartments to provide a substrate for egg laying and to protect embryos from cannibalism. Eggs were collected the following morning, and dead eggs were removed. Twenty-four hours later, 1 day postfertilization (DPF) embryos from each male were placed in 5 cm petri dishes with 15 ml aquarium water in densities of 13–30 for assessment of larval stress response phenotypes. All embryo dishes were incubated at approx. 26°C under a 12:12 light cycle.

2.5 | Larval exposure to alarm substance and physical stress

5-DPF offspring were tested for behavioural and physiological responses to stress. Two sets of offspring from each male were housed in petri dishes, and one was exposed to alarm substance (50 µl), while the other was given the same volume of aquarium water as a blank. Petri dishes were placed on a platform of frosted glass which was illuminated from below with LED lights to maximize contrast between fish and background. Dishes were video-recorded from above using a Panasonic HC-X920 digital camcorder beginning from 5 min prior to alarm or control treatment. A smaller number of sets of control and PCS larvae were exposed to physical stress, which was established by stirring the petri dish water by hand with a Pasteur pipette for 1 min (~2 rotations per second).

2.6 | Larval cortisol measurements

Fifteen minutes after exposure to stressors, larvae were immediately euthanized by immersion in ice water. For cortisol extraction, euthanized larvae were collected into 1.5-ml microfuge tubes and stored at –20°C until cortisol extraction. Pools of larvae were homogenized

in 2-ml tubes in 120 μ l distilled water using a pellet mixer. Cortisol was ethyl acetate-extracted from homogenates and quantified by ELISA, both following a previously published protocol (Yeh, Glöck, & Ryu, 2013).

2.7 | Larval group thigmotaxis

Thigmotaxis (edge preference behaviour) in response to alarm substance was recorded directly from videos using automated tracking software (Viewpoint® ZebraLab) which calculated the average number of animals present in a defined central zone (covering most of the dish area apart from the outer 6mm or approx. 1.5 larval standard lengths; Figure 1, inset) for every 10-s time interval. These values were subtracted from the total number of animals to calculate the percentage of animals in the peripheral zone (% thigmotaxis). These values were then summarized into 1-min time bins (i.e. the mean of six values). As the average thigmotaxis response appeared to reach a peak at around 5 min after exposure and subsequently began a gradual decrease, we considered only the first 10 min after exposure in the statistical analyses (i.e. 5 min either side of the observed peak).

2.8 | Statistical analyses of phenotype data

All statistical analyses were carried out in “R” 3.6.1 (R Core Team, 2018). We used linear mixed-effects models fit by restricted maximum likelihood (REML) with the LME4 package version 1.1.21 (Bates, Maechler, Bolker, & Walker, 2015) because of the hierarchical structure of the experimental design and the fact that experiments were carried out in batches. In all models, “Batch” was included as a random term. “Father ID” was included as a random effect (nested within batch) as the analyses included multiple samples from the same parent. “Dish ID” was included as a random effect (nested within Father ID) in the case of thigmotaxis as observations were made at multiple time points. “Assay” was included as a separate random effect in the case of cortisol analyses due to samples being run over three plates. Significance of fixed effect terms was evaluated using *F* tests and *t* tests with Kenward–Roger approximation of degrees of freedom, derived from summary tables obtained using the ANOVA() and summary() functions from the stats package in conjunction with LMERTEST version 3.1.1 (Kuznetsova, Brockhoff, & Christensen, 2017) and PBKRTTEST version 0.4.7 (Hakekoh & Hojsgaard, 2014) packages. Post hoc pairwise *t* tests of model terms were computed using the emmeans() function from the EMMEANS package version 1.4.3




FIGURE 1 Visual summary of larval behavioural and physiological stress responses. (a) Thigmotaxis (% of animals within 1.5 standard lengths of the dish edge) in 1-min time periods for control (circles, solid lines) and paternal chronic stress (PCS) offspring (triangles, dashed lines) exposed to alarm substance (black) or a “blank” (water; white) at 5 days postfertilization. The dashed vertical lines demarcate the period during which the alarm substance (or blank) was added to the dishes (between minutes 5 and 6). Inset: schematic representation of the measurement of thigmotaxis. (b) Cortisol levels in control and PCS offspring exposed to alarm substance (grey) or blank (white). (c) Cortisol levels from control and PCS offspring exposed to a stirring stimulus (grey) or blank (white). Each point represents a pool of 13–30 animals. Stars represent $p < .05$, derived from post hoc pairwise comparisons of mixed-effects models by estimated marginal means

(Lenth, 2019). To approximate normal distributions, the thigmotaxis response variable was subjected to a Gaussianization transformation using the `Gaussianize()` function from the `LAMBERTW` package version 0.6.4 (Goerg, 2016), while cortisol response variables were log-transformed. Details of the models, including model structure and sample sizes can be found in Table S2. The R code for the models is also provided in the Supporting information. Plots of phenotype data were generated using `GGPLOT2` version 3.2.1 (Wickham, 2011).

2.9 | Spermatozoal RNA isolation

Spermatozoa were collected from euthanized adult male zebrafish by disruption of dissected testes in ice-cold Hank's balanced salt solution (HBSS) to derive a testicular cell suspension, which was then centrifuged (800×g for 40 min) on top of a 40/80 Percoll gradient to separate spermatozoa from larger cells. After density gradient centrifugation, the upper layers containing nonsperm cells were removed and the sperm precipitate washed in ice-cold HBSS. Spermatozoa were then pelleted (1,000×g for 10 min), resuspended in Tri reagent and stored at -80°C until RNA extraction. Total RNA was extracted following the Tri reagent protocol with the addition of an overnight precipitation step with glycogen in 2-propanol at -20°C. RNA quantity was measured using the Qubit fluorometry kit, and size distribution was assessed using the Agilent Bioanalyser Pico kit. The effectiveness of the sperm enrichment protocol was indicated by the lack of prominent 18S and 28S ribosomal RNA in enriched samples, while a control sample which was not subjected to the enrichment protocol showed prominent rRNA peaks and a high RNA integrity number (Figure S1). Each sperm RNA sample comprised a pool of multiple individuals within the experimental batches such that each of six batches contributed one replicate pair (control and PCS), for a total of 12 biological samples. See Table S3 for the sperm RNA sampling scheme.

2.10 | RNA-seq library preparation and sequencing

Small RNA sequencing libraries were prepared using the SomaGenics (Santa Cruz) Real-Seq Biofluids kit following the manufacturers' instructions, using a minimum of 2.5 ng of input total RNA. Twenty PCR cycles were performed to amplify the reverse transcription product. The quality of the prepared libraries was examined using the Agilent Hi-sensitivity DNA chip prior to the individual libraries being pooled to a concentration of 4 nM for Illumina sequencing. Sequencing was carried out on an Illumina HiSeq 2,500 using SBS v3 chemistry and single end 65 base reads, producing on average 7.59 ± 2.22 million (mean \pm SD) reads per sample.

2.11 | RNA-seq data processing (quality control)

Reads were cropped to a maximum length of 50 nt and the adaptor sequence (TGGAATTCTCGGGTCCAAGG) removed using

trimmomatic (Bolger, Lohse, & Usadel, 2014) and read quality assessed using FASTQC (Andrews, Krueger, Seifert, Pichon, Biggins, & Wingett, 2015). Residual primer-derived contaminant sequences were also removed using trimmomatic following identification of possible primer contaminant sequences in FASTQC reports. Two library pairs were removed prior to running the final analysis workflow as they contained prominent outliers (apparent from a preliminary exploratory analysis; Figure S2), possibly due to lower read counts and/or higher amounts of possible primer contamination than other samples (Table S3). Excluding these library pairs, the average number of cleaned sRNA reads per sample was approx. 8.54 ± 1.92 million (mean \pm SD). Raw FASTQ files have been deposited to the NCBI sequence read archive under BioProject PRJNA635603.

2.12 | miRNA alignment and quantification

For miRNAs, reads in the length range of 18–24 nt were retrieved and aligned to miRbase mature zebrafish miRNA sequences using BOWTIE2 version 2.3.4.1 (Langmead & Salzberg, 2012) allowing no mismatches in the seed region but allowing the default maximum of two mismatches in the remainder of the alignment. Primary alignments were counted using SAMTOOLS version 1.7 (Li et al., 2009) with a mapping quality (MAPQ) filter of 10. To account for possible alternative miRNA isoforms derived from differential clipping of hairpin precursors (isomiRs), reads not aligning to mature miRNAs were recovered and aligned to known hairpin sequences retrieved from miRbase using the same Bowtie2 parameters. Primary alignments to hairpin sequences were also counted in the same manner, as we did not seek to identify specific isomiRs.

2.13 | piRNA alignment and quantification

For piRNAs, reads in the 25–32 nt range were aligned to predicted zebrafish piRNA cluster sequences obtained from the piRNA cluster database (<https://www.smallrnagroup.uni-mainz.de/piCdb/>; Rosenkranz, 2016). Alignment was carried out using Bowtie2 as above, and up to three alignments for each read were reported. As the MAPQ score produced by Bowtie2 is influenced by multiple alignments but we were interested only in cluster-level expression (and therefore not concerned with within-cluster multimapping), we did not consider MAPQ when counting alignments. Instead, alignments were filtered to retain those with alignment scores (AS) ≥ -11 (permitting one mismatch or one read gap of up to two bases), and for each read, the alignment with the highest AS was retained. Where reads had multiple alignments tied for the highest AS, these reads were kept if each alignment was within the same cluster and discarded if they aligned to multiple clusters. Multiple alignments were then assigned fractional values prior to counting the number of alignments to each cluster, such that the sum of all values counted equalled the total number of reads retained.

2.14 | tsRNA reference generation, alignment and quantification

Transfer RNA-derived small RNAs (tsRNAs) are typically derived from the 5' or 3' ends of mature tRNAs. Therefore, to obtain a reference database of putative spermatozoal tsRNAs, we pooled together 18–40 nt reads (which did not align to miRNAs/hairpins or piRNAs in previous steps) from eight biological samples (approx. 36 million reads) and aligned them (Bowtie2) to the first (5'ends) and last (3'ends) 40nt of mature tRNA sequences in the genomic tRNA database (Chan & Lowe, 2009). Perfect match primary alignments (i.e. no substitutions or indels; AS = 0) were extracted and further filtered to include only those that aligned to the edges of the reference sequences using custom R scripting. Unique sequences were considered as putative tsRNAs if they were represented by at least 4 reads. Due to many putative tsRNAs being redundant (i.e. could be completely encapsulated within another qualifying sequence) and so could not be pinpointed to a specific tsRNA, a clustering approach was adopted. Redundant sequences were removed, and nonredundant sequences were subjected to clustering using CD-HIT-EST from the CD-HIT software package version 4.8.1 (Li & Godzik, 2006) with a similarity threshold of 90%. Putative tsRNA sequences belonging to the same cluster were concatenated into single FASTA entries with each constitutive sequence separated by four Ns. Subsequently, the final reference FASTA contained both tsRNA sequence clusters and individual, nonredundant tsRNA sequences (i.e. "single-member" clusters).

18–40 nt reads from individual samples were then aligned to the putative tsRNA cluster reference using Bowtie2 as described for miRNAs. As a non-negligible portion of reads were expected to have multiple alignments within clusters, up to three alignments were reported per read which were subsequently filtered and counted using the same method described for piRNA clusters.

2.15 | Differential expression analysis

Differential expression analyses of counted miRNAs/hairpins, piRNA clusters and tsRNA clusters were carried out in R version 3.6.1 (R Core Team, 2018) using the EDGER package version 3.26.8 (McCarthy, Chen, & Smyth, 2012), with separate analyses carried out for each small RNA class. Counts matrices were filtered using the filterByExpr() function to retain RNAs with a minimum of five counts in the smallest library of at least one of the experimental groups. Counts were TMM-normalized using calcNormFactors(). The total counts in each sample were used as the library sizes for each of the respective small RNA classes. PCA plots based on multidimensional scaling were produced using the plotMDS() function, which calculated Euclidean distances between sample pairs using the log2 fold changes of all RNAs for all pairwise comparisons, and replotted using GGLOT2 version 3.2.1. Tagwise dispersion estimates were derived based on the trended dispersions, and statistical comparisons were performed using generalized linear models followed

by likelihood ratio tests. We allowed results with an FDR-corrected p -value of < 0.1 and a log2-fold change > 1 or < -1 to be considered differentially expressed. The TMM-normalized CPM values of all differentially expressed small RNAs were converted to a single set of Z-scores which were plotted as a heatmap with hierarchical clustering using the pheatmap() function from the PHEATMAP package version 1.0.12 with default parameters (<https://cran.r-project.org/web/packages/pheatmap/index.html>).

2.16 | MiRNA target analysis

Lists of predicted targets of differentially expressed mature miRNAs were obtained using the miRmap web interface (Vejnar, Blum, & Zdobnov, 2013) with a miRmap score cut-off of 95%, reflecting those in the top 5th percentile of predicted miRNA target repression strength. The obtained gene lists were filtered to retain only genes present at a minimum of 5 TMM-normalized counts per million in a previously published unfertilized egg transcriptome data set (Jiang et al., 2013) (GEO accession GSM1085060). Genes targeted by both up- and downregulated miRNAs were removed to obtain lists of genes targeted exclusively by either up- or downregulated miRNAs. To discern whether any pathways may be dysregulated as a result of differentially expressed miRNAs, the ENTREZ identifiers (ANNOTATIONDBI package version 1.46.1; Pages, Carlson, Falcon, & Li, 2018) of all genes targeted exclusively by up- or downregulated miRNAs were inputted into a single pathway overrepresentation analysis using the enrichKEGG() function from the CLUSTERPROFILER package version 3.12 (Yu, Wang, Han, & He, 2012) and KEGG pathway annotations for *D. rerio* (Kanehisa & Goto, 2000). The ENTREZ identifiers of all genes present in the egg transcriptome data set (≥ 5 CPM) were used as the background for pathway overrepresentation analysis. A network plot of miRNA–gene–pathway interactions was generated using the network() function from the NETWORK package version 1.16 (Butts, 2008) and the ggnet() function from the GGally GGLOT2 extension version 1.4 (Schloerke et al., 2018).

3 | RESULTS

3.1 | Offspring of males exposed to stressors have compromised stress responses

Both control and PCS offspring exhibited increased thigmotaxis in response to alarm cue (Figure 1a), but the response in the PCS larvae was markedly weaker, the overall mean dropping by 3.6 ± 2.5 (SEM) percentage points. A mixed-effects model (Table S4) revealed an overall significant effect of alarm substance on larval thigmotaxis (F test of linear mixed model with Kenward–Roger approximation of degrees of freedom, $F^{1,36} = 28$, $p < .001$), but no significant overall effect of PCS ($F^{1,32} = 1.61$, $p = .21$) or larval alarm \times PCS interaction, that is no effect of PCS on the larval response to alarm substance ($F^{1,36} = 2.1$, $p = .16$). However, post hoc pairwise comparisons




FIGURE 2 Sequencing of small RNAs in zebrafish spermatozoa. (a) Fluorescent live/dead staining image of spermatozoal sample enriched using density gradient centrifugation. Live cells are stained green with calcein-AM and dead cells are stained red with ethidium homodimer-1. (b) Sequence length distribution of spermatozoal RNA samples derived from small RNA sequencing. Two prominent peak regions < 30 nt corresponded with known size ranges of miRNA and piRNA, while the 18-40nt range is known to correspond with tsRNAs. (c) Sequence length distribution of putative tsRNA reads identified from a pool of eight biological samples. Putative tsRNAs were identified as 18-40nt reads with a perfect match alignment to either the 5' (5'tsRNAs, black) or 3' end (3'tsRNAs, grey) of mature tRNA sequences derived from the genomic tRNA database. (d) Sequence length distribution of putative unique 5'tsRNA (black bars) and 3'tsRNA sequences (grey bars), including redundant sequences

revealed that in the presence of alarm substance, thigmotaxis was significantly lower for PCS larvae ($T^{60.6} = 2.14$, $p = .036$).

As expected, both alarm substance and the physical stressor, stirring, induced a significant increase in cortisol in control offspring (estimated marginal means; alarm: $T^{30.6} = 2.69$, $p = .01$; stirring: $T^{10} = 3.05$, $p = .01$), but not in PCS offspring (alarm: $T^{30.14} = 1.27$, $p = .22$; stirring: $T^{10} = 0.56$, $p = .59$) (Figure 1b and c), although there were no significant interactions between paternal stress and either

of the larval stress treatments (alarm: $F^{1,30.1} = 0.97$, $p = .33$; stirring: $F^{1,10} = 3.1$, $p = .11$). Cortisol in response to stirring was markedly lower in PCS compared to control larvae, although the difference was not significant ($T^{13} = 1.19$, $p = .08$). Table S5 shows the results of the mixed-effects model.

PCS did not affect hatching success (Figure S3), suggesting that the observed effects on stress response were not a consequence of general developmental impairments.




FIGURE 3 Male stress alters small RNA expression in zebrafish spermatozoa. (a) Principal components analysis plots based on multidimensional scaling, showing Euclidean distances between samples according to differences in \log_2 FC of mature miRNAs, hairpin-derived sequences (putative isomiRs), piRNA clusters and tsRNA clusters in spermatozoa from paternal control (white points) and stress samples (black points), with experimental batch denoted by point shape. (b) Heatmap derived from hierarchical clustering of normalized expression values (Z-scores of counts per million) of 12 differentially expressed miRNAs, four hairpins exhibiting putative isomiR expression, six piRNA clusters and 12 differentially expressed putative tsRNAs or tsRNA clusters (FDR < 0.1). Putative tsRNAs are named after the mature tRNA sequence as well as the end (5' or 3') from which they are putatively derived. For clusters, the name of a representative tsRNA is shown, followed by the number of nonredundant tsRNAs in the cluster, in brackets. (c) Network plot of differentially expressed spermatozoal miRNAs, their predicted target genes (derived using miRmap) in the egg transcriptome and KEGG pathways enriched with miRNA targets (identified via pathway overrepresentation analysis). The plot is restricted to putative target genes that are included in enriched KEGG pathways (mitophagy and notch signalling) or those that are the predicted targets of at least three miRNAs (*socs5b*, *phyhipla* and *mf144aa*) and associated miRNAs. Target genes that were not included in any enriched pathways or which were targeted by fewer than three miRNAs are not shown. The two miRNAs in the dre-miR-723 family (dre-miR-723-5p and dre-miR-723-3p) were combined as one for this analysis

3.2 | Detection of miRNAs, piRNAs and tsRNAs in paternal spermatozoa

Following small RNA sequencing of spermatozoal samples (Figure 2a) from stressed and nonstressed males, $67.9 \pm 6.64\%$ (mean \pm SD) of total cleaned reads fell within the 18–40nt size range known to contain miRNAs, piRNAs and tsRNAs. Two distinct peak regions were identified in the sub-30nt range, which corresponded with known size ranges of miRNAs (18–24nt) and piRNAs (25–32nt), respectively (Figure 2b).

Approximately $15.5 \pm 3.3\%$ of reads in the 18–24 nt range (or $4.7 \pm 1.4\%$ of total cleaned sRNA reads) aligned to known mature miRNA sequences in miRbase, while a further $1.8 \pm 0.7\%$ (or $0.5 \pm 0.2\%$ of total cleaned sRNA reads) aligned to known hairpin sequences, constituting possible isomiRs.

Approximately $34 \pm 2.3\%$ of reads in the 25–32nt range (or $9.4 \pm 2.3\%$ of total cleaned sRNA reads) aligned to predicted piRNA cluster sequences derived from the piRNA cluster database, although

$31.6 \pm 1.8\%$ of piRNA-aligning reads with alignment score(s) of $>= -11$ had equally valid alignments to multiple clusters (these reads were not counted).

Following alignment of the remaining 18–40 nt reads to the 5' and 3' ends of mature tRNA sequences in the genomic tRNA database, we identified 610 and 142 unique sequences which we considered putative 5' and 3'tsRNAs, respectively. In terms of total read abundance, most 5'tsRNAs were 32–34 nt in length, while almost all 3'tsRNAs were 36 nt (Figure 2c). The distribution of unique sequences followed the same general pattern, albeit sequence lengths were more evenly distributed than for total reads (Figure 2d). When subjected to clustering by sequence similarity, 214 nonredundant tsRNA sequences (i.e. those which were not completely encapsulated by another sequence) formed 66 clusters. Re-alignment of 18–40nt reads (excluding miRNAs and piRNAs) from individual samples to clustered tsRNA sequences resulted in a $5.2 \pm 1.2\%$ alignment rate (or $2.8 \pm 0.7\%$ of total cleaned sRNA reads).

Together, miRNAs (including possible isomiRs), piRNAs and putative tsRNAs comprised approximately $17.3 \pm 2.3\%$ of cleaned sRNA reads.

3.3 | Differential expression of miRNAs, piRNAs and tsRNAs in paternal spermatozoa

Principal components analyses based on multidimensional scaling revealed that samples consistently segregated according to paternal treatment over the second principal component axis (Figure 3a), suggesting that male stress explained an appreciable proportion of variation in the populations of all three small RNA classes.

Out of 213 mature miRNAs that met the counts threshold for differential expression analysis, we detected 12 that were differentially expressed in response to stress (Figure 3b, Table 1), seven of which were downregulated and five upregulated in stressed males. The two topmost differentially expressed miRNAs, both downregulated, were let-7 family members: *dre-let-7d-5p* ($\log_2FC = -1.49$, $FDR = 0.024$) and *dre-let-7b* ($\log_2FC = -1.9$, $FDR = 0.024$). Among counts of 18–24nt reads which did not align to known mature miRNA sequences but which aligned to hairpin sequences, 79 such hairpins met the required counts threshold, of which four were found to be downregulated in response to stress (Figure 3b, Table 1). The topmost of these were *dre-mir-16a* ($\log_2FC = -2.3$, $FDR = 0.005$) and *dre-let-7b* ($\log_2FC = -2.84$, $FDR = 0.035$). Visual inspection of reads aligning to these hairpins using integrative genomics viewer (Robinson et al., 2011) indicated that most

TABLE 1 List of mature miRNAs differentially expressed in zebrafish spermatozoa in response to male stress, as well as hairpin sequences exhibiting additional differential expression arising from alternative miRNA isoforms (putative isomiRs)

Mature miRNA	logFC	logCPM	LR	FDR
<i>dre-let-7d-5p</i>	-1.493	8.372	13.721	0.024
<i>dre-let-7b</i>	-1.898	8.919	13.622	0.024
<i>dre-miR-723-3p</i>	1.260	13.989	11.273	0.056
<i>dre-miR-125a</i>	-2.075	7.760	10.509	0.058
<i>dre-miR-34b</i>	1.487	7.853	10.032	0.058
<i>dre-miR-17a-5p</i>	-1.205	8.702	9.560	0.058
<i>dre-miR-132-3p</i>	-1.313	8.151	9.503	0.058
<i>dre-miR-723-5p</i>	1.380	7.393	9.316	0.058
<i>dre-miR-19b-5p</i>	1.910	5.081	8.982	0.058
<i>dre-miR-200a-3p</i>	-1.811	5.540	8.825	0.058
<i>dre-miR-18c</i>	-2.104	4.826	8.813	0.058
<i>dre-miR-129-3-3p</i>	1.474	6.276	8.157	0.076
Hairpin sequence	logFC	logCPM	LR	FDR
<i>dre-mir-16a</i>	-2.312	9.494	19.267	< 0.001
<i>dre-let-7b</i>	-2.847	11.224	10.514	0.047
<i>dre-mir-733</i>	1.640	8.225	9.041	0.070
<i>dre-mir-194a</i>	-2.227	8.548	7.887	0.098

overlapped with the location of the known mature miRNA sequence, suggestive of isomiRs, as opposed to degraded fragments of hairpin precursors.

Out of 569 piRNA clusters that met the required counts threshold, six were differentially expressed, all of which were upregulated in stressed males (Figure 3b, Table S6). The topmost were cluster 252 located on chromosome 7 ($\log_2FC = 2.1$, $FDR < 0.0001$) and cluster 104 located on chromosome 4 ($\log_2FC = 2.84$, $FDR < 0.0001$).

Out of 55 putative tsRNAs or tsRNA clusters that met the required counts threshold, 12 were differentially expressed in response to male stress (Figure 3b, Table S7). Most of these were downregulated, including a single-member cluster comprising a nonredundant putative 3'tsRNA derived from tRNA-His-ATG-1-1 ($\log_2FC = -1.81$, $FDR = 0.002$), and a 4-member cluster of putative 5'tsRNAs represented by a sequence derived from tRNA-Ser-ACT-2-1 ($\log_2FC = -1.8$, $FDR = 0.002$).

3.4 | Mitophagy and Notch signalling pathways are enriched among predicted targets of differentially expressed spermatozoal miRNAs

Out of 643 total predicted miRNA targets present in the unfertilized egg transcriptome data set ($CPM \geq 5$), 301 genes were targeted exclusively by upregulated miRNAs, while 289 were targeted exclusively by downregulated miRNAs. Among genes targeted exclusively by upregulated miRNAs, none were targeted by more than two. Among genes targeted exclusively by downregulated miRNAs, we identified three—*socs5b*, *phyhipla* and *rnf144aa*—which were predicted targets of all three of *dre-miR-17a-5p*, *dre-miR-132-3p* and *dre-miR-200a* (Figure 3c). A pathway overrepresentation analysis of the putative targets of differentially expressed miRNAs revealed two KEGG pathways which were enriched with target genes: mitophagy (*dre04137*; 8 genes; $FDR = 0.08$) and notch signalling (*dre04330*; 11 genes; $FDR = 0.08$), the latter including *notch2* and *notch3* (Figure 3c).

4 | DISCUSSION

Although the role of paternal effects has been increasingly studied in the context of paternal investment postfertilization, such as brood pouch influences in seahorses and pipefish (Cunha, Berglund, Mendes, & Monteiro, 2018; Otero-Ferrer et al., 2020), and nest guarding in sticklebacks (Bell et al., 2016; Mcghee & Bell, 2014), paternal effects mediated by alterations to the sperm epigenome have hitherto not been well characterized in nonmammalian vertebrates such as fish. Here, we show that in a fish with no parental care, heightened paternal stress before fertilization results in offspring with a reduced tendency to exhibit a behavioural response to a natural danger signal (alarm cue). Paternal chronic stress (PCS) offspring furthermore did not exhibit an endocrine stress response in terms of increased cortisol levels following exposure to either alarm cue

or a physical stressor, suggesting that paternal experience disrupted offspring stress response physiology.

In terms of experimental framework and the context of paternal stress, the previous studies most comparable to the present study were carried out in rodent models of paternal stress, in which the production of corticosterone (Rodgers et al., 2013) and glucose (Gapp et al., 2014) was ameliorated in offspring from stressed fathers. Not only do our results mirror those of these studies on laboratory rodents, but they also add to a growing body of evidence from a range of organisms, including invertebrates (Crean, Dwyer, & Marshall, 2013; Guillaume et al., 2016; Tauffenberger & Parker, 2014; Zajitschek, Zajitschek, & Manier, 2017), fish (Evans, Lymbery, Wiid, Rahman, & Gasparini, 2017; Rodriguez Barreto et al., 2019; Zajitschek et al., 2014) and wild mammals (Weyrich et al., 2016) that the prefertilization paternal environment can be a crucial determinant of offspring epigenome and/or phenotype.

It is unclear whether the suppressed responses to stressors exhibited by PCS offspring has any adaptive significance, as suppression of behavioural response to a natural danger signal would more immediately appear to be maladaptive. However, in the context of neuroendocrine stress, such a suppression could also be seen as adaptive within higher-stress environments with the propensity to induce chronic stress, as it would ameliorate the risk of excessive glucocorticoid production causing dysregulation to other essential processes that are mediated by these hormones (Maestriperi & Hoffman, 2011; Rao & Androulakis, 2019). Furthermore, activation of neuroendocrine stress responses incurs an energetic cost while simultaneously suppressing foraging behaviour—a combination which can be highly detrimental to fitness (Clinchy et al., 2013). Indeed, wild songbirds have lower reproductive success under heightened predation threat, in part due to starvation of nestlings due to reduced parental foraging (Zanette, White, Allen, & Clinchy, 2011). Therefore, although weaker responses may be a disadvantage in the context of an isolated predator encounter, regular initiation of a strong response in the face of heightened predation risk may be ultimately unsustainable and detrimental to long-term fitness. Alternatively, it is possible that other mechanisms which rely less extensively on cortisol are employed to respond to challenges most adaptively (MacDougall-Shackleton et al., 2019). Regardless, intergenerational mechanisms to facilitate “fine-tuning” of the intensity or mode of stress response may be ultimately beneficial for long-term offspring fitness.

The differential expression of spermatozoal small RNAs in spermatozoa in response to stress shows that the molecular composition of spermatozoa is sensitive to environmental disruption and is further concordant with patterns in rodent models, suggesting that spermatozoal small RNAs comprise a form of heritable environmental information in teleosts. Two downregulated mature miRNAs found in the spermatozoa of stressed males (dre-let-7b and dre-let-7d-5p, the former of which also exhibited putative isomiR downregulation) belonged to the let-7 family, which is among the most evolutionarily conserved (Lee, Han, Kwon, & Lee, 2016). Interestingly, downregulation of spermatozoal let-7 miRNAs has been implicated in

intergenerational effects of nutritional stress (protein restriction) in a rodent model (Sharma et al., 2016), suggesting that the functions of intergenerational transmission of this miRNA family may be similarly conserved.

In addition to miRNAs, we detected increased relative expression of six piRNA clusters in the spermatozoa of stressed males. This particular class of small RNA has also been shown to play a role in intergenerational effects in other organisms, including maternal inheritance of an acquired trait in *Drosophila* (Grentzinger et al., 2012). Altered piRNA expression levels were also detected in the spermatozoa of male mice that had been stressed by maternal separation (Gapp et al., 2014). Altered levels of piRNA cluster expression in spermatozoa may reflect stress-induced changes to the expression of transposable elements (TEs) during spermatogenesis, in which piRNAs are employed extensively in the suppression of “opportunistic” TE expression during chromatin remodelling (Ernst, Odom, & Kutter, 2017).

As well as miRNAs and piRNAs, this study identified the presence of a novel class of small RNAs—tsRNAs—in zebrafish spermatozoa, which constituted a similar proportion of cleaned sRNA reads to mature miRNAs (2.8% compared to 4.7%). We identified clusters of similar tsRNAs which were mostly downregulated in response to male stress, mirroring the effect on sperm miRNAs which were also mostly downregulated. Recently, results from rodent models have implicated sperm tsRNAs in nongenetic inheritance of acquired metabolic syndrome (Chen, Yan, Cao, et al., 2016) and transgenerational influences of maternal overnutrition (Sarker et al., 2019). Our results suggest a similar role for sperm tsRNAs in fish, and therefore, tsRNAs may have evolutionarily conserved functions in conveying environmental information in the germ line. Indeed, tsRNAs are purported to represent some of the most primitive small RNA pathways from which more well-known RNA silencing pathways evolved (Garcia-Silva, Cabrera-Cabrera, Güida, & Cayota, 2012).

Although we identified differentially expressed spermatozoal RNAs in response to stress, their role in mediating phenotypic effects on the offspring is difficult to infer without functional experiments. However, it is likely that paternal miRNAs interact with maternal mRNAs to some degree immediately after fertilization, potentially influencing early developmental processes which are governed by maternal factors prior to zygotic genome activation (Wagner, Dosch, Mintzer, Wiemelt, & Mullins, 2004). We identified several putative targets of differentially expressed sperm miRNAs in the oocyte transcriptome, among which the mitophagy (mitochondrial destruction) and notch signalling pathways were notably enriched. Mitophagy, the coordinated destruction of paternal mitochondria by maternal factors, is one of the earliest events following fertilization (Al Rawi et al., 2011). The enrichment of this pathway with miRNA targets implies that this process could become disrupted as a result of changes to paternal miRNA, possibly causing further disruption to other early processes. Notch signalling is ubiquitous and crucial for differentiation and pattern formation as embryogenesis progresses. The function of maternal Notch signalling-related mRNA in very early embryos is not clear, although maternal mRNAs in general appear

to be important in early tissue specification in *Xenopus* embryos (Heasman, 2006).

Noncoding RNAs are believed play a role in chromatin remodelling during embryogenesis (Pauli, Rinn, & Schier, 2011), and mounting evidence suggests that piRNAs are capable of mediating DNA methylation (Calcagno et al., 2019). Therefore, paternally transmitted noncoding RNAs may influence the methylome of the early embryo. Indeed, zebrafish may be especially sensitive to paternal effects on methylation given that the methylation profile established in the early embryo recapitulates that of the sperm (Jiang et al., 2013; Potok, Nix, Parnell, & Cairns, 2013), although the mechanisms underlying this remain unclear. Further research is required to elucidate whether spermatozoal RNAs do indeed influence the embryonic methylome, thereby directing offspring phenotypes. Alternatively, the determination of phenotypes may be driven by other epigenetic mechanisms, of which the patterns of differential small RNA abundance observed here constitute mere by-products.

The potential adaptive importance of nongenetic inheritance is increasingly appreciated in regard to the responses of animals and plants to changing environments (Yin, Zhou, Lin, Li, & Zhang, 2019). However, despite the animals used in this study being a wild-type strain, our results must be interpreted with caution given the somewhat unique ecology of laboratory zebrafish. Namely, zebrafish kept in laboratory conditions can spawn all year round, in contrast to wild populations for which spawning events are purportedly confined to the monsoon season (Spence, Gerlach, Lawrence, & Smith, 2007). Nevertheless, given the fast spermatogenic cycle in zebrafish (Leal et al., 2009) and assuming frequent spawning opportunities, a male could potentially produce several different spermatozoal epigenotypes throughout the course of a single spawning season, thus transmitting a large degree of cumulative epigenetic variation to the next generation. Although some traits inherited epigenetically may be maladaptive to offspring, the range of possible phenotypic variation generated might be to the overall advantage of the father in increasing the overall likelihood of transmitting his genes to subsequent generations. In other words, epigenetic inheritance may constitute a form of genetically encoded plasticity by which the genome itself can enjoy protection from selective pressures.

To conclude, stress-related behavioural and physiological responses are sensitive to intergenerational perturbation via the paternal germ line in wild-type zebrafish. Our data support the idea that spermatozoal small RNAs are important in this process, as found in rodent models, suggesting that RNA-mediated paternal inheritance is a shared between mammals and fish. Equivalent studies on a broader range of species are needed, however, to determine whether this process is widespread among vertebrates and whether it can facilitate adaptation. Moreover, the mechanistic role of spermatozoal small RNAs requires validation in functional experiments, and the role of other epigenetic factors, particularly DNA methylation and histone modification, should be explored. Finally, the work demonstrates that the zebrafish could serve great utility as a model for the further exploration of paternal epigenetic effects.

ACKNOWLEDGEMENTS

The authors are grateful to Lynsey Gregory and Allison Blake for support with animal husbandry, Helen Eachus for advice with phenotyping methods, Rachel Tucker and Sarah Elliott for technical assistance, and Matthew Wyles for performing RNA library preparation and sequencing. The work benefited greatly from the Sheffield Advanced Research Computer (ShARC) HPC cluster. J.O. was supported by a University of Sheffield scholarship. The work received funding from the University of Sheffield and from the European Union's Horizon 2020 research and innovation programme under grant agreement No. 668989.

AUTHOR CONTRIBUTIONS

P.J.W., A.F. and J.O. developed the initial concept. J.O. designed experiments in consultation with P.J.W. and A.F. J.O. performed animal husbandry, phenotype experiments, RNA extraction and associated methodologies. P.R.H. oversaw RNA quality assessment, library preparation and sequencing, provided technical advice and contributed written methods. J.O. analysed phenotype data, performed all bioinformatic analyses and wrote the manuscript. All authors provided feedback on the written manuscript and approved the submitted manuscript.

ETHICAL APPROVAL

All animal work was carried out under a UK Home Office licence for the use of animals in scientific procedures (licence number 40/3704).

DATA AVAILABILITY STATEMENT

All data and scripts used for the phenotypic analyses (thigmotaxis and cortisol) as well as the complete workflow for the small RNA analyses have been deposited to Dryad (<https://doi.org/10.5061/dryad.hx3ffbgbr>). Raw FASTQ files have been deposited to the NCBI Sequence Read Archive (BioProject PRJNA635603; <https://www.ncbi.nlm.nih.gov/bioproject/PRJNA635603/>). The egg transcriptome data set used for miRNA target analysis (Jiang et al., 2013) is available in the Gene Expression Omnibus (GEO sample GSM1085060; <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSM1085060>; Jiang, 2013).

ORCID

James Ord  <https://orcid.org/0000-0001-6870-8927>

Alireza Fazeli  <https://orcid.org/0000-0003-0870-9914>

REFERENCES

- Al Rawi, S., Louvet-Vallee, S., Djeddi, A., Sachse, M., Culetto, E., Hajjar, C., ... Galy, V. (2011). Postfertilization autophagy of sperm organelles prevents paternal mitochondrial DNA transmission. *Science*, 334(6059), 1144–1147. <https://doi.org/10.1126/science.1211878>
- Andrews, S., Krueger, F., Secondos-Pichon, A., Biggins, F., & Wingett, S. (2015). FastQC. A quality control tool for high throughput sequence data. *Babraham Bioinformatics*. Retrieved from <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>.
- Barton, B. A. (2002). Stress in Fishes: A Diversity of Responses with Particular Reference to Changes in Circulating Corticosteroids.

- Integrative and Comparative Biology*, 42(3), 517–525. <https://doi.org/10.1093/icb/42.3.517>
- Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67(1), 1–48.
- Bell, A. M., McGhee, K. E., & Stein, L. R. (2016). Effects of mothers' and fathers' experience with predation risk on the behavioral development of their offspring in threespined sticklebacks. *Current Opinion in Behavioral Sciences*, 7, 28–32. <https://doi.org/10.1016/j.cobeha.2015.10.011>
- Best, C., Kurrasch, D. M., & Vijayan, M. M. (2017). Maternal cortisol stimulates neurogenesis and affects larval behaviour in zebrafish. *Scientific Reports*, 7, 40905. <https://doi.org/10.1038/srep40905>
- Bestion, E., Teyssier, A., Aubret, F., Clobert, J., & Cote, J. (2014). Maternal exposure to predator scents: Offspring phenotypic adjustment and dispersal. *Proceedings of the Royal Society B: Biological Sciences*, 281(1792), <https://doi.org/10.1098/rspb.2014.0701>
- Bizuayehu, T. T., & Babiak, I. (2014). MicroRNA in teleost fish. *Genome Biology and Evolution*, 6(8), 1911–1937. <https://doi.org/10.1093/gbe/evu151>
- Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics*, 30(15), 2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>
- Butts, C. T. (2008). network: A package for managing relational data in R. *Journal of Statistical Software*, 24(2), 1–36. <https://doi.org/10.18637/jss.v024.i02>
- Calcagno, D. Q., da Mota, E. R., Moreira, F. C., de Sousa, S. B. M., Burbano, R. R., & Assumpção, P. P. (2019). Role of PIWI-interacting RNA (piRNA) as epigenetic regulation. In V. B. Patel & V. R. Preedy (Eds.), *Handbook of nutrition, diet, and epigenetics*, Vol. 1 (pp. 187–209). Cham, Switzerland: Springer International Publishing. https://doi.org/10.1007/978-3-319-55530-0_77
- Champroux, A., Cocquet, J., Henry-Berger, J., Drevet, J. R., & Kocer, A. (2018). A decade of exploring the mammalian sperm epigenome: Paternal epigenetic and transgenerational inheritance. *Frontiers in Cell and Developmental Biology*, 6(50), 1–19. <https://doi.org/10.3389/fcell.2018.00050>
- Chan, P. P., & Lowe, T. M. (2009). GtRNAdb: A database of transfer RNA genes detected in genomic sequence. *Nucleic Acids Research*, 37(Database), D93–D97. <https://doi.org/10.1093/nar/gkn787>
- Chen, G. U., Wang, W., Ding, X., Yang, S., Wang, A., Yin, B., & Wei, W. (2018). Effects of maternal stress induced by predator odors during gestation on behavioral and physiological responses of offspring in Brandt's vole (*Lasiopodomys brandtii*). *Integrative Zoology*, 13(6), 723–734. <https://doi.org/10.1111/1749-4877.12355>
- Chen, Q., Yan, M., Cao, Z., Li, X., Zhang, Y., Shi, J., ... Zhou, Q. (2016). Sperm tsRNAs contribute to intergenerational inheritance of an acquired metabolic disorder. *Science*, 351(6271), 397–400. <https://doi.org/10.1126/science.aad7977>
- Chen, Q., Yan, W., & Duan, E. (2016). Epigenetic inheritance of acquired traits through sperm RNAs and sperm RNA modifications. *Nature Reviews Genetics*, 17(12), 733–743. <https://doi.org/10.1038/nrg.2016.106>
- Clinchy, M., Sheriff, M. J., & Zanette, L. Y. (2013). Predator-induced stress and the ecology of fear. *Functional Ecology*, 27(1), 56–65. <https://doi.org/10.1111/1365-2435.12007>
- Crean, A. J., Dwyer, J. M., & Marshall, D. J. (2013). Adaptive paternal effects? Experimental evidence that the paternal environment affects offspring performance. *Ecology*, 94(11), 2575–2582. <https://doi.org/10.2307/23597219>
- Cunha, M., Berglund, A., Mendes, S., & Monteiro, N. (2018). The 'Woman in Red' effect: Pipefish males curb pregnancies at the sight of an attractive female. *Proceedings of the Royal Society B: Biological Sciences*, 285(1885), 20181335. <https://doi.org/10.1098/rspb.2018.1335>
- Davis, E. P., & Sandman, C. A. (2010). The timing of prenatal exposure to maternal cortisol and psychosocial stress is associated with human infant cognitive development. *Child Development*, 81(1), 131–148. <https://doi.org/10.1111/j.1467-8624.2009.01385.x>
- Ernst, C., Odom, D. T., & Kutter, C. (2017). The emergence of piRNAs against transposon invasion to preserve mammalian genome integrity. *Nature Communications*, 8(1), 1411. <https://doi.org/10.1038/s41467-017-01049-7>
- Evans, J. P., Lymbery, R. A., Wiid, K. S., Rahman, M. M., & Gasparini, C. (2017). Sperm as moderators of environmentally induced paternal effects in a livebearing fish. *Biology Letters*, 13(4), 20170087. <https://doi.org/10.1098/rsbl.2017.0087>
- Feil, R., & Fraga, M. F. (2012). Epigenetics and the environment: Emerging patterns and implications. *Nature Reviews Genetics*, 13(2), 97–109. <https://doi.org/10.1038/nrg3142>
- Gapp, K., Jawaid, A., Sarkies, P., Bohacek, J., Pelczar, P., Prados, J., ... Mansuy, I. M. (2014). Implication of sperm RNAs in transgenerational inheritance of the effects of early trauma in mice. *Nature Neuroscience*, 17(5), 667–669. <https://doi.org/10.1038/nn.3695>
- Garcia-Silva, M. R., Cabrera-Cabrera, F., Güida, M. C., & Cayota, A. (2012). Hints of tRNA-Derived Small RNAs Role in RNA Silencing Mechanisms. *Genes*, 3(4), 603–614. <https://doi.org/10.3390/genes3040603>
- Godoy, L. D., Rossignoli, M. T., Delfino-Pereira, P., Garcia-Cairasco, N., & de Lima Umeoka, E. H. (2018). A comprehensive overview on stress neurobiology: basic concepts and clinical implications. *Frontiers in Behavioral Neuroscience*, 12, 127. <https://doi.org/10.3389/fnbeh.2018.00127>
- Goerg, G. M. (2016). *LambertW: Probabilistic Models to Analyze and Gaussianize Heavy-Tailed, Skewed Data*, R package version 0.6.4.
- Grentzinger, T., Armenise, C., Brun, C., Mugat, B., Serrano, V., Pelisson, A., & Chambeyron, S. (2012). piRNA-mediated transgenerational inheritance of an acquired trait. *Genome Research*, 22(10), 1877–1888. <https://doi.org/10.1101/gr.136614.111>
- Guillaume, A. S., Monro, K., & Marshall, D. J. (2016). Transgenerational plasticity and environmental stress: Do paternal effects act as a conduit or a buffer? *Functional Ecology*, 30(7), 1175–1184. <https://doi.org/10.1111/1365-2435.12604>
- Hakekoh, U., & Hojsgaard, S. (2014). A kenward-roger approximation and parametric bootstrap methods for tests in linear mixed models—the R package pbkrtest. *Journal of Statistical Software*, 59(9), 1–30.
- Heasman, J. (2006). Maternal determinants of embryonic cell fate. *Seminars in Cell & Developmental Biology*, 17(1), 93–98. <https://doi.org/10.1016/j.semcd.2005.11.005>
- Immler, S. (2018). The sperm factor: paternal impact beyond genes. *Heredity*, 121(3), 239–247. <https://doi.org/10.1038/s41437-018-0111-0>
- Jiang, L. (2013). egg.RNA-seq; Gene Expression Omnibus; GSM1085060. <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSM1085060>
- Jiang, L., Zhang, J., Wang, J.-J., Wang, L. U., Zhang, L. I., Li, G., ... Liu, J. (2013). Sperm, but not oocyte, DNA methylome is inherited by zebrafish early embryos. *Cell*, 153(4), 773–784. <https://doi.org/10.1016/j.cell.2013.04.041>
- Kanehisa, M., & Goto, S. (2000). KEGG: Kyoto encyclopedia of genes and genomes. *Nucleic Acids Research*, 28(1), 27–30. <https://doi.org/10.1093/nar/28.1.27>
- Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in Linear Mixed Effects Models. *Journal of Statistical Software*, 82(13), 1–26. <https://doi.org/10.18637/jss.v082.i13>
- Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature Methods*, 9(4), 357–359. <https://doi.org/10.1038/nmeth.1923>
- Leal, M. C., Cardoso, E. R., Nóbrega, R. H., Batlouni, S. R., Bogerd, J., França, L. R., & Schulz, R. W. (2009). Histological and stereological

- evaluation of zebrafish (*Danio rerio*) spermatogenesis with an emphasis on spermatogonial generations. *Biology of Reproduction*, 81(1), 177–187. <https://doi.org/10.1095/biolreprod.109.076299>
- Lee, H., Han, S., Kwon, C. S., & Lee, D. (2016). Biogenesis and regulation of the let-7 miRNAs and their functional implications. *Protein & Cell*, 7(2), 100–113. <https://doi.org/10.1007/s13238-015-0212-y>
- Lenth, R. (2019). *emmeans: Estimated Marginal Means. (Least-Squares Means)*. R package version 1.4.3.
- Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., ... Durbin, R. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, 25(16), 2078–2079. <https://doi.org/10.1093/bioinformatics/btp352>
- Li, W., & Godzik, A. (2006). Cd-hit: A fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics*, 22(13), 1658–1659. <https://doi.org/10.1093/bioinformatics/btl158>
- Lupien, S. J., McEwen, B. S., Gunnar, M. R., & Heim, C. (2009). Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nature Reviews Neuroscience*, 10(6), 434–445. <https://doi.org/10.1038/nrn2639>
- MacDougall-Shackleton, S. A., Bonier, F., Romero, L. M., & Moore, I. T. (2019). Glucocorticoids and “Stress” Are Not Synonymous. *Integrative Organismal Biology*, 1(1), obz017–. <https://doi.org/10.1093/IOB/OBZ017>
- Maestriperi, D., & Hoffman, C. L. (2011). Chronic stress, allostatic load, and aging in nonhuman primates. *Development and Psychopathology*, 23(4), 1187–1195. <https://doi.org/10.1017/S0954579411000551>
- Mathuru, A. S. (2016). Conspecific injury raises an alarm in medaka. *Scientific Reports*, 6, 36615. <https://doi.org/10.1038/srep36615>
- McCarthy, D., Chen, Y., & Smyth, G. (2012). Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation. *Nucleic Acids Research*, 40(10), 4288–4297. <https://doi.org/10.1093/nar/gks042>
- McEwen, B. S. (2004). Protection and damage from acute and chronic stress: Allostasis and allostatic overload and relevance to the pathophysiology of psychiatric disorders. *Annals of the New York Academy of Sciences*, 1032(1), 1–7. <https://doi.org/10.1196/annals.1314.001>
- McGheeK. E., Bell A. M. (2014). Paternal care in a fish: epigenetics and fitness enhancing effects on offspring anxiety. *Proceedings of the Royal Society B: Biological Sciences*, 281, (1794), 20141146 <http://dx.doi.org/10.1098/rspb.2014.1146>
- O’Dea, R. E., Noble, D. W. A., Johnson, S. L., Hesselson, D., & Nakagawa, S. (2016). The role of non-genetic inheritance in evolutionary rescue: Epigenetic buffering, heritable bet hedging and epigenetic traps. *Environmental Epigenetics*, 2(1), dvv014. <https://doi.org/10.1093/EEP/DVV014>
- Ord, J., Holmes, K. E., Holt, W. V., Fazeli, A., & Watt, P. J. (2020). Premature birth stunts early growth and is a possible driver of stress-induced maternal effects in the guppy *Poecilia reticulata*. *Journal of Fish Biology*, 96(2), 506–515. <https://doi.org/10.1111/jfb.14235>
- Otero-Ferrer, F., Lättelkivi, F., Ord, J., Reimann, E., Köks, S., Izquierdo, M., ... Fazeli, A. (2020). Time-critical influences of gestational diet in a seahorse model of male pregnancy. *Journal of Experimental Biology*, 223(3), jeb210302. <https://doi.org/10.1242/JEB.210302>
- Pages, H., Carlson, M., Falcon, S., & Li, N. (2018). AnnotationDbi: Annotation database interface. R Package Version 1.42.1. <https://doi.org/10.18129/B9.bioc.AnnotationDbi>
- Pauli, A., Rinn, J. L., & Schier, A. F. (2011). Non-coding RNAs as regulators of embryogenesis. *Nature Reviews. Genetics*, 12(2), 136–149. <https://doi.org/10.1038/nrg2904>
- Potok, M. E., Nix, D. A., Parnell, T. J., & Cairns, B. R. (2013). Reprogramming the maternal zebrafish genome after fertilization to match the paternal methylation pattern. *Cell*, 153(4), 759–772. <https://doi.org/10.1016/j.cell.2013.04.030>
- R Core Team (2018). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Rao, R., & Androulakis, I. P. (2019). Allostatic adaptation and personalized physiological trade-offs in the circadian regulation of the HPA axis: A mathematical modeling approach. *Scientific Reports*, 9(1), 11212. <https://doi.org/10.1038/s41598-019-47605-7>
- Räsänen, K., & Kruuk, L. E. B. (2007). Maternal effects and evolution at ecological time-scales. *Functional Ecology*, 21(3), 408–421. <https://doi.org/10.1111/j.1365-2435.2007.01246.x>
- Redfern, J. C., Cooke, S. J., Lennox, R. J., Nannini, M. A., Wahl, D. H., & Gilmour, K. M. (2017). Effects of maternal cortisol treatment on offspring size, responses to stress, and anxiety-related behavior in wild largemouth bass (*Micropterus salmoides*). *Physiology & Behavior*, 180, 15–24. <https://doi.org/10.1016/j.physbeh.2017.08.001>
- Robinson, J. T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E. S., Getz, G., & Mesirov, J. P. (2011). Integrative genomics viewer. *Nature Biotechnology*, 29(1), 24–26. <https://doi.org/10.1038/nbt.1754>
- Rodgers, A. B., Morgan, C. P., Bronson, S. L., Revello, S., & Bale, T. L. (2013). Paternal stress exposure alters sperm MicroRNA content and reprograms offspring HPA stress axis regulation. *Journal of Neuroscience*, 33(21), 9003–9012. <https://doi.org/10.1523/JNEUROSCI.0914-13.2013>
- Rodgers, A. B., Morgan, C. P., Leu, N. A., & Bale, T. L. (2015). Transgenerational epigenetic programming via sperm microRNA recapitulates effects of paternal stress. *Proceedings of the National Academy of Sciences of the United States of America*, 112(44), 13699–13704. <https://doi.org/10.1073/pnas.1508347112>
- Rodríguez Barreto, D., García de Leaniz, C., Verspoor, E., Sobolewska, H., Coulson, M., & Consuegra, S. (2019). DNA methylation changes in the sperm of captive-reared fish: A route to epigenetic introgression in wild populations. *Molecular Biology and Evolution*, 36(10), 2205–2211. <https://doi.org/10.1093/molbev/msz135>
- Rosenkranz, D. (2016). piRNA cluster database: A web resource for piRNA producing loci. *Nucleic Acids Research*, 44(D1), D223–D230. <https://doi.org/10.1093/nar/gkv1265>
- Sarker, G., Sun, W., Rosenkranz, D., Pelczar, P., Opitz, L., Efthymiou, V., ... Peleg-Raibstein, D. (2019). Maternal overnutrition programs hedonic and metabolic phenotypes across generations through sperm tsRNAs. *Proceedings of the National Academy of Sciences of the United States of America*, 116(21), 10547–10556. <https://doi.org/10.1073/pnas.1820810116>
- Schloerke, B., Crowley, J., Cook, D., Hofmann, H., Wickham, H., Briatte, F., ... & Larmarange, J. (2018). *GGally: Extension to 'ggplot2'*, R package version 1.4.0.
- Sharma, U., Conine, C. C., Shea, J. M., Boskovic, A., Derr, A. G., Bing, X. Y., ... Rando, O. J. (2016). Biogenesis and function of tRNA fragments during sperm maturation and fertilization in mammals. *Science*, 351(6271), 391–396. <https://doi.org/10.1126/science.aad6780>
- Spence, R., Gerlach, G., Lawrence, C., & Smith, C. (2007). The behaviour and ecology of the zebrafish, *Danio rerio*. *Biological Reviews*, 83(1), 13–34. <https://doi.org/10.1111/j.1469-185X.2007.00030.x>
- Steenbergen, P. J., Richardson, M. K., & Champagne, D. L. (2011). The use of the zebrafish model in stress research. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, 35(6), 1432–1451. <https://doi.org/10.1016/j.pnpb.2010.10.010>
- Stensmyr, M. C., & Maderspacher, F. (2012). Pheromones: Fish fear factor. *Current Biology*, 22(6), R183–R186. <https://doi.org/10.1016/j.cub.2012.02.025>
- Stephenson, J. F. (2016). Keeping eyes peeled: Guppies exposed to chemical alarm cue are more responsive to ambiguous visual cues. *Behavioral Ecology and Sociobiology*, 70, 575–584. <https://doi.org/10.1007/s00265-016-2076-4>
- Tauffmanberger, A., & Parker, J. A. (2014). Heritable transmission of stress resistance by high dietary glucose in *Caenorhabditis elegans*. *PLoS Genetics*, 10(5), e1004346. <https://doi.org/10.1371/journal.pgen.1004346>

- Vejnar, C. E., Blum, M., & Zdobnov, E. M. (2013). miRmap web: comprehensive microRNA target prediction online. *Nucleic Acids Research*, 41(W1), W165–W168. <https://doi.org/10.1093/nar/gkt430>
- Vitousek, M. N., Taff, C. C., Hallinger, K. K., Zimmer, C., & Winkler, D. W. (2018). Hormones and fitness: evidence for trade-offs in glucocorticoid regulation across contexts. *Frontiers in Ecology and Evolution*, 6, 42. <https://doi.org/10.3389/fevo.2018.00042>.
- Wagner, D. S., Dosch, R., Mintzer, K. A., Wiemelt, A. P., & Mullins, M. C. (2004). Maternal control of development at the midblastula transition and beyond: Mutants from the zebrafish II. *Developmental Cell*, 6(6), 781–790. <https://doi.org/10.1016/j.devcel.2004.04.001>
- Weyrich, A., Lenz, D., Jeschek, M., Chung, T. H., Rübensam, K., Göritz, F., ... Fickel, J. (2016). Paternal intergenerational epigenetic response to heat exposure in male Wild guinea pigs. *Molecular Ecology*, 25(8), 1729–1740. <https://doi.org/10.1111/mec.13494>
- Wickham, H. (2011). ggplot2. *Wiley Interdisciplinary Reviews: Computational Statistics*, 3(2), 180–185. <https://doi.org/10.1002/wics.147>
- Wienholds, E., Kloosterman, W. P., Miska, E., Alvarez-Saavedra, E., Berezikov, E., de Bruijn, E., Plasterk, R. H. A. (2005). MicroRNA Expression in Zebrafish Embryonic Development. *Science*, 309(5732), 310–311. <https://doi.org/10.1126/science.1114519>
- Yeh, C. M., Glöck, M., & Ryu, S. (2013). An optimized whole-body cortisol quantification method for assessing stress levels in larval zebrafish. *PLoS One*, 8(11), 1–8. <https://doi.org/10.1371/journal.pone.0079406>
- Yin, J., Zhou, M., Lin, Z., Li, Q. Q., & Zhang, Y. (2019). Transgenerational effects benefit offspring across diverse environments: A meta-analysis in plants and animals. *Ecology Letters*, 22(11), 1976–1986. <https://doi.org/10.1111/ele.13373>
- Yu, G., Wang, L. G., Han, Y., & He, Q. Y. (2012). ClusterProfiler: An R package for comparing biological themes among gene clusters. *OMICS A Journal of Integrative Biology*, 16(5), 284–287. <https://doi.org/10.1089/omi.2011.0118>
- Zajitschek, F., Zajitschek, S., & Manier, M. (2017). High-protein paternal diet confers an advantage to sons in sperm competition. *Biology Letters*, 13(2), <https://doi.org/10.1098/rsbl.2016.0914>
- Zajitschek, S., Herbert-Read, J. E., Abbasi, N. M., Zajitschek, F., & Immler, S. (2017). Paternal personality and social status influence offspring activity in zebrafish. *BMC Evolutionary Biology*, 17(1), 157. <https://doi.org/10.1186/s12862-017-1005-0>
- Zajitschek, S., Hotzy, C., Zajitschek, F., & Immler, S. (2014). Short-term variation in sperm competition causes sperm-mediated epigenetic effects on early offspring performance in the zebrafish. *Proceedings of the Royal Society B: Biological Sciences*, 281(1785), 20140422. <https://doi.org/10.1098/rspb.2014.0422>
- Zanette, L. Y., White, A. F., Allen, M. C., & Clinchy, M. (2011). Perceived predation risk reduces the number of offspring songbirds produce per year. *Science*, 334(6061), 1398–1401. <https://doi.org/10.1126/science.1210908>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Ord J, Heath PR, Fazeli A, Watt PJ. Paternal effects in a wild-type zebrafish implicate a role of sperm-derived small RNAs. *Mol Ecol*. 2020;00:1–14. <https://doi.org/10.1111/mec.15505>