White Rose Research Online

eprints@whiterose.ac.uk
https://eprints.whiterose.ac.uk
Universities of Leeds, Sheffield and York

Deposited via The University of Leeds.

White Rose Research Online URL for this paper:
https://eprints.whiterose.ac.uk/id/eprint/163212/

Version: Accepted Version

Article:
Butler, RWH, Lickorish, HW, Vinnels, J et al. (2020) Untangling the Annot sand fairway: structure and stratigraphy of the Eastern Champsaur Basin (Eocene-Oligocene), French Alps. Journal of the Geological Society, 177 (6). jgs2020-015. pp. 1197-1209. ISSN: 0016-7649

https://doi.org/10.1144/jgs2020-015

Reuse
Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown
If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

UNIVERSITY OF LEEDS
University of Sheffield
UNIVERSITY of York

revised text

Click here to access/download;text file;Champsaur_revised__FINAL.docx

Untangling the Annot sand fairway: structure and stratigraphy of the Eastern Champsaur Basin (Eocene-Oligocene), French Alps.

Robert W.H. Butler[1], Henry W. Lickorish[2], Jamie Vinnels[3] & William D. McCaffrey[2]

1 Geology and Geophysics, School of Geosciences, University of Aberdeen, Aberdeen AB24 3UE, UK
2 School of Earth and Environment, University of Leeds, Leeds LS2 9JT, UK
3 Equinor, 2107 City West Blvd, Houston, Texas, 77042, USA

Orchid: RWHB, 0000-0002-7732-9686
Correspondence: rob.butler@abdn.ac.uk

ABSTRACT

Early foredeep successions can yield insight on tectonic processes operating adjacent to and ahead of fledgling orogenic belts but are commonly deformed by the same orogens. We develop a workflow towards stratigraphic understanding of these deformed basins, applied to the Eastern Champsaur Basin of the French Alps. This contains a down-system correlative of the southern-sourced (Eocene-Oligocene) Annot turbidites. These strata are deformed by arrays of W-facing folds that developed beneath the Embrunais-Ubaye tectonic allochthon. The folds vary in geometry through the stratigraphic multilayer. Total shortening in the basin is around 4 km and the restored (un-decompacted) stratal thickness exceeds 980m. The turbidites are generally sand-rich and bed-sets can be correlated through the entire fold train. The succession shows onlap and differential thickening indicating deposition across palaeobathymetry that evolved during active basement deformation, before being over-ridden by the allochthon. The sand system originally continued over what is now the Ecrins basement massif that, while contributing to basin floor structure, only served to confine and potentially

focus further sediment transport to the North. Deformation ahead of the main Alpine orogen appears to have continued progressively, and the past definition of distinct “phases” (“pre-“ and “post-Nummulitic”) is an artefact of the stratigraphic record.

Keywords: turbidites, structural restoration, foreland basin,

Turbidite systems can offer important constraints on bathymetric continuity of arrays of sedimentary basins (e.g. Smith 2004; Puigdefabregas *et al.* 2004) and thus inform models of basin evolution. For ancient foredeep systems the challenge is to unravel useful stratigraphic information from successions that are involved in orogenic belts and then deformed. If stratigraphic information can be extracted from deformed systems, the insights so gained can then not only inform models of orogenic evolution but also impact on greater tectono-stratigraphic understanding of turbidite systems. This is especially important as many of the key ideas of turbidite sedimentation come from studies of successions in syn-orogenic sedimentary basins. The aims of this paper are two-fold. The first is to document a workflow for developing stratigraphic knowledge from highly folded successions. The second is to provide a linked stratigraphic and structural reconstruction of the Eastern Champsaur Basin – a deformed part of the Eocene-Oligocene Annot Sandstone system of SE France (the Grès d’Annot in Alpine literature: e.g. Sinclair 1997; Joseph & Lomas 2004) that has become incorporated into the Western Alpine orogen. While an extensive re-examination of the Annot system in the light of our reconstructions lies beyond the scope of this paper, we do explore some implications for the extent of the system, how crustal-scale deformation migrates through time and how tectonic evolution is represented in the stratigraphic record.

Geological context and motivation

The Annot system of SE France as a whole occupies a pivotal position in turbidite research, with outcrops providing test-beds for models of deep-water sedimentation and analogues for hydrocarbon reservoirs in the

subsurface. Bouma's pioneering descriptions of turbidite facies come from proximal parts of the Annot system (e.g. Bouma 1959; Stanley & Bouma 1964). This turbidite system was deposited in a broadly north-south trending foredeep basin, adjacent to the ancestral Alps and fed from eroding granitic basement to the south (Maures-Esterel massif, Fig. 1a) together with Corsica that, at the time, lay adjacent to the south coast of mainland France. Much of the early research is cited by Joseph & Lomas (2004) and papers that follow this introduction greatly expand upon these works. More recent studies include Mulder *et al.* (2010), Etienne *et al.* (2012), Salles *et al*. (2014), and Cunha *et al.* (2017).

Existing work has almost entirely focussed on sites that have experienced rather little deformation. The system is known to continue into more strongly deformed regions to the north and northeast (e.g. Sinclair 1997). By increasing understanding of the Annot system, insight is gained on the continuity and early deformation of the Alpine "foreland". As subaqueous gravity flows, turbidity currents seek low bathymetry. Therefore, the continuity of sand fairways, the geological record of the routes taken by their causative flows, provides evidence for relative bathymetry in front of the evolving Alpine orogen. The various relationships between the turbidites and their substrate chart not only the syndepositional deformation of the foredeep basin floor but also the deformation that predated basin subsidence. This understanding is important for evaluating the tectonic significance of unconformities associated with orogens and their implications for evaluating the progression of continental deformation. By increasing understanding of the Annot system, insight is gained on the continuity and early deformation of the Alpine "foreland".

Our case study here focusses on the south-eastern flank of the external Alpine basement massif of the Ecrins (Fig. 1a). The Eocene-Oligocene litho-stratigraphy of the area, in common with the main Annot system (Fig 2; e.g. Joseph & Lomas 2004), consists of three main lithofacies (Debelmas *et al.* 1980) that comprise the "Nummulitic trilogy". Collectively this represents the foredeep megasequence developed during this early stage of Alpine orogenesis. The megasequence oversteps a substrate of crystalline basement, continuous with the neighbouring Ecrins massif, together with

remnant patches of the Triassic-Jurassic cover to the massif (e.g. Debelmas *et al.* 1980). The eastern and southern margins to the basin are now buried beneath the tectonic allochthon of the Embrunais-Ubaye thrust sheets.

The first part of the Nummulitic trilogy is the Nummulitic Limestone, a shallow-water transgressive unit here of Priabonian age (Debelmas *et al.* 1980; Dumont *et al*. 2012). Locally, the Nummulitic Limestone is underlain by conglomerates derived from the adjacent crystalline basement, interpreted to be the fills to subaerially-incised palaeovalleys (e.g. Gupta 1997). The second part of the Nummulitic trilogy are the Blue Marls ("Marnes à Globigérines" on published maps, e.g. Debelmas *et al.* 1980). These calcareous mudstones chart palaeo-bathymetries increasing to several hundred metres depth. The final part of the trilogy are turbidites – the Champsaur Sandstone ("Grès du Champsaur" on published maps, e.g. Debelmas *et al.* 1980). Locally, there is a facies intermediate between these turbidites and the Blue Marls, the Brown Marls (e.g. Stanbrook & Clark 2004). These are generally inferred to have accumulated on local intra-basin highs from dispersed suspension clouds associated with the turbidity currents that deposited the Champsaur (and main Annot) Sandstone.

The Nummulitic trilogy is capped by a shaley olistostrome termed the "Schistes à Blocs" that lies directly under the Embrunais-Ubaye thrust sheets. It is classically inferred to have been shed off the advancing thrust sheets as the foredeep basin was closed (Kerckhove 1969). Detailed sedimentological investigation of the Schistes à Blocs in the Eastern Champsaur Basin is strongly inhibited by the intense penetrative deformation within this unit, which presumably reflects distributed shear deformation in the footwall to the overlying thrust sheets. Consequently, it is generally difficult to establish whether the unit represents a series of amalgamated debris flow deposits or is the product of a single submarine mass-wasting event.

Although given the single designator "Champsaur Sandstone", early Oligocene siliciclastic sandstones along the southern flank of the Ecrins massif have two distinct compositions and, by inference, provenances, lying in distinct depocentres (Debelmas *et al.* 1980). Following our previous usage (Vinnels *et al.* 2010; Butler 2017), we term these depocentres the Western and Eastern Champsaur Basins. They are separated by a zone of

136 deformation generally termed the Selle Fault (e.g. Tricart 1981; Ford 1996
137 and others since). To date, sedimentological research in the region has
138 mainly focussed on the westerly-derived, volcaniclastic turbidites that outcrop
139 in the Western Champsaur Basin, with detailed descriptions not only of
140 depositional architectures (e.g. Brunt *et al.* 2007) but also of the discrete fold-
141 thrust structures that have deformed the successions (Butler & McCaffrey
142 2004). Separated from this western basin by the Selle Fault Zone, the Eastern
143 Champsaur Basin is significantly more deformed than its western counterpart
144 (Fig. 3). It is characterised by folds on various wavelengths and amplitudes
145 that generally have tight hinges and straight limbs, approximating to chevron
146 geometry. This style of deformation is typical of successions with a
147 pronounced planar mechanical anisotropy such as is provided by inter-
148 bedded sandstones and shales (Ramsay 1974). The folds face westwards
149 with axial surfaces inclined moderately to the east. The structures were
150 described by Kerckhove *et al.* (1978) and studied in detail by Bürgisser & Ford
151 (1998). These studies show that the folds developed beneath, and locally
152 incorporate, the Embrunais Ubaye thrust sheets which overran both the
153 Eastern and Western Champsaur Basins. The thrust sheets provided tectonic
154 burial sufficient for peak temperatures in Eastern Champsaur of c 300 °C
155 (Bellanger *et al.* 2015).
156 Provenance of the Eastern Champsaur Sandstone has been disputed.
157 Sinclair (1997) depicts their turbidites to be the down-system continuation of
158 the Annot system otherwise preserved in the SW Alps of Provence. However,
159 in their regional overview, Joseph & Lomas (2004) indicate a northern
160 provenance, sourced from the an elevated, proto-Ecrins massif. The
161 argument is resolved by Vinnels *et al.* (2010) who present palaeocurrent data
162 to show a southerly derivation and document that the sandstones of the
163 Eastern Champsaur basin have the same composition as the Annot
164 Sandstone in its type area to the south. Thus Sinclair's (1997) view is
165 supported. Vinnels *et al.* (2010) note that, although deflected by subtle basin-
166 floor topography, the causative turbidity currents continued over what is now
167 the uplifted Ecrins basement massif. However, Vinnels *et al.* only focused on
168 the lower stratigraphic intervals, establishing facies variations and palaeoflow
169 deflections along the basal onlap surface of turbidites onto the Blue Marls that

drape uplifted substrate within the basin. The current paper focusses exclusively on the Eastern Champsaur Basin, building on our previous work (Vinnels *et al.* 2010). The regional structural setting is discussed extensively by Butler 2017). The stratigraphic relationships between the eastern and western Champsaur Basins and the role of the Selle Fault in partitioning distinct deep-water systems lie beyond the scope of the current paper and are reserved for later contributions.

A workflow

Geologists working in mountain belts have long had to grapple with the complexities of deformation when attempting to build stratigraphic knowledge. However, the methods used for creating this understanding are rarely documented, making it difficult to assess uncertainties in the larger-scale basin reconstructions. Furthermore, there has been a reluctance for many researchers investigating sedimentary basins associated with mountain belts to examine their more-deformed components. In presenting a workflow here we hope not only to promote research within these settings but also to encourage clearer documentation of how deformed stratigraphic sections are reconstructed.

The Champsaur area exhibits just under 2 km of topographic relief with cliff sections that are of limited accessibility, attributes that are typical of much of the Alpine mountain ranges and of young orogenic belts in general. A virtue of the relief in our Champsaur study area is that major hillsides trend east-west, perpendicular to the trend of fold axes. They therefore provide ideal cross-sections, akin to in-lines in grids of seismic reflection data. Accessible routes up these hillsides provide excellent vertical sections, equivalent to boreholes for subsurface examples. Consequently, we apply a workflow similar to that adopted for surface stratigraphic-structural mapping, tying detailed stratigraphic observations from vertical sections (boreholes) to remotely sensed cross-sections (in this case by direct observation from opposite valley-sides, equivalent to seismic profiles).

The first decision lies in selecting the best vertical section from which to build a stratigraphic succession. The west face of Le Piquet provides a strike

section, 3.5 km across, that runs from up from the base-Nummulitic unconformity to a small klippe of Embrunais-Ubaye thrust sheets (Fig. 5a). Only the upper part of this hillside provides continuous outcrop that is also amenable to direct observation – and this yields stratigraphic logs (Fig. 5a). Logged units include a c 380m interval that is sandstone-dominated (labelled A6 on Fig. 5a), capped by Schistes à Blocs. However, the broad bed-set characteristics of the underlying succession may be established by remote observation. These include shale-dominated levels that are more recessive on hillsides (A3 and A5 on Fig. 5a).

The boundaries between the bed-set units can be traced through the hillsides on Le Piquet and across the Dourmillouse Valley onto the dip section provided by its northern slopes (Fig. 5b). The upper part of the valley side, including the ridge line, provides near continuous, if largely inaccessible, outcrop within which the main folds can be identified. The fold axial traces can be mapped down dip into the Dourmillouse Valley. Then the bed-set stratigraphic units of the Champsaur sandstone can be traced through the fold train. This section is moderately accessible and locally provides accessible sections for detailed stratigraphic logging.

The axial traces of the folds provide the key linking structures to tie adjacent transects. These can be traced over the ridge line bounding the Dourmillouse Valley to the north and into the adjacent Fornel Valley (Fig. 4). The northern side of the ridge-line is much steeper than its southern counterpart so that accessibility is severely restricted. However, fold axial traces may be readily traced through the cliffs and the bed-set stratigraphy defined and similarly tracked through the fold array (Fig. 5c). Note that there is some polyharmonic folding but the axial surfaces for the main fold closures may be traced from the ridge-line down to the Blue Marls.

The bed-set stratigraphy

Following the workflow laid out above it is possible to map out not only the axial traces of the principal folds but also the bed-set stratigraphic units within the Champsaur sandstone (Fig. 4). Although structural deformation hinders identification of individual beds across the whole basin, thicker

intervals can be laterally traced continuously for several kilometres and probably extend much further. Vertical stratigraphic sections display significant heterogeneity in sandstone to shale ratio. These are well-illustrated by the cliff sections of Le Piquet (Fig. 5a),

As noted above, it is the variety in sandstone-shale abundance that provides a framework, on the bed-set scale, for dividing the basin stratigraphy into mappable units (Fig. 5). These units are illustrated on arrays of composite logs (Fig. 6). Principally, our sedimentary logs were measured from the fold limbs to avoid hinge areas, where beds show significant layer parallel shortening strain manifested by cleavage, and tracts of strongly overturned strata where bed thinning is likely. Strains in both of these structural settings are localised in the shales – sandstone beds retain unmodified sedimentary structures suggesting penetrative strains in these units are very low. We estimate an uncertainty of less than 10% on the normally-compacted thickness of sandstones and perhaps up to 20% for the shales on the values reported in Fig. 6, as we cannot rule out deformation even in the normal limbs of folds. The bed-set sequences may still be identified regardless of strain variations and we now use these to erect a basin-wide lithostratigraphy.

The Champsaur Sandstone onlaps its substrate towards the western basin margin (Vinnels *et al*. 2010 and references therein): the oldest preserved parts of the sandstone sequence are only present on the vegetated lower slopes of the far eastern exposures of the basin. In the Narreyroux valley (log location 4 on Fig. 4), a 670m thick section of Champsaur Sandstone (A4 on Fig. 6) is underlain to the east by considerably more stratigraphy but precise correlation has proven elusive. The higher parts of the stratigraphy are more tractable. Two levels are especially sandstone-poor and form key markers (A3 and A5 in Fig. 6). The lower of these two shaley intervals is approximately 50m thick and contains a distinctive couplet of 2 m thick sandstone beds that can be found in all sections that include this stratigraphic interval. There are a few thin sandstone beds present in the higher shale-dominated interval (A5 in Fig. 4) but these are laterally discontinuous. Between the two shaley sections lies an interval with abundant thick sandstone beds (A4 in Fig 4). The section below the lower shaley interval also has higher sandstone content; its upper portion contains

beds that are 3-5 m thick while the lower portion is bedded on the 50 cm – 1 m scale. These distinctions provide confirmations of stratigraphic correlations to elsewhere within the Eastern Champsaur district.

The lower shaley interval (A3) can be readily correlated between the Narreyroux, Fournel and Dourmillouse valleys (Fig. 4) to parts of the study area where more complete sections of the underlying sandstone-rich intervals are exposed. In these places we can separate a thicker-bedded upper component from a thinner-bedded lower component (A2 and A1 respectively in Fig. 6). The thicker-bedded interval (A2) is about 40 m thick. The logs from the Fournel and Dourmillouse valleys (Fig. 6) only demonstrate about 60 m of this lower portion. However, in the Narreyroux valley it achieves a thickness in excess of 150 m with a further 200 m represented by the poorly exposed and deformed terrain deeper in the section.

The upper shaley horizon defined in the Narreyroux valley (A5 on Fig. 6) can be correlated across the Dourmillouse valley (Fig. 4) to the cliffs of Le Piquet. This is the “Thick Shale” of Vinnels *et al.* (2010). Overlying this, 380 m of sandstone-rich turbidies (A6 on Fig. 4) are capped by the “Schistes a’ Blocs” that directly underlie the Embrunais-Ubaye thrust sheets. The sandstone rich interval (A6) can be traced through the fold structures of the upper Dourmillouse and Fournel Valleys, and constitute the upper 60m of the logged section in the Narreyroux valley (Fig. 4). In the upper Fournel Valley the thickness of this interval exceeds 400 m (log 5 on Fig. 6). Individual beds may be traced through folds and across the ridge line between the upper parts of the Dourmillouse and Fournel valleys. For example, a prominent shale interval (5-15 m thick; X on logs 5 and 6 on Fig. 6), at the base of interval A6b) can be traced between sections.

The Eastern Champsaur Basin continues over the watershed at the head of the Dourmillouse Valley into drainage of the Champoleon Valley (Fig. 4). Individual beds may be traced along the ridge line (Pointe des Estaris - Pointe des Pisses: log locations 8 and 9, Fig. 4) to complete the upper stratigraphy of this part of the Champsaur Sandstone up to the Schistes à Blocs (Fig. 6).

By combining the logs, we estimate that about 950m of Champsaur Sandstone stratigraphy (A6 – A7) overlies the prominent upper shaley horizon

306 (A5) in the ground between the upper Dourmillouse valley and Pointe des
307 Pisses (Fig. 4). The equivalent interval at Le Piquet is just 380m thick (log 1
308 on Fig. 6). Both sections are capped by the Schistes à Blocs so the thickness
309 difference is not due to post-depositional tectonic truncation. The difference is
310 also too great to be due to variations in distributed deformation or differential
311 compaction and there are significant differences in the bed-set stratigraphy
312 between these locations. We interpret the thickness variation to result from
313 differential stratigraphic growth across the Eastern Champsaur Basin.
314
315 Depositional architecture and growth
316 Although folding permits construction of substantial stratigraphic
317 sections and deformation has not been sufficient to prevent individual bed-
318 tracing through the high Alpine landscape, it does inhibit the recognition of
319 elements of depositional architecture that might be expressed at the km-scale
320 (such as the development of compensating turbidite lobes or, depending on
321 the scale, of erosional or constructional channels). In the Western Champsaur
322 basin, adjacent to our study area (but forming a distinct turbidite system),
323 extensive deep incision and former submarine canyons have been described
324 (e.g. Brunt *et al.* 2007). However, there is no evidence of incision within the
325 Eastern Champsaur Basin, beyond local scours (and associated layers of
326 mudstone rip-up clasts) and bed-scale amalgamation, nor of channel-levee
327 development. The 4-5 km strike section on the western face of Le Piquet
328 (located on Fig. 4, illustrated on Fig. 5a) reveals very low-angle discordances
329 within parts of the upper sandstone interval (A6) consistent with an expansion
330 of stratigraphic thickness from north to south (Vinnels *et al.* 2010). Individual
331 beds appear to thin northwards while bed-sets abut onto underlying beds –
332 relationships we interpret as internal onlap. Collectively these relationships
333 are plausibly interpreted as representing syn-depositional tilting within the
334 Eastern Champsaur Basin elevating Le Piquet area relative to the Pointe des
335 Pisses area during deposition of these upper sandstone intervals. Vinnels *et*
336 *al.* (2010) suggest that these thickness changes reflect differential growth of
337 an anticline in the basin floor. We infer that the intrabasinal slopes during
338 deposition of at least this part of the Champsaur Sandstone were relatively
339 gentle, there being no evidence for significant mass-wasting within the

succession. Therefore, evolving basin floor structures were largely swamped by concurrent sedimentation.

The post-depositional structure of the basin

The stratigraphic template of mappable bed-sets permits full correlation of units through the fold systems. These correlations are illustrated on our summary map (Fig. 4) and underpin two cross-sections through the Eastern Champsaur Basin (Fig. 7). The sections are necessarily simplified at the scale reproduced here, and no attempt is made here to illustrate detail of the thickness changes in the younger units (A6-A7), nor the precise geometry of polyharmonic folding deeper in the stratigraphic pile (A1).

The two cross-sections (Fig. 7) show a similar structural style: a train of asymmetric, west-facing folds with axial surfaces dipping towards the east. The folds detach downwards into the Blue Marls (Bürgisser & Ford 1998). Thrust faults are rare, show bed offsets of a few metres and are restricted to fold hinge zones. The attitude of the fold axial surfaces and fold interlimb angles vary through the region and, according to Bürgisser & Ford (1998), these variations reflect heterogenous simple shear strains distributed into the footwall of the Embrunais-Ubaye thrust sheets. Folds occur in clusters, separated by segments where stratigraphic sections are only weakly deformed. Within the clusters the folds have wavelengths of 200-500m and amplitudes of around 200m. The larger folds are broadly harmonic, with axial surfaces that can be traced through the visible succession of the Champsaur Sandstone. Prime examples are those folds that cross the ridge-line on the north side of the Dourmillouse valley near the summit of Pic Felix Neff (Figs 5b,c, 7). The distribution of these larger folds is consistent with a model that considers folds as ductile equivalents to imbricate thrusts (e.g. Pfiffner 1985; Butler 1992). Qualitatively, the general form of the folds appears to be influenced, at least in part, by variations in bed-thickness in the Champsaur Sandstone. The thicker, sandstone rich sequences (e.g. Fig. 2a; A6 on Figs 4 and 5) appear to act as “control units”, in the sense of Price & Cosgrove (1990), determining the wavelength of the main fold sets (Fig. 8). The older, thinner-bedded units (Fig. 2; A2-A4 on Figs 4 and 5) deform poly-harmonically

with respect to the younger, thick-bedded stratigraphy (e.g. unit A6), and are characterised by folding at short-wavelength. Together with this layer-dependence, the location of folds in the Champsaur Sandstone may relate to variations in the properties of the basal detachment zone in the Blue Marls, as is well-known in other detachment systems (e.g. Cotton & Koyi 2000 and many others). Less effective detachment would promote folding in the overlying strata. These aspects of the deformation await more detailed investigations.

The cross-sections (Fig. 7) demonstrate the regional discordance between the lithostratigraphic units in the Champsaur Sandstone and the Blue Marls. As noted by Vinnels *et al.* (2010, but well-known informally), this represents a regional onlap surface, albeit modified by the tectonic detachment. It charts deformation of the basin floor during the accumulation of the Nummulitic trilogy. Vinnels *et al.* (2010) treat the onlap angle qualitatively. To consider the angle of this onlap in profile, and to establish an estimate of tectonic shortening in the basin, we present a balanced and restored section (Fig. 8) for the Dourmillouse transect (Fig. 7b).

The restoration assumes that the sand-rich parts of the succession deformed largely by concentric folding, without appreciable changes in bed thickness by distortional strain. To track this assumption, we chart the thicknesses of bed-sets through the cross-section. These thicknesses are reported on both the final state and restored cross-sections (thin bed-perpendicular rulings). With the assumption of concentric folding, restorations can be performed simply by restoring the length of strata, conserving the sinuous lengths measured on the final state section. This is line-length restoration, as proposed by Dahlstrom (1969). Note that in this analysis we do not consider the impact of broad arching of the enveloping surfaces of the folds with in the Champsaur Sandstone. This longer-wavelength deformation couples with the underlying basement and it is the architecture of the basin fill that we wish to reconstruct.

The graphical restoration (Fig. 8b) is hung from the younger stratigraphic levels (A6), the top of which forms a horizontal datum, or “target horizon”, for determining the progress of deformation. This interval, together with those below, retain bed-length and bed-thickness from the deformed

state. A complete restoration might be expected to yield strata that are planar and parallel – features that are not evident here. There are various possible explanations for these discrepancies. First, distortional strains, especially within the mudstone-rich intervals, have not been considered in our method, which assumes ideally concentric folding. Certainly, these units are cleaved, implying that they have experienced such distortions. There may also be errors in the drafting of the cross-section, unavoidable given variations in outcrop quality and uncertainties in the precise position of stratigraphic boundaries. Errors arising from these effects amplify away from the "target horizon", creating significant short-wavelength relief on the base of section. These are regarded here to be restoration artefacts (selected examples identified in in Fig. 8b), as the top-basement surface is readily mapped and shows no such abrupt changes in relief where observed in the field.

Notwithstanding the short-wavelength artefacts, the restoration (Fig. 8b) can be used to calculate the angular relationship between the depositional units in the Champsaur Sandstone and the underlying Blue Marls. The restoration implies onlap of 980m of Champsaur Sandstone over 9.4 km – implying that the average angle of the onlap slope was 6 degrees. Note that this angle would have been significantly higher at the time of deposition as we have not allowed for vertical compaction during burial.

Tectonic shortening for the upper stratal levels (A6) is the difference in length between deformed and undeformed state, 4 km. However, the mismatches in restored lengths of the underlying strata indicate that the section is not balanced (the bed-lengths and implied longitudinal strain are not equal in layers, so that the restoration is not complete). We resolve this by inferring significant "top-to-the-west" shear distributed through the Champsaur Sandstone, as invoked by Bürgisser & Ford (1998). Variations in layer-parallel shortening can account for the open folding of underlying sandstone units and in the substrate.

Discussion – implications for tectonostratigraphic evolution

A palaeogeographic sketch is provided (Fig. 9), extending that of Joseph & Lomas 2004, their Fig. 5) which integrates our findings from the Eastern

Champsaur Basin into a semi-regional context. Following our earlier work (Vinnels *et al*. 2010), we consider its turbiditic basin fill to be a down-system continuation of the Annot sand fairway. We show the main Annot turbidite system to be derived from a southern source area and then routed along the sinuous Provencal basins defined by the underlying deformation within the broad foredeep basin (e.g. Salles *et al.* 2012). This system is distinct from a secondary turbidite system that feeds the Western Champsaur Basin.

Down-system continuity of the Annot Sand Fairway

The total (compacted) stratal thickness of the Champsaur Sandstone in the Western Champsaur Basin approaches 1km. Stratal patterns together with onlap at the base and within the Champsaur Sandstone indicate active deformation of the underlying basement during deposition. We propose that it is this deformation that provided the basin floor structure that served to confine and guide turbidity currents, facilitating sediment transport further down-system. The high proportion of coarse sandstone compared to finer-grained fractions, coupled with palaeocurrent data (Vinnels *et al*. 2010) implies that there was significant bypass through the Eastern Champsaur Basin. The Annot sand fairway therefore continued north from Champsaur, with no ponding behind the ancestral Ecrins massif (cf. Joseph & Lomas 2004; Fig. 9); turbidity currents are inferred here to have transited the region and fed down-system depocentres (Fig. 9). Certainly, the area of the Ecrins massif did not form a bathymetric high during deposition of the Annot system (cf. Ford *et al*., 1999) – it must have been deeper than the upstream Annot fairway. Future studies should address the sedimentology of the likely correlatives of the Annot turbidites further down-system, in the strongly deformed sections along the eastern part of the Ecrins and north in the Aiguilles d'Arves (Fig. 1).

Eastern confinement of the Annot system

For Joseph & Lomas (2004, their fig.5) and references therein) the eastern edge of the Annot sand system was confined by the Embrunais-Ubaye thrust sheets, an exotic tectonic allochthon emplaced from the internal Alpine domain onto the foreland domain (see also Ford *et al.* 2006). The

476 sheets were sufficiently thick to have buried the southern Ecrins district so
477 that it reached temperatures of c 300°C (Bellanger *et al.* 2015). Presumably
478 they created significant relief within the fordeep. Olistostromes were shed
479 from this relief during thrust sheet emplacement, manifest as the Schistes à
480 Blocs (e.g. Kerckhove 1969) which cap the turbidites in the Eastern
481 Champsaur Basin and elsewhere in the Annot system. Ford & Lickorish
482 (2004) suggest that it was the emplacement of the Embrunais-Ubaye thrust
483 sheets that closed off the Annot basin system. The implication is that this was
484 a progressive process, with the advance of thrust sheets gradually restricting
485 the pathways available for turbidity currents. However, the relationship
486 between the olistostomes and the Champsaur Sandstone is not consistent
487 with this implication. Muddy debris flows, such as carried the Schistes à
488 Blocs, would be expected to have had significant run-out distances, potentially
489 forming obstructions within the main flow paths for the Annot turbidity
490 currents. Deposits accumulating from axial flows in the case of the turbidity
491 currents, and from coeval basin flank-derived olistostromes, should be
492 interleaved. No such interleaving has been recognised by us in the Eastern
493 Champsaur Basin, nor elsewhere in the Annot System. Therefore, we deduce
494 that the olistostromes and Annot turbidites are not coeval – rather that the
495 Schistes à Blocs forms a distinct, younger depositional unit that overran (and
496 entrained) dark-shales. The corollary is that the flux of Annot turbidity currents
497 had terminated for reasons unrelated to the emplacement of the Embrunais-
498 Ubaye thrust sheets and that some other feature must have provided the
499 lateral (eastern) confinement to Annot flows.
500 If lateral confinement was not provided by the tectonic allochthon of the
501 Embrunais-Ubaye thrust sheets, then presumably it was provided by weakly
502 inverted parts of the ancestral rifted margin of Europe adjacent to the suture
503 of the closed Ligurian Tethys seaway. These may have formed additional
504 basement ridges, equivalent to the Ecrins but now buried beneath Alpine
505 thrust sheets. Alternatively, the Annot system may have abutted against
506 deformed Brianconnais and Sub-Brianconnais units, as illustrated on Fig. 9.
507 Both units contain sedimentary successions that are broadly time-equivalent
508 to the Annot turbidites – the so-called “Flysch Noir”. This is a succession of
509 pelites and thin, fine sandstones of mid-late Eocene age (e.g. Debelmas

1989). We tentatively suggest that the Flysch Noir represents an early part of the Annot turbidite system and that its pathway was elevated as its substrate of Brianconnais and Sub-Brianconnais units became deformed. Note that, by the early Oligocene, the eastern edge of the Brianconnais in this sector of the Alps had been incorporated into the Alpine orogen and was experiencing blue schist metamorphism, and maybe even greater burial (e.g. Michard *et al.* 2004; Dumont *et al.* 2012). Regardless of the nature of the eastern retaining margin to the Annot System, there is no documented evidence that it provided detritus into the sand fairway, until the emplacement olistostromes of the Schists à Blocs from the advancing Embrunais-Ubaye thrust sheets.

Structuring of the Eastern Champsaur Basin and the evolution of crustal shortening

The onlap of Champsaur Sandstone onto a tilted succession of Blue Marls and Nummulitic Limestone indicate that the floor to the Eastern Champsaur basin was actively deforming during deposition (Vinnels *et al.* 2010). Our studies confirm our earlier work and extend it – stratigraphic thickness variations across the Eastern Champsaur Basin (Fig. 6) strongly suggest that open folding continued to influence sand accumulation. However, sedimentation generally swamped the folds, with turbidites overstepping the anticline crests. These folds are cored by crystalline basement which outcrops in the Dourmillouse and Fournel valleys (Fig. 4) and it is this deep-rooting deformation style that is illustrated on Fig. 9. Basement-rooting deformation is also shown to provide the structural barrier that separated the distinct turbidite systems that characterise the Eastern and Western Champsaur Basins. Folds and thrusts in the adjacent crystalline basement now forming the Ecrins massif (located on Fig. 1a) are generally interpreted as resulting from tectonic inversion of half-graben inherited from Jurassic-aged continental rifting (e.g. Gillcrist *et al.* 1987; Dumont *et al.* 2008).

In Provencal basins (Fig. 9), active deformation during deposition of the Annot turbidites has been characterised as “wedge-top” (Salles *et al.* 2012 and references therein). The term harks back to idealised conceptualisations of orogenic margins that are divided into a foreland basin and an adjacent thin-skinned thrust system (e.g. De Celles & Giles 1996). In these models,

544 deposition can occur in the yet-to-deform foreland basin and in small basins
545 perched on the advancing thrust system (or wedge). However, such
546 distinctions appear inappropriate to us for the Champsaur district as
547 deformation of the basin floor during the turbidite deposition was thick-
548 skinned. Concepts of thrust wedges are unlikely to be directly relevant to
549 inversion tectonic systems such as the Ecrins or the initial deformation of the
550 Brianconnais as illustrated on Fig. 9.
551 In Provence, two distinct phases of folding are recognised – one
552 inferred to result from north-south contraction and traditionally related to
553 “Pyrenean-Provencal” orogeny (early Tertiary), and a subsequent SW-
554 directed thrust system (e.g. Siddans 1979; de Graciansky *et al.* 2011).
555 Likewise, deformation in the crystalline basement and the structural evolution
556 of the Ecrins basement massif has long been considered result from
557 punctuated tectonic activity - i.e., two distinct episodes of Alpine deformation,
558 separated by the unconformity that underlies the transgressive Nummulitic
559 Limestone (e.g. Gidon 1979). Ford (1996) correlates these with the two
560 tectonic episodes of Provence. Gupta & Allen (2000) argue that the southern
561 Ecrins was structured by folds and uplifted fault blocks to modulate the
562 Nummulitic transgression. It is a deduction that is consistent with the
563 punctuated model for tectonics in this part of the Alps. However, it seems
564 unlikely that initiation of a second episode of tectonic activity simply coincided
565 with the transgression of the Nummulitic megasequence across the
566 basement.
567 Recent radiometric dating of thrust zones in the Ecrins massif
568 (Bellanger *et al.* 2015) implies that deformation in the basement straddled
569 deposition of the post-Nummulitic successions of the Eastern Champsaur
570 Basin. Using these data, Butler (2017) proposes that crustal shortening in the
571 Ecrins and, by inference, beneath the Eastern Champsaur Basin was
572 continuous in time. Where the instantaneous syn-orogenic surface is sub-
573 areal, deformation is accompanied by erosion, marked by exhumation and
574 cooling of the basement. Where the instantaneous syn-orogenic surface
575 moves to below sea-level, deformation is accompanied by deposition. The
576 transition between these states is marked by transgression and its
577 diachroneity reflected by the sub-Nummulitic unconformity.

Dividing deformation episodes on the basis of an unconformity has a long tradition in orogenic geology (see discussion in Gray *et al.* 1997). However, is the interplay between uplift due to crustal shortening, with concomitant erosion, and subsidence due to long-wavelength orogenic loading of the lithosphere, with concomitant deposition, that defines the stratigraphic record of orogenesis. Focussing on unconformities, with their implications for missing time-sections, can create the false impression that deformation was punctuated. It is our proposal that the crust beneath the Champsaur district experienced deformation that was essentially continuous over geological time, through the late Eocene and into the Oligocene.

The arrival of the Embrunais-Ubaye Thrust Sheets in the late Rupelian (Dumont *et al.* 2008) marked a transition in the tectonic style to thin-skinned shearing. Deformation of the turbidites of the Eastern Champsaur Basin, dominantly by folding, happened beneath this over-riding allochthon. Shortening within the basin was approximately 4 km with differential shortening through the Champsaur Sandstone stratigraphy implying significant simple shear distributed penetratively through the multilayer (Bürgisser & Ford 1998). It is likely that thick-skinned deformation in the underlying crust continued, eventually arching the Embrunais-Ubaye thrust sheets and uplifting the adjacent Ecrins massif. The challenge facing further investigations of syn-tectonic sedimentation lies in unravelling deformed sedimentary basins to reveal their stratigraphy. These are endeavours that we hope will be encouraged by our study of the eastern Champsaur Basin.

Conclusions

Linked stratigraphic, sedimentological and structural studies in the Eastern Champsaur Basin, SE France, reveal:

1 – The Champsaur Sandstones of the basin represent a continuation of the late Eocene-early Oligocene Annot system. Therefore, the well-known type locations in Provence represent only a small part of this system and their causative turbidity currents continued up to, and presumably over and beyond what is now the Ecrins basement massif. There is no evidence of flow ponding in the Eastern Champsaur Basin, indeed bed character suggests

substantial sediment bypass continuing into more distal settings. These locations await further study.

2 – The continuity of sand fairways around the orogen can only arise where foredeep axial slopes and hence palaeobathymetry dipped monotonically in the direction of sediment dispersal. Therefore, during the early Oligocene, the Alpine foredeep became progressively deeper clockwise around the western Alpine arc in SE France. This implies greater tectonic subsidence, perhaps coupled with the building of depositional gradients in this direction, during the late Palaeogene. The proto-Ecrins basement massif did not lie at shallow bathymetric levels as suggested in some Alpine syntheses but, during the early Oligocene, lay under water depths significantly deeper than the Provencal sector of the foredeep.

3 – The floor to the foredeep basin within which the Annot turbidites were deposited was actively deforming. This has been inferred elsewhere for the type localities of the Annot system to the south of Champsaur where the foredeep megasequence rests on a gently folded substrate of Mesozoic strata. For the Eastern Champsaur, Mesozoic strata have been largely eroded so that the foredeep megasequence unconformably overlies basement. Small pockets of Triassic and Lower Jurassic strata are preserved, folded and thrust into basement beneath this unconformity, testifying to significant crustal deformation predating the deposition of the Nummulitic Limestone. We interpret the pre-Nummulitic deformation to form simply an early part of crustal shortening that progressed steadily through the late Eocene and early Oligocene.

4 – The Champsaur Sandstone succession is capped by the Schistes à Blocs, an olistostrome encased in dark shales, a harbinger of the advancing Embrunais-Ubaye thrust sheets. Notwithstanding the likely run-out distance of the causative debris flows ahead of the advancing thrust sheet, this olistostromal formation is not interbedded with the Champsaur Sandstone. Presumably these flows arrived on a basin floor that had largely been starved of sand supply and were not themselves the cause of this starvation.

5 - Deformation continued after the deposition of the sedimentary fill to the Eastern Champsaur Basin, with c. 4 km of shortening accommodated in the

645 footwall to the over-riding the tectonic allochthon represented by the
646 Embrunais-Ubaye thrust sheets.
647
648 Our study demonstrates that even in deformed parts of the Alpine orogen it is
649 still possible to establish detailed stratigraphies in turbidites, thereby opening
650 opportunities to extend tectono-stratigraphic investigations elsewhere in this
651 and other orogens. Such work may improve understanding not only of the
652 turbidite systems and their use as analogues for modern deep-water systems
653 but also of the tectonic relationships between orogens and their “forelands”.
654
655
656 Funding: This study was sponsored as part of the Turbidites Research Group
657 (2007-2010), funded by Anadarko, BG, BP, ConocoPhillips, Devon Energy,
658 Marathon, Maersk, Nexen, Petronas, StatoilHydro and Woodside. JV was
659 supported by BHP-Billiton.
660
661 Acknowledgements
662 We thank Mary Ford, Nigel Woodcock and an anonymous referee for
663 providing insightful comments on drafts of this paper. RWHB thanks Thierry
664 Dumont for discussions on Alpine geology and timings – although of course
665 the authors take sole responsibility for the information and interpretations
666 presented here.
667
668 **References**
669
670 Bellanger, M., Augier, R., Bellahsen, N., Jolivet, L., Monié, P., Baudin, T. &
671 Beyssax, O. 2015. Shortening of the European Dauphinois margin
672 (Oisans Massif, Western Alps): New insights from RSCM maximum
673 temperature estimates and $^{40}Ar/^{39}Ar$ in situ dating. *Journal of*
674 *Geodynamics*, **83**, 37-64.
675 Bouma, A. H. 1959. Some data on turbidites from the Alpes Maritimes
676 (France). *Geologie en Mijnbouw*, **21**, 223–227.
677 Brunt, R.L., McCaffrey, W.D. & Butler, R.W.H. 2007. Setting and architectural
678 elements of the Champsaur Sandstones, France. In: Nilsen, T.H., Shew,

679 R.D., Steffens, G.S. & Studlick, J.R.J. (eds), *Atlas of deep-water outcrops*
680 (American Association of Petroleum Geologists, Studies in Geology, **56**,
681 188-191
682 Bürgisser, J. & Ford, M. 1998. Overthrust shear deformation of a foreland
683 basin: structural studies south-east of the Pelvoux massif, SE France.
684 *Journal of Structural Geology,* **20**, 1455-1475.
685 Butler, R.W.H. 1992. Evolution of Alpine fold-thrust complexes: a linked
686 kinematic approach. In: Mitra, S. & Fisher, G. (eds) *Structural geology of fold*
687 *and thrust belts*. Johns Hopkins University Press, Baltimore, 29-44.
688 Butler, R.W.H. 2017. Basement-cover tectonics, structural inheritance and
689 deformation migration in the outer parts of orogenic belts: a view from the
690 western Alps. In: Law, R.D., Thigpen, R., Merschat, A., Stowell, H. &
691 Bailey, C. (eds) *Linkages and Feedbacks in Orogenic Process: A volume*
692 *in Honor of Robert D Hatcher Jr.* Memoir of the Geological Society of
693 America, **213**, 55-74.
694 Butler, R.W.H. & McCaffrey, W.D. 2004. Nature of thrust zones in deep water
695 sand-shale sequences: outcrop examples from the Champsaur sandstones
696 of SE France. *Marine and Petroleum Geology,* **21**, 911-921.
697 Cotton, J.T. & Koyi, H.A. 2000. Modeling of thrust fronts above ductile and
698 frictional detachments: Application to structures in the Salt Range and Potwar
699 Plateau, Pakistan. *Geological Society of America Bulletin*, **112**, 351-363.
700 Cunha, R.S., Tinterri, R. & Magalhaes, P.M. 2017. Annot Sandstone in the Peïra
701 Cava basin: An example of an asymmetric facies distribution in a confined
702 turbidite system (SE France). *Marine and Petroleum Geology*, **87**, 60-79.
703 Dahlstrom, C.D.A. 1969. Balanced cross-sections. *Canadian Journal of Earth*
704 *Sciences*, **6**, 743-757.
705 De Celles, P.G. & Giles, K.A. 1996. Foreland basin systems. *Basin Research*, **8**,
706 105-123.
707 Debelmas, J. 1989. On some key features of the evolution of the western Alps.
708 In: Sengor, A.M.C. (ed). *Tectonic Evolution of the Tethyan Region*. Springer,
709 Dordrecht. 23-42.
710 Debelmas, J., Durozoy, G., Kerckhove, C. Monjuvent, G., Mouterde, R. &
711 Pêcher, A. 1980. *Carte géologique de la France à 1:50,000: feuille*
712 *Orcières*. Bureau de Recherches Géologiques et Minières, Orleans.

713 Dumont, T., Champagnac, J-D., Crouzet, C. & Rochat, P. 2008. Multistage
714 shortening in the Dauphiné zone (French Alps): the record of Alpine
715 collision and implications for pre-Alpine restoration. *Swiss Journal of*
716 *Geosciences*, **101**, S89-S110.
717 Dumont, T., Schwartz, S., Guillot, S., Simon-Labric, T., Tricart, P. & Jourdan,
718 S. 2012. Structural and sedimentary records of the Oligocene revolution in
719 the Western Alpine arc. *Journal of Geodynamics*, **56-57**, 18-38.
720 Etienne, S., Mulder, T., Bez, M., Desaubliaux, G., Kwasniewski, A., Parize,
721 O., Dujoncquoy, E. & Salles, T. 2012. Multiple scale characterization of
722 sand-rich distal lobe deposit variability: examples from the Annot
723 Sandstones Formation, Eocene–Oligocene, SE France. *Sedimentary*
724 *Geology*, **273**, 1-18.
725 Ford, M. 1996. Kinematics and geometry of early Alpine, basement-involved
726 folds, SW Pelvoux Massif, SE France. *Eclogae Geologicae Helvetica*, **89**,
727 269-295.
728 Ford, M. & Lickorish, W.H. 2004. Foreland basin evolution around the western
729 Alpine Arc. In: Joseph, P. & Lomas, S. (eds) *Deep-water sedimentation in*
730 *the Alpine basin of SE France.* Special Publications of the Geological
731 Society, London, **221**, 39-63.
732 Ford, M., Lickorish, W.H. & Kusznir, N.J. 1999. Tertiary foreland sedimentation
733 in the Southern Subalpine Chains, SE France: a geodynamic appraisal.
734 *Basin Research*, **11**, 315-336.
735 Ford, M., Duchêne, S., Gasquet, D. & Vanderhaeghe, O. 2006. Two-phase
736 orogenic convergence in the external and internal SW Alps. *Journal of the*
737 *Geological Society*, **163**, 815-826.
738 Gidon, M. 1979. Le rôle des étapes successives de déformation dans la
739 tectonique alpine du massif du Pelvoux (Alpes occidentales). *Comptes*
740 *rendus de l'Académie des Sciences, Paris*, **288**, 803-806.
741 Gillcrist, R., Coward, M.P., & Mugnier, J-L. 1987. Structural inversion: examples
742 from the Alpine foreland and French Alps. *Geodinamica Acta*, **1**, 5-34.
743 de Graciansky, P-C., Roberts, D.G., and Tricart, P., 2011, *The Western Alps,*
744 *from rift to passive margin to orogenic belt*. Developments in Earth Surface
745 Processes, Elsevier, **14**, pp. 1-398.

746 Gray, D.R., Foster, D.A. & Bucher, M. 1997. Recognition and definition of
747 orogenic events in the Lachlan Fold Belt. *Australian Journal of Earth*
748 *Sciences*, **44**, 489-501.
749 Gupta, S. 1997. Tectonic control on paleovalley incision at the distal margin of
750 the early Tertiary Alpine foreland basin, southeastern France. *Journal of*
751 *Sedimentary Research*, **67***,* 1030-1043.
752 Gupta, S. & Allen, P.A. 2000. Implications of foreland paleotopography for
753 stratigraphic development in the Eocene distal Alpine foreland basin.
754 *Geological Society of America Bulletin*, **112**, 515-530.
755 Joseph, P. & Lomas, S.A. 2004. Deep-water sedimentation in the Alpine
756 foreland basin of SE France: New perspectives on the Grès d'Annot and
757 related systems — An introduction. In: Joseph, P. & Lomas, S. (eds).
758 *Deep-water sedimentation in the Alpine basin of SE France.* Special
759 Publications of the Geological Society, London, **221**, 1-16.
760 Kerckhove, C. 1969. La 'zone du flysch' dans les nappes de l'Embrunais-
761 Ubaye (Alpes Occidentales). *Géologie Alpine*, *Grenoble*, **45**, 5-204.
762 Kerckhove, C., Cochonat, P. & Debelmas, J. 1978. Tectonique du
763 soubassement parautochtone des nappes de l'Embrunais-Ubaye sur leur
764 bordure occidentale, du Drac au Verdon. *Géologie Alpine, Grenoble*, **54**,
765 67-82.
766 Michard, A., Avigad, D., Goffé, B. and Chopin, C., 2004. The high-pressure
767 metamorphic front of the south Western Alps (Ubaye-Maira transect,
768 France,Italy). *Schweizerische Mineralogische und Petrographische*
769 *Mitteilungen, **84***, 215-235.
770 Mulder, T., Callec, Y., Parize, O., Joseph, P., Schneider, J.L., Robin, C.,
771 Dujoncquoy, E., Salles, T., Allard, J., Bonnel, C. & Ducassou, E. 2010.
772 High-resolution analysis of submarine lobes deposits: Seismic-scale
773 outcrops of the Lauzanier area (SE Alps, France). *Sedimentary Geology*,
774 **229**, 160-191.
775 Pffifner, O.A. 1985. Displacements along thrust faults. *Eclogae Geologicae*
776 *Helvetica*, **78**, 313-333.
777 Price, N.J. & Cosgrove, J.W. 1990. *Analysis of Geological Structures*.
778 Cambridge University Press, Cambridge.

779 Puigdefabregas, C., Gjelberg, J. & Vaksdal, M. 2004. The Gres d'Annot in the
780 Annot syncline: outer basin-margin onlap and associated soft-sediment
781 deformation. In: Joseph, P. and Lomas, S. A. (eds). *Deep-Water*
782 *Sedimentation in the Alpine Basin of SE France: New Perspectives on the*
783 *Gres D'Annot and Related Systems*. Geological Society of London Special
784 Publication **221**, 367-388.
785 Ramsay, J.G. 1974. Development of chevron folds. *Geological Society of*
786 *America Bulletin*, **85**, 1741-1754.
787 Salles, L., Ford, M. & Joseph, P. 2014. Characteristics of axially-sourced
788 turbidite sedimentation on an active wedge-top basin (Annot Sandstone, SE
789 France): *Marine and Petroleum Geol*ogy, **56**, 305-323.
790 Schmid, S.M., Fügenschuh, B., Kissling, E. & Schuster, R. 2004. Tectonic map
791 and overall architecture of the Alpine orogeny. *Eclogae geologicae*
792 *Helvetiae*, **97**, 93-117.
793 Siddans, A.W.B. 1979. Arcuate fold and thrust patterns in the Subalpine chains
794 of Southeast France. *Journal of Structural Geology*, **1**, 117-126.
795 Sinclair, H.D. 1997. Tectonostratigraphic model for underfilled peripheral
796 foreland basins: An Alpine perspective: *Geological Society of America*
797 *Bulletin*, **109**, 324-346.
798 Smith, R. 2004. Silled sub-basins to connected tortuous corridors: sediment
799 distribution systems on topographically complex sub-aqueous slopes. In:
800 Lomas, S.A. & Joseph, P. (eds) *Confined turbidite systems*. Special
801 Publications of the Geological Society, London, **222**, 23-43.
802 Stanbrook, D.A. and Clark, J.D., 2004. The Marnes Brunes Inférieures in the
803 Grand Coyer remnant: characteristics, structure and relationship to the Grès
804 d'Annot. . In: Joseph, P. & Lomas, S. (eds). *Deep-water sedimentation in*
805 *the Alpine basin of SE France.* Special Publications of the Geological
806 Society, London, **221**, 285-300.
807 Stanley, D. J. & Bouma, A. H. 1964. Methodology and paleogeographic
808 interpretation of Flysch formations: a summary of studies in the Maritime
809 Alps. In: A. H. Bouma and A. Brouwer (eds.), *Turbidites*. Developments in
810 Sedimentology **3**. Elsevier, Amsterdam, 34–64.

811 Tricart, P. 1981. Les marqueurs des mouvements verticaux du socle aux
812 abords du massif du Pelvoux (Alpes occidentales). *Revue de géologie*
813 *dynamique et de géographie physique Paris*, **23**, 289-300.
814 Vinnels, J.S., Butler, R.W.H., McCaffrey, W.D. & Lickorish, W.H. 2010.
815 Sediment distribution and architecture around a bathymetrically complex
816 basin: Eastern Champsaur Basin, SE France. *Journal of Sedimentary*
817 *Research*, **80**, 216-235.
818
819

820 **Figures**

821

822 Fig. 1. a) Location map of the Eastern Champsaur Basin (boxed area is Fig 4)
823 in the context of the western Alps (modified after Schmid *et al*. 2004). b) the
824 continuity of the Annot sand system in SE France, in the context of other
825 Eocene-Oligocene turbidite units in the external Western Alps (modified after
826 Joseph & Lomas 2004). The grey area represents thrust sheets, including the
827 Embrunais-Ubaye and lateral equivalents that have over-ridden the Annot
828 sand system, which are shown as inferred at depth. SF – Selle Fault.

829

830 Fig. 2. Idealised Nummulitic stratigraphy (modified after Joseph & Lomas
831 2004)

832

833 Fig. 3. Fold geometries, all of which face west, in various parts of the
834 Champsaur Sandstone (see Fig. 4 for place names). a) looking NNW onto Pic
835 Felix Neff, to illustrate the broadly harmonic and relatively long-wavelength
836 folding of the upper stratigraphic packages (unit A6 on Figs 4 and 6). b)
837 looking SE onto polyharmonic folding in sandstone units (unit A3 on Figs 4
838 and 6) directly above the regional detachment zone of the Blue Marls. The
839 visible cliff height is 100m. c) looking SSW onto chevron-style folding in the
840 lower units of the thinly bedded Champsaur sandstone (Unit A1 on Figs 4 and
841 6). The bottom of the image is at an altitude of c 2200m, the skyline reaches
842 up to 3085m.

843

844 Fig. 4. Geological map of the Eastern Champsaur Basin showing the
845 distribution and structure of the mappable units within the Champsaur
846 Sandstone as defined by this study (Fig. 4). The location of lines of section in
847 Fig 5a, b are annotated (X-X' and Y-Y' respectively), along with logged
848 stratigraphic sections (circled numbers, 1-9; Fig. 6) and field photographs of
849 Fig. 5.

850

851 Fig. 5. Selected key panoramas that serve to trace the bed-set stratigraphy of
852 the Champsaur Sandstone, with varying accessibility. The viewpoints are
853 shown on Fig. 4. Stratigraphic boundaries are shown as yellow lines while

fold axial traces are shown by red dashed lines. a) illustrates the west face of Le Piquet, a strike-section, together with the location and content of two sedimentary logs (1 and 2 on Fig. 4) that characterise the upper part of the Champsaur Sandstone. b) Is an oblique view of the north side of the Dourmillouse Valley, which provides a dip-section through the fold belt. This hillside is partly accessible permitting ground-truthing of units and completion of measured sections. c) illustrates part of the steep south side of the Fournel Valley. This provides exceptional exposures for the stratigraphy and folds but has limited accessibility.

Fig. 6. A selection of lithostratigraphic logs with defined mappable units (A1 – A6-A7), correlated through the basin. The locations are shown on Fig. 4. These sections were constructed with direct bed-measurement and field observations – their location being restricted by accessibility in the terrain.

Fig. 7. Cross-sections through the Eastern Champsaur Basin. The lines of section are shown in Fig. 2. a) the northern basin; b) the Dourmillouse transect.

Fig. 8. A line-length restoration of the Dourmillouse cross-section through the basin. The thicknesses of bed-sets are shown (intra-formation rulings), and retained between the final state section (a) and its restored version (b). Selected bed-set levels are illustrated (in colour) together with the top basement (in red).

Fig. 9. A schematic representation of the extent of the Annot turbidite system, connecting its well-studied proximal areas in Provence to the almost un-studied, more distal regions that include the Aiguilles d'Arves basin, via the Eastern Champsaur Basin described here. The diagram is inspired by an equivalent perspective provided by Joseph & Lomas (2004). Evidence for deformation and its timing beneath the sand fairway within the Ecrins and beneath the Aiguilles d'Arves sector is discussed by Butler (2017).

figure 1 new

Click here to access/download;figure;Fig 1 maps NEW.pdf



figure 2

Click here to access/download;figure;Fig 2 strat NEW.pdf



figure 3

Click here to access/download;figure;Fig 3 folds new.jpg

figure 4 new

Click here to access/download;figure;Fig 4 MAP NEW.pdf



figure 5

Click here to access/download;figure;Fig 5_panoramas.jpg









Click here to access/download;figure;Fig 6_logsNEW.pdf



figure 7 new

Click here to access/download;figure;Fig 7_ cross sections.pdf

WSW

Selle Valley

Narreyroux valley

ENE

X

X'

a)

A6
A2 - A5
A1
Champsaur Sandstone

Nummulitic Limestone and Blue Marls

Jurassic

basement

2km

Pic Felix Neff

Crete de Martinet

Tete de Dourmillouse

W

E

Y

Y'

b)



Click here to access/download;figure;Fig 8_restoration.pdf

figure 9 new

Click here to access/download;figure;Fig 9_pgeog.pdf