



This is a repository copy of *The hypercomplex genome of an insect reproductive parasite highlights the importance of lateral gene transfer in symbiont biology.*

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/164886/>

Version: Published Version

Article:

Frost, C.L., Siozios, S., Nadal-Jimenez, P. et al. (4 more authors) (2020) The hypercomplex genome of an insect reproductive parasite highlights the importance of lateral gene transfer in symbiont biology. *mBio*, 11 (2). e02590-19. ISSN 2150-7511

<https://doi.org/10.1128/mbio.02590-19>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:
<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>



The Hypercomplex Genome of an Insect Reproductive Parasite Highlights the Importance of Lateral Gene Transfer in Symbiont Biology

Crystal L. Frost,^a  Stefanos Siozios,^a Pol Nadal-Jimenez,^a Michael A. Brockhurst,^b Kayla C. King,^c  Alistair C. Darby,^a  Gregory D. D. Hurst^a

^aInstitute of Integrative Biology, University of Liverpool, Liverpool, United Kingdom

^bDepartment of Animal and Plant Sciences, University of Sheffield, Sheffield, United Kingdom

^cDepartment of Zoology, University of Oxford, Oxford, United Kingdom

Crystal L. Frost and Stefanos Siozios contributed equally to this project. Author order is alphabetical.

ABSTRACT Mobile elements—plasmids and phages—are important components of microbial function and evolution via traits that they encode and their capacity to shuttle genetic material between species. We here report the unusually rich array of mobile elements within the genome of *Arsenophonus nasoniae*, the son-killer symbiont of the parasitic wasp *Nasonia vitripennis*. This microbe's genome has the highest prophage complement reported to date, with over 50 genomic regions that represent either intact or degraded phage material. Moreover, the genome is predicted to include 17 extrachromosomal genetic elements, which carry many genes predicted to be important at the microbe-host interface, derived from a diverse assemblage of insect-associated gammaproteobacteria. In our system, this diversity was previously masked by repetitive mobile elements that broke the assembly derived from short reads. These findings suggest that other complex bacterial genomes will be revealed in the era of long-read sequencing.

IMPORTANCE The biology of many bacteria is critically dependent on genes carried on plasmid and phage mobile elements. These elements shuttle between microbial species, thus providing an important source of biological innovation across taxa. It has recently been recognized that mobile elements are also important in symbiotic bacteria, which form long-lasting interactions with their host. In this study, we report a bacterial symbiont genome that carries a highly complex array of these elements. *Arsenophonus nasoniae* is the son-killer microbe of the parasitic wasp *Nasonia vitripennis* and exists with the wasp throughout its life cycle. We completed its genome with the aid of recently developed long-read technology. This assembly contained over 50 chromosomal regions of phage origin and 17 extrachromosomal elements within the genome, encoding many important traits at the host-microbe interface. Thus, the biology of this symbiont is enabled by a complex array of mobile elements.

KEYWORDS bacteriophage evolution, endosymbionts, genomics, plasmids

Phages and plasmids play important roles in the ecology and evolution of bacteria (1). Both types of elements have the capacity to promote the lateral transfer of traits. Temperate phages commonly carry genes that encode phenotypes of benefit to bacterial survival, including key determinants of the ability to thrive in association with hosts (2–4). Plasmids are also key determinants of phenotype, most importantly acting as shuttles for antibiotic resistance (5). Plasmids also carry a range of traits important for association with hosts. For instance, both essential amino acid synthesis in the aphid

Citation Frost CL, Siozios S, Nadal-Jimenez P, Brockhurst MA, King KC, Darby AC, Hurst GDD. 2020. The hypercomplex genome of an insect reproductive parasite highlights the importance of lateral gene transfer in symbiont biology. *mBio* 11:e02590-19. <https://doi.org/10.1128/mBio.02590-19>.

Editor Nicole Dubilier, Max Planck Institute for Marine Microbiology

Copyright © 2020 Frost et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Gregory D. D. Hurst, g.hurst@liv.ac.uk.

Received 4 October 2019

Accepted 2 March 2020

Published 24 March 2020

symbiont *Buchnera* and the reproductive parasitic phenotype of male-killing *Spiroplasma* in *Drosophila melanogaster* are encoded on plasmids (6–7).

Recent advances in long-read sequencing technologies (read lengths of >20 kb) allow closure of prophage-rich bacterial genomes whose repetitive nature prevented them from being completed with previous technologies. Our own attempts to complete the genome of *Arsenophonus nasoniae*, the male-killing endosymbiont of *Nasonia* and other chalcid wasps (8), represents an instructive case study. This wasp species parasitizes filth fly (calliphorid) pupae. *Arsenophonus nasoniae* passes from a female into the fly pupa on wasp oviposition, from where it infects larvae through feeding, ultimately establishing as an extracellular infection in the wasp ovipositor (9). The microbe kills male hosts as embryos and may also exhibit pathological relationship in diapausing (overwintering) wasp larvae.

Our first attempt to sequence this genome used standard and paired-end libraries with 454 sequencing, resulting in a fragmented assembly with 665 contigs (143 scaffolds and 261 sequencing gaps) (10). We have since used Illumina, PacBio, and, lately, Oxford Nanopore reads to produce an improved reference genome assembly for *A. nasoniae* (strain FIN). Hybrid assembly using PacBio (15- to 20-kb-size-selected library, ~4-kb median read length, ~12× coverage) and Nanopore long reads (~10-kb median read length, 252-kb maximum read length, 9,631 reads of >20 kb, ~169× coverage) with subsequent Illumina polishing resulted in a closed circular genome (for details of strain isolation and sequencing methods, see <https://doi.org/10.6084/m9.figshare.11842425>).

This closed genome revealed a 3.9-Mb main chromosome with abundant phage-derived chromosomal islands (Fig. 1). PHASTER (11) estimates the presence of 27 phage-derived regions, of which 18 are classed as complete and may represent intact prophage, 3 are classed as unsure, and 6 are classed as incomplete and probable relics. The sizes of these chromosomal phage-derived regions range between 4.3 and 101.7 kbp (Fig. 1; see also <https://doi.org/10.6084/m9.figshare.11845134>). This density of phage-derived elements (6.9/Mb) is almost double that of *Lactococcus lactis* subsp. *cremoris* MG1363, the most prophage-rich genome recorded in previous systematic analyses (12). Of the 27 phage-derived elements inferred in the assembly, 26 are confirmed through individual reads that cover both flanking regions. All were confirmed through tiled long reads with unique overlapping sequences (Fig. 1).

Phage-derived elements classed as potentially complete prophage by PHASTER were examined in more detail (<https://doi.org/10.6084/m9.figshare.11845182>). Flanking attachment sites were observed in all cases. However, these putative prophage elements varied in their ranges of core phage functions identified through homology. Two of 18 putative prophage elements lacked genes predicted to function in lysis, 4 lacked predicted packaging genes, and 3 lacked structural genes. From these data, we conclude either (i) that some of these elements are not autonomous (excision would require complementation by other phage-derived elements) or (ii) that the missing components are novel and within the unannotated portion of the putative prophage element.

Syntenic mapping using Sibelia (13) shows that the phage-derived regions within the *A. nasoniae* main chromosome are not identical but do share many repetitive elements (Fig. 1). This mosaicism of repetitive sequences breaks the assembly when the read length is insufficient to fully span the repetitive region(s) in any given phage-derived region, and these elements are the primary cause of assembly breakage in previous sequencing efforts, with break points commonly occurring in phage-derived regions.

The assembly also predicted a complex complement of extrachromosomal DNA, with at least 17 extrachromosomal DNA elements (15 circular, 1 linear, 1 unclear) (Fig. 2). Patterns of coverage (<https://doi.org/10.6084/m9.figshare.11845263>), combinations of long-read pairs that overlap and form a circle, and the presence of open reading frames (ORFs) predicted to function in plasmid maintenance (retention systems and/or replication initiation proteins) support their status as plasmids (<https://doi.org/10.6084/m9.figshare.11845197> and <https://doi.org/10.6084/m9.figshare.11845305>). The




FIG 1 Prophage-rich content of *A. nasoniae* strain FIN'13's main chromosome. Inwards, the first circle represents the complete and closed chromosome generated from the hybrid assembly of the Nanopore, PacBio, and Illumina data. The different colors depict intact (green), questionable (blue), and incomplete (red) phage elements, as estimated by PHASTER, while black bars represent the seven rRNA operons. The second circle depicts the transposable elements identified in the *A. nasoniae* genome. The third circle shows uniquely mapped long Nanopore reads spanning the prophage regions. Only the top five largest reads spanning each region are shown. The fourth circle shows the corresponding assembly using only the Illumina data. The innermost circle represents patterns of GC skew. Colored ribbons in the interior connect the different syntenic blocks (>5 kb). The scale is in kilobase pairs.

predicted plasmid diversity exceeds the 11 recorded in *Marinovum algicola*, which the authors considered “unprecedented for proteobacteria” (14). Outside of the proteobacteria, a similar number of extrachromosomal elements (up to 21) was found in *Borrelia burgdorferi* (15). This parasitic spirochete, like *A. nasoniae* (7), has two different host species in its life cycle (15). There are 1,426 predicted genes on the plasmids, of which 320 code for accessory genes and 419 are annotated as hypothetical (<https://doi.org/10.6084/m9.figshare.11857836>). ORFs present are predicted to encode diverse functions, including the capacity to induce apoptosis in host cells, toxin elements and




FIG 2 Plasmid maps of *A. nasoniae* in linearized form. Predicted intact, questionable, and incomplete prophage elements are shown in green, blue, and red, respectively. Colored ribbons connect the different synteny blocks (>5 kb). The assignment O indicates that the element is predicted to be circular. The scale is in kilobase pairs.

transporters, type III secreted effectors that manipulate eukaryotic cell physiology, and proteins that allow microbes to adhere to and invade eukaryotic cells.

The extrachromosomal DNA is predicted to harbor 28 further phage-derived elements (PHASTER, 11 complete, 5 unsure, and 12 incomplete). Four of the extrachromosomal elements consisted entirely of circularized prophage DNA (<https://doi.org/10.6084/m9.figshare.11845305>) and were classified as plasmids on the basis of their Rep-encoding genes or addiction systems. Synteny mapping indicates that extrachromosomal elements share many repetitive elements with each other, and these regions are primarily of phage origin (Fig. 2). In contrast, only one 5-kb window from the

extrachromosomal genome compartment was observed to show sequence similarity to the main chromosome (with a threshold of 90% of the nucleotide sequence) (<https://doi.org/10.6084/m9.figshare.11857794>).

Arsenophonus nasoniae represents the most-prophage-rich genome documented to date. With 55 predicted phage-derived regions in total, 27% of the main chromosome and 43% of the total genome are phage derived. Phage-derived regions on the main chromosome are predicted to encode 1,490 protein sequences. Of the genes, around 250 are not associated with core phage functions, and an additional 491 genes were annotated as encoding hypothetical proteins that lack domains recognized by Pfam. ORFs present are predicted to encode diverse functions, including the capacity to induce apoptosis in host cells, toxin elements and transporters, type III secreted effectors that manipulate eukaryotic cell physiology, and proteins that allow microbes to adhere to and invade eukaryotic cells (<https://doi.org/10.6084/m9.figshare.11857836>). However, they do not contain ORFs of obvious eukaryotic origin, as typified in the eukaryote association modules of *Wolbachia* phage (16).

The ORFs present in phage-derived regions highlight the potentially rich capacity for phages to drive lateral transfer of important genes between bacterial lineages. Previous work indicated shared phage-derived elements between *Arsenophonus* spp. and the aphid symbiont *Hamiltonella defensa* (17). We examined likely sources of the phage-derived elements of *Arsenophonus nasoniae* using Krona charts, examining the closest allied matches to ORFs within phage-derived regions (<https://doi.org/10.6084/m9.figshare.11857860>). Of the genes in *A. nasoniae* phage-derived regions, the best matches were in the genus *Arsenophonus*, other closely related genera, and other insect-associated gammaproteobacteria. In contrast, with regard to the genes not present within the predicted phage-derived regions, the best matches were most commonly to genes in the *Arsenophonus* symbiont from *Nilaparvata lugens*.

We complemented this approach with phylogenetic analysis of three prophage-encoded proteins: portal protein, lysozyme, and capsid protein (<https://doi.org/10.6084/m9.figshare.11857869>). Predicted capsid proteins and predicted phage portal proteins from *A. nasoniae* prophage-derived elements were most closely related to the respective gene in prophage elements from *Xenorhabdus*, *Proteus*, *Providencia*, and *Morganella*, genera closely related to *Arsenophonus*. In contrast, predicted lysozyme proteins from *A. nasoniae* prophage-derived regions had a distinct pattern, with most closely related genes being from other *Arsenophonus* strains, and this cluster then allied to ones derived from prophages in the aphid symbiont *Regiella insecticola*.

Overall, there is no consistent pattern supporting direct lateral transfer of an intact phage element from any currently sampled taxon. Rather, the elements observed are chimeric, predominantly with nearest matches from closely allied genera, but with some components deriving from more distantly related insect-associated gammaproteobacteria. Past work examining phage transfer in *Wolbachia* has emphasized the importance of coinfection as an arena for transfer (18). As a gammaproteobacterium that exists both as an endosymbiont and external to hosts, *Arsenophonus* likely encounters a broad range of microbes, and this establishes opportunities for lateral transfer. *Arsenophonus nasoniae*, for instance, exists alongside both *Proteus* and *Providencia* within the *Nasonia* gut (19).

Phage transfers have occurred despite the presence of a type I CRISPR-Cas system in the *A. nasoniae* genome. The main chromosome of *A. nasoniae* carries a CRISPR-Cas system most similar in organization to that found in *Yersinia pseudotuberculosis* (type I-F) (<https://doi.org/10.6084/m9.figshare.11857887>). A notable difference is that in *A. nasoniae*, the *cas2-cas3* fusion gene, a typical characteristic of the type I-F CRISPR-Cas system (20), is interrupted by a degenerated insertion element of the IS91 family and by a small gene predicted to encode a hypothetical protein. Diverse spacer elements are found adjacent to the CRISPR-Cas cluster. Eight of the 11 spacers show homology to either prophage or plasmid sequences found in the *A. nasoniae* genome; the other three are of unknown origin and potentially evidence elements that were historically present. It is possible that the CRISPR system suppresses prophage reactivation in *A.*

nasoniae, as observed in other systems (21). It is currently unclear if the intact prophage elements retain the capacity to establish a lytic cycle.

This genome was challenging to assemble because of its structural complexity. It is notable that many of the reported closed bacterial chromosomes with high predicted prophage content were obtained using methods developed prior to next-generation sequencing (NGS), in which cosmid- or fosmid-based scaffolding processes are employed (<https://doi.org/10.6084/m9.figshare.11857893>). These long-range scaffolding tools make assemblies relatively robust for long repetitive elements, such as prophages. Conversely, closed bacterial chromosomes completed using short-read sequencing methods tend not to contain large numbers of phage-derived elements. Short-read sequencing methods may have thus led to an underrepresentation of closed bacterial genomes with complex architectures. Because the causes of assembly failure (repetitive elements like phages [22]) are biologically important, these broken assemblies obscure important properties. A full understanding of the role of prophages in the genome evolution of bacteria will require more widespread application of long-read sequencing. Without this approach, we will continue to underestimate the complexity and diversity of host-associated microbes.

Data availability. Raw sequences have been submitted to the NCBI SRA under accession numbers [SRR8797496](https://www.ncbi.nlm.nih.gov/sra/SRR8797496), [SRR8797497](https://www.ncbi.nlm.nih.gov/sra/SRR8797497), and [SRR8797498](https://www.ncbi.nlm.nih.gov/sra/SRR8797498) for PacBio, Illumina, and Nanopore data, respectively. Data can be found under BioProject number [PRJNA529362](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA529362). Supplementary material may be downloaded from <https://doi.org/10.6084/m9.figshare.c.4853367.v1>.

ACKNOWLEDGMENTS

We thank Esa Lehikoinen, for collecting and providing spent birds' nests, from which *Nasonia* and the *A. nasoniae* isolates were obtained. We also thank Steve Paterson for commenting on the manuscript and Seth Bordenstein and an anonymous reviewer for constructive critiques.

Genome sequencing was provided in part by MicrobesNG (<http://www.microbesng.uk>), which is supported by the BBSRC (grant BB/L024209/1). This project has received funding from the European Union's Horizon 2020 research and innovation program under Marie Skłodowska-Curie grant agreements 704382 and 708232 (to C.L.F. and P.N.-J., respectively), as well as funding from the NERC (grant NE/101067X/1 to G.D.D.H., M.A.B., and K.C.K.).

This project was conceived by G.D.D.H., M.A.B., A.C.D., and K.C.K. Microbial isolation, culture, DNA isolation, and sequencing were performed by C.L.F., S.S., P.N.-J., and K.C.K. Genome assembly and analysis were performed by C.L.F., S.S., A.C.D., and G.D.D.H. The paper was written by C.L.F., S.S., and G.D.D.H., with comments from P.N.-J., M.A.B., K.C.K., and A.C.D. All authors approved the final draft of the manuscript and can confirm that there are no potential conflicts of interest arising from this paper.

REFERENCES

- Hall JPJ, Brockhurst MA, Harrison E. 2017. Sampling the mobile gene pool: innovation via horizontal gene transfer in bacteria. *Philos Trans R Soc Lond B Biol Sci* 372:20160424. <https://doi.org/10.1098/rstb.2016.0424>.
- Davies EV, Winstanley C, Fothergill JL, James CE. 2016. The role of temperate bacteriophages in bacterial infection. *FEMS Microbiol Lett* 363:fnw015. <https://doi.org/10.1093/femsle/fnw015>.
- LePage DP, Metcalf JA, Bordenstein SR, On J, Perlmutter JI, Shropshire JD, Layton EM, Funkhouser-Jones LJ, Beckmann JF, Bordenstein SR. 2017. Prophage WO genes recapitulate and enhance *Wolbachia*-induced cytoplasmic incompatibility. *Nature* 543:243–247. <https://doi.org/10.1038/nature21391>.
- Weldon SR, Strand MR, Oliver KM. 2013. Phage loss and the breakdown of a defensive symbiosis in aphids. *Proc Biol Sci* 280:20122103. <https://doi.org/10.1098/rspb.2012.2103>.
- San Millan A. 2018. Evolution of plasmid-mediated antibiotic resistance in the clinical context. *Trends Microbiol* 26:978–985. <https://doi.org/10.1016/j.tim.2018.06.007>.
- Harumoto T, Lemaitre B. 2018. Male-killing toxin in a bacterial symbiont of *Drosophila*. *Nature* 557:252–255. <https://doi.org/10.1038/s41586-018-0086-2>.
- Wernegreen JJ, Moran NA. 2001. Vertical transmission of biosynthetic plasmids in aphid endosymbionts (*Buchnera*). *J Bacteriol* 183:785–790. <https://doi.org/10.1128/JB.183.2.785-790.2001>.
- Gherna RL, Werren JH, Weisburg W, Cote R, Woese CR, Mandelco L, Brenner DJ. 1991. *Arsenophonus nasoniae* gen. nov., sp. nov., the causative agent of the son-killer trait in the parasitic wasp *Nasonia vitripennis*. *Int J Syst Evol Microbiol* 41:563–565. <https://doi.org/10.1099/00207173-41-4-563>.
- Nadal-Jimenez P, Griffin JS, Davies L, Frost CL, Marcello M, Hurst G. 2019. Genetic manipulation allows in vivo tracking of the life cycle of the son-killer symbiont, *Arsenophonus nasoniae*, and reveals patterns of

- host invasion, tropism and pathology. *Environ Microbiol* 21:3172–3182. <https://doi.org/10.1111/1462-2920.14724>.
10. Darby AC, Choi J-H, Wilkes T, Hughes MA, Werren JH, Hurst GDD, Colbourne JK. 2010. Characteristics of the genome of *Arsenophonus nasoniae*, son-killer bacterium of the wasp *Nasonia*. *Insect Mol Biol* 19:75–89. <https://doi.org/10.1111/j.1365-2583.2009.00950.x>.
 11. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44:W16–W21. <https://doi.org/10.1093/nar/gkw387>.
 12. Touchon M, Bernheim A, Rocha EP. 2016. Genetic and life-history traits associated with the distribution of prophages in bacteria. *ISME J* 10: 2744–2754. <https://doi.org/10.1038/ismej.2016.47>.
 13. Minkin I, Patel A, Kolmogorov M, Vyahhi N, Pham S. 2013. Sibelia: a scalable and comprehensive synteny block generation tool for closely related microbial genomes, p 215–229. In Darling A, Stoye J (ed), *Algorithms in bioinformatics*. Springer, Berlin, Germany.
 14. Frank O, Göker M, Pradella S, Petersen J. 2015. Ocean's twelve: flagellar and biofilm chromids in the multipartite genome of *Marinovum algicola* DG898 exemplify functional compartmentalization. *Environ Microbiol* 17:4019–4034. <https://doi.org/10.1111/1462-2920.12947>.
 15. Casjens SR, Mongodin EF, Qiu W-G, Luft BJ, Schutzer SE, Gilcrease EB, Huang WM, Vujanovic M, Aron JK, Vargas LC, Freeman S, Radune D, Weidman JF, Dimitrov GI, Khouri HM, Sosa JE, Halpin RA, Dunn JJ, Fraser CM. 2012. Genome stability of Lyme disease spirochetes: comparative genomics of *Borrelia burgdorferi* plasmids. *PLoS One* 7:e33280. <https://doi.org/10.1371/journal.pone.0033280>.
 16. Bordenstein SR, Bordenstein SR. 2016. Eukaryotic association module in phage WO genomes from *Wolbachia*. *Nat Commun* 7:e13155. <https://doi.org/10.1038/ncomms13155>.
 17. Duron O. 2014. *Arsenophonus* insect symbionts are commonly infected with APSE, a bacteriophage involved in protective symbiosis. *FEMS Microbiol Ecol* 90:184–194. <https://doi.org/10.1111/1574-6941.12381>.
 18. Kent BN, Salichos L, Gibbons JG, Rokas A, Newton IL, Clark ME, Bordenstein SR. 2011. Complete bacteriophage transfer in a bacterial endosymbiont (*Wolbachia*) determined by targeted genome capture. *Genome Biol Evol* 3:209–218. <https://doi.org/10.1093/gbe/evr007>.
 19. Brucker RM, Bordenstein SR. 2013. The hologenomic basis of speciation: gut bacteria cause hybrid lethality in the genus *Nasonia*. *Science* 341: 667–669. <https://doi.org/10.1126/science.1240659>.
 20. Koonin EV, Makarova KS, Zhang F. 2017. Diversity, classification and evolution of CRISPR-Cas systems. *Curr Opin Microbiol* 37:67–78. <https://doi.org/10.1016/j.mib.2017.05.008>.
 21. Edgar R, Qimron U. 2010. The *Escherichia coli* CRISPR system protects from λ lysogenization, lysogens, and prophage induction. *J Bacteriol* 192:6291–6294. <https://doi.org/10.1128/JB.00644-10>.
 22. Schmid M, Frei D, Patrignani A, Schlapbach R, Frey JE, Remus-Emsermann MNP, Ahrens CH. 2018. Pushing the limits of de novo genome assembly for complex prokaryotic genomes harboring very long, near identical repeats. *Nucleic Acids Res* 46:8953–8965. <https://doi.org/10.1093/nar/gky726>.