

This is a repository copy of *The predictors of cigarette smoking, smokeless tobacco consumption and use of both forms in adolescents in South Asia:a secondary analysis of the Global Youth Tobacco Surveys*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/166412/>

Version: Accepted Version

Article:

Mishu, Masuma Pervin orcid.org/0000-0002-6545-9117, Siddiqui, Faraz orcid.org/0000-0002-2253-3911, Shukla, Radha et al. (3 more authors) (2021) The predictors of cigarette smoking, smokeless tobacco consumption and use of both forms in adolescents in South Asia:a secondary analysis of the Global Youth Tobacco Surveys. *Nicotine and tobacco research*. pp. 956-965. ISSN 1469-994X

<https://doi.org/10.1093/ntr/ntaa202>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Supplementary Figure 1: Expected association of anti- and pro-tobacco factors with tobacco use

