

This is a repository copy of *Paleotsunami research along the Nankai Trough and Ryukyu Trench subduction zones—Current achievements and future challenges*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/166432/>

Version: Accepted Version

Article:

Fujiwara, Osamu, Goto, Kazuhisa, Ando, Ryosuke et al. (1 more author) (2020)
Paleotsunami research along the Nankai Trough and Ryukyu Trench subduction zones—
Current achievements and future challenges. *Earth-Science Reviews*. 103333. ISSN 0012-
8252

<https://doi.org/10.1016/j.earscirev.2020.103333>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

1 Paleotsunami research along the Nankai Trough and Ryukyu Trench subduction zones
2 – Current achievements and future challenges –
3
4

5 **Authors:**

6 Osamu Fujiwara^{1*}, Kazuhisa Goto², Ryosuke Ando² and Ed Garrett^{3,4}
7

8 **Affiliations:**

- 9 1. Geological Survey of Japan, National Institute of Advanced Industrial Science and
10 Technology, AIST Central 7, 1-1-1 Higashi, Tsukuba, 305-8567, Japan.
11 2. Department of Earth and Planetary Science, The University of Tokyo, 7-3-1 Hongo,
12 Bunkyo-ku, Tokyo, 113-0033, Japan.
13 3. Department of Environment and Geography, University of York, Wentworth Way,
14 Heslington, York, YO10 5NG, United Kingdom
15 4. Geological Survey of Belgium, Royal Belgian Institute of Natural Sciences, Brussels,
16 Belgium
17

18 *Corresponding Author: Osamu Fujiwara; Email: o.fujiwara@aist.go.jp.
19

20 **Keywords:**

21 tsunami deposit, paleoseismology, earthquake forecasting, Japanese islands, Maximum possible
22 tsunami
23
24

25 **1. Introduction**

26 Facing the convergent plate boundary along the northwestern Pacific margin, the Japanese
27 islands have repeatedly suffered from great subduction zone earthquakes and related tsunamis.
28 As it is a mountainous country, the large population and industrial activity are concentrated on
29 small coastal plains, making them particularly vulnerable to these natural hazards. The most
30 recent example is the 2011 Tōhoku earthquake and tsunami, which occurred along the northern
31 Japan Trench, where the Pacific Plate subducts beneath the Japanese archipelago.

32 One area highly likely to generate a future magnitude (M) 8.5+ subduction zone earthquake
33 and tsunami is the Nankai Trough-Ryukyu Trench subduction zone, where the Philippine Sea
34 Plate descends beneath the Japanese archipelago (e.g., HERP, 2013). Forecasting of the next
35 earthquake and tsunami in these regions is an urgent issue for the Japanese government's
36 national disaster prevention measures. To address this issue, many research projects have sought
37 to identify the locations, timing, and size of past great earthquakes, based on historical records
38 along with geological evidence. Documented Nankai Trough earthquakes and related tsunamis
39 date back to the 7th century CE and suggest that M8-class great earthquakes have occurred at an
40 interval of 100-200 years in this region (HERP, 2013), with 75 years having passed since the
41 last events, which occurred in 1944 and 1946 CE.

42 On the other hand, documented records of great earthquakes and tsunamis along the
43 Ryukyu Trench are limited mainly to the 17th century CE and later. It was once thought that no
44 great earthquakes occur along the Ryukyu Trench; however, results from seismology and
45 tsunami deposit studies since 2011 CE have overturned this theory and have highlighted the
46 importance of estimating the magnitude and location of possible future great earthquakes in this
47 region (Ando et al., 2009, 2012a, 2018).

48 Both the historical and geological records are inevitably biased by age and region as
49 archives for paleoseismic events. Furthermore, geological records tend to be biased towards
50 larger events. However, the geological record has the greatest advantage of all: the length of
51 time it covers. Geological and archeological studies have been conducted to supplement the
52 imperfect documented earthquake and tsunami history and to extend the record back to earlier
53 periods. The first line of these studies is "Earthquake archaeology", established in the late 1980s,
54 which seeks to identify the past earthquakes based on evidence of strong ground shaking, such
55 as liquefaction features, found in archeological sites (Sangawa, 1992). These studies suggested
56 the possibility of subduction zone earthquakes, such as the 887 and 684 CE Tōkai earthquakes
57 along the eastern Nankai trough, which had not been confirmed in the documentary record.
58 Several earlier Nankai Trough earthquakes dating back to 1800-2000 years ago have been also
59 proposed from the archaeological data (e.g., Sangawa, 2007). However, this method generally

60 does not provide information on tsunamis, which are a key characteristic of major subduction
61 zone earthquakes.

62 A second line of research, paleotsunami studies, commenced along the Ryukyu Trench in
63 the 1960s, and full-scale research along the Nankai Trough began in the 1990s. Tsunamis
64 following two M9-class earthquakes that occurred in close succession in the 21st century, the
65 2004 CE Indian Ocean tsunami and 2011 CE Tōhoku tsunami, have provided a clear incentive
66 to study the record of past tsunamis and incorporate paleotsunami data into forecasts of future
67 events. Results of paleotsunami studies along the Nankai and Sagami Troughs up to 2007 CE
68 are summarized in Komatsubara and Fujiwara (2007). Garrett et al. (2016) provides a
69 comprehensive review of the recurrence history of great Nankai Trough earthquakes in the
70 Middle to Late Holocene, including not only tsunami deposits and liquefaction features on land,
71 but also marine and lacustrine turbidites.

72 The Geological Society of Japan published a special issue summarizing the results of
73 paleotsunami research along the Japanese archipelago as one of the commemorative
74 publications celebrating its 125th anniversary (*Journal of the Geological Society of Japan* 123,
75 No. 10). Wallis et al. (2018) describes the changes in the Japanese government's policy
76 concerning tsunami countermeasures following the Tōhoku event, and the changes in the
77 paleotsunami community's research direction reflecting the new policy. The policy (CDMC,
78 2011) reflected on the overreliance on geodetic observation data (~130 year-long) and “recent
79 and reliable” historical documents (covering the last ~400 years) in the long-term earthquake
80 prediction for Japan, and stipulated that all information from various different sources including
81 geological data should be considered in the new earthquake and tsunami countermeasures (see
82 summary in Goto et al., in this issue). As a response to this situation, in Japan, many
83 paleotsunami studies have been conducted and some are in progress under the auspices of the
84 national and local governments.

85 Tsunami deposit studies along the Nankai and Ryukyu subduction zones have been
86 independently reviewed (e.g., Fujiwara and Tanigawa, 2017; Goto, 2017), but there is no
87 comprehensive review comparing the two or summarizing the latest research. In this review, we
88 intend to address this point. For example, differences in the climate and geology of the two
89 regions create differences in the occurrence and composition of typical tsunami deposits: sandy
90 tsunami deposits for the Nankai and tsunami boulders (mainly coral blocks) for the Ryukyu
91 subduction zone. Additionally, as the two regions comprise a continuous subduction zone, the
92 possibility of a giant earthquake spanning both regions must carefully be assessed (Furumoto
93 and Ando, 2009). For this reason, synthesizing the earthquake and tsunami histories of both
94 subduction zones is particularly important.

95 Earthquake and tsunami forecasts operate at various timescales ranging from earthquake
96 early warnings seconds before shaking is felt to long-term probability predictions over decades
97 where geological data is predominant (e.g., Satake and Atwater, 2007). Every case requires the
98 specification of three factors: where, when, and how big. From this perspective, this paper
99 reviews the results of research on onshore tsunami deposits, their limitations, and existing
100 knowledge gaps, and discusses measures to address the remaining issues. When and where
101 earthquakes occurred is reconstructed based mainly on tsunami deposits studies with the help of
102 other geological evidence. A comprehensive earthquake recurrence model is still incomplete for
103 these subduction zones. This paper therefore also addresses the role that tsunami deposit
104 research plays in the development of this recurrence model. The magnitude of earthquakes that
105 occur along a subduction zone varies widely (e.g., Satake and Atwater, 2007); this paper
106 discusses the variation in the size of earthquakes inferred from tsunami deposit research.

107 The possibility of the occurrence of an unforeseen or unexpected large earthquake and
108 tsunami is a major point of reflection following the 2011 Tōhoku event. Learning from this, an
109 estimation of the “maximum possible earthquake and tsunami” that could affect the Japanese
110 coast has been included in disaster prevention measures in Japan (CDMC, 2011; Cabinet Office,
111 2012a). The CDMC (2011) stated that future tsunami countermeasures will basically require the
112 assumption of two possible levels of tsunamis. Level one reflects the largest tsunamis known to
113 occur on centennial timescales and forms the basis for designing the height of seawalls and
114 other features. Level two is the maximum possible tsunami, which occurs at a much lower
115 frequency than the first level but is considered to be more destructive. The hypothetical size of a
116 level two tsunami is based on studies of tsunami deposits and crustal movements. The 2011
117 Tōhoku tsunami is an example of a level two tsunami. Countermeasures against a level two
118 tsunami could include relocating settlements to higher ground and building huge seawalls, but
119 specific measures are difficult to implement. Complicating attempts to mitigate future impacts,
120 it is unknown whether such extremely large earthquakes and tsunamis have occurred in the past
121 along the Nankai and Ryukyu subduction zones. This paper refers to the plausibility of this
122 scenario from the standpoint of the geological evidence.

123

124 **2. Tectonic setting**

125 **2.1. Plate subduction and related geomorphology**

126 The Nankai and Ryukyu subduction zones mark the northwestern margin of the Philippine
127 Sea Plate (Fig. 1), but the nature of plate subduction, geological structures, and occurrence
128 mode of earthquakes are largely different between the two. The Palau-Kyushu Ridge forms the
129 boundary between the two subduction zones. The Nankai Trough extends for 700 km from the
130 southwest of Shikoku Island in the west to Suruga Bay in the east, where it is also called the

131 Suruga Trough. The Ryukyu Islands consist of numerous small islands extending over 1000 km
132 along the Ryukyu Trench, which belong to the subtropical climate zone and hence the
133 depositional environment is very different from other parts of Japan. The convergence rate is
134 generally 40-50 mm a⁻¹ along the Nankai subduction zone and 50-63 mm a⁻¹ along the Ryukyu
135 subduction zone, respectively. The slope of the descending slab, with some local variations, is
136 generally steeper in the Ryukyu subduction zone.

137 The Amami Plateau, Daito Ridge, Oki-Daito Ridge and basins between them extend in
138 E-W or NW-SE directions on the Philippine Sea Plate. Western extensions of these topographic
139 irregularities subduct beneath the Ryukyu arc (e.g., Kato, 1993). The Ryukyu and Nankai
140 subduction zones are represented by old (40-49 Ma), cold slab that originated from the West
141 Philippine Basin and young (<20 Ma), hot slab that originated from the Shikoku Basin,
142 respectively (e.g., Ishizuka et al., 2011). A thick accretionary prism characterizes the Nankai
143 Trough, in which high-angle reverse faults (splay faults) branching off the plate boundary
144 megathrust may move together with the main thrust during a great earthquake and enhance the
145 tsunami (Baba et al., 2006; Moore et al., 2007).

146 The Nankai Trough fault region has been empirically divided into five fault planes (rupture
147 zones), which denoted as segments A to E from west to east, respectively (Fig. 1) (Ando, 1975a,
148 b; Ishibashi, 1976, 1981). This classification is mainly based on studies of the source
149 mechanisms of historical earthquakes and the forearc submarine topography, which reflects the
150 seismotectonic activities along this region. Each of the five segments has a length of 100-150
151 km along the trough axis and almost corresponds to the distribution of forearc basins (e.g.,
152 Awata and Sugiyama, 1989; Sugiyama, 1990). In general, plate boundary earthquakes
153 (M8-class) that are considered to have source areas in areas A to B and C to E are called Nankai
154 and Tōkai earthquakes, respectively.

155 Allocating the source areas of past earthquakes to any of segments A to E is based on the
156 implicit assumption that a large earthquake recurs on its own fault plane. The necessity to
157 criticize and review this assignment has recently been pointed out (e.g., Seno, 2012). The
158 Hyūga-nada area (Figs. 1, 2), where M7-class earthquakes occur separately from the Nankai and
159 Tōkai earthquakes, may be another seismic zone denoted as Z (e.g., Wells et al., 2003). This
160 empirical segmentation of rupture zones has not been applied to the Ryukyu Trench as historical
161 and seismological information is scarce.

162 Flight of marine terraces, mainly from mid-Pleistocene to Holocene in age, are distributed
163 on the peninsulas, capes and islands of the Nankai and Ryukyu subduction zones (Koike and
164 Machida, 2001), and there have been many discussions about their formation process in relation
165 to sea-level changes and crustal movements (e.g. Yoshikawa et al., 1964; Webster et al., 1998).
166 They show a wide regional variation in average uplift rates. For example, the terraces of marine

167 oxygen isotope stage (MIS) 5e are 40-50 m high in many areas, but reach nearly 190 m at Cape
168 Muroto, the boundary between segments A and B. Coseismic uplift and subsidence during
169 historical earthquakes characterize the tips of the peninsulas and their inland plains, respectively
170 (e.g., Shikoku; Geographical Survey Institute, 1952; Tokai; Ishibashi, 1981). In these area,
171 reverse sense crustal deformation is observed in the interseismic period; gradual subsidence of
172 the peninsulas and uplift of the lowland, respectively (Geographical Survey Institute, 2008).

173 Kikai Island, located at the northern margin of the Amami Islands on the subducting
174 Amami Plateau, is the closest island to the Ryukyu Trench axis. It shows an extremely high
175 uplift rate, with the height of the MIS 5e marine terrace exceeding 200 m. On this island,
176 Holocene marine terraces consisting of coral reefs record several-meter-scale uplift events that
177 recurred with 1500-year intervals (Webster et al., 1998). The other islands along the Ryukyu
178 Trench do not show remarkable crustal deformation, but the coastal notch in Ishigaki Island
179 records a cumulative uplift of ~2 m in the last 2000 years (Kawana, 1989). Some of these
180 marine terraces distributed along the Nankai and Ryukyu coasts have provided the theoretical
181 basis for time-predictable recurrence models for subduction zone earthquakes (e.g., Shimazaki
182 and Nakata, 1980).

183

184 **2.2. Seismicity**

185 Understanding the state of inter-plate coupling and seismogenesis in the subduction zones
186 provides basic information for estimating the region's potential of great earthquake and tsunami.
187 Especially, in the Hyūga-nada region, connecting the Nankai and Ryukyu subduction zones, this
188 information is the key to identifying the geological and seismological differences between the
189 two.

190

191 **Geodetic deformation and inferences of inter-plate coupling along the Nankai Trough and** 192 **Ryukyu Trench subduction zones**

193 The source areas of large earthquakes are usually expected to present strong inter-plate
194 coupling during inter-seismic periods (e.g., Lay and Kanamori, 1981). The degree of the
195 coupling can vary over fault areas, reflecting differences in the fault's frictional properties. If a
196 fault creeps steadily, it has a low degree of coupling, while high coupling results from a
197 seismogenic fault that is stuck during inter-seismic periods. The degree of the coupling is
198 quantified in geodesy (Savage, 1983) by the slip-deficit rate (SDR) (or back-slip rate), which is
199 the rate of upper plate displacement along with the plate interface driven by the subduction
200 motion of the lower plate. The coupling coefficient is also determined as the SDR divided by
201 the long-term average rate of the relative motion between the two plates. When the fault is
202 completely coupled, the SDR is equal to the long-term average relative plate motion rate and the

203 coupling coefficient is unity. SDRs are estimated from geodetic data by applying slip inversion
204 analysis (Savage, 1983).

205 Several geodetic inferences of inter-plate coupling have been recorded along the Nankai
206 Trough subduction zone (Miyazaki and Heki, 2001; Ito and Hashimoto, 2004; Nishimura and
207 Hashimoto, 2006; Wallace et al., 2009; Loveless and Meade, 2010; Yokota et al., 2016), while
208 only a few studies have been conducted along the Ryukyu Trench (Ando et al., 2009; Tadokoro
209 et al., 2018) due to limited spatial coverage of onshore GNSS observations in this island area.
210 Along the Nankai Trough, spatial distributions of SDR have been inferred and certain
211 heterogeneous structures have been identified (Fig. 1), owing to a dense onshore GNSS network
212 called GEONET and the recent development of seafloor geodetic observation networks by the
213 Japan Coast Guard (Yokota et al., 2016). The seafloor observation sites deployed near the
214 trough axis contribute greatly to resolving the previously unknown coupling at shallow depths
215 of the plate interface (Sagiya and Thatcher, 1999), which caused an ambiguity in the estimation
216 of the maximum earthquake size in the area. Lack of knowledge of shallow coupling was clearly
217 illustrated along the Japan Trench subduction zone, which hosted the 2011 Tōhoku-oki
218 earthquake (Nishimura et al., 2000; Loveless and Meade, 2011).

219 All of the SDR estimates indicate that the Nankai Trough subduction interface is strongly
220 coupled overall, with Yokota et al. (2016) further resolving regional heterogeneity, with three
221 areas of relatively stronger coupling with higher SDRs inferred offshore Shikoku Island, the Kii
222 Peninsula and the region from the Atsumi Peninsula to the Omaezaki Peninsula (Fig. 1). The
223 first two areas seem to coincide with the source areas of the 1946 CE Nankai (M8.0) and the
224 1944 CE Tōnankai (M7.9) earthquakes, and the third area seems to be included in the source
225 area of the 1854 CE Ansei-Tōkai earthquake (Ando, 1975b). Given such correlations, the
226 observed heterogeneous distribution of SDR might have played a role in defining the
227 segmentation of historical earthquakes. However, some earthquakes can involve multiple
228 segments, as in the 1707 Hōei (M8.6) earthquake, which is considered to have ruptured all three
229 of the high SDR segments (Ando, 1975b), and the earthquake recurrence pattern is not always
230 characteristic (Seno, 2012).

231 Near the southwestern end of the Nankai Trough subduction zone, the degree of coupling
232 offshore of southern Kyushu Island becomes rather ambiguous due to poorer observational
233 coverage; however, geodetic inferences still tend to suggest weaker coupling in the southern
234 half of the Hyūga-nada region (Nishimura and Hashimoto, 2006; Wallace et al., 2009; Loveless
235 and Meade, 2010) (see Fig.1), reflecting the rotation of the displacement vectors observed over
236 southern Kyushu. The location of this weak coupling is suggested to be correlated with the
237 subducting Kyushu-Palau ridge (Wallace et al., 2009; Yamamoto et al., 2013). To the south,

238 inferences of inter-plate coupling become more difficult due to the much poorer observational
239 coverage.

240 Along the Ryukyu Trench subduction zone, inter-plate coupling has been evaluated
241 qualitatively due to the sparseness of onshore observational stations, with generally weak
242 coupling inferred (Nishimura et al., 2004). However, more quantitative geodetic inferences have
243 been conducted with the addition of ocean bottom geodetic observations. Seafloor displacement
244 data has recently been retrieved from the area offshore the Okinawa Islands, in the center of the
245 Ryukyu Trench, and almost complete coupling of the plate interface is inferred by Tadokoro et
246 al. (2018), in contrast to the onshore-based estimation. It is also suggested that this highly
247 coupled area corresponds to the source area of the 1791 CE M 8 inter-plate earthquake.

248

249 **Seismological inferences of inter-plate coupling using repeating earthquakes**

250 Repeating earthquakes are moderate-sized earthquakes that are considered to occur
251 repeatedly at the same points on faults due to the similarity in their seismic waveforms. They
252 are used to infer the degree of inter-plate coupling in a similar fashion to creep meters (Nadeau
253 and Johnson, 1998). While the absence of repeating earthquakes characterizes both the end
254 members of the fully coupled and the fully decoupled fault, once repeating earthquakes are
255 observed, inter-plate coupling can be estimated, with higher production rates reflecting weaker
256 coupling and more frequent occurrence of fault slip. The use of repeating earthquakes can
257 supplement geodetic inferences due to its higher detection capability, reflecting the weaker
258 geometrical spreading nature of the seismic waves than the geodetically-observed static
259 displacement. Nevertheless, both observations are inevitably limited in island areas.

260 Igarashi (2010) inferred inter-plate coupling by using repeating earthquakes and mapped its
261 spatial variation on the plate interfaces off the Japanese islands. The background seismic
262 activity of the Nankai Trough is generally low reflecting the strong coupling and repeating
263 earthquakes are rarely found except the shallowest portion of the plate interface near the trough
264 axis. However, it is evident that inter-plate coupling becomes weaker from the Hyūga-nada
265 region towards the Ryukyu Trench to the south, matching to the inference from geodetic
266 observations. Yamashita et al. (2012) increased the detection of repeating earthquakes, focusing
267 on the Hyūga-nada region, and mapped a more detailed spatial distribution of inter-plate
268 coupling (Fig. 1). In this region, the coupling appears to be weakest along the band where the
269 Kyushu-Palau ridge is being subducted (Yamamoto et al., 2013), with the repeating earthquakes
270 accommodating most of the long-term relative plate motion at shallower (<~20 km) and deeper
271 (>40 km) depths. However, to the south, the coupling increases in the area off Tanegashima
272 Island, corresponding to the northern end of the Ryukyu Trench, where repeating earthquakes
273 accommodate approximately half the relative plate motion.

274 In the Hyūga-nada region, the history of moderate- and large-sized earthquakes appears to
275 be matched with the inferred coupling distribution (Yamashita et al., 2012). In the northern area
276 of the Hyūga-nada, M 6 and 7-class earthquakes have repeatedly occurred, while in the southern
277 part, the 1996 CE M 6.5 earthquake occurred. Between these regions, major earthquakes are not
278 known, while moderate-sized earthquakes ($5 < M < 6.5$) have been observed in the relatively
279 strongly coupled area.

280 Inter-plate coupling along the Ryukyu Trench seems to show a systematic variation where
281 the northern region – from Tanegashima Island to Amami Island – exhibits relatively strong
282 coupling and the southern region around the Sakishima Islands exhibits weaker coupling.
283 However, the coupling state may not be definitively identified solely based on repeating
284 earthquakes in these island areas; in the central region, while repeating earthquakes suggest
285 moderate coupling, seafloor observations suggest almost full coupling (Tadokoro et al., 2018).

286

287 **Spatial distribution of shallow and deep slow earthquakes; implications for the state of** 288 **inter-plate coupling**

289 Weak inter-plate coupling does not necessarily mean that faults creep steadily. Weak
290 coupling can reflect intermittent slip events on the plate interfaces. Such slip events are
291 generally called slow earthquakes and include slow slip events (SSEs), low-frequency
292 earthquakes and (non-volcanic) tremor. Slow earthquakes are thought to occur along the
293 transition zones of the seismogenic and aseismic zones of plate interfaces (Ando et al., 2012b;
294 Obara and Kato, 2016) both at the up-dip (~ 5 km) and down-dip ends (~ 30 km) of the
295 seismogenic zones that generate the huge earthquakes.

296 The family of slow earthquakes has been discovered over the past twenty years along the
297 Nankai Trough by seismological (e.g., Obara, 2002) and geodetic (e.g., Hirose and Obara, 2005)
298 observations, and their correlation with inferred inter-plate coupling states are confirmed in
299 great detail. Along the plate interface at depths from ~ 30 to ~ 40 km, tremor and short-term
300 SSEs (S-SSEs) are generally observed with recurrence periods of ~ 6 months (Obara, 2002;
301 Hirose and Obara, 2005), while long-term SSEs with M 7 have recurred at ~ 10 -year intervals
302 particularly beneath Lake Hamana in Tōkai region (Suito and Ozawa, 2009 and references
303 therein). For the Ryukyu Trench, seismological observations suggest that slow earthquakes,
304 called very low frequency earthquakes, probably occur on the plate interface over the entire
305 Ryukyu Trench (Ando et al., 2012a), and geodetic observations identify seemingly
306 corresponding short-term SSEs (S-SSEs) and long-term SSEs (L-SSEs) (Heki and Kataoka,
307 2008; Nishimura, 2014). Nishimura (2014) suggested that the occurrence of S-SSEs at shallow
308 depths may be a signature of incomplete inter-plate coupling. Tadokoro et al. (2018) suggested

309 the inferred strong coupling area does not overlap with the SSE areas, and heterogeneity in the
310 areas of SSE occurrence should exist along the Ryukyu Trench.

311 The Hyūga-nada region appears to present a transitional character in terms of the spatial
312 distributions of slow earthquakes. In the other areas of the Nankai Trough, shallow slow
313 earthquakes and deep S-SSEs and tremors are both observed, while deep S-SSE/tremor is not
314 observed in the Hyūga-nada region (Obara and Kato, 2016). The occurrence of shallow slow
315 earthquakes has been characterized in great detail owing to the recent deployment of ocean
316 bottom seismometers (Yamashita et al., 2015). Tremor was observed to migrate in the shallow
317 depths of the plate interface, and the lateral extent of the migration was delimited by the
318 possible area of the subducting Kyushu-Palau ridge (Yamamoto et al., 2013). The down-dip
319 limit of the migration was not reached until ~ 20 km, corresponding to the hypocentral depths of
320 M 7-class earthquakes in this region, suggested that it stopped at the shallower end of the
321 seismogenic zone (Yamashita et al., 2015). This implies that the intermediate depths of the plate
322 interface are still coupled and have the potential to nucleate a major earthquake. The
323 geophysical observations clearly show that the inter-plate coupling of this area is relatively low
324 overall, but these observations still leave the possibility that the rupture of this coupled zone
325 could be triggered by neighboring large earthquakes, or the rupture of this area could facilitate
326 interactions between neighboring large earthquakes, if their timings are synchronized.

327 While the above-mentioned geophysical observations have clarified the spatial
328 characteristics of the inter-plate properties, we should still remember that our experience is
329 limited by the short history of geophysical observations. The states of inter-plate coupling can
330 slowly fluctuate even over the course of several years (Uchida et al., 2016), and limited
331 snapshots can fail to reveal critical properties. Integration of paleoseismological evidence is still
332 critically important to capture phenomena spanning long periods of time.

333

334 **3. Record of tsunamis and tsunami deposits**

335 **3.1. Nankai Trough region**

336 **3.1.1. Historical Tōkai and Nankai earthquakes and tsunamis**

337 The Nankai Trough region, which is close to the political and cultural center of Japan, the
338 Kyoto-Nara-Osaka area, has the longest and most complete documented record for the
339 recurrence of subduction zone earthquakes and tsunamis in the world. According to this
340 1300-year long record, earthquakes have occurred in a variety of modes. Some earthquakes
341 have ruptured almost the entire subduction zone at the same time, while at other times, two
342 earthquakes, referred to as Tōkai and Nankai earthquakes, have occurred consecutively,
343 separated by short intervals (Fig. 2). Nankai earthquakes that are historically confirmed include
344 the 684 CE Hakuho (M8.0-8.5), 887 CE Nin-na (M8.0-8.5), 1361 CE Kōan (M8.2-8.5), 1707

345 CE Hōei, 1854 CE Ansei-Nankai (M8.4; only 32 hours after the 1854 Ansei-Tōkai earthquake),
346 and 1946 CE Showa-Nankai (M8.0) earthquakes, with further unconfirmed events in 1099 and
347 1614 CE. Tōkai earthquakes confirmed by documented records include the 1096 CE Eichō
348 (M8.0-8.5), 1498 CE Meiō (M8.2-8.4), 1707 CE Hōei (M8.6), 1854 Ansei-Tōkai (M8.4), and
349 1944 CE Tōnankai (M7.9) earthquakes, and a further rupture may have occurred in 1614 CE.
350 The 1707 earthquake was a simultaneous rupture of the Tōkai and Nankai fault segments
351 (segments A to D, and perhaps also including the southern part of E) and was the largest
352 earthquake in Japanese history until the occurrence of the 2011 Tōhoku earthquake.

353 There are two possibilities for a great earthquake that occurred along the Nankai Trough in
354 the early 17th century. The first is the 1605 CE Keichō earthquake, which generated a large
355 tsunami from Kanto to Kyushu. This event has been considered as a “tsunami earthquake”,
356 because only weak shaking accompanied the large tsunami (e.g., Ishibashi and Satake, 1998).
357 Ando and Nakamura (2013) suggested that the rupture occurred along a shallow portion of the
358 plate interface. An alternative hypothesis is that the 1605 earthquake occurred along the
359 Izu-Ogasawara Trench, to the east of Izu-Bonin Arc (Harada et al., 2013). Instead of 1605 CE
360 event, Ishibashi (2014) proposed the 1614 CE earthquake as a candidate for a 17th century
361 Nankai Trough earthquake, based on records of strong ground shaking in Kyoto and tsunamis at
362 the Kii Peninsula. A Nankai earthquake paired with the 1498 CE Meiō Tōkai earthquake has not
363 been confirmed; nevertheless, liquefaction features in western Japan suggest the occurrence of a
364 Nankai earthquake around 1498 CE (Sangawa, 2001, 2007). Ishibashi (2014) proposed four
365 earthquakes that occurred between 1498 and 1512 CE as candidates, but there is no solid record
366 of tsunami inundation during this period.

367 Documentary records confirming the occurrence of a Tōkai earthquake paired with the
368 1361 CE Nankai earthquake have not yet been discovered. As a candidate for this earthquake,
369 Ishibashi (1998, 2014) proposed an earthquake that featured strong shaking in Kyoto, Nara, and
370 Kumano (north of Shiono-misaki; boundary between the segments B and C) on July 24th 1361
371 (2 days before the Nankai earthquake). Late 14th century coseismic uplift around Cape
372 Omaezaki (Fig. 2, site 27) shown by emerged boring bivalves (Kitamura et al., 2018c) and
373 marine terrace (Fujiwara et al., 2010) support Ishibashi’s hypothesis.

374 The occurrence of the 1099 CE earthquake was originally inferred from documents
375 suggesting strong ground shaking in the Kyoto, Nara, and Osaka region (but less than 1096 CE
376 earthquake) and coastal subsidence in Kochi (around site 10 in Fig. 2); which matches general
377 characteristics of other Nankai earthquakes (e.g., Ishibashi, 1999). However, the occurrence of
378 this earthquake is uncertain due to the unreliability of underlying evidence, especially the lack
379 of reports of strong ground shaking and tsunami inundation in the area facing segments A and B.
380 Ishibashi (2016) suggested that the 1099 CE earthquake did not actually exist and that the 1096

381 CE earthquake was instead a full-length rupture of the Tōkai and Nankai segments. Ishibashi
382 (2014) also suggested the possibility that the 1099 CE Nankai earthquake did occur but was
383 smaller in magnitude than other Nankai earthquakes.

384 The possibility that a Tōkai earthquake occurred at the same time as the Nankai earthquake
385 on 22nd August, 887 CE has been suggested from a description in the Nihon Sandai Jitsuroku
386 (日本三代実録, "The true history of three reigns of Japan"; completed in 901 CE) (e.g.,
387 Ishibashi, 1999, 2014). The chronicle's record for that day shows that strong ground shaking
388 was felt in wide area including the Tōkai region along with the occurrence of an earthquake and
389 tsunami in the Nankai region. The occurrence of the 684 CE Nankai earthquake is documented
390 in the Nihon Shoki (日本書紀, The Chronicles of Japan, completed in 720 CE), with a report of
391 tsunami and coastal submergence in Kōchi, but there are no records of a corresponding
392 earthquake in the Tōkai region. Only liquefaction features in archaeological sites in the Tōkai
393 region have suggested as possible evidence of a Tōkai earthquake corresponding to the 684 CE
394 Nankai earthquake (e.g., Sangawa, 2001, 2007). However, given the lack of evidence for
395 tsunami to determine the occurrence of the Tōkai earthquake in either case, the possibility of an
396 inland earthquake from an active fault cannot be ruled out. These imperfections in the historical
397 earthquake record have been a hindrance in discussing the occurrence time of Tōkai and Nankai
398 earthquakes and their linkage.

399 The 1498 CE Meiō tsunami might have been somewhat higher than 1707 and 1854 CE
400 tsunamis, ~ 8 m along the Enshu-nada coast, which includes the sites 19 to 27 in Fig. 2 (Hatori,
401 1975). The height of the 1605 CE tsunami was generally 4 - 6 m and up to 13 m in Shikoku
402 (Murakami et al., 1996). The 1707 CE tsunami generally reached 5 - 8 m-high along the Tōkai
403 and Nankai coasts, over 10 m in some locations on the Kii Peninsula and Shikoku, and 2 - 4 m
404 along the Kyushu coast (Murakami et al., 1996, Watanabe, 1998). The 1854 CE Ansei-Tōkai
405 tsunami had a height of 5 - 8 m, with some isolated cases of ~20 m at the tips of promontories
406 (Watanabe, 1998). The 1854 CE Ansei-Nankai tsunami generally reached 5 - 6 m-high,
407 occasionally over 8 m along the Kii Peninsula and Shikoku (Murakami et al., 1996, Watanabe,
408 1998), and 2 - 4 m along the Kyushu coast (Cabinet Office, 2012b). The 1944 CE tsunami was
409 3 - 7 m-high along the eastern coast of Kii Peninsula and 0.9 - 2 m-high along the Enshu-nada
410 coast (Watanabe, 1998). The height of the 1946 CE Showa-Nankai tsunami was 2 - 4 m in
411 general, up to 6 m on the Kii and Shikoku coasts and 1 - 1.6 m on the Kyushu coast (Murakami et
412 al., 1996, Watanabe, 1998). These tsunamis generally caused greater human and economic
413 damages than resulted from ground shaking during historical Nankai Trough earthquakes (e.g.,
414 Yata, 2009, 2018; Ishibashi, 2014).

415

416 **3.1.2. Tsunami deposits**

417 Nankai Trough coasts have few extensive wetlands suitable for reconstructing the tsunami
418 inundation area, which characterize the major paleotsunami study fields in the world, such as
419 Hokkaido (e.g., Nanayama et al., 2003), Cascadia (e.g., Peters et al., 2007) and Chile (e.g.
420 Cisternas et al., 2005). Artificial disturbance of surface sediments reduces the chances of
421 finding tsunami deposits from this region, along with the difficulties of distinguishing tsunami
422 deposits from the other washover (mainly storm) deposits and river-flooding deposits.
423 Consequently, coastal back-barrier lakes and ponds, and infilled valleys have mainly been
424 targeted as paleotsunami study sites. The barriers between these sites and the sea are mainly
425 composed of sand dunes and beach ridges formed during the last seven to six thousand years,
426 after the mid Holocene sea level high stand in Japan (e.g., Sato et al., 2016a). Many of these
427 sites have gradually subsided through the Holocene, and stably accumulated fine-grained
428 sediments, sometimes rich in organic matter (e.g., Okamura and Matsuoka, 2012). They have a
429 high potential for preserving marine overwash event deposits. Coring surveys using hand corers,
430 piston corers, and geoslicers (Nakata and Shimazaki, 1997) have been the main methods to
431 reveal the tsunami deposits, while their depositional ages have mainly been determined by
432 radiocarbon dating. Due to the very limited distribution of widespread Holocene tephtras along
433 the Nankai and Ryukyu subduction zones, it is difficult to estimate the age of tsunami deposits
434 based on volcanic ash stratigraphy and to compare them between regions, as has been done in
435 the Hokkaido (e.g., Nanayama et al., 2003) and Tōhoku (e.g., Sawai et al., 2012) regions.

436 Tsunami and possible tsunami deposits have been reported from a total of 23 study sites
437 along the Nankai Trough coast (Fig. 2, sites 7 to 20, 22 to 26, 28, 29, 31 and 32). Adjacent
438 points that cannot be distinguished on the map, such as some points in the southern part of the
439 Kii Peninsula, are counted as one. Garrett et al. (2016) recalibrated all available published
440 radiocarbon ages to take advantage of the latest radiocarbon calibration curves, updated
441 estimates of local marine radiocarbon reservoir effects, and Bayesian age modeling approaches.
442 As a result, some of the originally reported correlations between sedimentary layers and
443 historical earthquakes have been revised. We follow Garrett et al. (2016) for the ages of tsunami
444 deposits summarized here.

445 Most of the reported tsunami and possible deposits are washover sand deposits intercalated
446 in muddy or peaty deposits, with two exceptions of the tsunami boulders found from the
447 southern Kii peninsula (Fig. 2, site 14) and the southern Izu peninsula (Fig. 2, site 32). Sandy
448 tsunami deposits generally show a tapering shape with a fining landward trend and sometimes
449 include remains of marine organism. They may be a single layer, but often consist of multiple
450 layers, each of which shows a graded structure. Sedimentary structures indicating deposition
451 from a tractive flow, such as basal erosion surface, cross-lamination, and inverse grading also
452 characterize many cases.

453

454 **Historical tsunami deposits**

455 Examples of tsunami deposits that can be accurately correlated to historical earthquakes are
456 still limited in the Nankai Trough region (Fig. 2). Limited numbers of measurements and
457 chronological uncertainties inherent in the radiocarbon approach make it difficult to distinguish
458 individual historical tsunami deposits that formed at short intervals, often of less than 100 years,
459 and compare them between distant sites.

460 The easternmost examples are reported by Kitamura and Kobayashi (2014a) and Kitamura
461 et al. (2014) from Shimoda, southern Izu Peninsula (Fig. 2, Site 32). The former is a laminated
462 sand bed about 10 cm thick, with rip-up clasts that is estimated to have been deposited between
463 430-290 cal. BP and 1950 CE, possibly due to the 1854 CE Ansei-Tōkai or 1707 CE Hōei
464 tsunami. The latter is a 32-tonne boulder located on a coastal plateau, which is attributed to the
465 1854 CE Ansei-Tōkai earthquake based on the radiocarbon ages of sessile marine fossils
466 attached to the boulder.

467 On the Ita Lowlands (Fig. 2, site 31), one possible sandy tsunami deposit – with diatoms
468 suggesting a marine incursion – was found from a sedimentary sequence covering the last 2500
469 years (Sawai et al., 2016). Radiocarbon dating limited the age of the tsunami bed to 1200-1320
470 CE or 1150-1330 CE and Sawai et al. (2016) attributed it to the 1096, 1099, or 1361 CE Nankai
471 Trough earthquake or the 1293 CE Kantō earthquake, centered on the Sagami Trough, east of
472 Izu Peninsula. On the Shimizu Plain, Kitamura and Kobayashi (2014b) reported the deposition
473 of gravelly sand beds associated with the erosion of the muddy bay bottom between 670–600
474 and 250–50 cal. BP (Fig. 2, Site 29). They inferred that the sand beds may be attributed to the
475 tsunamis from 1498 CE Meiō or 1707 CE Hōei earthquakes.

476 Komatsubara et al. (2008) conducted paleotsunami research in a back-marsh at Shirasuka
477 (Fig. 2, site 20) using a 4 m-long geoslicer and found exotic sand beds within the muddy
478 sedimentary sequence. Comparing the radiocarbon ages obtained from the cores and historical
479 documents, five sand beds, each of which is 10 - 20 cm thick, were attributed to tsunamis
480 associated with the 1498, 1605, 1707, 1854 CE earthquakes and the 1680 or 1699 CE typhoon.
481 This hypothesis was verified with new age models developed using radiocarbon ages (Garrett et
482 al., 2018) and optically stimulated luminescence (OSL) ages (Riedesel et al., 2018) from
483 subsequently excavated sediment cores from the same site as the original study. The sand unit A
484 of Komatsubara et al., (2008), located beneath the 1498 CE tsunami deposit and originally
485 interpreted as a slope failure deposit, was recognized as a possible tsunami deposit based on the
486 occurrence of marine and brackish diatoms (Garrett et al. 2018). Both radiocarbon and OSL age
487 models suggest an age for this tsunami deposit of the late 14th century and the layer may be
488 attributed to the 1361 CE Tōkai earthquake (Garrett et al., 2018; Riedesel et al., 2018).

489 Fujino et al. (2018) conducted an array coring survey on the Shima Lowland (Fig. 2, site
490 17), a Holocene drowned valley, and found 10 exotic sand beds within the muddy marsh
491 sequence. A tsunami origin for these beds is inferred from sedimentary features common in
492 modern tsunami deposits, including marine bioclast-rich sand, alternating sub-layers of sand and
493 silt, rip-up clasts, and upward-fining structures. The youngest three tsunami beds were
494 tentatively attributed to 1498, 1096, and 684 CE Nankai Trough earthquakes.

495 A group of possible tsunami boulders, consisting of hundreds of angular boulders weighing
496 up to 100 tons (Namegaya et al., 2011), are found on a wave-cut bench near Cape
497 Shiono-misaki (Fig. 2, site 14). These boulders were derived from the Middle Miocene
498 quartz-porphry dyke penetrating the basement sedimentary rock, which is located seaward of
499 the boulder group. The dyke and boulders are famous as the tourist spot of “Hashigui-iwa”,
500 which means large rocks (iwa) look like bridge (Hashi) piles (gui or kui) in Japanese. Excluding
501 smaller ones < 1m in diameter, these boulders were not moved by the 1946 CE Showa-Nankai
502 tsunami or subsequent storms (Shishikura, 2013). Shishikura (2013) recognized two major
503 boulder transport events based on the radiocarbon ages of sessile marine fossils, the 12-14th
504 century and the 17-18th century CE, and attributed them to the 1361 and 1707 CE tsunamis,
505 respectively.

506 Tōkai earthquakes corresponding to the 887 and 684 CE Nankai earthquakes have
507 remained unconfirmed for a long time, but have recently been substantiated by the discovery of
508 tsunami deposits on the Otagawa Lowland (Fig. 2, site 25) in western Shizuoka Prefecture
509 (Fujiwara et al., 2020). Excavation walls exposed by river improvement work in the lowland
510 revealed a ~1-km long coast-normal and 4 to 5 m-deep cross section of the strand plain (Fig.
511 3A). Fujiwara et al. (2020) mapped four sandy tsunami deposits along the excavation walls,
512 each of which shows inundation over 2 km inland from the coast at the time of tsunami, with
513 landward thinning and fining trends. Radiocarbon ages link the younger two tsunami deposits to
514 the 1498 and 1096 CE Tōkai earthquakes. The older two deposits indicate the occurrence of
515 Tōkai earthquakes near to the end of the 7th and 9th centuries CE. The late 9th century tsunami
516 deposit, integrated with the description of strong ground shaking in the Nihon Sandai Jitsuroku,
517 demonstrates that the 887 CE earthquake was a full-length rupture of the Tōkai and Nankai
518 segments (Fujiwara et al., 2020). As the late 7th century tsunami deposit lacks data to identify
519 the precise date of its occurrence, it is not known whether a Tōkai earthquake is coincident with
520 the 684 CE Nankai earthquake.

521 For Shikoku and northern Kyushu, in addition to the summary provided by Okamura and
522 Matsuoka (2012) from 26 ponds, Baranes et al. (2016) conducted paleotsunami research in three
523 ponds. Among these 29 sites, historical tsunami deposits have been found in four locations. At
524 Ryujin-ike in northern Kyushu (Fig.2, site 7), separated by a 17 m high and 100 m wide sand

525 dune, a narrow channel is the only access to the Pacific. According to historical documents, the
526 1707 CE Hōei tsunami, over 10 m-high, was much higher than that of 1854 and 1946 CE
527 tsunamis around Ryujin-ike (e.g., Chida and Nakaue, 2007). Among 40 sand beds intercalated
528 within the 5.4 m long cored humic mud sequence, eight prominent beds, each 5 - 20 cm-thick,
529 are interpreted as washover deposits due to their thinning landward trend and the inclusion of
530 beach sand and marine bioclasts (Okamura and Matsuoka, 2012). Based on radiocarbon dating,
531 Okamura and Matsuoka (2012) linked the youngest three beds to the 1707, 1361 and 684 CE
532 tsunamis.

533 Baranes et al. (2016) reported a total of nine washover sand beds from the back-barrier
534 lakes facing the Bungo Channel in western Shikoku (Fig. 2, site 8). A 9.3 m-deep sediment core
535 from Lake Ryuuō, consisting of mainly clay beds and covering ~2500-year-long record of
536 deposition, shows an environmental change from a brackish lake to freshwater conditions that
537 occurred around 1000 CE (about 6 m below the lake bottom). Nine washover sand beds occur in
538 the freshwater clay between the depths of 595 and 271 cm, accumulated from 1000 to 1800 CE.
539 Although with substantial chronological uncertainties, their depth-to-age model suggests the
540 possibility that four washover sand beds represent the 1099, 1361, 1605, and 1707 CE tsunamis.
541 There are two sand layers deposited around 1500 CE, one of which may be due to a Nankai
542 earthquake that is unknown from historical records and paired with the 1498 Meiō Tōkai
543 earthquake. Baranes et al. (2016) concluded that the 1707 CE tsunami, which left the thickest
544 (25 cm) coarse-grained sand bed, was one of the most significant floods over the last
545 millennium in the region.

546 Fourteen sandy tsunami deposits ranging from 4800 to 1300 cal BP in age are reported
547 from Tadasu-ike (Fig.2, site 9), located 800 m inland from the modern coast and behind a 5 m
548 high beach ridge (Okamura and Matsuoka, 2012). The sand beds range from 5 to 27 cm in
549 thickness. The pond was inundated by the 1707 and 1946 CE tsunamis, however the record of
550 the last 1300 years is missing in this pond due to artificial disturbance; the youngest bed was
551 linked to the 684 CE tsunami (Okamura and Matsuoka, 2012). Kaniga-ike in Shikoku (Fig.2,
552 site 10), located about 400 m inland from the modern coast and fringed by 6 m-high coastal
553 dunes, records six sandy tsunami deposits in the last 2000 years (Okamura and Matsuoka, 2012).
554 The tsunami beds range from 5 to 75 cm in thickness. Based on historical documents and
555 limited radiocarbon ages (Cabinet Office, 2011a), the youngest two can be correlated with the
556 1854 CE Ansei-Nankai and 1707 CE Hōei earthquakes, while the third and fourth ones from the
557 top may be attributed to the 1361 and 684 CE Nankai earthquakes, respectively.

558

559 **Prehistoric tsunami deposits**

560 Tsunami deposits before the oldest documented Nankai Trough earthquake, the 684 CE
561 Hakuho earthquake, have been recorded as far back as 6000 years ago. Kitamura and Kobayashi
562 (2014a) reported four possible tsunami sand beds from a muddy shallow bay sequence in the
563 Shimizu Plain (Fig. 2, site 29), which are dated to 6180–6010 to 5700–5580, 5700–5580 to
564 5520–5320, 4335–4125 to 4250–4067, and 3670–3540 to 3500–3360 cal. BP, respectively.
565 Kitamura et al. (2013) identified two possible tsunami sand beds from a back-marsh sequence in
566 the Shizuoka Plain covering over the past 4000 years (Fig. 2, site 28).

567 Drowned valleys on the Shima Peninsula archive a long and numerous record of possible
568 paleotsunamis. Okahashi et al. (2005) reported up to 12 washover beds deposited from 6,000 to
569 1,600 cal BP at Osatsu (Fig. 2, site 18). At Shima (Fig. 2, site 17), Fujino et al. (2018) reported
570 seven washover sand beds beneath the three historical tsunami beds. Their depth-to-age model,
571 based on radiocarbon dates, indicates that these ten tsunami beds were deposited from about
572 4,500 to 500 cal BP, with an interval of 100 to 600 years.

573 Lake Hamana, a large brackish lagoon on the Enshu-nada coast (Fig. 2, site 23), has also
574 been targeted for paleotsunami studies by several research groups. Tsuji et al. (1998) analyzed
575 cores of up to 2 m in length from the flood-tide delta and interpreted eight exotic coarse-grained
576 beds with gravel and marine shells as possible tsunami deposits. The four older deposits formed
577 between about 4,800 and 3,200 cal BP; storm surges and tidal channel migration also remain
578 possible explanations for them. The four youngest deposits are linked with tsunami inundation
579 in 1854 or 1707, 1498 CE, the 13th century and 1096 CE by Tsuji et al. (1998); however, more
580 recent recalibration of radiocarbon data highlights difficulties with ascribing particular historical
581 tsunamis to these deposits (Garrett et al., 2016). Sato et al. (2016b) identified a marine incursion
582 event around 4790 – 4420 cal. BP based on fossil diatom analyses of a sediment core obtained
583 from the central Lake Hamana.

584 The western Hamamatsu Plain (Fig. 2, site 24) has a maximum width of about 4 km and
585 six rows of beach ridges (Fig. 4A). Possible tsunami sand beds intercalated with mud and peat
586 were reported from two separate swales (Sato et al., 2016a) and an incised valley (Fujiwara et
587 al., 2013a). Sato et al. (2016a) does not state that the sand beds are tsunami deposits, but
588 additional evidence suggests a possible tsunami origin for them. Each sand bed, generally less
589 than 20 cm thick, shows an erosive base and upward fining trend, and records the occurrence of
590 sediment flow in the back-marsh. A rapid increase in marine and brackish diatoms recognized
591 within and just after the deposition of some of the sand beds and suggests coastal subsidence
592 coincident with the marine incursion (Fig. 4D). The uppermost tsunami deposits are
593 progressively younger with decreasing distance from the present coastline. In a
594 landward swale (Fig. 4B) and in the Rokkengawa lowland (Fig. 4E), 4 km inland from the
595 modern coast, the uppermost deposit dates to just before the fall of the Kawagodaira pumice

596 (1210-1187 cal BCE, Tani et al., 2013). In a more seaward swale, 2 km from the present coast
597 (Fig. 4C), the uppermost deposit dates to 775-895 CE.

598 In Shikoku and Kyushu, five tsunami deposits have been recognized from Ryujin-ike (Fig.
599 2, site 7) between 3300 and 1600 years ago, thirteen from Tadasu-ike (Fig. 1, site 9) between
600 3300 and 1600 years ago, and two from Kaniga-ike (Fig. 2, site 10) between 1000 BCE and 600
601 CE (Okamura and Matsuoka, 2012). Tanigawa et al. (2018) reported four sandy washover
602 deposits tapering landward from sediment cores drilled in a small coastal lowland in Nankoku,
603 Kochi Prefecture (Fig. 2, site 11). The lowland is separated from the Pacific by dunes 450 m
604 wide and 13 m high. The sand beds are laterally extensive through marine-brackish clay and
605 overlying freshwater clay and peat. Tanigawa et al. (2018) interpreted the sand beds as having
606 been deposited by tsunamis or unusually large storm surges. The possible tsunami sand beds
607 were found in sediments with ages between 5970 and 2440 cal BP. They attributed the lack of
608 marine incursions after 2440 cal BP to the development of beach ridges that protected the site
609 from tsunamis and storm surges.

610 Shimada et al. (2019) reported five tsunami inundations between 5581 and 3640 cal BP
611 from a small drowned valley in Mugi (Fig. 2, site 12), eastern Shikoku. Array coring along
612 the valley axis and diatom analyses of the cored sediments revealed the deposition of at least
613 nine exotic sand and gravel beds in the muddy brackish-to-fresh wetland sequence.
614 Environmental changes inferred from diatom assemblages suggest the occurrence of coastal
615 subsidence associated with the deposition of three of the sand beds. Subsidence of 0.3 m or
616 more was reported around the study site during the 1946 CE Showa-Nankai earthquake (Japan
617 Coast Guard, 1948). Shimada et al. (2019) tentatively attributed two other sand beds to tsunami
618 events, while acknowledging the possibility of deposition from a storm surge or far-field
619 tsunami. Reported tsunami deposits can be traced up to 240 m or more inland from the current
620 shoreline, and they range in thickness from 5 to 20 cm with some exceptions up to 75 cm.

621

622 **3.2. Transitional boundary of Nankai Trough and Ryukyu Trench (Hyūga-nada)**

623 **3.2.1. Historical earthquakes and tsunamis**

624 The Hyūga-nada region covers the western end of Nankai Trough and the northern end of
625 Ryukyu Trench. Seismic activity in this region is relatively high and small tsunamis were
626 detected following the 1941, 1961, 1968, and 1984 CE earthquakes (Hatori, 1985). Historically,
627 large tsunamis occurred in 1662 and 1769. The 1662 CE earthquake was the largest historical
628 event, with a magnitude of 7.6 (HERP, 2004) and a maximum tsunami height of <5 m at
629 Miyazaki Plain (Hatori, 1985). Hatori (1985) estimated that the 1662 CE tsunami could have
630 inundated the lowland of Miyazaki Plain via the large rivers, while tsunamis might not have

631 been able to overtop the coastal sand dunes because of their height (~10 to 15 m, Nagaoka et al.,
632 1991).

633

634 **3.2.2. Paleotsunami deposits**

635 Paleotsunami studies, both historical and prehistoric, are very scarce in this region; a major
636 reason is probably the lack of suitable lowlands. Although the Miyazaki Plain is one candidate
637 site for paleotsunami research, the coastal dunes are too high and the rivers and floodplains are
638 large. Therefore, it is not straightforward to locate tsunami deposits in such a depositional
639 environment. Nevertheless, Ichihara et al. (2015) reported a possible tsunami deposit at around
640 2000 years ago on this plain.

641 Further south at Kushima City (Fig. 2, site 6), Yamada et al. (2020) conducted paleotsunami
642 research and found only one tsunami deposit at 4600 years ago during the period between 2000
643 and 5000 years ago. Based on this result, they suggested that tsunamigenic earthquakes
644 probably only occur at low frequencies in the Hyūga-nada region.

645

646 **3.3. Paleotsunamis on the Ryukyu Islands**

647 **3.3.1. Historical earthquakes and tsunamis**

648 Most islands in the Ryukyu Islands are surrounded by fringing reefs; these are typically
649 wider in the south of the island chain (~1.5 km wide) and narrower in the north (a few hundred
650 meters wide). Based on the geographical distribution and paleotsunami records, we divide these
651 islands into three groups: the Amami, Okinawa and Sakishima Islands. The three groups face
652 the northern, central, and southern Ryukyu Trench, respectively (Fig. 1). In between Kyushu
653 and the Amami Islands, there are several islands which we group as the Ōsumi and Tokara
654 Islands. Since no paleotsunami generated by subduction zone earthquake have been reported,
655 we do not discuss this region in this study. The Ryukyu Islands align parallel to the trench and
656 hence each island has an approximately equal chance of being affected by tsunamis if
657 earthquakes are generated along the trench.

658 In contrast to Nankai Trough region, historical records from the Ryukyu Islands are only
659 available for the last ~400 years. Historical and instrumental data show that earthquake and
660 tsunami occurrence in this region is remarkably low in frequency. Indeed, the only earthquakes
661 with magnitudes >8 during the past ~400 years were in 1911 CE in the northern Ryukyu Trench
662 (Mw=8.0, e.g., Tsuji, 1997), in 1791 CE in the central Ryukyu Trench (Mw=8.0 - 8.3,
663 Nakamura and Kinjou, 2013; Tadokoro et al., 2018), and in 1771 CE in the southern Ryukyu
664 Trench (Mw>8.0, e.g., Nakamura, 2009). However, because there is no record of ground
665 shaking accompanying the 1791 CE tsunami, it is uncertain whether this resulted from a
666 near-field tsunamigenic earthquake (Nakamura and Kinjou, 2013; Tadokoro et al. 2018) or was

667 potentially a far-field tsunami generated somewhere in the Pacific Ocean (Matu'ura, in press).
668 The 1771 CE tsunami was exceedingly large with maximum run-up heights of approx. 30 m
669 (e.g., Goto et al., 2010a). However, based on detailed historical records, it is also well known
670 that the tsunami-affected area was spatially limited to a narrow range of the Sakishima Islands
671 (Goto et al., 2010a).

672 Lowland plains suitable for paleotsunami research using sandy tsunami deposits are
673 extremely scarce because of the fringing reef environment. On the other hand, numerous coral
674 boulders are scattered on the reefs surrounding many islands and they are considered as useful
675 geological evidence of past extreme wave events such as tsunamis and storm waves (Fig. 3B,
676 e.g., Kawana and Nakata, 1994; Goto et al., 2010a).

677 In this section, we summarize paleotsunami research on the Ryukyu Islands based on the
678 comprehensive review by Goto (2017) with additional recent works. As a brief summary, the
679 paleotsunami history of the Amami and Okinawa Islands can be characterized by scarce (or
680 even no) evidence of large tsunamis over the last few thousand years, while the Sakishima
681 Islands can be characterized by frequent large tsunamis with intervals of a few hundred years.

682

683 **3.3.2. Paleotsunami deposits on the Amami and Okinawa Islands**

684 On the Amami Islands, Kawana and Nakata (2003) estimated the possible occurrence of
685 tsunamis in 360-510 cal BP, 1390-1770 cal BP, and 2100-2110 cal BP, based on dating coral
686 boulders on the reef. Similarly, Kawana (2006) and Iwai and Kawana (2008) suggested possible
687 large tsunamis around 3400 years ago on the Okinawa Islands, based on the dating of beach
688 rocks and coral boulders. However, following studies by Goto et al. (2009, 2013), field surveys
689 and remote sensing analyses revealed that all of the coral boulders deposited on the reefs
690 fringing the Amami and Okinawa Islands can be explained by transport during storms despite
691 their heavy weight (~160 tons). The boulders are deposited close to the source area (i.e. around
692 the reef edge) and have not been transported a long distance. Since the reef was formed and the
693 boulders were emplaced on the reef at least 2300 years ago (Kawana and Nakata, 2003), the
694 present clast size and spatial distribution of boulders reflects the cumulative impacts of many
695 storm waves since at least 2300 years ago. If large tsunamis had occurred during this period,
696 boulders originally deposited on the reef by storm waves should be transported further inland, as
697 is observed in the Sakishima Islands (discussed below). Therefore, Goto et al. (2013) concluded
698 that no tsunamis large enough to significantly modify the characteristic distribution of storm
699 wave boulders have affected these islands in the last 2300 years. Indeed, using numerical
700 modeling, Minamidate et al. (2020) confirmed that boulders in the Okinawa Islands can be
701 explained by the realistic size of storm waves generated by past typhoons.

702 Another important paleoseismological aspect, especially in the Amami Islands, is a rapid
703 uplift of Kikai Island (e.g., Webster et al., 1998). At Kikai Island, the terrace of MIS 5e is at an
704 elevation of 224 m (e.g., Ota and Omura, 2000). At least 4 emerged Holocene coral terraces are
705 observed and each terrace is associated with emergence of 1 to 4 m (Webster et al., 1998;
706 Sugihara et al., 2003; Hongo, 2010). Because of these geomorphological features, it is thought
707 that Kikai Island was intermittently uplifted by large earthquakes (Nakata et al., 1978). On the
708 other hand, considering the absence of tsunami boulders but presence of storm boulders both on
709 Kikai and Amami-Oshima Islands (Goto et al., 2013), earthquakes that might have caused the
710 uplift of the island may not have generated large tsunamis. The uplift might be explained by slip
711 on an intra-plate fault, such as a splay fault, rather than slip on the subduction interface (Goto,
712 2017). This idea may better fit with the fact that nearby Amami-Oshima Island (~40 km to the
713 west) has not been uplifted significantly compared with Kikai Island (Ikeda, 1977) and hence
714 these islands are tilted within a narrow area. Alternatively, Shikakura (2014) numerically
715 modeled crustal movements and the formation of the coral terraces at Kikai Island and
716 suggested that large earthquakes are not mandatory requirements, but steady uplift with some
717 intermittent earthquakes may alternatively be enough to explain the uplift as well as the
718 formation of multiple coral terraces.

719

720 **3.3.3. Paleotsunamis on the Sakishima Islands**

721 Because of historical records that describe the movement of huge boulders during the 1771
722 CE tsunami and the presence of numerous boulders deposited on the reef and on land,
723 identification of historically described 1771 CE tsunami boulders has proved an attractive topic.
724 With this objective, geological and historical studies of the 1771 CE tsunami and boulders
725 began earlier than tsunami studies in other regions, with the earliest work by Makino (1968).
726 Makino (1968, 1981) interpreted that all boulders deposited on land were transported by the
727 1771 CE tsunami. However, following studies cautioned the need for careful identification
728 because many of the boulders are composed of the Pleistocene Ryukyu Limestone that forms
729 the island itself and may have been sourced from higher elevations and moved by gravity (Kato
730 and Kimura, 1983; Kawana and Nakata, 1994). Nevertheless, there are numerous coral boulders
731 with fresh Holocene coral skeletons that must have been transported from the sea possibly by
732 large tsunamis during Holocene (Kato and Kimura, 1983; Kawana and Nakata, 1994).

733 Radiocarbon dating of coral boulders has been performed since early 1980's and boulders
734 with ages around 1771 CE have been discovered, suggesting their origin from this tsunami (e.g.,
735 Kato and Kimura, 1983; Kawana and Nakata, 1994). However, many studies have revealed that
736 there are boulders with much older ages than 1771 CE, suggesting that these boulders are likely

737 to have been deposited by repeated tsunamis during the few thousand years before 1771 CE
738 (Kawana and Nakata, 1994; Omoto, 2012; Araoka et al., 2013).

739 Problems associated with pre-2010 CE studies on these boulders include 1) discrimination
740 of tsunami boulders from storm boulders, and 2) dating accuracy and reliability of tsunami age
741 estimation. Regarding the first issue, Goto et al. (2010b, 2013) investigated clast size and the
742 spatial distribution of boulders on the Sakishima Islands and suggested that coral and reef
743 boulders can be classified into two groups. One group has a characteristic distribution of storm
744 wave boulders such as a narrow range in spatial distribution from the reef edge and an
745 exponentially fining landward trend in clast-size distribution. The boulders of this group can be
746 considered as of storm wave origin (Goto et al., 2010b, 2013). On the other hand, the other
747 group can be characterized by boulders that are deposited along the coast and on land. They are
748 deposited far beyond the possible landward limit of storm wave boulders (i.e. ~300 m from the
749 reef edge, Goto et al., 2009). Also, there is no clear clast size distribution indicative of storm
750 deposition. Based on these features, Goto et al. (2010b, 2013) identified the boulders in this
751 group as of tsunami origin. Later, Watanabe et al. (2016) numerically confirmed the tsunami
752 origin of the boulders in the latter group, identifying that extremely long wavelengths are
753 required to explain the deposition of boulders along the coast.

754 Regarding the second issue, Omoto (2012, 2019) (Fig. 2, site 3) and Araoka et al. (2013)
755 (Fig. 2, site 2) carefully selected samples for dating. They limited measurements to the boulders
756 that meet tsunami boulder identification criteria proposed by Goto et al. (2010b, 2013) and that
757 were composed of single massive corals (i.e. *Porites* sp., Fig. 3B). Based on this approach,
758 Araoka et al. (2013) estimated a tsunami recurrence interval of about 150-400 years on the
759 Sakishima Islands. It should be noted, however, that they measured ages of various sizes of
760 boulders so it is uncertain whether the paleotsunamis that cast the boulders ashore were large or
761 small.

762 There are some studies that focus on sandy tsunami deposits from the Sakishima Islands.
763 Possible sandy tsunami deposits, especially relating to 1771 CE, were previously reported by
764 archaeologists (e.g., Yamamoto, 2008; Kugai, 2011; Nakaza, 2017) and coastal engineers
765 (Nakaza et al., 2013) following the work done by Kawana and Nakata (1994) but their detail
766 history had remained uncertain. Recently, Ando et al. (2018) performed a 120 m-long trench
767 survey and found that paleotsunamis may have occurred at approximately 600-year intervals
768 over the last 2000 years (Fig. 2, site 1). This observation is consistent with numerical modeling
769 of the movement of a very large boulder at southeastern Ishigaki Island by Hisamatsu et al.
770 (2014) who suggested at least 2 large tsunamis equivalent to or even larger than the 1771
771 tsunami occurred after 2000 years ago but before 1771 CE. Interestingly, the recurrence interval
772 of paleotsunamis estimated from sandy tsunami deposits (Ando et al., 2018; Kitamura et al.,

2018a, b) is longer than that estimated from boulders (Araoka et al., 2013). This discrepancy may be explained by coastal boulders being more sensitive in recording tsunamis because even small tsunamis can transport small boulders to the coast. Therefore, evidence of small to medium tsunamis are included in the estimation of tsunami recurrence intervals from boulders. Meanwhile, sandy tsunami deposits that extended far inland may only be suitable to recognize large events. Alternatively, this discrepancy might also reflect the small number of studies carried out on sandy deposits in this region so further research is required. Although some researchers have suggested the absence of a predecessor to the 1771 CE tsunami (Nakaza et al., 2013, 2015), geological evidence of both boulders and sandy deposits provides crucial evidence that either small or large tsunamis have repeatedly occurred over the last few thousand years around the Sakishima Islands.

784

785 **4. When and where: the history of great earthquakes**

786 This section summarizes the spatial and temporal distribution of paleotsunami evidence
787 along the Nankai and Ryukyu subduction zones. Table 1 summarizes when the historical
788 earthquakes occurred in these regions.

789

790 **4.1. Nankai subduction zone**

791 Great earthquakes occurred in the Tōkai region nine times in total, including the
792 controversial 1614 CE event, from the end of the 7th century until 1944 CE. On the other hand,
793 great earthquakes occurred in the Nankai region eight times from 684 to 1946 CE. The 1099 CE
794 earthquake was recorded with an incorrect date, and may have actually occurred in 1096 CE
795 (Ishibashi, 2016). The Tōkai and Nankai earthquakes always occur in pairs within a few days to
796 several years, except for the 1498 CE Meiō earthquake. A Nankai earthquake paired with the
797 1498 CE Meiō Tōkai earthquake is likely to have occurred, but we have not been able to
798 definitively ascribe any of the multiple candidates from the historical record. The 1707 CE Hōei
799 and 887 CE Nin-na earthquakes were full-length ruptures of both the Tōkai and Nankai fault
800 segments.

801 Some questions remain about the recurrence pattern of Nankai Trough earthquakes. First,
802 the recurrence intervals of earthquakes in the Tōkai and Nankai regions appears to be different
803 before and after the 1361 CE earthquake (Table 1). Even if the 1614 CE earthquake did exist, it
804 seems unsuitable to include it within a series of great Nankai Trough earthquakes because the
805 seismic intensity and tsunami during the 1614 CE event were much smaller than other
806 earthquakes. If this earthquake is included, the recurrence interval from the 1361 CE earthquake
807 to the 1854 CE earthquake were 137 years (1361 - 1498 CE), 116 years (1498 - 1614 CE), 93
808 years (1614 - 1707 CE), and 147 years (1707 - 1854 CE). On the other hand, before 1361CE,

809 recurrence intervals were much longer; 203 years (684 - 887CE), 209 years (887 - 1096 CE),
810 and 265 years (1361 - 1096 CE). It is unlikely that the process of plate boundary earthquakes
811 suddenly changed at some point and, while it is possible that the apparent change is an artifact
812 of the small sample number, other explanations may also be considered.

813 It is possible that other great earthquakes may be missing from the documented record due
814 to a range of circumstances (Koyama, 1999). However, if M8-class earthquakes repeated at
815 intervals of ~100 years before 1361 CE, due to the large Medieval population, it would be
816 surprising if no historical documents recorded their occurrence. Reflecting the sensitivity with
817 which historical records from this period document earthquakes, in the period between 1200 and
818 1260 CE, there are over 100 earthquakes detailed in documents from Kyoto (Ishibashi, 2014).
819 Nevertheless, liquefaction features suggest the occurrence of a Tōkai earthquake between 1096
820 and 1361 CE and evidence for a Nankai earthquake between 1099 and 1361 CE has been found
821 from archaeological sites in southern Osaka and the southern Kii Peninsula (Sangawa, 2007).
822 Further searches for tsunami deposits are necessary to identify whether there are additional
823 historical Nankai Trough earthquakes other than those shown in Table 1.

824

825 **4. 2. Ryukyu subduction zone**

826 In contrast to the Nankai Trough region, there seems to be little evidence of large
827 earthquakes ($M_w > 8$) and tsunamis from the Hyūga-nada or the northern and central Ryukyu
828 Trench regions over the last few thousand years. However, more data needs to be collected,
829 especially from the Hyūga-nada region.

830 Large tsunamis have repeatedly occurred around the southern Ryukyu Islands. If
831 submarine landslides have contributed to enhancing tsunami magnitudes (Miyazawa et al.,
832 2012; Okamura et al., 2018), it is difficult to estimate earthquake magnitudes. However, it is
833 probably reasonable to infer from the paleotsunami record that $M_w > 8$ earthquakes probably
834 occurred, with some potentially triggering submarine landslides (Hisamatsu et al., 2014).

835

836 **5. Variation in size and recurrence time**

837 The size of earthquakes and tsunamis occurring on the Nankai and Ryukyu subduction
838 zones shows a wide variation. However, lesser tsunamis may not leave a trace in the geological
839 record. Of course, the creation and preservation potential of tsunami deposits can change
840 depending on the geological and geomorphological conditions of the place where the tsunami
841 hit. For example, intervals between 10 tsunami deposits left in the Shima Lowlands (Fig. 2, site
842 17) between 4500 and 500 years ago show variations from 100 to 600 years (Fujino et al., 2018),
843 which are longer than expected from the historical Nankai Trough record (Fig. 2). The lack of
844 intervening tsunami deposits may simply be due to the lesser size of those tsunamis. Only the

845 larger tsunamis that reached the study site with sufficient stream power would be recorded in
846 the sediment record.

847 In that sense, it is interesting that traces of the 1361 CE tsunami were not found between
848 the 1096 and 1498 CE tsunami deposits in the Ota River Lowland, where a careful investigation
849 was conducted by Fujiwara et al. (2020) on a large outcrop over a length of 1 km (see chapter
850 3.1.2). This probably indicates that the 1361 CE tsunami was too small to inundate the study
851 area, or that, if the tsunami reached the area, its stream power was too weak to form an
852 identifiable tsunami deposit.

853 To enhance the contribution of the geological record, it is necessary to expand the search
854 for paleotsunami deposits to sediments that have not been targeted so far. For example, the 2011
855 Tōhoku tsunami widely left fine-grained deposits, mainly clay beds, inland from the area where
856 sandy deposits are distributed (e.g., Goto et al., 2011; Fujiwara, 2015). If the mud supply is
857 sufficient, such a mud layer may be left behind after the tsunami. However, muddy
858 paleotsunami deposits are likely to have been overlooked in previous studies. Development of
859 criteria and techniques for identifying such muddy tsunami deposits in the geological record
860 will help to expand the range of available sites.

861 Analyses of seawater components (Chagué-Goff et al., 2017) and marine biomarkers in the
862 sediment (e.g., Shinozaki et al., 2015) are among several promising new ways to solve this
863 problem. While efforts should be made to develop approaches for identifying tsunami deposits,
864 geological methods alone have their limitations. For example, identification of the difference
865 between a tsunami and a storm surge by sediment transport modeling is also expected
866 (Watanabe et al., 2018). Although many studies have performed paleotsunami surveys on
867 low-lying plains or coastal lakes, it is important to expand the survey area to the other regions
868 such as narrow valleys (e.g., Fujiwara, 2015; Abe et al., 2020) in order to increase number of
869 potential sites in the region.

870 There are only a few "treasure troves" where many tsunami deposits are recorded. Methods
871 for accurately dating the sediments and estimating the tsunami size from such records may have
872 not been sufficiently developed. Recent advances in coring, non-destructive testing technology
873 (Falvard and Paris, 2017, Paris et al., 2020), chemical analysis methods (e.g. Chagué-Goff et al.,
874 2017) and geophysical sensing techniques (e.g., Obrocki et al., 2020) will drive the
875 development of more complete tsunami records from such key sites. Advances in radiocarbon
876 dating and calibration techniques will enable more accurate estimation of the occurrence timing
877 of great earthquakes and tsunamis (Ishizawa et al., 2020).

878 Progress in methods for quantifying the tsunami size from deposits are needed to elucidate
879 the variations in earthquake size. Most lowlands along the Nankai Trough coast are small, with
880 mountains and hills close to the coast. In such a situation, tsunamis reach the landward margin

881 of the lowland irrespective of their size, and it is not possible to compare the differences in
882 inundation distances. The tsunami height could, however, be estimated using sediment transport
883 modeling, using parameters such as grain size, the thickness of tsunami deposits, and the height
884 of coastal barrier (cf. Baranes et al., 2016). Tsunami boulders may be a useful tool for
885 estimating local tsunami size especially in the Ryukyu Islands (e.g., Watanabe et al., 2016).
886 However, the future improvement of models and development of methodologies to collect field
887 data about the boulder source and shape are required for further high accuracy modeling.

888

889 **6. How big?**

890 As the 2004 Sumatra-Andaman and 2011 Tōhoku earthquakes demonstrated, an
891 earthquake much larger than any in the region's instrumental or documentary history may occur
892 in the future. Paleoseismological evidence shows that these extraordinary events have occurred
893 repeatedly along subduction zones around the world, including the Kuril Trench (e.g.,
894 Nanayama et al., 2003), northern Japan Trench (e.g., Sawai et al., 2012; Sawai, in press),
895 Peru-Chile Trench (e.g. Cisternas et al., 2005), and Sunda-Java Trench (e.g., Malik et al., 2011).
896 These giant earthquakes have much longer (~300-500 years or longer) recurrence intervals than
897 those of great earthquakes (M8-class) that have occurred in each subduction zone. According to
898 the available paleotsunami evidence, the Nankai and Ryukyu subduction zones may not be
899 exceptions.

900

901 **6.1. Nankai subduction zone**

902 An updated fault model for the 1707 CE Hōei earthquake based on crustal deformation
903 inferred from the GPS network and evidence of tsunami deposits at Ryujin Pond (Fig. 2, site 7)
904 suggest that its source area extended 70 km further to the west of the 1854 earthquake
905 (Furumura et al., 2011). Okamura and Matsuoka (2012) suggested that earthquakes of similar
906 magnitude to the 1707 event could occur every 300-500 years along this subduction zone. The
907 number of tsunami deposits varies from site to site and they are generally less frequent than
908 expected from the frequency of historical earthquakes. This may be the result of only larger
909 tsunamis overwashing the coastal barriers and leaving deposits in back-barrier ponds. The 684,
910 1361 and 1707 CE tsunamis selectively recorded at the Ryujin, Ryuuō, and Kaniga-ike Pond
911 (Fig. 2, sites 7, 8 and 10) may therefore be relatively larger than other historical tsunamis
912 (Okamura and Matsuoka, 2012; Baranes et al., 2016). If larger tsunamis result from larger
913 earthquakes, this indicates that larger earthquakes with 300 - 500-year recurrence intervals and
914 smaller earthquakes with shorter recurrence intervals coexist in the Nankai Trough region,
915 suggesting the recurrence pattern of this region is not periodic, but may still be time predictable
916 or slip predictable (Fig. 5A).

917 Okamura and Matsuoka (2012) also pointed out that the historical period of 1300 years
918 may not be enough to consider the maximum-size earthquake and tsunami of this region. The
919 discovery of tsunami deposits thicker than those from the tsunami that followed the 1707 CE
920 Hōei earthquake, the largest in the Nankai region's history, from Tadasu-ike (Fig. 2, site 9),
921 Kaniga-ike (Fig. 2, site 10), Kamoda-oike (Fig.2, site 13) and Sugari-oike (Fig. 2, site 15), all of
922 which were aged around 2000 years ago, led Okamura and Matsuoka (2012) to hypothesize the
923 occurrence of a giant tsunami. However, it is unreasonable to discuss the size of a tsunami
924 based on the thickness of tsunami deposits alone. The thickness of different tsunami deposits
925 within a site may be influenced by the state of the tide and the time of year, which controls
926 vegetation cover and therefore sediment transport. Furthermore, changes in coastal
927 geomorphology over time may change a site's susceptibility to accumulating and preserving
928 tsunami deposits (Garrett et al., 2016). Underdeveloped low dunes and beach ridges can allow
929 large amounts of overflow, even during lesser tsunamis, resulting in the formation of thicker
930 tsunami deposits in back-barrier ponds (Fujiwara, 2015). Additionally, the estimated ages for
931 the thick tsunami bed in each of Okamura and Matsuoka's (2012) ponds has substantial
932 uncertainties of hundreds of years. Consequently, it is ambiguous whether the thick tsunami bed
933 in each of the ponds was formed by one extremely large tsunami or series of closely spaced
934 large tsunamis.

935 Deposition of a thick tsunami deposits around 2000 years ago is not seen at Nankoku (Fig.
936 2, site 11) (Tanigawa et al., 2018) or Ryuuō Pond (Fig. 2, site 8) (Baranes et al., 2016). In
937 addition, at Ryūjin Pond in northern Kyushu, a sand bed deposited around 3300 years ago is the
938 thickest, with only a minor sand bed deposited around 2000 years ago. Thick tsunami deposits
939 formed around 2000 years ago have not been found from the Tōkai region (Fujiwara, 2013;
940 Fujiwara and Tanigawa, 2017). The late 7th century tsunami deposits in the Otagawa Lowland
941 is thick, over 40 cm more than 2 km inland from the coast at the time of tsunami occurrence, but
942 its distribution is limited to the topographic lows in the lowland (Fujiwara et al., 2020).
943 Fujiwara et al. (2020) interpreted that the tsunami was not so large and inundation and sand bed
944 deposition was limited to the topographic depressions.

945 Kitamura (2016) summarized the spatio-temporal distribution of reported tsunami
946 deposits from the eastern coast of the Nankai Trough, including sites 23, 24, 26, 28, 29, and 32
947 in Figure 2, and concluded that no tsunami deposits with a wide regional distribution that might
948 suggest the occurrence of "level 2" tsunami over the past 4000 years have been found in this
949 area. However, due to the limitation in the age control of the tsunami deposits, it would be
950 difficult to deny the existence of a giant tsunami on this basis alone.

951

952 **6.2. Hyūga-nada and the Ryukyu subduction zone**

953 Historical and geological evidence is too scarce to evaluate whether large tsunamigenic
954 earthquakes have been generated in the Hyūga-nada region. Historical records suggest no large
955 earthquakes with magnitude >8 have occurred over the last 400 years in this region. Geological
956 evidence from the Pacific coast of central and southern Kyushu is probably still insufficient to
957 discuss paleoseismicity in this region, although recent works suggest the rare occurrence of
958 tsunamigenic earthquakes (Ichihara et al., 2015; Yamada et al., 2020).

959 M8-class earthquakes can be generated elsewhere along the Ryukyu Trench based on
960 historical and instrumental records; however, the Ryukyu Trench is generally considered as an
961 aseismic zone in terms of the potential occurrence of giant earthquakes of magnitude ~9,
962 although potential coupling of the plate boundary and the consequent occurrence of extremely
963 large earthquakes along the Ryukyu Trench has been suspected by some researchers (e.g., Ando
964 et al., 2009, 2012a; Aydan and Tokashiki, 2018; Hsu et al., 2012). Geological evidence shows
965 that tsunamigenic earthquakes may be rare or even nonexistent in the northern and central
966 Ryukyu Trench (Goto et al., 2013). Even though Kikai Island in the northern Ryukyu Trench
967 has been uplifted quickly and the repeated occurrence of large earthquakes has been suspected
968 (e.g., Nakata et al., 1978), relatively small earthquakes just beneath the island (Goto, 2017) or
969 stable uplift without large earthquakes (Shikakura, 2014) may also explain this record.

970 Nakamura and Sunagawa (2015) studied the distribution of very low frequency
971 earthquakes (VLFs) along entire the Ryukyu Trench. They suggested that occurrence of
972 VLFs is high in the northern and central Ryukyu Trench but low in the southern Ryukyu
973 Trench. Therefore, it is likely that the southern Ryukyu Trench has potential to generate large
974 earthquakes, unlike the northern and central regions (Nakamura and Sunagawa, 2015).
975 Nishimura (2014) also suggested that short-term slow slip events (SSEs) cluster near the areas
976 where large earthquakes have occurred historically (e.g., the 1911 and 1771 CE events).
977 Nevertheless, the sources of these large earthquakes do not overlap with the clusters of
978 short-term SSEs and the reason remains uncertain.

979 Arai et al. (2016) and Nakamura (2017) suggested there is a strongly coupled zone at
980 shallower plate boundary depths in the southern Ryukyu Trench. Therefore, localized
981 tsunamigenic earthquakes at the southern Ryukyu Trench inferred from tsunami boulders (Goto
982 et al., 2013) may be explained by the heterogeneity of the coupling along the Ryukyu Trench.
983 Nevertheless, the source model of the 1771 CE tsunami is still controversial with proposed
984 models including an intra-plate earthquake plus a submarine landslide (Imamura et al., 2008;
985 Miyazawa et al., 2012), a tsunami earthquake (Nakamura, 2009), splay faulting (Hsu et al.,
986 2013), and a large submarine landslide near the trench axis (Okamura et al., 2018). By assuming
987 a large subduction zone earthquake along the Ryukyu Trench, Ando et al. (2018) estimated a
988 seismic coupling coefficient of 20%. Alternatively, if a submarine landslide played an important

989 role in the generation of the 1771 CE tsunami and predecessors, then a special tectonic state
990 capable of generating large earthquakes only at the southern Ryukyu Trench may not need to be
991 considered. Relationships among VLFs, short-term SSEs, and large earthquakes ($M_w \sim 8$) are
992 still uncertain, so further seismological research with reference to paleotsunami results is
993 required along the Ryukyu Trench.

994

995 **7. Knowledge gaps and future work**

996

997 **7.1. Earthquake recurrence model**

998 Since Kanamori and McNally (1982) explained a series of different size earthquakes off
999 the Colombian coast as the rupture of different combinations of multiple asperities, similar
1000 phenomena have been reported from subduction zones around the world. Some examples show
1001 that giant (M_9 -class) and large (M_{7-8} class) earthquakes may share fault plane asperities, and
1002 the former have much longer recurrence intervals than the latter (Cisternas et al., 2005, 2017).
1003 The Nankai and Ryukyu subduction zones may be no exception. This may indicate that the
1004 recurrence pattern of subduction zone earthquakes is “nonpredictable” (Fig.5A).

1005 The recurrence of Nankai Trough earthquakes has generally been explained as the repeated
1006 rupture of all or part of characteristic fault planes (segments) A, B, C, D and E (Fig. 1 and Table
1007 1). Observations that the amount of uplift during the 1707, 1854, and 1946 CE Nankai
1008 earthquakes at Cape Muroto was proportional to the length of the interval preceding each
1009 earthquake led to the proposal of the time-predictable recurrence model (Fig. 5A) by Shimazaki
1010 and Nakata (1980). Kumagai (1996) applied this model to Nankai Trough earthquakes using the
1011 time series of great earthquakes since 684 CE. However, according to current knowledge,
1012 Kumagai’s series of historical earthquakes contains some uncertainties, and the reliability of the
1013 recurrence model is ambiguous. For example, the 1230 CE earthquake and the 1498 CE Meiō
1014 earthquake in the Nankai region (as opposed to the undisputed Tōkai event) in his series are still
1015 undetermined. Results of paleotsunami research in the Otagawa Lowland (Fujiwara et al., 2020)
1016 does not support the time-predictable recurrence model. According to this study, the 1361 CE
1017 Tōkai earthquake has the longest pre-earthquake interval (265 years following the 1096 CE
1018 earthquake), but its tsunami seems to have been less extensive than others, suggesting a smaller
1019 magnitude of earthquake (see chapter 5).

1020 Seno (2012) proposed a new earthquake recurrence model for Nankai Trough earthquakes
1021 that challenged the traditional idea of the characteristic earthquake model, which partitions the
1022 Nankai Trough seismogenic zone into ruptures A-E and assign the epicenters of historical
1023 earthquakes to them. He assumed two complementary earthquakes that have different
1024 non-overlapping asperities mainly based on the spatial distribution of seismic intensities, while

1025 also considering tsunami heights and coastal deformation estimated from documented records
1026 and liquefaction features in archaeological sites (Fig. 5B). According to his idea, historical great
1027 Nankai Trough earthquakes are classified into two types; “Hōei-type” earthquakes (887, 1361,
1028 1707 and 1944+1946 CE) and the “Ansei-type” earthquakes (684, 1096+1099, 1489,
1029 1854+1854 CE), which alternately occur with recurrence intervals of around 350 and 400 years,
1030 respectively. Seno’s (2012) model seems to explain the variations in the recurrence intervals
1031 and extent of source area of each earthquake in Nankai Trough; however, the reliability of
1032 model is reduced by the decrease in the number of reliable documentary records with increasing
1033 age, especially before the 1096 CE earthquake.

1034 In contrast to the Nankai Trough region, the establishment of an earthquake recurrence
1035 model along the Ryukyu Trench is not straightforward because of limited historical and
1036 geological data. Presently available data suggest that earthquakes with magnitudes of around 8
1037 have occurred patchily in space and time from the northern to the central Ryukyu Trench. The
1038 repeated occurrence of these earthquakes is uncertain. On the other hand, in the southern
1039 Ryukyu Trench, the repeated occurrence of large tsunamis is suggested (Omoto, 2012; Araoka
1040 et al., 2013; Ando et al., 2018). Therefore, the future occurrence of large earthquakes in this
1041 region is likely.

1042 The amount of reported paleo-tsunami data is still insufficient to update the earthquake
1043 recurrence model in these regions (Figure 2). Updating the model also requires improved
1044 accuracy in the dating and the identification of tsunami deposits, which is a particularly
1045 important issue with sites surveyed in the early stages of the development of the field.
1046 Re-examination of these sites with current knowledge and technology, as discussed in chapter 5,
1047 may reveal some rules in the recurrence mode of the earthquakes in these regions.

1048 Coastal deformation data will enhance the revision of earthquake recurrence models (e.g.,
1049 Garrett et al, 2016). In particular, uplift along the western Suruga Bay coast, including Cape
1050 Omaezaki (representing the rupture of segment E), and the southern Kii Peninsula (representing
1051 the rupture of segment C) is key to proving the hypothesis of Ansei- and Hōei-type earthquakes
1052 (Fig. 5B). Kitamura et al. (2019) suggested a possible seismic uplift occurred around 400 CE
1053 from the Shimizu Plain (Fig. 2, site 29).

1054 Integration of offshore paleoseismic features, such as turbidite sequences, with onshore
1055 tsunami deposits also will help us to consider which recurrence model is plausible. In fact, the
1056 integrated study of coastal deformation and onshore tsunami deposits, and offshore
1057 seismo-turbidite sequences in the Cascadia subduction zone and northern San Andreas fault
1058 systems has revealed the recurrence mode of great earthquakes in this region (e.g., Goldfinger et
1059 al., 2003 a, b). These studies also contribute to regional earthquake and tsunami
1060 countermeasures through underpinning tsunami simulations (e.g., Priest et al., 2017). Although

1061 there are some initial studies on seismo-turbidites in the Nankai (Garrett et al., 2016 and
1062 references therein) and the Ryukyu subduction zone (e.g., Ujiie et. al. 1997; Ikehara et al., 2017),
1063 at present, coordination of onshore and offshore paleoseismic research is an area where future
1064 progress is expected.

1065

1066 **7.2. Maximum possible earthquake and tsunami**

1067 As a response to the occurrence of unexpectedly huge 2011 Tōhoku earthquake, the
1068 Cabinet Office (2011b) attempted to quantify the maximum possible earthquake and tsunami for
1069 the Nankai Trough subduction zone (“maximum scenario” in Fig. 1). The earthquake source
1070 consists of the strongly coupled zone of 10 to 20 km depth, which has been considered to be the
1071 main seismogenic zone, with the addition of down-dip weakly coupled zones at ~30 to 40 km
1072 depth, which are the location of the occurrence of SSE and deep low-frequency tremor. Its
1073 western end is delimited by the Kyushu-Palau ridge. The up-dip zone from 10 km to the trough
1074 axis, which may also generate large tsunamis, is assumed as an additional tsunami source area.
1075 According to the worst scenario by the Cabinet Office (2012a), the height of the tsunami
1076 occurring from this M9 earthquake could exceed 10 m over a wide area along the Pacific coast
1077 of western Japan, and reach in excess of 30 m in some places. However, no geological evidence,
1078 such as tsunami deposits, was taken into account for this catastrophic earthquake and tsunami
1079 scenario (e.g., Wallis et al., 2018). Determining when the most recent outside earthquake and
1080 tsunami occurred and how large it was will help to address this concern.

1081

1082 **Largest tsunami in the area estimated from geological data**

1083 The “2 ka giant tsunami hypothesis” by Okamura and Matsuoka (2012), derived from the
1084 discovery of extremely thick tsunami deposits, played an important role in making people aware
1085 of the possibility of a giant earthquake and tsunami in the Nankai Trough region and raising
1086 awareness of disaster management. Nevertheless, it has not been tested. First, it is necessary to
1087 ensure that multiple seismic events are not confused with a single major event. To solve this
1088 issue, the extremely thick tsunami deposits should be re-examined by increasing the number of
1089 reliable radiocarbon ages and using new age modeling methods (e.g., Bronk Ramsey, 2009;
1090 Lougheed and Obrochta, 2016, 2019).

1091 Other causes of the thick tsunami deposit in individual ponds should also be considered.
1092 Earthquake-induced submarine landslides are a candidate source for locally thick tsunami
1093 deposits. They generate much larger, but regionally limited, tsunamis than would be expected
1094 based on the size of concurrent earthquakes (e.g., Tappin et al., 2008; Harbitz et al., 2013;
1095 Kawamura et al., 2017). In fact, many submarine landslides have been found on the submarine
1096 slopes along the Nankai Trough and some of them have areas of $\geq 10 \text{ km}^2$ (Moriki et al., 2017).

1097 The study of the evolutionary history of both coastal and submarine landforms may be the key
1098 to solving this problem.

1099 Reconstruction of the inundation area is the most important indicator for determining the
1100 size of a tsunami from its deposit (e.g., Fujiwara, 2015). In the case of eastern Hokkaido, the
1101 discovery of prehistoric tsunami deposits showing inundation areas several times larger than
1102 those of historical M8-class earthquakes revealed that unusually large tsunamis repeatedly
1103 occurred on the Kuril Trench (e.g., Nanayama et al., 2003). Similarly, the 869 CE Jōgan
1104 tsunami deposit showed much larger inundation areas than those of other historical earthquakes
1105 that occurred in the northern Japan Trench, suggesting an unprecedentedly large earthquake
1106 (Satake et al., 2008; Namegaya et al., 2010, Sawai et al., 2012).

1107 The Hamamatsu Plain (Fig. 2, site 24) is the first candidate site where we can suggest that
1108 no giant tsunami has occurred in the eastern Nankai Trough over the latter half of the Holocene
1109 based on the tsunami inundation area shown by the tsunami deposits. The strand plain has
1110 prograded seaward over the last 6000 years following the Holocene sea-level high stand (e.g.,
1111 Sato et al., 2011, 2016a), as evidenced by preservation of rows of sandy beach ridges. This
1112 progradation process can be seen in the vertical change of diatom assemblages in swale deposits,
1113 which shows a change from brackish-marine to freshwater conditions (Fig. 4B). In harmony
1114 with the progradation of the coast, older washover deposits, probably including tsunami
1115 deposits, are distributed inland and younger ones are found in more seaward locations (Fig. 4 B,
1116 C). Additionally, at each site, younger tsunami deposits tend to be thinner (Fig. 4 B, C).
1117 Assuming tsunamis (or other washover events) of similar size over time, the progradation of the
1118 coast would make it less likely that each successive tsunami would reach the inland areas, and
1119 the observed sequence of tsunami deposits would be left on the Hamamatsu Plain (Fig. 6). If an
1120 extremely large tsunami occurred over the past few thousand years, its deposit would have
1121 extended from the coast to further inland, as shown at the top of the Fig. 6. However, no such
1122 trace has yet been found (Fujiwara, 2013; Fujiwara and Tanigawa, 2017).

1123 Along the Ryukyu subduction zone, the 1771 CE tsunami at the Sakishima Islands is the
1124 largest known event. Based on numerical modeling of boulder transport Hisamatsu et al. (2014)
1125 suggested that at least one prehistoric tsunami event was even larger than the 1771 CE tsunami.
1126 However, since information is limited, further geological data is required to clarify the size of
1127 prehistoric tsunamis.

1128

1129 **Crustal movements suggesting outsized earthquakes along the Nankai Trough**

1130 In discussing the possibility of a giant earthquake, it is necessary to assess a wide range of
1131 crustal deformation patterns occurring at different temporal and spatial scales. In the case of
1132 eastern Hokkaido, where the recurrence of giant tsunami along the Kuril Trench is shown by

1133 tsunami deposits, there is a contradiction between the coastal deformation pattern shown by
1134 geological and geodetic data. In this area, 60 to 80 km above the top of the subducting plate,
1135 marine terraces showing 20 to 50 m of net uplift in the past 125 ka (Okumura, 1996) coexist
1136 with tide gauge and bench mark-inferred subsidence of up to 1 m in the past 100 years
1137 (Shimazaki, 1974a; Ozawa et al., 1997). Little, if any, uplift is known from the M8-class
1138 earthquakes in this region's written history during the past 200 years (Shimazaki, 1974b;
1139 Kasahara, 1975; Kasahara and Kato, 1980; Yamanaka and Kikuchi, 2003).

1140 Micropaleontological research on marsh deposits in this area (Sawai et al., 2004) revealed
1141 post-seismic uplift associated with unusually large earthquakes, which was induced by greater
1142 creep further down the plate boundary than during any of the Kuril subduction zone's historical
1143 events. The last uplift event of up to 2 m, which might resolve some of the contradiction
1144 between geologic and geodetic data, occurred over the first few decades following an outsized
1145 17th-century earthquake.

1146 A similar phenomenon can be seen around Lake Hamana. Middle Pleistocene (MIS 9 to
1147 MIS 5) terraces, mainly of marine and partly of fan origin, are found in this area (Sugiyama,
1148 1991; Nakashima et al., 2008), showing that uplift has been dominant over 10^4 to 10^5 -year
1149 scales. The MIS-9 marine terrace reaches 78 m in elevation and is highest on the west coast of
1150 Lake Hamana, decreasing in height to the north and west. Subsidence of 0.1-0.6 m during the
1151 1854 CE Ansei-Tōkai earthquake is estimated around the Lake Hamana area (Ishibashi, 1981;
1152 Sato and Fujiwara, 2017). The northern coast of the lake is also known to have subsided during
1153 the 1707 CE Hōei earthquake (Yata, 2013). Coseismic subsidence between 7000 and 5700 years
1154 ago, similar to the 1707 and 1854 CE events, was reported from Shinjo Lowland (Fig. 2, site 21)
1155 on the west coast of Lake Hamana (Sato and Fujiwara, 2017). Multiple active reverse faults
1156 (The research group for active submarine faults off Tokai, 1999) and normal faults (Arai et al.,
1157 2006) parallel to the trough axis are distributed off the Hamana area. However, they cannot
1158 explain the coastal deformation in the area.

1159 Major differences between this area and eastern Hokkaido include the inter-seismic uplift
1160 after the 1854 CE Ansei-Tōkai earthquake and the short-term uplift associated with SSEs.
1161 Inter-seismic uplift, averaging ~6 mm/y between 1901 and 2008 CE (Geographical Survey
1162 Institute, 2008), may compensate for the coseismic subsidence during "ordinal" Tōkai
1163 earthquakes; however, other factors are needed to explain the formation of high and tilted
1164 terraces. This average uplift rate includes the effect of SSEs (see chapter 2.2). SSEs seem to
1165 occur with decadal recurrence intervals and are high frequency uplift events on a geological
1166 time scale, but associated uplift seems too small to form the terraces. For example, a Tōkai slow
1167 slip event (Ozawa et al., 2002; Miyazaki et al., 2006) occurred under Lake Hamana from
1168 January 2000 to July 2005, with a cumulative moment magnitude of ~7.1, and generated ~5 cm

1169 uplift in this area (Suito and Ozawa, 2009). At present the importance of SSEs for terrace
1170 formation remains unclear and their contribution may need to be considered further.

1171 To elucidate the possibility of a giant earthquake, a unified explanation is required for
1172 geodetic and geologic phenomena on different timescales, including SSEs with 10^0 to 10^1 -year
1173 scales, inter-seismic uplift with a 10^2 -year scale, and tilting uplift of the Pleistocene terraces
1174 over 10^4 to 10^5 -year scales. This requires further investigation of Holocene crustal movements
1175 on a 10^3 -year scale, which connects the two.

1176 The possibility of eastward rupture propagation from the Nankai Trough to the
1177 Fujikawa-kako fault zone (FKZ) must also be assessed in the context of defining the maximum
1178 possible earthquake size. The FKZ is an eastern extension of the Nankai or Suruga Trough and
1179 might rupture in combination with future great Nankai Trough earthquakes (HERP, 2010).
1180 Nevertheless, the recurrence history is still unclear at present. Marsh deposits on the
1181 Ukishima-gahara Lowlands (Fig. 2, site 30) record a 1500-year history of repeated coastal
1182 subsidence suggesting the activity of the FKZ (Fujiwara et al., 2016). Further research is needed
1183 to elucidate the relationship between episodes of coastal subsidence and historical earthquakes.

1184

1185 **Possibility of full-length rupturing event of the Nankai and Ryukyu subduction zone**

1186 The Nankai and the Ryukyu subduction zones make up one plate boundary through the
1187 Hyūga-nada region, and both may have the potential to generate M9-class earthquakes.
1188 Therefore, it is important to understand the possible risk of the occurrence of extremely large
1189 earthquakes crossing this boundary, as suggested by some researchers (e.g., Furumoto and Ando,
1190 2009). As reviewed in this paper, the occurrence of such an extremely large earthquake and
1191 tsunami in the past few thousand years is not supported by the presently available historical and
1192 geological data in this region. However, the spatio-temporal correlation of earthquakes that
1193 occurred on the Nankai and Ryukyu subduction zones or the possibility of a full-length
1194 mega-earthquake linking both seismogenic zones has not been evaluated well. From this point
1195 of view, research integrating various scientific fields such as seismology, paleoseismology,
1196 geology, and geodesy is also an issue for the future.

1197

1198 **8. Conclusions**

1199 This review illustrates the difference in the type of tsunami deposits and the progress of
1200 research between the Nankai and Ryukyu subduction zones. We also summarize the recurrence
1201 mode of great earthquakes and tsunamis in these regions based on geological evidence. The
1202 volume of information currently available about paleoearthquakes and tsunamis varies greatly
1203 different between the two subduction zones, despite the fact they comprise a contiguous
1204 northwestern boundary of the subducting Philippine Sea Plate beneath the Japanese archipelago.

1205 Information from the Ryukyu subduction zone is very limited compared to the Nankai
1206 subduction zone.

1207 The model describing the recurrence mode of great earthquakes in this region is still
1208 incomplete, and updating the model requires future studies to gather more paleoseismic data
1209 from both on land and offshore. Reconstruction of the largest event ever in the region based on
1210 geological evidence will help discussions about the “real” size of the maximum possible
1211 earthquake in this region. This requires a new conceptual model that can unify geological
1212 processes that occur on various time scales. Differentiation of extremely large earthquakes and
1213 piecemeal ruptures of lesser magnitude in the geological record is also important for the
1214 accurate reconstruction of past events and corresponding forecasts of future earthquake.
1215 Progress in age estimation techniques will help to diminish this enduring problem. The potential
1216 of giant earthquakes that rupture both the Nankai and Ryukyu subduction zones remains
1217 unknown. Paleoseismological studies in Hyūga-nada region that connects the two subduction
1218 zones are key to solving this question.

1219

1220

1221 **Acknowledgement**

1222 The part of the review concerning Nankai subduction zone paleotsunami deposits led by
1223 OF and EG was conducted as contribution to the QuakeRecNankai project supported by the
1224 Belgian Science Policy Office (BELSPO BRAIN-be BR/121/A2).

1225

1226

1227 **References**

1228

1229 Abe, T., Goto, K. and Sugawara, D., 2020. Spatial distribution and sources of tsunami deposits
1230 in a narrow valley setting - insight from 2011 Tohoku-oki tsunami deposits in northeastern
1231 Japan. *Progress in Earth and Planetary Science*. doi.org/10.1186/s40645-019-0318-6.

1232 Ando, M., 1975a. Possibility of a major earthquake in the Tokai district, Japan and its
1233 pre-estimated seismotectonic effects. *Tectonophysics* 25, 69-85.
1234 doi.org/10.1016/0040-1951(75)90011-6.

1235 Ando, M., 1975b. Source mechanisms and tectonic significance of historical earthquakes along
1236 the Nankai Trough, Japan. *Tectonophysics* 27, 119-140.
1237 doi.org/10.1016/0040-1951(75)90102-X.

1238 Ando, M., Kitamura, A., Tu, Y., Ohashi, Y., Imai, T., Nakamura, M., Ikuta, R., Miyairi Y.,
1239 Yokoyama, Y. and Shishikura, M., 2018. Source of high tsunamis along the southernmost
1240 Ryukyu trench inferred from tsunami stratigraphy. *Tectonophysics* 722, 265–276.
1241 doi.org/10.1016/j.tecto.2017.11.007.

1242 Ando, M. and Nakamura, M., 2013. Seismological evidence for a tsunami earthquake recorded
1243 four centuries ago on historical documents. *Geophysical Journal International* 195, 1088–
1244 1101. doi.org/10.1093/gji/ggt270

1245 Ando, M., Nakamura, M., Matsumoto, T., Furukawa, M., Tadokoro, K. and Furumoto, M., 2009.
1246 Is the Ryukyu subduction zone in Japan coupled or decoupled? -The necessity of seafloor
1247 crustal deformation observation, *Earth Planets Space* 61, 1031-1039, doi:
1248 10.1186/Bf03352954.

1249 Ando, M., Tu, Y., Kumagai, H., Yamanaka, Y. and Lin C. H., 2012, Very low frequency
1250 earthquakes along the Ryukyu subduction zone. *Geophysical Research Letters* 39,
1251 L04303. doi.org/10.1029/2011GL050559.

1252 Ando, R., Takeda, N. and Yamashita, T., 2012b. Propagation dynamics of seismic and aseismic
1253 slip governed by fault heterogeneity and Newtonian rheology. *Journal of Geophysical*
1254 *Research* 117, B11308. doi:10.1029/2012jb009532.

1255 Arai, K., Okamura, Y., Ikehara, K., Ashi, J., Soh, W., Kinoshita, M., 2006. Active faults and
1256 tectonics on the upper forearc slope off Hamamatsu City, central Japan. *Journal of the*
1257 *Geological Society of Japan* 112, 749-759. (in Japanese with English abstract). doi:
1258 https://doi.org/10.5575/geosoc.112.749.

1259 Arai, R., Takahashi, T., Kodaira, S., Kaiho, Y., Nakanishi, A., Fujie, G., Nakamura, Y.,
1260 Yamamoto, Y., Ishihara, Y., Miura, S. and Kaneda, Y., 2016. Structure of the tsunamigenic
1261 plate boundary and low-frequency earthquakes in the southern Ryukyu Trench. *Nature*
1262 *Communications*. 7, 12255. DOI: 10.1038/ncomms12255.

1263 Araoka, D., Yokoyama, Y., Suzuki, A., Goto, K., Miyagi, K., Miyazawa, K., Matsuzaki, H. and
1264 Kawahata, H., 2013. Tsunami recurrence revealed by Porites coral boulders in the southern
1265 Ryukyu Islands, Japan. *Geology* 41, 919–922. doi.org/10.1130/G34415.1.

1266 Awata, Y. and Sugiyama, Y., 1989. Geological structure due to the right–lateral reverse faulting
1267 accompanied with the large earthquakes along the Nankai Trough). *Zishin (Journal of the*
1268 *Seismological Society of Japan)*, 2nd ser., **42**, 231–233.

1269 Aydan, O. and Tokashiki, N., 2018. Tsunami boulders and their implications on the potential for
1270 a mega-earthquake along the Ryukyu Archipelago, Japan. *Bulletin of Engineering Geology*
1271 *and the Environment* 78, 3917-3925. doi.org/10.1007/s10064-018-1378-3.

1272 Baba, T., Cummins, P.R. Hori, T. and Kaneda, Y., 2006. High precision slip distribution of the
1273 1944 Tonankai earthquake inferred from tsunami waveforms: Possible slip on a splay fault.
1274 *Tectonophysics* 426, 119–134. doi:10.1016/j.tecto.2006.02.015

1275 Baranes, H.E., Woodruff, J.D., Wallace, D.J., Kanamaru, K. and Cook, T.L., 2016.
1276 Sedimentological records of the C.E. 1707 Hōei Nankai Trough tsunami in the Bungo
1277 Channel, southwestern Japan. *Natural Hazards* 84, 1185-1205. DOI
1278 10.1007/s11069-016-2498-3.

1279 Bronk Ramsey, C., 1995. Radiocarbon calibration and analysis of stratigraphy; the OxCal
1280 program. *Radiocarbon* 37, 425–430.

1281 Bronk Ramsey, C., 2008. Deposition models for chronological records. *Quat. Sci. Rev.* 27, 42–
1282 60.

1283 Bronk Ramsey, C., 2009. [Bayesian analysis of radiocarbon dates](#). *Radiocarbon* 51, 337–360.

1284 Cabinet Office, Government of Japan, 2011a. Information provided by members at the second
1285 meeting of the Committee for evaluating the rupture model of megathrust earthquakes
1286 along the Nankai Trough. <http://www.bousai.go.jp/jishin/nankai/model/2/pdf/1.pdf>.

1287 Cabinet Office, Government of Japan, 2011b. Interim report of the Committee for evaluating
1288 the rupture model of megathrust earthquakes along the Nankai Trough.
1289 http://www.bousai.go.jp/jishin/nankai/model/pdf/chukan_matome.pdf (in Japanese, last
1290 accessed 2020/2/21)

1291 Cabinet Office, Government of Japan, 2012a. Second report of the Committee for evaluating the
1292 rupture model of megathrust earthquakes along the Nankai Trough: tsunami fault moles.
1293 http://www.bousai.go.jp/jishin/nankai/model/pdf/20120829_2nd_report01.pdf. (in
1294 Japanese, last accessed 2020/2/11)

1295 Cabinet Office, Government of Japan, 2012b. Second report of the Committee for evaluating
1296 the rupture model of megathrust earthquakes along the Nankai Trough: tsunami fault
1297 moles. Reference: Basic concept for tsunamis with high frequency of occurrence in the

1298 Nankai Trough. http://www.bousai.go.jp/jishin/nankai/model/pdf/20120829_2nd_report04.pdf
1299 pdf (in Japanese, last accessed 2020/2/11).

1300 CDMC (Central Disaster Management Council), 2011. Report of the committee for technical
1301 investigation on countermeasures for earthquakes and tsunamis based on the lessons
1302 learned from the “2011 off the Pacific coast of Tohoku Earthquake”
1303 <http://www.bousai.go.jp/kaigirep/chousakai/tohokukyokun/pdf/Report.pdf> (last accessed
1304 2020/2/11).

1305 Chagué-Goff, C., Szczuciński, W. and Shinozaki, T. 2017. Applications of geochemistry in
1306 tsunami research: A review. *Earth-Science Reviews* 165, 203-244.
1307 <https://doi.org/10.1016/j.earscirev.2016.12.003>

1308 Chida, N., and Nakaue, K., 2007. Time series analysis of the tsunamis caused by Hiei and
1309 Ansei Nankai earthquakes in Yonouzu, Oita prefecture, east central Kyushu, Japan. The
1310 research bulletin of the Faculty of Education and Welfare Science, Oita University, 29, 69–
1311 80.

1312 Cisternas, M., Atwater, B.F, Torrejon, F., Sawai, Y., Machuca, G., Lagos, M., Eipert, A.,
1313 Youlton, C., Salgado, I., Kamataki, T., Shishikura, M., Rajendran, C.P., Malik, J.K., Rizal,
1314 Y. and Husni, M., 2005. Predecessors of the giant 1960 Chile earthquake. *Nature* 437:
1315 404–407. DOI: 10.1038/nature03943

1316 Cisternas, M., Garrett, E., Wesson, R., Dura, T., Ely, L.L., 2017. Unusual geologic evidence of
1317 coeval seismic shaking and tsunamis shows variability in earthquake size and recurrence in
1318 the area of the giant 1960 Chile earthquake. *Marine Geology* 385, 101–113.

1319 Falvard, S. and Paris, R., 2017. X-ray tomography of tsunami deposits: Towards a new
1320 depositional model of tsunami deposits. *Sedimentology* 64, 453–477. doi:
1321 10.1111/sed.12310.

1322 Fujino, S., Kimura, H., Komatsubara, J., Matsumoto, D., Namegaya, Y., Sawai, Y., Shishikura,
1323 M., 2018. Stratigraphic evidence of historical and prehistoric tsunamis on the Pacific coast
1324 of central Japan: Implications for the variable recurrence of tsunamis in the Nankai Trough.
1325 *Quaternary Science Reviews* 201, 147–161.
1326 <https://doi.org/10.1016/j.quascirev.2018.09.026>.

1327 Fujiwara, O., 2013. Earthquakes and tsunamis along the Nankai Trough, inferred from geology
1328 and geomorphology: examples in Tokai Region., *GSJ Chishitsu News* 2, 197-200 (in
1329 Japanese)

1330 Fujiwara, O., 2015. The science of tsunami deposits. University of Tokyo Press, 283p (in
1331 Japanese).

1332 Fujiwara, O., Aoshima, A., Irizuki, T., Ono, E., Obrochta, S.P., Sampei, Y., Sato, Y. and
1333 Takahashi, A., 2020. Tsunami deposits refine great earthquake rupture extent and

1334 recurrence over the past 1300 years along the Nankai and Tokai fault segments of the
1335 Nankai Trough, Japan. *Quaternary Science Reviews* 277,
1336 doi.org/10.1016/j.quascirev.2019.105999.

1337 Fujiwara, O., Fujino, S., Komatsubara, J., Morita, Y. and Namegaya, Y. (2016) Paleoeological
1338 evidence for coastal subsidence during five great earthquakes in the past 1500 years along
1339 the northern onshore continuation of the Nankai subduction zone. *Quaternary International*
1340 397, 523-540. doi.org/10.1016/j.quaint.2015.11.014.

1341 Fujiwara, O. Hirakawa, K., Irizuki, T, Hasegawa, S., Hase, Y., Uchida, J., Abe, K., 2010.
1342 Millennium-scale recurrent uplift inferred from beach deposits bordering the eastern
1343 Nankai Trough, Omaezaki area, central Japan. *Island Arc* 19, 374–388.
1344 <https://doi.org/10.1111/j.1440-1738.2010.00729.x>

1345 Fujiwara, O., Ono, E., Satake, K., Sawai, Y., Umitsu, M., Yata, T., Abe, K., Ikeda, T., Okamura,
1346 Y., Sato, Y., Aung, T.T., Uchida, J., 2007. Trace of the AD1707 Hiei earthquake from the
1347 coastal lowland, Shizuoka Prefecture, central Japan. *Annual Report on Active Fault and*
1348 *Paleoearthquake Researches*, Geological Survey of Japan, AIST, No. 7, 157–171 (in
1349 Japanese with English abstract).

1350 Fujiwara, O., Sato, Y., Ono, E. and Umitsu, M. 2013a. Researches on tsunami deposits using
1351 sediment cores: 3.4 ka tsunami deposit in the Rokken-gawa Lowland near Lake Hamana,
1352 Pacific coast of central Japan. *Journal of Geography (Chigaku Zasshi)* 122, 308-322. (in
1353 Japanese with English abstract). doi:10.5026/jgeography.122.308

1354 Fujiwara, O., Ono, E., Yata, T., Umitsu, M., Sato, Y. and Heyvaert, V.M.A., 2013b. Assessing
1355 the impact of 1498 Meio earthquake and tsunami along the Enshu-nada coast, central Japan
1356 using coastal geology. *Quaternary International* 308, 4–12.
1357 <https://doi.org/10.1016/j.quaint.2012.12.009>.

1358 Fujiwara, O. and Tanigawa, K., 2017. Researches on the tsunami deposits along the Nankai
1359 Trough: Their achievements and upcoming challenges. *Journal of the Geological Society*
1360 *of Japan* 123, 831-842. (in Japanese with English abstract).
1361 doi.org/10.5575/geosoc.2017.0061

1362 Furumoto, M. and Ando, M. 2009. Possibility of super large earthquake from southwest Japan
1363 to Ryukyu Islands*, *Chikyu Monthly* 31, 197–202, in Japanese.

1364 Furumura, T., Imai, K. and Maeda, T., 2011. A revised tsunami source model for the 1707 Hiei
1365 earthquake and simulation of tsunami inundation of Ryujin Lake, Kyushu, Japan. *Journal*
1366 *of Geophysical Research* 116, B02308. doi:10.1029/2010JB007918, 2011

1367 Garrett, E., Fujiwara, O., Garrett, P., Heyvaert, V.M.A., Shishikura, M., Yokoyama, Y.,
1368 Hubert-Ferrari, A., Brückner, H., Nakamura, A., De Batist, M., the QuakeRecNankai team,
1369 2016. A systematic review of geological evidence for Holocene earthquakes and tsunamis

1370 along the Nankai-Suruga Trough, Japan. *Earth-Science Reviews* 159, 337–357
1371 <https://doi.org/10.1016/j.earscirev.2016.06.011>.

1372 Garrett, E., Fujiwara, O., Riedesel, S., Deforce, W. J., Yokoyama, Y., Schmidt, S., Brückner, H.,
1373 De Batist, M., Heyvaert, V.M.A., the QuakeRecNankai team., 2018. Historical
1374 Nankai-Suruga megathrust earthquakes recorded by tsunami and terrestrial mass
1375 movement deposits on the Shirasuka coastal Lowlands, Shizuoka Prefecture, Japan. *The*
1376 *Holocene* 28, 968–983 <https://doi.org/10.1177/0959683617752844>.

1377 Geographical Survey Institute, 1952. Re-survey of the south-western part of Japan after the
1378 great Nankaido earthquake of 1946. *Bulletin of the Geographical Survey Institute* 3,
1379 31-118.

1380 Geographical Survey Institute, 2008. Crustal Movements in the Tokai District. Report of the
1381 Coordinating Committee for Earthquake Prediction, Japan 79, 80p. (in Japanese).

1382 Goldfinger, C., Nelson, C.H., Johnson, J.E., and the Shipboard Scientific Party, 2003a.
1383 Holocene earthquake records from the Cascadia subduction zone and northern San Andreas
1384 Fault based on precise dating of offshore turbidites. *Annual Review of Earth and Planetary*
1385 *Sciences* 31, 555-577, doi: 10.1146/annurev.earth.31.100901.141246.

1386 Goldfinger, C., Nelson, C.H., Johnson, J.E., and the Shipboard Scientific Party, 2003b,
1387 Deep-water turbidites as Holocene earthquake proxies: the Cascadia subduction zone and
1388 Northern San Andreas Fault systems. *Annals OF Geophysics* 46, 1169-1194. DOI:
1389 <https://doi.org/10.4401/ag-3452>

1390 Goto, K., 2017. Paleotsunami researches along the Ryukyu Trench. *Journal of the Geological*
1391 *Society of Japan* 123, 843–855 (in Japanese with English abstract)
1392 doi.org/10.5575/geosoc.2017.0046

1393 Goto, K., Chagué-Goff, C., Fujino, S., Goff, J., J., Jaffe, B. Nishimura, Y., Richmond, B.,
1394 Sugawara, D., Szczuciński, W., Tappin, D.R., Witter, R.C. and Yulianto, E., 2011. New
1395 insights of tsunami hazard from the 2011 Tohoku-oki event. *Marine Geology* 290, 46-50.
1396 doi.org/10.1016/j.margeo.2011.10.004

1397 Goto, K., Ishizawa, T., Ebina, Y., Imamura, F., Sato, S., Udo, K., submitted. Ten years after the
1398 2011 Tohoku-oki tsunami: Aftermath of environmental and social effects. *Earth-Science*
1399 *Reviews*.

1400 Goto, K., Kawana, T. and Imamura, F., 2010a. Historical and geological evidences of boulders
1401 deposited by tsunamis, southern Ryukyu Islands, Japan. *Earth-Science Reviews* 102, 77–
1402 99. doi.org/10.1016/j.earscirev.2010.06.005.

1403 Goto, K., Miyagi, K., Kawamata, H. and Imamura, F., 2010b. Discrimination of boulders
1404 deposited by tsunamis and storm waves at Ishigaki Island, Japan. *Marine Geology* 269, 34–
1405 45. doi.org/10.1016/j.margeo.2009.12.004.

1406 Goto, K., Miyagi, K. and Imamura, F., 2013. Localized tsunamigenic earthquakes inferred from
1407 preferential distribution of coastal boulders on Ryukyu Islands, Japan. *Geology* 41, 1139–
1408 1142.

1409 Goto, K., Okada, K. and Imamura, F., 2009. Characteristics and hydrodynamics of boulders
1410 transported by storm waves at Kudaka Island, Japan. *Marine Geology* 262, 14–24.
1411 doi.org/10.1016/j.margeo.2009.03.001.

1412 Harada, T., Ishibashi, K., Satake, K., 2013. Tsunami numerical simulation for hypothetical
1413 giant/great earthquakes along the Izu-Bonin Trench. Programme and Abstracts, the
1414 Seismological Society of Japan, 2013 Fall Meeting, 186 (in Japanese).

1415 Harbitz, C. B., Løvholt, F. and Bungum, H., 2013. Submarine landslide tsunamis: how extreme
1416 and how likely? *Natural Hazards* 72, 1341-1374. DOI 10.1007/s11069-013-0681-3.

1417 Hatori, T., 1974. Sources of large tsunamis in southwest Japan. *Zishin* (Journal of the
1418 Seismological Society of Japan), 2nd ser., 27, 10-24. (in Japanese with English abstract).

1419 Hatori, T., 1975. Sources of large tsunamis generated in the Boso, Tokai and Nankai regions in
1420 1498 and 1605. *Bulletin of the Earthquake Research Institute, University of Tokyo* 50,
1421 171-185 (in Japanese with English abstract).

1422 Hatori, T., 1976. Documents of tsunami and crustal deformation in Tokai district associated
1423 with the Ansei Earthquake of Dec. 23, 1854. *Bulletin of the Earthquake Research Institute,*
1424 *University of Tokyo* 51, 13–28. (in Japanese with English abstract).

1425 Hatori, T., 1985. Field investigation of historical tsunamis along the east coast of Kyushu, west
1426 Japan. *Bulletin of the Earthquake Research Institute, University of Tokyo* 60, 439–459. (in
1427 Japanese).

1428 Heki, K., and Kataoka, T., 2008. On the biannually repeating slow-slip events at the Ryukyu
1429 Trench, southwestern Japan. *Journal of Geophysical Research* 113(B11).
1430 doi:10.1029/2008JB005739.

1431 HERP (Headhunters for Earthquake Research Promotion), 2004. Long-term evaluation of
1432 seismic activity around the Hyuga Nada and the Nansei-shoto Trench.
1433 https://www.jishin.go.jp/main/chousa/kaikou_pdf/hyuganada.pdf (last accessed
1434 2020/3/12).

1435 HERP, 2010. Partial revision of long-term evaluation of Fujikawa Estuary fault zone.
1436 http://www.jishin.go.jp/main/chousa/10oct_fujikawa/ (In Japanese, last accessed 19 Feb,
1437 2020.).

1438 HERP, 2013. Evaluation of long-term earthquake activity along the Nankai Trough. 2nd edn.,
1439 HERP, Japan. https://www.jishin.go.jp/main/chousa/kaikou_pdf/nankai_2.pdf (in Japanese,
1440 last acceded 10 February 2020).

1441 Hirose, H., and Obara, K., 2005. Repeating short- and long-term slow slip events with deep
1442 tremor activity around the Bungo channel region, southwest Japan, *Earth Planets Space* 57,
1443 961-972.

1444 Hisamatsu, A., Goto, K. and Imamura, F., 2014. Local paleo-tsunami size evaluation using
1445 numerical modeling for boulder transport at Ishigaki Island, Japan. *Episodes* 37, 265–276.
1446 DOI: 10.18814/epiiugs/2014/v37i4/006

1447 Hongo, C., 2010. Sea-level standstill and dominant hermatypic coral from the Holocene raised
1448 reef terraces at the Kikai Island, Ryukyu Islands. *Journal of Geography* 119, 860–871. (In
1449 Japanese with English abstract)

1450 Hsu, Y. J., Ando, M., Yu, S. B. and Simons, M., 2012. The potential for a great earthquake
1451 along the southernmost Ryukyu subduction zone. *Geophysical Research Letters* 39,
1452 L14302. doi.org/10.1029/2012GL052764

1453 Hsu, S. K., Yeh, Y. C., Sibuet, J. C., Doo, W. B. and Tsai, C. H., 2013. A mega-splay fault
1454 system and tsunami hazard in the southern Ryukyu subduction zone. *Earth and Planetary
1455 Science Letters* 362, 99–107. doi.org/10.1016/j.epsl.2012.11.053.

1456 Ichihara, N., Kuroki, A. and Ijyuuin H., 2015. Tsunami event deposits in Miyazaki Plain.
1457 Abstract of 122nd Annual Meeting of Geological Society of Japan.
1458 https://doi.org/10.14863/geosocabst.2015.0_188 (in Japanese).

1459 Igarashi, T., 2010. Spatial changes of inter-plate coupling inferred from sequences of small
1460 repeating earthquakes in Japan. *Geophysical Research Letters* 37, L20304. doi:
1461 10.1029/2010GL044609.

1462 Ikeda, M., 1977. Coastal terraces and their deformations in Amami-Oshima, Ryukyu Islands,
1463 Southern Japan. *Journal of Geography (Chigaku Zasshi)* 86, 43-49. (in Japanese with
1464 English abstract).

1465 Ikehara, K., Kanamatsu, T., Usami, K. and Hsiung, K.H., 2017. Depositional processes and
1466 recurrence of seismo-turbidites along the southern Ryukyu Trench. EGU General
1467 Assembly Conference Abstracts, 2017 EGUGA.19.34511.

1468 Imamura, F., Goto, K. and Ohkubo, S., 2008. A numerical model for the transport of a boulder
1469 by tsunami. *Journal of Geophysical Research, Oceans* 113, C01008.
1470 doi.org/10.1029/2007JC004170.

1471 Ishibashi, K., 1976. Re-examination of a great earthquake expected in the Takaki district,
1472 central Japan: Possibility of the “Suruga Bay earthquake”. Abstract of the Seismological
1473 Society of Japan, 1976, No.2, 30–34 (In Japanese).

1474 Ishibashi, K., 1981. Specification of a soon-to-occur seismic faulting in the Tokai district,
1475 central Japan, based upon seismotectonics. *Earthquake Prediction: An international review*
1476 4, Maurice Ewing Series, 297–332. <https://doi.org/10.1029/ME004p0297>

- 1477 Ishibashi, K., 1998. Inferred Tokai earthquake paired with the 1361 Shohei Nankai earthquake.
1478 Programme and abstracts, the Seismological Society of Japan, 1988 Fall Meeting, 125 (in
1479 Japanese). <https://historical.seismology.jp/ishibashi/archive/1361Ko-anTokai98.pdf>
- 1480 Ishibashi, K., 1999. Great Tokai and Nankai, Japan, earthquakes as revealed by historical
1481 seismology: 1. Review of the events until the mid-14th century. *Journal of Geography*, 108,
1482 399–423 (in Japanese with English abstract).
- 1483 Ishibashi, K., 2014. Nankai Trough great earthquake, History, Science and Society. Iwanami
1484 Shoten, Publishers, 205p (in Japanese).
- 1485 Ishibashi, K., 2016. Possibility that the 1099 Jotoku (Kowa) Nankai earthquake is unreal and the
1486 1096 Kaho (Eicho) earthquake was an “entire-Nankai Trough” earthquake: examination of
1487 *Kansenji-an*, a historical document describing an earthquake in Shikoku. *Rekishi-jisihin*
1488 (Historical Earthquakes) 31, 81–88(in Japanese with English abstract).
- 1489 Ishibashi, K. and Satake, K., 1998. Problems on forecasting great earthquakes in the subduction
1490 zones arund Japan by means of paleoseismology. *Zisin (Journal of the Seismological*
1491 *Society of Japan)*, 2nd ser. 50, 1–21 (in Japanese with English abstract).
- 1492 Ishizawa, T., Goto, K., Yokoyama, Y. and Goff, J., 2020. Dating tsunami deposits: Present
1493 knowledge and challenges. *Earth-Science Reviews*, 200, 102971.
1494 doi.org/10.1016/j.earscirev.2019.102971.
- 1495 Ishiuka, O., Taylor, R.N., Yuasa, M. and Ohara, Y., 2011., Making and breaking an island arc:
1496 A new perspective from the Oligocene Kyushu–Palau arc, Philippine Sea. *Geochemistry,*
1497 *Geophysics, Geosystems* 12, <https://doi.org/10.1029/2010GC00344>
- 1498 Ito, T., and Hashimoto, M., 2004. Spatiotemporal distribution of interplate coupling in southwest
1499 Japan from inversion of geodetic data. *Journal of Geophysical Research* 109, B02315,
1500 [doi:10.1029/2002JB002358](https://doi.org/10.1029/2002JB002358).
- 1501 Iwai, K. and Kawana, T., 2008. Physiographical studies on locations of archaeological sites in
1502 Okinawa Island, the Ryukyus, Japan, during past several thousand years. *Okinawa J. Geog.*
1503 *Stud.* 8, 25–36. (in Japanese).
- 1504 Japan coast guard, 1948. Research report of the 1946 Nankai earthquake -part of ground
1505 deformation and damage-. *Hydrographic bulletin* 201, 139 p. (in Japanese)
- 1506 Kanamori H, McNally K., 1982. Variable rupture mode of the subduction zone along the
1507 Ecuador-Colombia coast. *Bulletin of Seismological Society of America* 72, 1241–1253.
- 1508 Kasahara, K., 1975. Aseismic Faulting Following the 1973 Nemuro-oki Earthquake, Hokkaido,
1509 Japan (a Possibility), *Pure and applied Geophysics* 113, 127–139.
1510 [doi:10.1007/BF01592905](https://doi.org/10.1007/BF01592905)

1511 Kasahara, K. and Kato, T. (1980) Aseismic faulting following the 1973 Nemuro-oki earthquake,
1512 Hokkaido, Japan (a retrospective study). *Pure and Applied Geophysics* 119, 392-403. Doi:
1513 10.1007/BF00877774.

1514 Kato, Y., 1993. Geological structure and topography of the Amami Plateau. Report of
1515 Hydrographic Researches 29, 51-63 (in Japanese with English abstract).

1516 Kato, Y. and Kimura, M., 1983. Age and origin of so-called “tsunami-ishi”, Ishigaki Island,
1517 Okinawa Prefecture. *Journal of the Geological Society of Japan* 89, 471–474. (in
1518 Japanese).

1519 Kawamura, K., Kanamatsu, T. and Yamada, Y., 2017. Submarine slides and marine geohazards:
1520 the study results so far and current problems. *Journal of Geological Society of Japan* 123,
1521 999-1014. (in Japanese with English abstract).

1522 Kawana, T., 1989. Quaternary crustal movement in the Ryukyu Islands. *Earth Mon.* 11, 618–
1523 630 (in Japanese, original title translated).

1524 Kawana, T., 2006. Invasion of about 3400 cal BP large wave in the southeastern Okinawa
1525 Island and the surroundings, the Ryukyus, Japan, as deduced from coralline deposits. *Bull.*
1526 *Faculty Educ. Univ. Ryukyus* 68, 265–271. (in Japanese).

1527 Kawana, T. and Nakata, T., 1994. Timing of late Holocene tsunamis originated around the
1528 southern Ryukyu Islands, Japan, deduced from coralline tsunami deposits. *Journal of*
1529 *Geography (Chigaku Zasshi)* 103, 352–376. (in Japanese with English abstract).

1530 Kawana, T. and Nakata, K., 2003. Calibrated calendar ages of the paleotsunamis attacked
1531 eastern part of the Kasari Peninsula of Amami Island and the northern part of Kikai Island,
1532 the north Ryukyus, Japan. *Okinawa Geog. Soc. Bull.* 6, 33–40. (in Japanese).

1533 Kitamura, A., 2016. Examination of the largest-possible tsunamis (Level 2) generated along the
1534 Nankai and Suruga troughs during the past 4000 years based on studies of tsunami deposits
1535 from the 2011 Tohoku-oki tsunami. *Progress in Earth and Planetary Science* 3:12. DOI
1536 10.1186/s40645-016-0092-7.

1537 Kitamura, A., Fujiwara, O., Shinohara, K., Akaike, S., Masuda, T., Ogura, K., Urano, Y.,
1538 Kobayashi, K., Tamaki, C. and Mori, H., 2013. Identifying possible tsunami deposits on
1539 the Shizuoka Plain, Japan and their correlation with earthquake activity over the past 4000
1540 years. *The Holocene* 23, 1684–1698. doi.org/10.1177/0959683613505345.

1541 Kitamura, A., Ina, T., Suzuki, D., Tsutahara, K., Sugawara, D., Yamada, K. and Aoshima, A.,
1542 2019. Geologic evidence for coseismic uplift at ~AD 400 in coastal lowland deposits on
1543 the Shimizu Plain, central Japan. *Progress in Earth and Planetary Science* 6,
1544 Article number 57. Doi:10.1186/s40645-019-0305-y

1545 Kitamura, A., Ito, M., Sakai, S., Yokoyama, Y. and Miyairi, Y., 2018a. Identification of tsunami
1546 deposits using a combination of radiometric dating and oxygen-isotope profiles of

1547 articulated bivalves. *Marine Geology* 403, 57–61.
1548 <https://doi.org/10.1016/j.margeo.2018.04.003>.

1549 Kitamura, A., Ito, M., Ikuta, R. and Ikeda, M., 2018b. Using molluscan assemblages from
1550 paleotsunami deposits to evaluate the influence of topography on the magnitude of late
1551 Holocene mega-tsunamis on Ishigaki Island, Japan. *Prog. Earth Planet. Sci.* 5, 41. DOI:
1552 10.1186/s40645-018-0200-y.

1553 Kitamura, A. and Kobayashi, K., 2014a. Geologic record of middle-late Holocene
1554 paleotsunamis and paleo-earthquakes on the Shizuoka Plain and coastal lowland of the
1555 southern Izu Peninsula, central Japan. *Journal of Geography (Chigaku Zasshi)* 122, 813–
1556 834 (in Japanese with English abstract).

1557 Kitamura, A. and Kobayashi, K., 2014b. Geologic evidence for prehistoric tsunamis and
1558 coseismic uplift during the AD 1854 Ansei-Tokai earthquake in Holocene sediments on the
1559 Shimizu Plain, central Japan. *The Holocene* 24, 814–827. Doi:
1560 [10.1177/0959683614530447](https://doi.org/10.1177/0959683614530447)

1561 Kitamura, A., Ohashi, Y., Miyairi, Y., Yokoyama, Y. and Yamaguchi, T., 2014. Discovery of a
1562 tsunami boulder along the coast of Shimoda, Shizuoka, central Japan. *The Quaternary*
1563 *Research (Daiyonki-Kenkyu)* 53, 259–264 (In Japanese with English abstract).

1564 Kitamura, A., Seki, Y., Kitamura, Y., Haga, T., 2018c. The discovery of emerged boring
1565 bivalves at Cape Omaezaki, Shizuoka, Japan: Evidence for the 1361 CE Tokai earthquake
1566 along the Nankai Trough. *Marine Geology* 405, 114–119.
1567 <https://doi.org/10.1016/j.margeo.2018.08.006>

1568 Koike, K. and Machida, H. eds., 2001. Atlas of Quaternary marine terraces in the Japanese
1569 islands. University of Tokyo Press, 105pp. with 3 CD-ROMs and 2 maps. (in Japanese).

1570 Komatsubara, J. and Fujiwara, O. (2007) Overview of Holocene tsunami deposits along the
1571 Nankai, Suruga, and Sagami Troughs, southwest Japan. *Pure and Applied Geophysics*, 164,
1572 493-507.

1573 Komatsubara, J. Fujiwara, O., Takada, K., Sawai, Y., Than T. Aung and Kamataki, T. (2008)
1574 Historical tsunamis and storms recorded in a coastal lowland, Shizuoka prefecture, along
1575 the Pacific Coast of Japan. *Sedimentology*, 55, 1703-1716.

1576 Koyama, M., 1999. A review of historical seismology in Japan: analyses and proposals for next
1577 stage of study of historical documents. *Journal of Geography* 108, 346-369. (in Japanese
1578 with English abstract) https://doi.org/10.5026/jgeography.108.4_346

1579 Kugai, M., 2011. Discovery of the foraminifera sedimentation at “Tomori-motojima” and
1580 “Sunagawa-motojima”*. *Bull. Miyakojima City Mus.**, 15, 65–75. (in Japanese).

1581 Kumagai, H., 1996. Time sequence and the recurrence models for large earthquakes along the
1582 Nankai Trough revisited. *Geophysical Research Letters*, 23, 1139-1142.
1583 <https://doi.org/10.1029/96GL01037>

1584 Lay, T., and Kanamori, H., 1981. An asperity model of great earthquake sequences, in
1585 *Earthquake prediction - An international review*, AGU Geophys. Mono., edited by D. W.
1586 Simpson and P. G. Richards, pp. 579-592, Amer. Geophys. Union, Washington, D.C.

1587 Lougheed, B.C., Obrochta, S.P., 2016. MatCal: open source bayesian ¹⁴C age calibration in
1588 MatLab. *J. Open Res. Software* 4. <http://doi.org/10.5334/jors.130>.

1589 Lougheed, B.C. and Obrochta, S.P., 2019. A Rapid, deterministic age-depth modelling routine
1590 for geological sequences with inherent depth uncertainty: *Paleoceanography and*
1591 *Paleoclimatology* 34, 122-133. doi.org/10.1029/2018PA003457

1592 Loveless, J. P., and Meade, B. J., 2010. Geodetic imaging of plate motions, slip rates, and
1593 partitioning of deformation in Japan. *Journal of Geophysical Research* 115, B02410.
1594 doi.org/10.1029/2008JB006248

1595 Loveless, J. P., and Meade, B. J., 2011. Spatial correlation of interseismic coupling and
1596 coseismic rupture extent of the 2011 Mw=9.0 Tohoku-oki earthquake. *Geophysical*
1597 *Research Letters* 38, L17306, doi:10.1029/2011GL048561.

1598 Makino, K., 1968. Meiwa tsunami at Yaeyama*. Published by author, 447p. (in Japanese).

1599 Makino, K., 1981. Meiwa tsunami at Yaeyama -revision and enlargement-*. Published by
1600 author, 462p. (in Japanese).

1601 Malik, J.N., Shishikura, M., Echigo, T., Ikeda, Y., Satake, K., Kayane, H., Sawai, Y., Murty,
1602 C.V.R. and Dikshit, O. 2011. Geologic evidence for two pre-2004 earthquakes during
1603 recent centuries near Port Blair, South Andaman Island, India. *Geology* 39, 559-562. DOI:
1604 10.1130/G31707.1

1605 Matsu'ura, R. S., in press, The cause of the tsunami in 1791 at the main island of Okinawa.
1606 *Historical Earthquakes* (in Japanese).

1607 Miyazaki, S. and Heki, K., 2001. Crustal velocity field of southwest Japan: Subduction and
1608 arc-arc collision. *Journal of Geophysical Research* 106(B3), 4305-4326.
1609 doi.org/10.1029/2000JB900312.

1610 Minamidate, K., Goto, K., Watanabe, M., Roeber, V., Toguchi, K., Sannoh, M., Nakashima, Y.,
1611 Kan, H., 2020. Millennial scale maximum intensities of typhoon and storm wave in the
1612 northwestern Pacific Ocean inferred from storm deposited reef boulders. *Scientific reports*.
1613 10, 7218. <https://doi.org/10.1038/s41598-020-64100-6>.

1614 Miyazaki, S., Segall, P., McGuire, J.J., Kato, T. and Hatanaka, Y., 2006. Spatial and temporal
1615 evolution of stress and slip rate during the 2000 Tokai slow earthquake. *Journal of*
1616 *Geophysical Research* 111, B03409.doi:10.1029/2004JB003426.

1617 Miyazawa, K., Goto K., and Imamura, F., 2012. Re-evaluation of the 1771 Meiwa Tsunami
1618 source model, southern Ryukyu Islands, Japan. *In* Submarine Mass Movements and Their
1619 Consequences, *Adv. Nat. Tech. Hazards Res.* 31, 497-506. DOI:
1620 10.1007/978-94-007-2162-3_44.

1621 Moore, G.F., Bangs, N.L., Taira, A., Mumamoto, S., Pangborn, E. and Tobin, H.J., 2007.
1622 Three-dimensional splay fault geometry and implication for tsunami generation. *Science*
1623 318, 1128-1131. DOI: 10.1126/science.1147195

1624 Moriki, H., Kumamoto, T., Nakata, T., Goto, H., Izumi, N. and Nishizawa, A., 2017.
1625 Identification of landslide and its characteristics on the seafloor around Japan using
1626 anaglyph images. *Report of Hydrographic and Oceanographic Researches No.54*, 1-16. (in
1627 Japanese with English abstract).

1628 Murakami, H., Shimada, T., Itoh, S., Yamamoto, N. and Ishizuka, J., 1996. Reexamination of
1629 the heights of the 1605, 1707 and 1854 Nankai tsunamis along the coast of Shikoku island.
1630 *Journal of Japan Society for Natural Disaster Science* 15, 39-52 (In Japanese with English
1631 abstract)

1632 Nadeau, R. M. and Johnson, L. R., 1998. Seismological studies at Parkfield VI: Moment release
1633 rates and estimates of source parameters for small repeating earthquakes. *Bulletin of*
1634 *Seismological Society of America* 88(3), 790-814.

1635 Nagaoka, S., Maemoku, H. and Matsushima, Y., 1991. Evolution of Holocene coastal landforms
1636 in the Miyazaki Plain, southern Japan. *The Quaternary Reach (Daiyonki Kenkyu)* 30,
1637 59-78 (in Japanese with English abstract).

1638 Nakamura, M., 2009. Fault model of the 1771 Yaeyama earthquake along the Ryukyu Trench
1639 estimated from the devastating tsunami. *Geophysical Research Letters* 36, L19307,
1640 doi:10.1029/2009GL039730.

1641 Nakamura, M., 2017. Distribution of low frequency earthquakes accompanying the very low
1642 frequency earthquakes along the Ryukyu Trench. *Earth Planets. Space* 69, 49, DOI
1643 10.1186/s40623-017-0632-4.

1644 Nakamura, M. and Kinjou, A., 2013. Fault model of the 1768 and 1791 tsunamis in the central
1645 Ryukyu. Abstract for the 2013 AGU Fall Meeting, T13E-2574.

1646 Nakamura, M. and Sunagawa, N., 2015. Activation of very low frequency earthquakes by slow
1647 slip events in the Ryukyu Trench. *Geophysical Research Letters* 42, 1076–1082.
1648 <https://doi.org/10.1002/2014GL062929>.

1649 Nakashima, R., Hori, N., Miyazaki, K. and Nishioka, Y., 2008. Geology of the Toyohashi and
1650 Tahara districts. *Quadrangle Series, 1: 50,000, Geological Survey of Japan, AIST*, 113 p. (in
1651 Japanese with English abstract 3 p.)

- 1652 Nakata, T. and Shimazaki, K., 1997. Geo-slicer, a newly invented soil sampler, for
1653 high-resolution active fault studies. *Journal of Geography*, 106, 59-69. (in Japanese with
1654 English abstract). doi.org/10.5026/jgeography.106.59.
- 1655 Nakata, T., Takahashi, T. and Koba, M., 1978. Holocene-emerged coral reefs and sea-level
1656 changes in the Ryukyu Islands. *Geographical Review of Japan* 51, 87-108. (in Japanese
1657 with English abstract).
- 1658 Nakaza, H., 2017. What is Shiraho-Saonetabaru Cave Site? Outline of survey and main results.
1659 *Kagaku* 87, 532–536. (in Japanese).
- 1660 Nakaza, E., Iribe, T., Tokuhisa, R., Miyazato, N., Inagaki, K. and Savou, R., 2013. Prehistorical
1661 and historical tsunamis of Ryukyu Islands estimated through tsunami depositions. *Journal*
1662 *of the Japan Society of Civil Engineering, Ser. B3 (Ocean Eng.)* 69, 515–520. (in Japanese
1663 with English abstract).
- 1664 Nakaza, E., Tokuyama, R., Inagaki, K., 2015. Original locations of huge boulders moved by
1665 tsunami and its generation mechanism. *Journal of the Japan Society of Civil Engineers, Ser.*
1666 *B2 (Coastal Eng.)*, 71, 193–198. (in Japanese with English abstr).
1667 doi.org/10.2208/kaigan.71.I_193
- 1668 Nishimura, T., 2014. Short-term slow slip events along the Ryukyu Trench, southwestern Japan,
1669 observed by continuous GNSS. *Progress in Earth and Planetary Science* 1, 22.
1670 <http://www.proearthplanetsci.com/content/1/1/22>.
- 1671 Namegaya, Y., Maemoku, H., Shishikura, M., Echigo, T. and Nagai, A., 2011. Factors causing
1672 scattered boulders located around Hashigui-iwa, the southernmost of Kii peninsula, Japan.
1673 Abstract of Japan Geoscience Union Meeting 2011, AAA035-12 (in Japanese).
- 1674 Namegaya, Y., Satake, K. and Yamaki, S., 2010. Numerical simulation of the AD 869 Jogan
1675 tsunami in Ishinomaki and Sendai plains and Ukedo river-mouth lowland. *Annual Report*
1676 *on Active Fault and Paleoeearthquake Researches, Geological Survey of Japan, AIST, No.*
1677 *10, 1-21* (in Japanese with English abstract).
- 1678 Nanayama, F., Satake, K., Furukawa, R., Shimokawa, K., Atwater, B.F., Shigeno, S., Yamaki,
1679 S., 2003. Unusually large earthquakes inferred from tsunami deposits along the Kuril
1680 trench. *Nature* 424, 660–663, DOI: 10.1038/nature01864.
- 1681 Nishimura, T., 2014. Short-term slow slip events along the Ryukyu Trench, southwestern Japan,
1682 observed by continuous GNSS. *Progress in Earth and Planetary Science* 1, 22,
1683 doi:10.1186/s40645-014-0022-5.
- 1684 Nishimura, S. and Hashimoto, M., 2006. A model with rigid rotations and slip deficits for the
1685 GPS-derived velocity field in Southwest Japan. *Tectonophysics* 421, 187-207.
1686 doi:10.1016/j.tecto.2006.04.017.

- 1687 Nishimura, S., Hashimoto, M. and Ando, M., 2004. A rigid block rotation model for the GPS
1688 derived velocity field along the Ryukyu arc. *Physics of the Earth and Planetary Interiors*
1689 142(3-4), 185-203, doi:10.1016/j.pepi.2003.12.014.
- 1690 Nishimura, T., Miura, S., Tachibana, K., Hashimoto, K., Sato, T., Hori, S., Murakam, E., Kono,
1691 T., Nida, K., Mishina, M., Hirasawa, T. and Miyazaki, S., 2000. Distribution of seismic
1692 coupling on the subducting plate boundary in northeastern Japan inferred from GPS
1693 observations. *Tectonophysics* 323, 217-238. doi.org/10.1016/S0040-1951(00)00108-6.
- 1694 Obara, K., 2002. Nonvolcanic deep tremor associated with subduction in southwest Japan.
1695 *Science*, 296(5573), 1679-1681. 10.1126/science.1070378.
- 1696 Obara, K., and Kato, A., 2016. Connecting slow earthquakes to huge earthquakes. *Science*,
1697 353(6296), 253-257, doi:10.1126/science.aaf1512.
- 1698 Obrocki, L., Vött, A., Wilken, D., Fischer, P., Willershäuser, T., Koster, B., Lang, F.,
1699 Papanikolaou, I., Rabbel, W. and Reicherter, K., 2020. Tracing tsunami signatures of the
1700 AD 551 and AD 1303 tsunamis at the Gulf of Kyparissia (Peloponnese, Greece) using direct
1701 push *in situ* sensing techniques combined with geophysical studies. *Sedimentology* 67,
1702 1274-1308. doi: 10.1111/sed.12555.
- 1703 Okahashi, H., Yasuhara, M., Mitamura, M., Hirose, K., Yoshikawa, S., 2005. Event deposits
1704 associated with tsunamis and their sedimentary structure in Holocene marsh deposits on
1705 the east coast of the Shima Peninsula, central Japan. *J. Geosci. Osaka City Univ.* 48,
1706 143-158.
- 1707 Okumura, K., 1996. Tephrochronology, correlation and deformation of marine terraces in
1708 eastern Hokkaido, Japan. *Geographical Reports of Tokyo Metropolitan University* 31,
1709 19-26.
- 1710 Okamura, M. and Matsuoka, H., 2012. Nankai earthquake recurrences from tsunami sediment.
1711 *Kagaku*, 82, 182-191. (in Japanese)
- 1712 Okamura, Y., Nishizawa, A., Fujii, Y. and Yanagisawa, H., 2018. Accretionary prism collapse:
1713 a new hypothesis on the source of the 1771 giant tsunami in the Ryukyu Arc, SW Japan.
1714 *Scientific Reports* 8, 13620. DOI:10.1038/s41598-018-31956-8.
- 1715 Omoto, K., 2012. Distribution of coral boulders beached by huge tsunamis and calibrated
1716 radiocarbon ages of fossil *Porites* sp. Of Maibahabama, SE of Miyako Island, Okinawa
1717 Prefecture. *Journal of Geography (Chigaku Zasshi)* 121, 1043–1051. (in Japanese with
1718 English abstract).
- 1719 Omoto, K., 2019. ¹⁴C ages of large coral boulders washed ashore on the southeast coast of
1720 Miyako Island, Okinawa Prefecture. *Quarterly Journal of Geography* 71, 82-88. (In
1721 Japanese with English abstract)

1722 Ota, Y. and Omura, A., 2000. Review of coral reef terrace studies at Kikai Island, Ryukyu
1723 Islands, southwestern Japan. *The Quaternary Research (Daiyonki-Kenkyu)* 39, 45–53 (in
1724 Japanese with English abstract).

1725 Ozawa, S., Hashimoto, M. and Tada, T., 1997. Vertical crustal movements in the coastal areas
1726 of Japan estimated from tidal observations. *Bull. Geog. Surv. Inst.* 43, 1-21.

1727 Ozawa, S., Murakami, M., Kaidzu, M., Tada, T., Sagiya, T., Hatanaka, Y., Yurai, H. and
1728 Nishimura T., 2002. Detection and monitoring of ongoing aseismic slip in the Tokai region,
1729 central Japan. *Science* 298, 1009–1012. doi:10.1126/science.1076780.

1730 Paris, R, Falvard, S., Chagué, C., Goff, J., Etienne, S. and Doumalin, P., 2020. Sedimentary
1731 fabric characterized by X-ray tomography: A case-study from tsunami deposits on the
1732 Marquesas Islands, French Polynesia. *Sedimentology* 67, 1207-1229. doi:
1733 10.1111/sed.12582.

1734 Peters, R., Jaffe, B. and Gelfenbaum, G., 2007. Distribution and sedimentary characteristics of
1735 tsunami deposits along the Cascadia margin of western North America. *Sedimentary*
1736 *Geology*, 200, 372-386. doi.org/10.1016/j.sedgeo.2007.01.015

1737 Priest, G.R., Witter, R.C., Zhang, Y.J., Goldfinger, C., Wang, K., Allan, J.C., 2017, New
1738 constraints on coseismic slip during southern Cascadia subduction zone earthquakes over
1739 the past 4,600 years implied by tsunami deposits and marine turbidites, *Natural Hazards*
1740 DOI: 10.1007/s11069-017-2864-9.

1741 Riedesel, S., Brill, D., Roberts, H.M., Duller, G.A.T., Garrett, E., Zander, A.M., King, G.E.
1742 Tamura, T., Burow, C., Cunningham, A., Seeliger, M., De Batist, M., Heyvaert, V.M. A.,
1743 Fujiwara, O., Brückner, H., the QuakeRecNankai team, 2018. Single-grain feldspar
1744 luminescence chronology of historical extreme wave event deposits recorded in a coastal
1745 lowland, Pacific coast of central Japan. *Quaternary Geochronology* 45, 37–49.
1746 <https://doi.org/10.1016/j.quageo.2018.01.006>.

1747 Ryan, W.B.F., S.M. Carbotte, J.O. Coplan, S. O'Hara, A. Melkonian, R. Arko, R.A. Weissel, V.
1748 Ferrini, A. Goodwillie, F. Nitsche, J. Bonczkowski, and R. Zemsky (2009), *Global*
1749 *Multi-Resolution Topography synthesis*, *Geochem. Geophys. Geosyst.*, 10, Q03014,
1750 doi:10.1029/2008GC002332.

1751 Sagiya, T., and Thatcher, W., 1999. Coseismic slip resolution along a plate boundary
1752 megathrust: The Nankai Trough, southwest Japan. *Journal of Geophysical Research*
1753 104(B1), 1111-1129. doi.org/10.1029/98JB02644.

1754 Saito, E., 2017. GIS data of the depth of the oceanic plate surfaces beneath the Japanese
1755 archipelago. *GSJ Open-File Report*, no. 647, Geological Survey of Japan, AIST. (in
1756 Japanese).

- 1757 Sangawa, A., 1992. Earthquake archaeology: Earthquake history revealed from archaeological
1758 site). *Chuōkōron-shin sha*, 251p. (in Japanese)*.
- 1759 Sangawa, A., 2001. Recent results of paleoseismological study based on earthquake traces
1760 excavated at archaeological sites. *Annual Report on Active Fault and Paleoearthquake*
1761 *Researches*, Geological Survey of Japan, AIST, No. 1, 287–300 (in Japanese with English
1762 abstract).
- 1763 Sangawa, A., 2007. Japanese History Looking Back from Large Earthquakes. *Chu-ko Shinsho*,
1764 268pp (in Japanese) *.
- 1765 Sateke, K. and Atwater, B.F., 2007. Long-term perspectives on giant earthquakes and tsunamis
1766 at subduction zones. *Annual Review of Earth and Planetary Sciences* 35, 349-374.
1767 <https://doi.org/10.1146/annurev.earth.35.031306.140302>.
- 1768 Satake, K., Namegaya, Y. and Yamaki, S., 2008. Numerical simulation of the AD 869 Jogan
1769 tsunami in Ishinomaki and Sendai plains. *Annual Report on Active Fault and*
1770 *Paleoearthquake Researches*, Geological Survey of Japan, AIST, No. 8, 71-89 (in Japanese
1771 with English abstract).
- 1772 Sato, Y. and Fujiwara, O., 2017. Microfossil evidence for recurrent coseismic subsidence
1773 around Lake Hamana, near the Nankai-Suruga trough, central Japan. *Quaternary*
1774 *International* 456, 39-52. <http://dx.doi.org/10.1016/j.quaint.2017.08.040>
- 1775 Sato, Y., Fujiwara, O. and Ono, E., 2016a. Late Holocene geomorphological development of
1776 beach ridges in western Hamamatsu strand plain, central Japan. *The Quaternary Research*
1777 55, 17-35 (in Japanese with English abstract)
- 1778 Sato, Y., Fujiwara, O., Ono, E., Umitsu, M., 2011. Environmental change in coastal lowlands
1779 around the Lake Hamana during the middle to late Holocene. *Geographical Review of*
1780 *Japan Series A*, 84, 258-273. (in Japanese with English abstract).
- 1781 Sato, Y., Matsuoka, H., Okamura, M., Kashima, K., 2016b. Late Holocene environmental
1782 changes of coastal lagoon inferred from a fossil diatom analysis of sediment core from
1783 Lake Hamana, central Japan. *Quat. Int.* 397, 317–329.
- 1784 Savage, J. C., 1983. A dislocation model of strain accumulation and release at a subduction zone.
1785 *Journal of Geophysical Research* 88(Nb6), 4984-4996, doi:10.1029/JB088iB06p04984.
- 1786 Sawai, Y., in press. Subduction zone paleoseismology along the Pacific coast of northeast Japan
1787 – progress and remaining problems. *Earth-Science Reviews*.
- 1788 Sawai, Y., Namegaya, Y., Okamura, Y., Satake, K. and Shishikura, M., 2012, Challenges of
1789 anticipating the 2011 Tohoku earthquake and tsunami using coastal geology. *Geophysical*
1790 *Research Letters*, **39**, doi: 10.1029/2012GL053692.
- 1791 Sawai, Y., Satake, K., Kamataki, T., Nasu, H., Shishikura, M., Atwater, B.F., Horton, B. P.,
1792 Kelsey, H.M., Nagumo, T. and Yamaguchi, M., 2004. Transient uplift after a 17th-century

1793 earthquake along the Kuril subduction zone. *Science* 306, 1918-1920. DOI:
1794 10.1126/science.1104895

1795 Sawai, Y., Tanigawa, K., Tamura, T. and Namegaya, Y., 2016. Medieval coastal inundation
1796 revealed by a sand layer on the Ita lowland adjacent to the Suruga Trough, central Japan.
1797 *Natural Hazards* 80, 505–519. DOI 10.1007/s11069-015-1980-7.

1798 Seno, T., 2012. Great earthquakes along the Nankai Trough – A new idea for their rupture mode
1799 and time series–. *Zisin (Journal of the Seismological Society of Japan)*, 2nd ser. 64, 97–
1800 116 (in Japanese with English abstract).

1801 Shikakura, Y., 2014. Marine terraces caused by fast steady uplift and small coseismic uplift and
1802 the time-predictable model: Case of Kikai Island, Ryukyu Islands, Japan. *Earth Planet. Sci.*
1803 *Lett.*, 404, 232–237. <https://doi.org/10.1016/j.epsl.2014.08.003>

1804 Shimada Y, Fujino S, Sawai Y, Koichiro T, Matsumoto D, Momohara A,
1805 Saito-Kato M, Yamada M, Hirayama E, Suzuki T, Chagué C., 2019.
1806 Geological record of prehistoric tsunamis in Mugi Town, facing the Nankai
1807 Trough, western Japan. *Progress in Earth and Planetary Science*, 6 (33).
1808 <https://doi.org/10.1186/s40645-019-0279-9>.

1809 Shimazaki, K., 1974a. Pre-seismic crustal deformation caused by an underthrusting oceanic
1810 plate, in eastern Hokkaido, Japan. *Physics of the Earth and Planetary Interiors* 8, 148-157
1811 [https://doi.org/10.1016/0031-9201\(74\)90127-7](https://doi.org/10.1016/0031-9201(74)90127-7).

1812 Shimazaki, K., 1974b. Nemuro-Oki earthquake of June 17, 1973: A lithospheric rebound at the
1813 upper half of the interface. *Physics of the Earth and Planetary Interiors*, 9, 314-327.
1814 [https://doi.org/10.1016/0031-9201\(74\)90059-4](https://doi.org/10.1016/0031-9201(74)90059-4).

1815 Shimazaki, K. and Nakata, T., 1980. Time□predictable recurrence model for large earthquakes.
1816 *Geophysical Research Letters*, 7, 279-282. <https://doi.org/10.1029/GL007i004p00279>

1817 Shinozaki, T., Fujino, S., Ikehara, M. Sawai, Y., Tamura, T., Goto, K., Sugawara, D. and Abe,
1818 T., 2015. Marine biomarkers deposited on coastal land by the 2011 Tohoku-oki tsunami.
1819 *Natural Hazards* 77, 445–460. doi.org/10.1007/s11069-015-1598-9.

1820 Shishikura, M., 2013. Earthquake and tsunamis along the Nankai Trough, inferred from geology
1821 and geomorphology -examples in Nankai region-. *GSJ Chishitsu News*, 2, 201-204. (in
1822 Japanese)

1823 Shishikura, M., Echigo, T., Maemoku, H., Ishiyama, T., 2008a. Height and ages of uplifted
1824 sessile assemblage distributed along the southern coast of the Kii Peninsula, southcentral
1825 Japan – reconstruction of multi-segment earthquake history along the Nankai Trough.
1826 *Annual Report on Active Fault and Paleoequake Researches*, Geological Survey of
1827 Japan, AIST, No. 8, 267–280 (in Japanese with English abstract).

- 1828 Sugihara, K., Nakamori, T., Iryu, Y., Sasaki, K. and Blanchon, P., 2003. Holocene sea-level
1829 change and tectonic uplift deduced from raised reef terraces, Kikai-jima, Ryukyu Islands,
1830 Japan. *Sedimentary Geology* 159, 5-25. [doi.org/10.1016/S0037-0738\(03\)00092-7](https://doi.org/10.1016/S0037-0738(03)00092-7).
- 1831 Sugiyama, Y., 1990. Seismotectonics of the Suruga Bay and Enshu-nada region. *Zishin (Journal*
1832 *of the Seismological Society of Japan)*, 2nd ser., 43, 439–442. (in Japanese)
- 1833 Sugiyama, Y., 1991. The Middle Pleistocene deposits in the Atsumi Peninsula and along the
1834 east coast of Lake Hamana, Tokai district -sedimentary cycles formed by the
1835 glacio-eustatic sea-level change and their correlations to the contemporaneous deposits in
1836 the Kanto and Kinki districts-. *Bulletin of the Geological Survey of Japan* 42, 75-109. (in
1837 Japanese with English abstract).
- 1838 Suito, H. and Ozawa, S., 2009. Transient crustal deformation in the Tokai district —The Tokai
1839 slow slip event and post seismic deformation caused by the 2004 off southeast Kii
1840 Peninsula earthquake —. *Zishin (Journal of the Seismological Society of Japan)*, 2nd ser.,
1841 61, 113-135 (in Japanese with English abstract).
- 1842 Tadokoro, K., Nakamura, M., Ando, M., Kimura, H., Watanabe, T. and Matsuhiro, K., 2018.
1843 Interplate coupling state at the Nansei-Shoto (Ryukyu) Trench, Japan, deduced from
1844 seafloor crustal deformation measurements. *Geophysical Research Letters* 45, 6869–6877.
1845 <https://doi.org/10.1029/2018GL078655>.
- 1846 Takada, K., Satake, K., Sangawa, A., Shimokawa, K., Kumagai, H., Goto, K., Haraguchi, T.,
1847 2002. Survey of tsunami deposits at an archaeological site along the eastern Nankai trough.
1848 Annual report on active fault and paleoearthquake researches, Geological Survey of Japan,
1849 AIST, No. 2, 235-243. (In Japanese with English abstract)
- 1850 Tani, S., Kitagawa, H., Hong, W., Park, J.H. Sung, K.S. and Park G., 2013. Age determination
1851 of the Kawagodaira volcanic eruption in Japan by ¹⁴C wiggle-matching. *Radiocarbon* 55,
1852 748-752. DOI: 10.2458/azu_js_rc.55.16312.
- 1853 Tanigawa, K., Shishikura, M., Fujiwara, O., Namegaya, Y., and Matsumoto, D., 2018. Mid-to
1854 late-Holocene marine inundations inferred from coastal deposits facing the Nankai Trough
1855 in Nankoku, Kochi Prefecture, southern Japan. *The Holocene* 28, 867–878
1856 <https://doi.org/10.1177/0959683617752837>.
- 1857 Tanioka, Y., Satake, K. 2001a. Detailed coseismic slip distribution of the 1944 Tonankai
1858 earthquake estimated from tsunami waveforms. *Geophys. Res. Lett.* 28, 1075–1078.
1859 [doi:10.1029/2000GL012284](https://doi.org/10.1029/2000GL012284).
- 1860 Tanioka, Y., Satake, K., 2001b. Coseismic slip distribution of the 1946 Nankai earthquake and
1861 aseismic slips caused by the earthquake. *Earth, Planets and Space* 53, 235–241.
1862 <https://doi.org/10.1186/BF03352380>.

1863 Tappin, D. R., Watts, P. and Grilli, S. T., 2008. The Papua New Guinea tsunami of 17 July
1864 1998: anatomy of a catastrophic event. *Nat. Hazards Earth Syst. Sci.* 8, 243–266.

1865 The research group for active submarine faults off Tokai(ed.), 1999. active submarine faults off
1866 Tokai -Results from the Japan-France KAIKO-TokaiProject. University of Tokyo Pres,
1867 151p. (In Japanese with English abstract)

1868 Tsuji, Y., 1997 Damage to villages on Amami Archipelago due to the 1995
1869 Amami-Oshima-Kinkai earthquake-tsunami. *Journal of Geography (Chigaku Zasshi)* 106,
1870 486–502 (in Japanese with English abstract).

1871 Tsuji, Y., Okamura, M., Matsuoka, H., Goto, T., Han, S.S., 2002. Prehistorical and historical
1872 tsunami traces in lake floor deposits, Oike Lake, Owase City and Suwaike Lake,
1873 Kii-Nagashima City, Mie Prefecture, central Japan. *Chikyu Mon.* 24, 743–747 (In
1874 Japanese)

1875 Tsuji, Y., Okamura, M., Matsuoka, H., Murakami, Y., 1998. Study of tsunami traces in lake
1876 floor sediment of the Lake Hamanako. *Hist. Earthq.* 14, 101–113 (In Japanese)

1877 Uchida, N., Iinuma, T., Nadeau, R. M., Burgmann, R. and Hino, R., 2016. Periodic slow slip
1878 triggers megathrust zone earthquakes in northeastern Japan. *Science*, 351(6272), 488-492,
1879 doi:10.1126/science.aad3108.

1880 Ujiie, H., Nakamura, T., Miyamoto, Y., Park, J.O., Hyun, S., and Oyakawa, T., 1997.
1881 Holocene turbidite cores from the southern Ryukyu Trench slope: suggestions of periodic
1882 earthquakes. *The Journal of the Geological Society of Japan* 103, 590-603.
1883 <https://doi.org/10.5575/geosoc.103.590>.

1884 Wallace, L. M., Ellis, S., Miyao, K., Miura, S., Beavan, J. and Goto, J., 2009. Enigmatic, highly
1885 active left-lateral shear zone in southwest Japan explained by aseismic ridge collision.
1886 *Geology*, 37, 143-146, doi:10.1130/G25221a.1.

1887 Wallis, S.R., Fujiwara, O. and Goto, K. (2018) Geological studies in tsunami research since the
1888 2011 Tohoku earthquake. *Geological Society London Spec. Publication*, 456, 39-53.


1889 Watanabe, H., 1998. Comprehensive list of tsunamis to hit the Japanese Islands. Tokyo: Tokyo
1890 University Press, 238 pp (in Japanese).

1891 Watanabe, M., Goto, K., Imamura, F. and Hongo, C., 2016. Numerical identification of tsunami
1892 boulders and estimation of local tsunami size at Ibaruma reef of Ishigaki Island, Japan.
1893 *Island Arc* 25, 316–332. doi.org/10.1111/iar.12115.

1894 Watanabe, M., Goto, K., Bricker, J. D. and Imamura, F., 2018. Are inundation limit and
1895 maximum extent of sand useful for differentiating tsunamis and storms? An example from
1896 sediment transport simulations on the Sendai Plain, Japan. *Sedimentary Geology*, 364,
1897 204-216. doi.org/10.1016/j.sedgeo.2017.12.026.

- 1898 Webster, J. M., Davies P. J. and Konishi, K., 1998. Model of fringing reef development in
1899 response to progressive sea level fall over the last 7000 years - (Kikai-jima, Ryukyu
1900 Islands, Japan). *Coral Reefs* 17, 289–308.
- 1901 Wells R. E., Blakely, R. J., Sugiyama, Y., Scholl, D. W. and Dinterman, P. A., 2003, Basin-
1902 centered asperities in great subduction zone earthquakes: A link between slip, subsidence,
1903 and subduction erosion? *J. Geophys. Res.*, **108**, doi: 10.1029/2002JB002072.
- 1904 Yamada, M., Fujino, S., Chiba, T., Goto, K. and Goff, J., 2020. Redeposition of volcanoclastic
1905 sediments by a tsunami 4600 years ago at Kushima City, south-eastern Kyushu, Japan.
1906 *Sedimentology*. doi.org/10.1111/sed.12547.
- 1907 Yamamoto, M., 2008. Trace of earthquake and tsunami in Ishigaki Island, Yaeyama
1908 Archipelago. *Archaeol. J.* 577, 9–14. (in Japanese).
- 1909 Yamamoto, Y., Obana, K., Takahashi, T., Nakanishi, A., Kodaira, S. and Kaneda, Y., 2013.
1910 Imaging of the subducted Kyushu-Palau Ridge in the Hyuga-nada region, western Nankai
1911 Trough subduction zone. *Tectonophysics* 589, 90-102, doi:10.1016/j.tecto.2012.12.028.
- 1912 Yamanaka, Y. and Kikuchi, M., 2003. Source process of the recurrent Tokachi-oki earthquake
1913 on September 26, 2003, inferred from teleseismic body waves. *Earth Planets Space* 55,
1914 e21-e24. doi:10.1186/BF03352479.
- 1915 Yamashita, Y., Shimizu, H. and Goto, K., 2012. Small repeating earthquake activity, interplate
1916 quasi-static slip, and interplate coupling in the Hyuga-nada, southwestern Japan subduction
1917 zone. *Geophysical Research Letters* 39, doi:Artn L08304 10.1029/2012gl051476.
- 1918 Yamashita, Y., Yakiwara, H., Asano, Y., Shimizu, H., Uchida, K., Hirano, S., Umakoshi, K.,
1919 Miyamachi, H., Nakamoto, M., Fukui, M., Kamizono, M., Kanehara, H., Yamada, T.,
1920 Shinohara, M. and Obara, K., 2015. Migrating tremor off southern Kyushu as evidence for
1921 slow slip of a shallow subduction interface. *Science* 348, 676-679. DOI:
1922 10.1126/science.aaa4242.
- 1923 Yata, T., 2009. Great earthquakes in Japanese medieval times. *Yoshikawa kobunkan*, 203p (in
1924 Japanese)*.
- 1925 Yata, T., 2018. Great earthquakes in Japanese early modern times. *Yoshikawa kobunkan*, 248p.
1926 (in Japanese)*.
- 1927 Yata. T., 2013. Subsidence of the northern part of Lake Hamana during the 1707 Hoei
1928 Earthquake. *Shiryogaku-kenkyu* (Researches on historical materials), No. 10, 1-14 (in
1929 Japanese)*.
- 1930 Yokota, Y., Ishikawa, T., Watanabe, S., Tashiro, T. and Asada, A., 2016. Seafloor geodetic
1931 constraints on interplate coupling of the Nankai Trough megathrust zone. *Nature* 534,
1932 374-377, doi:10.1038/nature17632.

1933 Yoshikawa, T., Kaizuka, S. and Ota, Y., 1964. Mode of crustal movement in the late Quaternary
1934 on the southeast coast of Shikoku. southwestern Japan. Geographical Review of Japan 37,
1935 627-648. (in Japanese with English abstract) <https://doi.org/10.4157/grj.37.627>
1936 Zang, S.X., Chen, Q.Y., Ning, J.Y., Shen, Z.K. and Liu, Y.G., 2002. Motion of the Philippine
1937 Sea plate consistent with the NUVEL-1A model. Geophysical Journal International 150,
1938 809-819. <https://doi.org/10.1046/j.1365-246X.2002.01744.x>
1939
1940 *English translation from the original written in Japanese.
1941
1942




1944

1945 **Fig. 1. Map showing the outline of the seismotectonic setting around the Nankai and**
 1946 **Ryukyu subduction zones.**

1947 Tsunami source areas for the 1707 CE (green line) and two 1854 CE earthquakes (orange
 1948 solid and dashed lines), as modified from Hatori (1974, 1976). Earthquake rupture areas of the
 1949 1944 CE Tōnankai and 1946 CE Showa-Nankai earthquakes (blue shaded areas) are simplified
 1950 from Tanioka and Satake (2001a, b). Plate movement vector from Zang et al. (2002).
 1951 Classification of rupture zones, A to E and Z, follows Ando, 1975 (a, b), Ishibashi, 1976, and
 1952 Wells et al. (2003). Contour lines showing the depth of the Philippine Sea plate surface, 10
 1953 km-interval, modified from Saito (2017). Distribution of high interpolate slip-deficit rate area
 1954 (>5 cm yr⁻¹) along the Nankai Trough is simplified from Yokota et al. (2016). Spatial
 1955 distribution of the averaged quasi-static slip rate in the Hyūga-nada region (from May 1994 to
 1956 May 2011) is reproduced from Fig. 2 b of Yamashita et al. (2012) with the permission of
 1957 publisher (John Wiley and Sons; license number 4775371178223). Source area of “the

1958 maximum possible large earthquake” in Nankai subduction zone modified from Cabinet Office
 1959 (2011b).
 1960



1961

1962 **Fig. 2. Spatiotemporal compilation of representative paleotsunami evidence along the**
 1963 **Ryukyu and Nankai subduction zones**

1964 Horizontal lines indicate Nankai megathrust earthquake rupture zones, following Fujiwara
 1965 et al. (2020); dashed where uncertain, dotted to indicate the debated 1099 CE earthquake. We
 1966 include four sites that record evidence for coseismic deformation (numbers 14, 21, 27 and 30).

1967 Site numbers: 1. Ibaruma, Ishigaki Island; 2. Sakishima Islands; 3. Miyako Islands[†]; 4.
 1968 Okinawa Islands; 5. Amami Islands; 6. Kushima City[†]; 7. Ryūjin-ike; 8. Ryuuoo-ike; 9.

1969 Tadasu-ike*; 10. Kaniga-ike; 11. Nankoku[†]; 12. Mugi Town[†]; 13. Kamoda-oike; 14. Kii
1970 Peninsula* (including Kuchiwabuka, Ameshima, Shionomisaki, Izumozaki, Arafunezaki,
1971 Ikeshima, Yamamibana, Taiji, and Suzushima); 15. Ōike (Sagari) Pond*; 16. Suwa Pond*; 17.
1972 Shima Lowlands[†]; 18. Ōsatsu Town*; 19. Nagaya Moto-Yashiki*; 20. Shirasuka; 21. Shinjo
1973 Lowlands[†]; 22. Arai*; 23. Lake Hamana*; 24. Western Hamamatsu strand plain[†]; 25. Ōtagawa
1974 Lowlands[†]; 26. Yokosuka Lowlands*; 27. Omaezaki[†]; 28. Oya Lowlands*; 29. Shimizu Plain*;
1975 30. Ukishima-ga-hara*; 31. Ita Lowlands*; 32. Shimoda*.

1976 Sites with age recalibrated or modelled in Garrett et al. (2016) marked *, sites with age
1977 ranges recalibrated or modelled in this publication marked [†]. We modeled age ranges using the
1978 OxCal program v.4.2 (Bronk Ramsay, 2009) using P_Sequence and Sequence models (Bronk
1979 Ramsay, 1995, 2008, 2009), following the approach detailed in Garrett et al. (2016). For the
1980 Miyako Islands, we interpreted peaks in a cumulative probability distribution as indicating the
1981 timing of boulder movement events, following the approach of Araoka et al. (2013).

1982 Basemap uses Global Multi-Resolution Topography (Ryan et al., 2009) through
1983 www.GeoMapApp.org.


1984

1985 Main references cited in Garrett et al. (2016) but not discussed further in this paper are as
1986 follows.

1987 Site 14: Shishikura et al. (2008a), Site 15, 16: Tsuji et al., (2002), Site 19: Takada et al., 2002,
1988 Site 22: Fujiwara et al. (2013b), Site 26: Fujiwara et al. 2007, Site 28: Kitamura et al. (2013),
1989 Site 32: Kitamura et al. (2014).

1990

1991



1992

1993

Fig. 3. Photographs showing examples of tsunami deposits

1994

A) Overview of four historical tsunami deposits at the Ōtagawa Lowlands, facing the Tōkai segment of the Nankai Trough. Stratified sandy tsunami deposits are distinguished as horizontal color bands, which serve as an aquifer, in back-marsh mud.


1995




1996


1997

1998 B) A *Porites* tsunami boulder with microatoll shape deposited on the eastern coast of Ishigaki
1999 Island, Japan. The scale at lower right is 30 cm.
2000

2001







2004

2005 **Fig 4. Paleotsunami deposits in the western Hamamatsu Plain**

2006 A) Geomorphic classification map of Hamamatsu-Lake Hamana area showing the main
 2007 coring sites B to E. Map modified from Sato et al. (2011) with additional air photo
 2008 analyses and land survey data.

2009 B) Geological columnar section and results of diatom analyses from Site B. Occurrence of
 2010 possible tsunami sand beds is limited beneath the Kg tephra (1210-1187 cal BCE, Tani et
 2011 al., 2013). Up-core changes in diatom assemblages suggest increasingly freshwater
 2012 conditions in the back marsh. Increased occurrences of marine-brackish diatoms
 2013 accompany the deposition of some sand beds. Figure captions are common for Fig. B to
 2014 Fig. D.


2015 C) Geological columnar section from Site C. The occurrence of possible washover sand beds is
 2016 limited to the period older than 1175-1055 cal BP (775-895 CE).

2017 D) Geological columnar section and results of diatom analyses from Site D. An increase of
 2018 brackish-marine diatom species suggests the deposition of a tsunami sand bed ~3700-3300
 2019 cal BP following the coastal subsidence.

2020 E) Geological cross-section at the site E. A tsunami sand bed showing a landward fining and
 2021 thinning trend over 500 m along the incised valley axis. The occurrence of tsunami beds
 2022 and other sand beds are limited below the Kg tephra.

2023

2024 Figs. 4B to 4D and Fig. 4E are modified from Sato et al. (2016a) and Fujiwara et al. (2013a),
 2025 respectively.



B


	Ansei-type						Hoei-type					
	Z	A	B	C	D	E	Z	A	B	C	D	E
Showa								1946		1944		
Ansei		1854			1854							
Hoei								1707				
Meio		?			1498							
Shohei								1361		1361		
Kowa/Eicho		1099			1096							
Ninna									887			
Hakuho		684			?							

2026

2027 Fig. 5 Schematic illustration of proposed recurrence models for subduction zone earthquakes

2028

A) modified from Satake and Atwater (2007), B) simplified from Seno (2012).



2029

2030 Fig. 6. Conceptual model illustrating the reconstruction of paleotsunami size from the spatial
 2031 distribution of tsunami deposits considering the geomorphic evolution of a strand plain

2032 This figure was originally shown as Fig. 3 in Fujiwara (2013) and quoted in Fujiwara and
 2033 Tanigawa (2017).

		Rupture zone									
		Sakishima	Okinawa	Amami	Osumi and Tokara	Z	A	B	C	D	E
Length of history		Ryukyu Trench ~400 years				Hyuga-nada ?	Nankai Trough ~1300 years				
historical event (Mw>8.0)	1946 Showa										
	1944 Showa										
	1911 Kikai										
	1854 Ansei										
	1854 Ansei										
	1791 event										
	1771 Meiwa										
	1707 Hoei										
	1614 Keicho										
	1498 Meio							△	△		
	1361 Shohei										
	1361 Shohei										
	1099 Kowa										
	1096 Eicho										
	887 Ninna										
684 Hakuho											
	latest 7th century								△	△	
Seismological evidence	slow slip	not studied	not studied	not studied	not studied	○	○	○	○	○	○
	coupled or decoupled	⊙	○	not studied	not studied	not studied	⊙	○	○	⊙	○
Paleotsunami evidence	late Holocene paleotsunami evidence	Y	N	N	not studied	very rare?	Y	Y	Y	Y	Y
	paleotsunami interval (years)	150-400	-	-	not studied		90-265				
Geomorphological evidence	co-seismic crustal deformation	-	-	Y	-	-	Y	Y	Y	Y	Rare
	...										

2034

	Shaking+tsunami+coastal deformation	△	Liquefaction
	Shaking+tsunami		tsunami
	Shaking+coastal deformation		

2035

2036

Table 1. Paleoearthquake and tsunami evidence and related seismotectonic data along the Nankai and Ryukyu subduction zones.

2037

2038

2039

2040