

This is a repository copy of *In anti-CCP+ at-risk individuals, radiographic bone erosions* are uncommon and are not associated with the development of clinical arthritis.

White Rose Research Online URL for this paper: https://eprints.whiterose.ac.uk/166595/

Version: Accepted Version

Article:

Di Matteo, A orcid.org/0000-0003-0867-7051, Mankia, K, Nam, JL et al. (5 more authors) (2020) In anti-CCP+ at-risk individuals, radiographic bone erosions are uncommon and are not associated with the development of clinical arthritis. Rheumatology. ISSN 1462-0324

https://doi.org/10.1093/rheumatology/keaa761

© 2020, Oxford University Press. This is an author produced version of an article published in Rheumatology. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/

Baseline	Hands/feet CR-BE distribution	SENS score (BE)	SvdH score (BE)	Radiologist's diagnosis	Follow-up	Hands/feet CR-BE distribution	SENS score (BE)	SvdH score (BE)	Status
	R radiocarpal and 1 st MCP joints	2	4	IA		R radiocarpal and 1 st MCP joints	2	4	Ongoing
	L 5 th DIP joint	0	0	OA		L 5 th DIP joint	0	0	Progressed
	L 3 rd DIP joint and R 5 th DIP joints, bilateral 1 st MTP joints, bilateral 5 th MTP joints	4	12	IA		L 3 rd DIP and R 5 th DIP joints, R 1 st DIP joint, bilateral 1 st MTP joints, bilateral 5 th MTP joints	4	14	Progressed
	R 2 nd and 3 rd MCP joints, R 2 nd DIP joint, L 5 th MCP and 5 th PIP joints, bilateral 1 st MTP joints, L 1 st IP joint (foot)	7	13	IA		R 2 nd and 3 rd MCP joints, R 2 nd DIP joint, bilateral 4 th PIP joints, L 5 th PIP joint, bilateral 1 st MTP joints, L 1 st IP joint (foot)	8	17	Ongoing
	L 3 rd MCP joint, R 2 nd PIP joint, bilateral 3 rd PIP joints, L 4 th PIP joint, R 4 th PIP joint, bilateral 5 th PIP joints, bilateral 1 st MTP joints, 5 th R MTP joint	11	17	Erosive OA		L 3 rd MCP joint, L 2 nd PIP joint, bilateral 2 nd , 3 rd , 4 th PIP joints, bilateral 1 st MTP joints, R 5 th MTP joint	11	22	Ongoing
	L 5 th MTP joint	1	3	IA		R 2 nd and 3 rd DIP joints, L 5 th MTP joint	1	3	Progressed
	L 3 rd MTP joint, bilateral 1 st IP joints (foot)	3	4	IA		L 3rd MTP joint, bilateral 1st IP joints (foot)	3	12	Ongoing
	R 1 st MTP joint	1	6	OA		R 1 st MTP joint	1	9	Ongoing
	Bilateral 2 nd tarso-metatarsal joints	0	0	OA		R 2 nd DIP joint, bilateral 2 nd tarso-metatarsal joints	0	0	Ongoing
	R 2 nd MCP joint, R 5 th MCP joint	2	5	IA		R 2 nd MCP joint, R 5 th MCP joint	2	4	Ongoing

Table 4. Repeat CR in 10 out of 17 CCP+ at risk individuals with baseline bone erosions

Legend. BE: bone erosions, CR: conventional radiography, DIP: distal interphalangeal, IA: inflammatory arthritis, IP: interphalangeal, L: left, MCP: metacarpophalangeal, MTP: metatarsophalangeal, OA: osteoarthritis, PIP: proximal interphalangeal, R: right, SENS: Simple Erosions Narrowing Score, SvdH: Sharp van der Heijde.