



This is a repository copy of *Methods matter : pilot and feasibility studies in sports medicine*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/166911/>

Version: Accepted Version

Article:

Campbell, M.J. orcid.org/0000-0003-3529-2739, Mansournia, M.A. and Lancaster, G. (2020) *Methods matter : pilot and feasibility studies in sports medicine*. *British Journal of Sports Medicine*, 54 (22). pp. 1309-1310. ISSN 0306-3674

<https://doi.org/10.1136/bjsports-2020-102631>

© 2020 The Author(s). This is an author-produced version of a paper subsequently published in *British Journal of Sports Medicine*. Campbell MJ, Mansournia MA, Lancaster G., *Methods matter: pilot and feasibility studies in sports medicine*. *British Journal of Sports Medicine* Published Online First: 14 July 2020. doi: 10.1136/bjsports-2020-102631

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Pilot and feasibility studies in Sports Medicine

Michael J. Campbell¹ , Mohammed A Mansournia²

Gillian A. Lancaster³

Address:

SCHARR, University of Sheffield (address 91 Stumperlowe Hall Rd, Sheffield, UK S10 3QT.)

¹Corresponding Authors: Michael J Campbell and Mohammad Mansournia

Email: m.j.campbell@sheffield.ac.uk +44 114 230 1837

² Mohammad affiliation

³Professor of Medical Statistics, School of Primary, Community and Social Care and Keele Clinical Trials Unit (CTU)

Over recent years, there has been an increasing emphasis on the importance of preliminary work prior to the organisation of large-scale, publicly funded randomised controlled trials. Many large public funding bodies now expect substantial work to have been done prior to the main bid for money to fund a large trial or indeed any large study. The value of preliminary work is now recognised and researchers are encouraged to carry out feasibility and pilot studies before a major trial and to publish both the protocol and the results of their work in advance of the main trial. There is a new journal, *Pilot and Feasibility Studies*, which publishes the results of such studies.

Similar to a randomised controlled trial, a pilot trial should have clear objectives and a priori criteria for success. It should be designed, conducted and reported using the same standards as any high-quality randomized controlled trial. Pilot and feasibility studies are important in that they can ensure that scarce research money and researcher efforts are being invested in efficient trials that can provide definitive answers to important research questions. Guidelines for reporting pilot trials as part of an extension of the CONSORT statement have been published [1,2]

A recent review of physical activity trials comments on some of the feasibility issues that are more common in this area [3]. These include: the challenge of blinding of group allocation, retaining interest of the comparison group, participant and instructor fidelity, and being able to recruit people who are not already active enough, but interested enough in becoming active to sign up for the study. A second review in sports medicine reported on five journals which published over five articles which related to randomized pilot trials between 2012 and 2015 [4] The authors found that reporting and understanding of pilot and feasibility studies in studies of physical activity was poor, with few providing sufficient explanation. Over half of the studies did not have feasibility objectives which corresponded with previous research findings, so that these findings were not being disseminated effectively to researchers in the field of physical activity. The low standard of reporting across most of the reviewed articles

and the fact that the extended CONSORT 2010 statement was ignored by the journal editors highlights the need to actively disseminate these guidelines to ensure their impact. The authors also reported that journal editors in this area will rarely publish pilot and feasibility studies, but this might be because the studies are generally of poor quality and an objective of this paper is to encourage better design and reporting in the area of physical activity trials.

Current Definitions

There are several papers that clarify the various distinctions in pilot and feasibility studies and these are summarised in the Box. [2,5,.6]

Box here

Researchers should view feasibility as an overarching concept, with all studies done in preparation for a main study open to being called feasibility studies, and with pilot studies as a subset of feasibility studies. Authors should describe their feasibility objectives and methods well and in accordance with the guidelines [1], reporting clearly if their study is in preparation for a future RCT to evaluate the effect of an intervention or therapy. A pilot study should address the main uncertainties that have been identified in the development work in order to plan a main trial or other major piece of research.

The CONSORT statement and the Horne review [1,4] only apply to *randomised external* pilot trials (See Box). An internal pilot study will have a trial monitoring committee which will make suggestions as to whether the trial should be abandoned at that stage or extended either in length or eligibility of participants and possibly suggest revising the sample size upward to accommodate unanticipatedly large variability in the outcome variable. An external pilot trial may suggest, for example, that an individually randomised trial is not possible and so the team may consider a cluster trial as the main trial. The results from the pilot may suggest the main outcome variable may be difficult to measure accurately, so other outcomes measures might be proposed. Note that changing the outcome variable from what is proposed in the protocol to that which appears in the main trial report often highlights major generalisability

issues with the trial and so it is better to identify the main outcome variable whilst planning the trial. [7]

Examples

Haines aimed to describe the feasibility of an exercise intervention (reduced-exertion, high-intensity interval training REHIT) in nondiabetic hyperglycaemia patients [8]. The study intended to recruit 40 participants but achieved only 6. It revealed several issues including patient eligibility, challenges to recruitment, patient consent, and poor clinician engagement. The process of accurately screening and case finding eligible patients was problematic. The author concluded that a trial in its current form is not feasible, but made suggestions as to changes that could occur in future studies.

An excellent trial in sports medicine that, nevertheless, might have benefited from piloting is the INSPIRE (INtervention Study on Prevention of Injuries in Runners at Erasmus MC) Trial [9]. This study with 2378 participants aimed to reduce running injuries in recreational runners by providing evidence-based online advice on modifying known risk factors. It not only failed to show an effect, but the point estimates were in the wrong direction to that expected. It also had a high drop-out rate (28% compared to 5% expected). A feasibility or pilot study might have highlighted the drop-out rate, giving the authors a chance to try and make completion of the outcome questionnaire more attractive. It may also have suggested that the intervention as it stands would be unlikely to be beneficial to the participants, leading to an opportunity for modifying the intervention. However, it is good that such a large, negative trial was published, because *a priori* the hypothesis was worth testing.

We hope that this article will encourage more researchers to think about a pilot or feasibility study before the main trial, and to publish the results. Even if the authors themselves do not plan on a subsequent major study, the information will help other researchers in the field to plan future studies.

Acknowledgements

We thank members of the CONSORT team on Pilot and Feasibility trials for helpful discussion (Sandra Eldridge, Christine Bond, Sally Hopewell, Claire Chan and Lehana Thabane)

References

1. Eldridge SM, Chan CL, Campbell MJ, Bond CM, Hopewell S, Thabane L, Lancaster GA on behalf of the PAFS consensus group. CONSORT 2010 statement: extension to randomised pilot and feasibility trials *BMJ* 2016;355:i5239
2. Eldridge SM, Lancaster GA, Campbell MJ, Thabane L, Hopewell S, Coleman CL, Bond CM Defining feasibility and pilot studies in preparation for randomised controlled trials: development of a conceptual framework. *PLoS One* 2016;11:3
3. El-Kotob R, Giangregorio LM. Pilot and feasibility studies in exercise, physical activity, or rehabilitation research. *Pilot and feasibility studies* 2018;4(1):1-7
4. Horne E, Lancaster GA, Matson R, Cooper A, Ness A, Leary S. Pilot trials in physical activity journals: A review of reporting and editorial policy *Pilot and feasibility studies*; 2018; 4(1):125.
5. Whitehead AL, Sully BG, Campbell MJ. Pilot and feasibility studies: is there a difference from each other and from a randomised controlled trial?. *Contemporary clinical trials*; 2014; 38(1):130-3.
6. Arain M, Campbell MJ, Cooper CL and Lancaster GA What is a pilot or feasibility study? A review of current practice and editorial policy *BMC Medical Research Methodology* 2010; 10:67
7. Kirkham JJ et al. The impact of outcome reporting bias in randomised controlled trials on a cohort of systematic reviews *BMJ* 2010; 340 doi: <https://doi.org/10.1136/bmj.c365>
8. Haines M. Feasibility of procedures for a randomised pilot study of reduced exertion, high-intensity interval training (REHIT) with non-diabetic hyperglycaemia patients. *Pilot and Feasibility Studies* 2020; 6(1):1-3.
9. Fokkema T, de Vos R-J, van Ochten JM, Verhaar JAN, Davis IS, et al. Online multifactorial prevention programme has no effect on the number of running-related injuries: a randomised controlled trial *Br J Sports Med*. 2019;53:1479–1485.

Box : Feasibility and Pilot Studies

A *feasibility* study asks whether something can be done, should the investigator proceed with it, and if so, how.

A *pilot* study asks the same questions but also has a specific design feature. A pilot study replicates a future study, or part of one conducted on a smaller scale.

An *internal* pilot study is simply the first part of the main trial. For an internal pilot trial, the design is fixed and decisions will be made (after a year say) as to whether recruitment to the full trial is possible in the time postulated.

In an *external* pilot trial, the subjects are not expected to be included in the main trial, and more radical changes to the design may be made, such as changing the outcome variable.

Feasibility studies include all preliminary work prior to a main study, and pilot studies are a subset of this.

To facilitate their identification, these studies should be clearly identified using the terms 'feasibility' or 'pilot' as appropriate, including feasibility studies that are largely qualitative.

Investigators should report appropriate objectives and methods related to feasibility; and give clear confirmation whether their study is in preparation for a future randomised controlled trial designed to assess the effect of an intervention.