



This is a repository copy of *Spin fluctuation anisotropy as a probe of orbital-selective hole-electron quasiparticle excitations in detwinned Ba(Fe_{1-x}Cox)₂As₂*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/167067/>

Version: Published Version

Article:

Tian, L., Liu, P., Xu, Z. et al. (7 more authors) (2019) Spin fluctuation anisotropy as a probe of orbital-selective hole-electron quasiparticle excitations in detwinned Ba(Fe_{1-x}Cox)₂As₂. *Physical Review B*, 100 (13). 134509. ISSN 2469-9950

<https://doi.org/10.1103/physrevb.100.134509>

© 2019 American Physical Society. Reproduced in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Spin fluctuation anisotropy as a probe of orbital-selective hole-electron quasiparticle excitations in detwinned $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$

Long Tian¹, Panpan Liu¹, Zhuang Xu¹, Yu Li², Zhilun Lu³, H. C. Walker⁴, U. Stuhr⁵,
Guotai Tan¹, Xingye Lu¹, and Pengcheng Dai^{2,1,*}


¹Center for Advanced Quantum Studies and Department of Physics, Beijing Normal University, Beijing 100875, China

²Department of Physics and Astronomy, Rice University, Houston, Texas 77005, USA

³Helmholtz-Zentrum Berlin für Materialien und Energie GmbH, Berlin 14109, Germany

⁴ISIS Facility, Rutherford Appleton Laboratory, Chilton, Didcot, Oxfordshire OX11 0QX, United Kingdom

⁵Laboratory for Neutron Scattering and Imaging, PSI, CH-5232 Villigen, Switzerland

 (Received 27 July 2019; revised manuscript received 14 September 2019; published 21 October 2019)

We use inelastic neutron scattering to study spin excitation anisotropy in mechanically detwinned $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ with $x = 0.048$ and 0.054 . Both samples exhibit a tetragonal-to-orthorhombic structural transition at T_s , a collinear static antiferromagnetic order at wave vector $\mathbf{Q}_1 = \mathbf{Q}_{\text{AF}} = (1, 0)$ below the Néel temperature T_N , and superconductivity below T_c ($T_s > T_N > T_c$). In the high-temperature paramagnetic tetragonal phase ($T \gg T_s$), spin excitations centered at \mathbf{Q}_1 and $\mathbf{Q}_2 = (0, 1)$ are gapless and have fourfold (C_4) rotational symmetry. On cooling to below T_N but above T_c , spin excitations become highly anisotropic, developing a gap at \mathbf{Q}_2 but still are gapless at \mathbf{Q}_1 . Upon entering into the superconducting state, a neutron spin resonance appears at \mathbf{Q}_1 with no magnetic scattering at \mathbf{Q}_2 . By comparing these results with those from angle-resolved photoemission spectroscopy experiments, we conclude that the anisotropic shift of the d_{yz} and d_{xz} bands in detwinned $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ below T_s is associated with the spin excitation anisotropy, and the superconductivity-induced resonance arises from the electron-hole Fermi surface nesting of quasiparticles with the d_{yz} orbital characters.

DOI: [10.1103/PhysRevB.100.134509](https://doi.org/10.1103/PhysRevB.100.134509)

I. INTRODUCTION

Unconventional superconductors such as copper oxides, iron pnictides, and heavy fermions are interesting because superconductivity in these materials is derived from their long-range antiferromagnetic (AF) ordered parent compounds [1]. Although there is no consensus on the microscopic origin of superconductivity, there is increasing evidence that electron pairing in these superconductors is mediated by spin fluctuations (excitations) [1–5]. In particular, superconductivity is intertwined with magnetic degrees of freedom, and forms a state coexisting with the static AF order in the underdoped regime [1–5]. Therefore, to understand the fundamental interactions that lead to unconventional superconductivity, it is important to investigate how magnetism interacts with superconductivity in the coexisting regime of unconventional superconductors.

In the case of Co-underdoped iron pnictide superconductors such as $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ with $0.03 < x < 0.065$, they exhibit a tetragonal-to-orthorhombic structural transition at T_s , a collinear static AF order below the Néel temperature T_N , and superconductivity below T_c ($T_s > T_N > T_c$) as shown in Fig. 1(a) [6–11]. As a function of decreasing temperature, a collinear static AF order is established below T_N at wave vector $\mathbf{Q}_1 = \mathbf{Q}_{\text{AF}} = (1, 0)$ [inset in Fig. 1(a)]. On further cooling across T_c , the static ordered moment decreases below T_c accompanied by the formation of a neutron spin

resonance coupled to superconductivity [9–12]. For optimally and overdoped $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$, where their T_s and T_N are suppressed and the system is in the paramagnetic tetragonal state, neutron spin resonance occurs at wave vectors $\mathbf{Q}_1 = (1, 0)$ and $\mathbf{Q}_2 = (0, 1)$, and therefore obeys fourfold rotational (C_4) symmetry of the underlying tetragonal lattice [13–16]. While superconductivity clearly competes with static AF order in underdoped $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ [9–11], much is unclear concerning how the superconductivity-induced neutron spin resonance interacts with spin waves from the AF-ordered phase. Although the collinear AF order and associated low-energy spin waves should have twofold rotational (C_2) symmetry below T_N , the observed resonance and spin waves have the C_4 symmetry from the presence of twin domains of the orthorhombic phase below T_s [9–11]. Therefore, to understand the interplay between spin waves associated with static AF order and the neutron spin resonance connected with superconductivity, one must carry out inelastic neutron scattering experiments on detwinned samples with static AF order and superconductivity.

In this paper, we report comprehensive inelastic neutron scattering experiments designed to study spin excitations in detwinned $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ with coexisting AF order and superconductivity. From the phase diagram of $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ in Fig. 1(a) established from our own and previous transport work [11], we know that static AF order decreases with increasing Co doping, and competes with superconductivity, which increases with increasing Co doping [9–11]. To study the interplay of spin waves associated with the static AF order and resonance connected

*pdai@rice.edu




FIG. 1. (a) The phase diagram of $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ with the arrows indicating the Co-doping concentrations of our samples ($x = 0.048$ and 0.054). The left-hand inset shows a schematic of the collinear AF ordering of the Fe spins in real space and the applied uniaxial pressure direction is marked by vertical arrows. The right-hand inset is the corresponding reciprocal space map showing the AF ordering wave vector \mathbf{Q}_1 (green) and wave vector \mathbf{Q}_2 (red). The filled hexagonal and circular points represent T_s and T_N , respectively, obtained from resistance measurements. The filled and open square points mark T_c determined from the resistance and magnetic susceptibility measurements, and from Ref. [11]. Schematic Fermi surfaces of underdoped $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ in (b) paramagnetic tetragonal state and (c) nematic state below T_s . The red, green, and blue colors represent d_{xz} , d_{yz} , and d_{xy} orbitals, respectively. The arrows mark nesting wave vectors \mathbf{Q}_1 and \mathbf{Q}_2 between the Z and X/Y points. Our definition of the X/Y is switched from that of the ARPES work [29]. The normalized temperature-dependent resistance data indicate the superconducting transition temperatures of (d) $x = 0.048$ and (e) $x = 0.054$.

with superconductivity, one must judiciously choose the Co-doping concentrations where the strength of the spin waves is comparable with the superconductivity-induced resonance [17,18]. For this purpose, we prepared single crystals of $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ with $x = 0.048$ and 0.054 [7–9]. At zero external uniaxial pressure [19–22], the $x = 0.048$ samples have $T_c = 18$ K, $T_N = 52$ K, and $T_s = 63$ K [Fig. 1(d)], and




FIG. 2. (a) Elastic rocking curve scans of the sample angle (A_3) around $\mathbf{Q}_1 = (1, 0, 1)$ and $\mathbf{Q}_2 = (0, 1, 1)$ in uniaxial strained $\text{Ba}(\text{Fe}_{0.952}\text{Co}_{0.048})_2\text{As}_2$. The double peaks show the sample has two major domains separated by about 4.5° . (b) Temperature differences of the transverse scans around \mathbf{Q}_1 and \mathbf{Q}_2 in uniaxial strained $\text{Ba}(\text{Fe}_{0.946}\text{Co}_{0.054})_2\text{As}_2$.

0.054 crystals have $T_c = 21$ K, $T_N = 38$ K, and $T_s = 53$ K [Fig. 1(e)]. Upon detwinning these crystals using a device similar to previous work with about 40 MPa uniaxial pressure [23,24], the system no longer has a clean T_s because the C_4 rotational symmetry in the tetragonal phase is already broken by the applied pressure and T_N increases several K under pressure consistent with the earlier work [19–22]. We carried out inelastic neutron scattering measurements on nearly 100% detwinned $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$. In the normal state above T_c but below T_N , spin excitations are gapless and increase with increasing energy at the AF ordering wave vector $\mathbf{Q}_1 = (1, 0)$, but have a ~ 12 meV gap at $\mathbf{Q}_2 = (0, 1)$ [see inset in Fig. 1(a)], showing strong magnetic anisotropy. On cooling to below T_c , a neutron spin resonance at E_r and a spin gap at energies below E_r are formed at $\mathbf{Q}_1 = (1, 0)$, but there is no superconductivity-induced magnetic scattering at $\mathbf{Q}_2 = (0, 1)$. On warming to temperatures slightly above the finite pressure T_N , the spin excitations are still anisotropic, showing much strong scattering at $\mathbf{Q}_1 = (1, 0)$. Finally, on warming to temperatures well above T_N and zero pressure T_s , spin excitations become the same at $\mathbf{Q}_1 = (1, 0)$ and $\mathbf{Q}_2 = (0, 1)$, and obey the C_4 rotational symmetry. By comparing these results with those from angle-resolved photoemission spectroscopy (ARPES) experiments [25–30], we conclude that the anisotropic shift of the d_{yz} and d_{xz} bands of the electron Fermi




FIG. 3. (a) Constant- \mathbf{Q} scans at \mathbf{Q}_1 below and above T_c in $\text{Ba}(\text{Fe}_{0.952}\text{Co}_{0.048})_2\text{As}_2$. The resonance is seen as intensity gain below T_c around $E_r \approx 5.5$ meV. The gray data points represent the background scattering. (b) Identical scans at \mathbf{Q}_2 . (c), (d) Background subtracted constant- \mathbf{Q} scans below and above T_c at the \mathbf{Q}_1 and \mathbf{Q}_2 points. The vertical arrow indicates the position of the neutron spin resonance.

pockets at the X and Y points in detwinned $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ below T_s is associated with the spin excitation anisotropy, and the superconductivity-induced resonance arises from the electron-hole Fermi surface nesting of quasiparticles with the d_{yz} orbital characters along the \mathbf{Q}_1 direction as shown schematically in Fig. 1(c).




FIG. 4. Constant-energy scans in $\text{Ba}(\text{Fe}_{0.952}\text{Co}_{0.048})_2\text{As}_2$ below and above T_c with $E_r = 5.5$ meV along the $[H, 0, 1]$ and $[0, K, 1]$ directions around (a) \mathbf{Q}_1 and (b) \mathbf{Q}_2 .




FIG. 5. Two-dimensional images of the spin excitations of $\text{Ba}(\text{Fe}_{0.946}\text{Co}_{0.054})_2\text{As}_2$ within the $[H, K]$ plane below and above T_c at energies of (a), (b) $E = 5 \pm 1$ meV; (c), (d) $E = 8 \pm 1$ meV; (e), (f) $E = 12 \pm 1$ meV. The vertical color bars indicate scattering intensity in arbitrary units.

II. EXPERIMENTAL RESULTS

Our neutron scattering experiments were carried out on the FLEXX cold neutron three-axis spectrometer at Helmholtz Zentrum Berlin, Germany, and the MERLIN neutron time-of-flight (TOF) chopper spectrometer at ISIS, Rutherford-Appleton Laboratory, UK [31]. The detwinning ratio for $\text{Ba}(\text{Fe}_{0.946}\text{Co}_{0.054})_2\text{As}_2$ samples used at ISIS was measured on the EIGER thermal neutron three-axis spectrometer at Paul Scherrer Institute, Switzerland. Sizable single crystals of $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ were grown by self-flux method and cut along the a - and b -axis directions of the orthorhombic lattice below T_s [4]. Each cut sample was mounted on a specially designed aluminum-based sample holder with uniaxial pressure applied along the b -axis direction [23,24]. The total mass of our samples is ~ 2.5 g for the $x = 0.048$ used for FLEXX experiment, ~ 3.6 g for the $x = 0.054$ used for MERLIN and EIGER experiments. The momentum transfer \mathbf{Q} in three-dimensional reciprocal space is defined as $\mathbf{Q} = H\mathbf{a}^* + K\mathbf{b}^* + L\mathbf{c}^*$, where H , K , and L are Miller indices and $\mathbf{a}^* = \hat{a}2\pi/a$, $\mathbf{b}^* = \hat{b}2\pi/b$, $\mathbf{c}^* = \hat{c}2\pi/c$ with $a = 5.615$ Å, $b = 5.573$ Å, and $c = 12.95$ Å in the low-temperature orthorhombic state [9–11]. In this notation, the AF order occurs at the in-plane wave vector $\mathbf{Q}_{\text{AF}} = (1, 0)$, and there should be no elastic magnetic scattering at wave vector $(0, 1)$.




FIG. 6. Two-dimensional images of magnetic scattering in $\text{Ba}(\text{Fe}_{0.946}\text{Co}_{0.054})_2\text{As}_2$ along the $[1, K]$ and $[H, 1]$ directions at temperatures (a), (b) 4.6 K; (c), (d) 23 K; (e), (f) 60 K; (g), (h) 100 K. The incident beam energy is $E_i = 25$ meV and the partial detwinning ratio of the sample has been corrected for.

For measurements on three-axis spectrometers, we aligned the samples in the $[1, 0, 1] \times [0, 1, 1]$ scattering plane where we can measure the static magnetic order and excitations at both $\mathbf{Q}_{\text{AF}} = \mathbf{Q}_1 = (1, 0, 1)$ and $\mathbf{Q}_2 = (0, 1, 1)$ simultaneously [23]. The fixed final neutron energies are $E_f = 5$ and 14.7 meV for FLEXX and EIGER experiments, respectively. For experiments on TOF spectrometer MERLIN, the direction of the incident beam is parallel to the c axis. Using multi- E_i mode with a primary incident neutron beam energy of $E_i = 80$ meV and Fermi chopper frequency of $\omega = 250$ Hz, we were able to measure with two additional incident energies of 25 and 12 meV, thus allowing spin excitations up to $E < 70$ meV to be probed.

Figure 1(a) shows the phase diagram of electron-doped $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ as determined from our transport measurements. Consistent with previous work [7], we find that the ratio between the actual and nominal Co-doping level is about 0.74. For the experiments, we chose $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ with Co-doping levels $x = 0.048$ and 0.054 as marked by vertical arrows in Fig. 1(a). Figures 1(b) and 1(c) show Fermi surfaces of underdoped $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ with coexisting AF order and superconductivity above and below the zero pressure T_s , respectively, as obtained from ARPES experiments on uniaxial pressure detwinned samples [25–30]. The temperature dependence of the normalized resistance for $\text{Ba}(\text{Fe}_{0.952}\text{Co}_{0.048})_2\text{As}_2$ and $\text{Ba}(\text{Fe}_{0.946}\text{Co}_{0.054})_2\text{As}_2$ reveals superconducting transition temperatures of $T_c = 18$ and 21 K, respectively [Figs. 1(d) and 1(e)].

In order to carry out inelastic neutron scattering experiments on detwinned $\text{Ba}(\text{Fe}_{0.952}\text{Co}_{0.048})_2\text{As}_2$ and $\text{Ba}(\text{Fe}_{0.946}\text{Co}_{0.054})_2\text{As}_2$, one must mount crystals in a uniaxial detwinning device and apply uniaxial pressure along one axis of the orthorhombic lattice to detwin the

samples [see inset of Fig. 1(a)]. For fully detwinned samples, one would expect to observe magnetic Bragg intensity at \mathbf{Q}_1 but no magnetic signal at \mathbf{Q}_2 . Figures 2(a) and 2(b) show the background-subtracted rocking curve elastic scans around \mathbf{Q}_1 and \mathbf{Q}_2 for $\text{Ba}(\text{Fe}_{0.952}\text{Co}_{0.048})_2\text{As}_2$ and $\text{Ba}(\text{Fe}_{0.946}\text{Co}_{0.054})_2\text{As}_2$, respectively. While the $\text{Ba}(\text{Fe}_{0.952}\text{Co}_{0.048})_2\text{As}_2$ sample is $\sim 100\%$ detwinned, there is a weak peak at $\mathbf{Q}_2 = (0, 1, 1)$ for $\text{Ba}(\text{Fe}_{0.946}\text{Co}_{0.054})_2\text{As}_2$. Defining the detwinning ratio as $\eta = (I_{10} - I_{01}) / (I_{10} + I_{01})$, where I_{10} and I_{01} are magnetic Bragg scattering at \mathbf{Q}_1 and \mathbf{Q}_2 , respectively [32,33], we find that $\text{Ba}(\text{Fe}_{0.946}\text{Co}_{0.054})_2\text{As}_2$ has a detwinning ratio of $\eta \approx 85\%$. Using the measured η , we can estimate the intrinsic magnetic scattering at \mathbf{Q}_1 and \mathbf{Q}_2 for different energy transfers, and thus determine the energy dependence of the magnetic scattering at these wave vectors and the related dynamic magnetic susceptibility $\chi''(\mathbf{Q}, E)$ [32,33].

We first present our inelastic neutron scattering results for $\text{Ba}(\text{Fe}_{0.952}\text{Co}_{0.048})_2\text{As}_2$. Figures 3(a) and 3(b) show the constant- \mathbf{Q} scans at \mathbf{Q}_1 and \mathbf{Q}_2 below and above T_c , as well as background scattering at wave vectors transversely rotated $\sim 15^\circ$ from \mathbf{Q}_1 and \mathbf{Q}_2 . At \mathbf{Q}_1 , the scattering increases with increasing energy, and superconductivity induces a resonance at $E_r \approx 5.5$ meV below T_c [Fig. 3(a)]. Figure 3(c) shows background-subtracted scattering, suggesting that superconductivity opens a spin gap below about 2 meV. At \mathbf{Q}_2 , we find no discernible signal above the background scattering both below and above T_c , suggesting the presence of a large spin gap in the normal state and that superconductivity does not induce any magnetic intensity [Fig. 3(b)]. The background-subtracted scattering in Fig. 3(d) suggests a spin gap of ~ 7 meV in both the normal and superconducting states, with detectable magnetic scattering above 8 meV. To further confirm these




FIG. 7. The energy/ c -axis wave-vector dependence of the spin excitations around the in-plane wave vector \mathbf{Q}_1 (green) and \mathbf{Q}_2 (red) positions at temperatures (a) 4.6 K, (b) 23 K, (c) 60 K, (d) 100 K. The incident beam energy is $E_i = 25$ meV, and the vertical arrows indicate the energy values with integer L . The partial detwinning ratio has been corrected for.

results, we show in Figs. 4(a) and 4(b) constant-energy scans at $E_r = 5.5$ meV at \mathbf{Q}_1 and \mathbf{Q}_2 , respectively. While there is a clear magnetic peak centered at \mathbf{Q}_1 that enhances the intensity below T_c , no discernible peak is seen at \mathbf{Q}_2 below or above T_c .

The two-dimensional (2D) magnetic scattering images of $\text{Ba}(\text{Fe}_{0.946}\text{Co}_{0.054})_2\text{As}_2$ in the (H, K) plane for different energy transfers below ($T = 4.6$ K) and above ($T = 23$ K) T_c are shown in Figures 5(a), 5(c), 5(e) and 5(b), 5(d), 5(f), respectively. At $E = 5 \pm 1$ meV, the scattering is centered

around \mathbf{Q}_1 and clearly enhances below T_c , and there is no scattering at \mathbf{Q}_2 [Figs. 5(a) and 5(b)]. On increasing energy to $E = 8 \pm 1$ meV, the situation is similar at \mathbf{Q}_1 but there may be some scattering at \mathbf{Q}_2 [Figs. 5(c) and 5(d)]. On increasing energy to $E = 12 \pm 1$ meV, superconductivity has little effect on spin excitations at \mathbf{Q}_1 and there is weak magnetic signal at \mathbf{Q}_2 [Figs. 5(e) and 5(f)].

Figure 6 summarizes the energy dependence of the spin excitations at \mathbf{Q}_1 and \mathbf{Q}_2 as a function of increasing temperature. The energy dependence of the scattering is obtained by integrating wave vectors $0.9 < H < 1.1$ around \mathbf{Q}_1 in Figs. 6(a), 6(c), 6(e), 6(g) and $0.9 < K < 1.1$ in Figs. 6(b), 6(d), 6(f), and 6(h) around \mathbf{Q}_2 . The effect of the partial detwinning ratio was corrected using the method developed in Ref. [33]. Consistent with Figs. 4 and 5, we find that superconductivity induces a broad resonance (or two resonances) around $E_r \approx 5$ meV at \mathbf{Q}_1 [Figs. 6(a) and 6(c)] [17,34–37], but has no effect at \mathbf{Q}_2 [Figs. 6(b) and 6(d)] [38]. On warming to 60 K, which is above the zero pressure T_N and T_S , there is still clear magnetic excitation anisotropy [Figs. 6(e) and 6(f)]. Finally, on warming to 100 K, the spin excitations at \mathbf{Q}_1 and \mathbf{Q}_2 become essentially identical with no observable anisotropy.

To accurately determine the dynamic magnetic susceptibility anisotropy, which is associated with spin nematic order [39–45] and may be important for superconductivity [46,47], we cut the data in Fig. 6 along the TOF direction, which couples the energy transfer of the spin excitations with L modulation. Figures 7(a)–7(d) show the temperature dependence of the magnetic scattering at \mathbf{Q}_1 and \mathbf{Q}_2 . Consistent with earlier work [13,14], we find that spin excitations at \mathbf{Q}_1 have a strong L modulation with high magnetic intensity at $L = 1, 3, \dots$ [Figs. 7(a)–7(c)]. The spin excitations have a




FIG. 8. (a), (b) The energy dependence of the resonance along the $\mathbf{Q} = [1, K]$ and $[H, 1]$ directions in $\text{Ba}(\text{Fe}_{0.946}\text{Co}_{0.054})_2\text{As}_2$. (c)–(h) The wave-vector dependence of the spin resonance at $E = 5 \pm 1, 9 \pm 1, 13 \pm 1$ meV around \mathbf{Q}_1 . The vertical and horizontal dashed lines indicate the in-plane momentum integration range used in (a) and (b).




FIG. 9. (a) Energy dependence of the resonance at \mathbf{Q}_1 and \mathbf{Q}_2 obtained by taking temperature differences of the constant- \mathbf{Q} cuts below and above T_c . The vertical dashed lines indicate the energy integration range in Figs. 8(c)–8(h). (b) In-plane wave-vector dependence of the magnetic scattering near the resonance energy around \mathbf{Q}_1 and \mathbf{Q}_2 .

gap of about 8 meV at \mathbf{Q}_2 below 23 K [Fig. 7(b)], and become gapless and similar to those at \mathbf{Q}_1 around 100 K [Fig. 7(d)].

The impact of superconductivity on the spin excitations at \mathbf{Q}_1 and \mathbf{Q}_2 can be further evaluated by the temperature differences plot below and above T_c . Figures 8(a) and 8(b) confirm that superconductivity induces a broad resonance at \mathbf{Q}_1 and has no effect at \mathbf{Q}_2 . The in-plane wave-vector dependence of the resonance at \mathbf{Q}_1 in Figs. 8(c), 8(e), and 8(g) reveals no strong evidence of an incommensurate dispersive resonance as seen in electron-doped $\text{Ba}(\text{Fe}_{0.963}\text{Ni}_{0.037})_2\text{As}_2$ [48] and hole-doped $\text{Ba}_{0.67}\text{K}_{0.33}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ with $x = 0$ and 0.08 [49], possibly due to the fact that $\text{Ba}(\text{Fe}_{0.946}\text{Co}_{0.054})_2\text{As}_2$ is still not close to optimal superconductivity, or our measurements have insufficient in-plane wave-vector resolution to resolve the expected transverse incommensurate scattering [48]. Figure 9(a) shows the energy dependence of the resonance at \mathbf{Q}_1 and \mathbf{Q}_2 , confirming the results of Figs. 8(a) and 8(b). The wave-vector dependence of the resonance at different energies is plotted in Fig. 9(b), which again reveals no evidence of incommensurate scattering.

In previous work on nearly optimally doped $\text{BaFe}_{1.9}\text{Ni}_{0.1}\text{As}_2$ [32] and BaFe_2As_2 [24], spin excitation anisotropy associated with the spin-driven Ising-nematic phase was defined as $\delta = (I_{10} - I_{01}) / (I_{10} + I_{01})$, where I_{10} and I_{01} are the inelastic magnetic scattering at \mathbf{Q}_1 and \mathbf{Q}_2 , respectively. Here δ is not related to the detwinning ratio η ,




FIG. 10. Temperature dependence of the $E = 5$ meV spin excitations in $\text{Ba}(\text{Fe}_{0.946}\text{Co}_{0.054})_2\text{As}_2$ measured with $E_i = 25$ meV at (a) 46 K, (b) 60 K, (c) 88 K, (d) 100 K. The scattering intensity has been corrected for the Bose population factor to obtain $\chi''(\mathbf{Q}, E)$.

which is determined from elastic magnetic Bragg peaks at \mathbf{Q}_1 and \mathbf{Q}_2 in a partially detwinned sample, but a measure of inelastic spin excitation intensity differences between \mathbf{Q}_1 and \mathbf{Q}_2 for a fully detwinned sample. For BaFe_2As_2 , the spin excitation anisotropy δ extends to about $E = 180$ meV at 7 K (at zero uniaxial pressure, $T_N \approx 138$ K), reduces to ~ 120 meV at 145 K (around pressure induced T_N) [22] and ~ 40 meV at 170 K (well above T_N under pressure), and finally becomes isotropic at 197 K [24]. In the case of $\text{BaFe}_{1.9}\text{Ni}_{0.1}\text{As}_2$, the spin excitation anisotropy δ is nonzero below about ~ 60 meV at 5 K (where $T_N \approx T_s \approx 30 \pm 5$ K) and unchanged on warming to 35 K [32]. Figure 10 shows the temperature dependence of the magnetic scattering at the resonance energy of $E_r \approx 5$ meV. On warming from 45 to 100 K, the scattering profile clearly changes from C_2 to C_4 symmetric and becomes isotropic.

To quantitatively determine the temperature/energy dependence of the spin excitation anisotropy δ in $\text{Ba}(\text{Fe}_{0.946}\text{Co}_{0.054})_2\text{As}_2$, we estimate the energy dependence of the dynamic susceptibility $\chi''(\mathbf{Q}, E)$, which can be calculated by $\chi''(\mathbf{Q}, E) \propto (1 - e^{-E/k_B T})I(\mathbf{Q}, E)$, at \mathbf{Q}_1 and \mathbf{Q}_2 as a function of increasing temperature. Figure 11(a) shows the energy dependence of $\chi''(\mathbf{Q}_1, E)$ and $\chi''(\mathbf{Q}_2, E)$ at $T = 4.5$ K, revealing magnetic anisotropy below about 30 meV. Each point of $\chi''(\mathbf{Q}, E)$ is obtained by integrating the magnetic scattering over wave vectors $-0.05 < H < 0.05$ and $-0.05 < K < 0.05$ around \mathbf{Q}_1 or \mathbf{Q}_2 . On warming to $T = 23$ K [Fig. 11(b)], 46 K [Fig. 11(c)], 60 K [Fig. 11(d)], 78 K [Fig. 11(e)], and 100 K [Fig. 11(f)], the spin excitation anisotropy gradually decreases and finally vanishes at 100 K. Figure 11(g) shows the energy dependence of the spin excitation anisotropy δ at different temperatures. Since $\text{Ba}(\text{Fe}_{0.946}\text{Co}_{0.054})_2\text{As}_2$ has $T_N \approx 38$ K similar to nearly optimal electron-doped $\text{BaFe}_{1.9}\text{Ni}_{0.1}\text{As}_2$ with $T_N \approx 30$ K [32], one would expect similar spin excitation anisotropy in $\text{Ba}(\text{Fe}_{0.946}\text{Co}_{0.054})_2\text{As}_2$ and $\text{BaFe}_{1.9}\text{Ni}_{0.1}\text{As}_2$, as confirmed by




FIG. 11. Energy dependence of $\chi''(E)$, where the in-plane momentum transfers are integrated around \mathbf{Q}_1 and \mathbf{Q}_2 , at temperatures (a) 4.6 K, (b) 23 K, (c) 46 K, (d) 60 K, (e) 78 K, (f) 100 K. The first two data points at low energies in each figure are measured with incident energy $E_i = 25$ meV and the rest are from $E_i = 80$ meV. The values of $\chi''(E)$ are obtained by fitting the transverse cuts with one Gaussian and linear background. The Gaussian intensity above background was then corrected by the magnetic form factor, Bose factor, and the partial detwinning ratio. (g) Temperature dependence of the spin excitation anisotropy between \mathbf{Q}_1 and \mathbf{Q}_2 . The first two data points collected using $E_i = 25$ meV are plotted together. The dashed lines are guides to the eye.

comparing Fig. 11(g) and Fig. 4 of Ref. [32]. The magnetic anisotropy is clearly present above the pressure induced T_N .

III. DISCUSSION AND CONCLUSION

To understand our neutron scattering results in terms of the electron-hole Fermi-surface nesting picture [50,51], we consider electron and hole Fermi surfaces of uniaxial pressure detwinned $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ above and below T_s as determined from ARPES measurements [25,26,29,30]. In the paramagnetic tetragonal state above T_s , the hole Fermi surfaces near Γ and Z points are composed of d_{xy} and degenerate d_{xz}/d_{yz}

orbitals, respectively. If low-energy spin excitations arise from quasiparticle excitations between the hole-electron Fermi surfaces as suggested in an itinerant picture of magnetism and superconductivity [50–52], the electron-hole Fermi surface nesting of the d_{yz} and d_{xz} orbital quasiparticles are along the \mathbf{Q}_1 and \mathbf{Q}_2 directions, respectively. Since a d_{xy} orbital has C_4 symmetry, it cannot by itself induce any anisotropic magnetic scattering through the hole-electron Fermi surface nesting along the \mathbf{Q}_1 and \mathbf{Q}_2 directions.

When we cool $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ below T_s in the orthorhombic nematic phase, the d_{yz} band of the electron Fermi surface at X/Y goes up in energy, while the d_{xz} band goes down in energy. Since these are electron pockets, the green part (the d_{yz} band) of the Fermi surface near the Fermi level will shrink in size, while the red part (the d_{xz} band) of the Fermi surface will expand, resulting in different shaped Fermi surfaces as shown in Fig. 1(c). At the Z point, which has hole pockets, the changes for the d_{yz} and d_{xz} bands are opposite with much smaller amplitude [29,30]. So, we can basically assume that the holelike Fermi surfaces are not modified much below T_s . If spin fluctuations arise from intraorbital but interband quasiparticle excitations between hole and electron Fermi surfaces [53], spin fluctuations along \mathbf{Q}_1 should arise mostly from the d_{yz} band scattering between the Z and X points. Similarly, one would expect spin fluctuations along the \mathbf{Q}_2 direction to arise mostly from the d_{xz} band scattering between the Z and Y points [Figs. 1(b) and 1(c)]. In the high-temperature paramagnetic tetragonal phase, the d_{yz} and d_{xz} orbital Fermi surfaces are degenerate, resulting in identical shapes for the electron Fermi pockets at the X and Y points, and an isotropic hole Fermi surface at the Z point [Fig. 1(b)]. The quasiparticle scattering across the hole-electron Fermi pockets along the \mathbf{Q}_1 and \mathbf{Q}_2 directions and associated spin fluctuations therefore have the same scattering intensity and behave identically.

On cooling to below T_s , the lifting of the d_{yz} band makes the electron Fermi pocket at the X point to be better matched with the d_{yz} orbital in the hole pocket, and the reduction in the d_{xz} band enhances the oval shape of the electron Fermi pocket at Y point as shown in Fig. 1(c). At the Z point, the hole Fermi surfaces also change line shape due to the rising d_{xz} band and the reduction of the d_{yz} band, but to a much smaller extent compared with the shifts in Fermi surfaces at the X/Y points [Fig. 1(c)] [29,30]. Therefore, the major effect of the tetragonal-to-orthorhombic lattice distortion and associated nematic phase is to change the shapes of the electron Fermi pockets at the X and Y points as shown in Fig. 1(c). From a pure hole-electron Fermi surface nesting point of view [50], the nesting condition along the \mathbf{Q}_1 direction improves below T_s because of the better matched hole-electron Fermi surfaces of the d_{yz} band [Fig. 1(c)]. On the other hand, the ~ 30 meV downward shift of the d_{xz} band below T_s at the Y point enlarges the electron pocket along the \mathbf{Q}_1 direction [25] and thus makes the d_{xz} - d_{xz} hole-electron Fermi surface nesting along the \mathbf{Q}_2 direction less favorable.

If we assume that low-energy spin fluctuations in $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ arise from quasiparticle excitations between hole-electron Fermi pockets at the Z and X/Y points, spin fluctuations along the \mathbf{Q}_1 and \mathbf{Q}_2 directions should be sensitive to the nesting condition associated with the splitting

energy between the d_{xz} and d_{yz} bands below T_s , and become C_4 rotational symmetric above T_s . Since the splitting energy between the d_{xz} and d_{yz} bands is around 60 meV in undoped BaFe_2As_2 , decreases to about 30 meV in $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ with $x = 0.045$, and vanishes around optimal superconductivity [25], the energy scale of the spin fluctuation anisotropy δ should decrease with increasing x and vanish near optimal superconductivity. This is qualitatively consistent with the spin anisotropy results on BaFe_2As_2 [24], $\text{BaFe}_{1.9}\text{Ni}_{0.1}\text{As}_2$ [32], and $\text{Ba}(\text{Fe}_{0.946}\text{Co}_{0.054})_2\text{As}_2$ [Fig. 11(g)], suggesting that low-energy spin fluctuations at the wave vector \mathbf{Q}_1 have a strong d_{yz} orbital character and arise from the d_{yz} - d_{yz} hole-electron Fermi surface quasiparticle excitations. Since superconductivity-induced neutron spin resonance in underdoped $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ only appears at \mathbf{Q}_1 (Figs. 3, 8, and 9), it is tempting to argue that superconductivity in these materials arises mostly from electrons with d_{yz} orbital characters [54]. However, such a picture is strictly only true within the itinerant model of magnetism and superconductivity in iron pnictides [54].

The above discussion centers on the assumption that energy splitting of the d_{xz} and d_{yz} bands originates from orbital/nematic ordering below T_s . In a recent ARPES work on BaFe_2As_2 [30], it was argued that the splitting of the d_{xz} and d_{yz} bands is induced not by orbital/nematic order at T_s , but by static AF order occurring at a temperature T_N just below T_s [55]. In this scenario, spin fluctuations occurring at \mathbf{Q}_1 in the AF-ordered state originate from spin waves of static ordered moments. The presence of a large spin gap at \mathbf{Q}_2 [24] and effective magnetic exchange coupling anisotropy [56] can be well understood by including a biquadratic coupling term in the local moment Heisenberg Hamiltonian [57–59]. In the underdoped regime where superconductivity coexists with AF order, the broad (or double) resonance mode seen in neutron scattering experiments of twinned iron pnictides [17,34–37] may arise from interacting spin waves with itinerant electrons [60]. In this picture, the resonance associated with the AF order should exclusively appear at \mathbf{Q}_1 , while the resonance associated with itinerant electrons and simple nested Fermi surfaces should appear at both \mathbf{Q}_1 and \mathbf{Q}_2 [60]. Our results in Figs. 8 and 9 clearly disagree with this picture.

Alternatively, the neutron spin resonance [17,34–37] can arise from orbital-selective pairing-induced superconducting gap anisotropy [61]. Here, the broadening of the resonance is a consequence of anisotropic superconducting gap in the electron pockets at the X and Y points. Below T_s , the unfavorable nesting condition of the d_{xz} band along the \mathbf{Q}_2 means low-energy spin excitations are gapped at \mathbf{Q}_2 . Therefore,

the appearance of the resonance exclusively at \mathbf{Q}_1 suggests that superconducting electrons have mostly the d_{yz} orbital characters below the nematic ordering temperature T_s . In a recent work on detwinned FeSe [33], which has no static AF order below T_s [62], we again find that the superconductivity-induced resonance only appears at \mathbf{Q}_1 , consistent with orbital-selective Cooper pairing in FeSe seen by scanning tunneling microscopy experiments [63]. This further supports the notion that orbital order and the nematic phase below T_s induce the energy splitting of the d_{xz} and d_{yz} bands in electron pockets, which in turn modifies the Fermi-surface nesting condition and associated spin fluctuations along the \mathbf{Q}_1 and \mathbf{Q}_2 directions.

In conclusion, our inelastic neutron scattering experiments on mechanically detwinned $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ with $x = 0.048$ and 0.054 , which has coexisting AF order and superconductivity, reveal highly anisotropic spin fluctuations with large magnetic scattering intensity at the AF ordering wave vector \mathbf{Q}_1 and weak scattering at \mathbf{Q}_2 at temperatures below the tetragonal-to-orthorhombic structural transition T_s . On cooling to a temperature above T_c but below T_N , a large spin gap appears at the \mathbf{Q}_2 point and spin fluctuations are mostly centered at the \mathbf{Q}_1 point. Upon entering the superconducting state, a neutron spin resonance appears at the \mathbf{Q}_1 point with no magnetic scattering at the $\mathbf{Q}_2 = (0, 1)$ point. By comparing these results with those from ARPES experiments, we conclude that the anisotropic shift of the d_{yz} and d_{xz} electronlike bands in detwinned $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ below T_s is associated with the spin excitation anisotropy, and the superconductivity-induced resonance arises from itinerant electrons with the d_{yz} orbital characters. Therefore, low-energy spin fluctuations in underdoped $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$ are highly orbital selective below T_s , suggesting that the orbital order and the nematic phase are correlated with spin fluctuations and superconductivity in underdoped iron pnictide superconductors.

ACKNOWLEDGMENTS

We are grateful to M. Yi for helpful discussions and detailed explanation of the ARPES results on $\text{Ba}(\text{Fe}_{1-x}\text{Co}_x)_2\text{As}_2$. We thank Y. Song for a critical reading of the paper and S. Li for allowing us to use his laboratory's equipment where some of the single crystals were grown. The work at Beijing Normal University is supported by the Fundamental Research Funds for the Central Universities (Grants No. 310432101 and No. 2014JJCB27) and the National Natural Science Foundation of China (Grant No. 11734002). The neutron scattering work at Rice is supported by the U. S. NSF Grant No. DMR-1700081 (P.D.).

-
- [1] D. J. Scalapino, *Rev. Mod. Phys.* **84**, 1383 (2012).
 [2] E. Fradkin, S. A. Kivelson, and J. M. Tranquada, *Rev. Mod. Phys.* **87**, 457 (2015).
 [3] M. Eschrig, *Adv. Phys.* **55**, 47 (2006).
 [4] P. C. Dai, *Rev. Mod. Phys.* **87**, 855 (2015).
 [5] M. Kenzelmann, *Rep. Prog. Phys.* **80**, 034501 (2017).

- [6] G. R. Stewart, *Rev. Mod. Phys.* **83**, 1589 (2011).
 [7] N. Ni, M. E. Tillman, J.-Q. Yan, A. Kracher, S. T. Hannahs, S. L. Bud'ko, and P. C. Canfield, *Phys. Rev. B* **78**, 214515 (2008).
 [8] C. Lester, J.-H. Chu, J. G. Analytis, S. C. Capelli, A. S. Erickson, C. L. Condon, M. F. Toney, I. R. Fisher, and S. M. Hayden, *Phys. Rev. B* **79**, 144523 (2009).

- [9] D. K. Pratt, W. Tian, A. Kreyssig, J. L. Zarestky, S. Nandi, N. Ni, S. L. Bud'ko, P. C. Canfield, A. I. Goldman, and R. J. McQueeney, *Phys. Rev. Lett.* **103**, 087001 (2009).
- [10] A. D. Christianson, M. D. Lumsden, S. E. Nagler, G. J. MacDougall, M. A. McGuire, A. S. Sefat, R. Jin, B. C. Sales, and D. Mandrus, *Phys. Rev. Lett.* **103**, 087002 (2009).
- [11] S. Nandi, M. G. Kim, A. Kreyssig, R. M. Fernandes, D. K. Pratt, A. Thaler, N. Ni, S. L. Budko, P. C. Canfield, J. Schmalian, R. J. McQueeney, and A. I. Goldman, *Phys. Rev. Lett.* **104**, 057006 (2010).
- [12] H. Luo, X. Lu, R. Zhang, M. Wang, E. A. Goremychkin, D. T. Adroja, S. Danilkin, G. Deng, Z. Yamani, and P. Dai, *Phys. Rev. B* **88**, 144516 (2013).
- [13] M. D. Lumsden, A. D. Christianson, D. Parshall, M. B. Stone, S. E. Nagler, G. J. MacDougall, H. A. Mook, K. Lokshin, T. Egami, D. L. Abernathy, E. A. Goremychkin, R. Osborn, M. A. McGuire, A. S. Sefat, R. Jin, B. C. Sales, and D. Mandrus, *Phys. Rev. Lett.* **102**, 107005 (2009).
- [14] S. Chi, A. Schneidewind, J. Zhao, L. W. Harriger, L. Li, Y. Luo, G. Cao, Z. Xu, M. Loewenhaupt, J. Hu, and P. C. Dai, *Phys. Rev. Lett.* **102**, 107006 (2009).
- [15] S. Li, Y. Chen, S. Chang, J. W. Lynn, L. Li, Y. Luo, G. Cao, Z. Xu, and P. C. Dai, *Phys. Rev. B* **79**, 174527 (2009).
- [16] M. Wang, C. Zhang, X. Lu, G. Tan, H. Luo, Y. Song, M. Wang, X. Zhang, E. A. Goremychkin, T. G. Perring, T. A. Maier, Z. Yin, K. Haule, G. Kotliar, and P. Dai, *Nat. Commun.* **4**, 2874 (2013).
- [17] M. Wang, M. Yi, H. L. Sun, P. Valdivia, M. G. Kim, Z. J. Xu, T. Berlijn, A. D. Christianson, S. Chi, M. Hashimoto, D. H. Lu, X. D. Li, E. Bourret-Courchesne, P. Dai, D. H. Lee, T. A. Maier, and R. J. Birgeneau, *Phys. Rev. B* **93**, 205149 (2016).
- [18] M. Wang, H. Luo, M. Wang, S. Chi, J. A. Rodriguez-Rivera, D. Singh, S. Chang, J. W. Lynn, and P. Dai, *Phys. Rev. B* **83**, 094516 (2011).
- [19] C. Dhital, Z. Yamani, W. Tian, J. Zeretsky, A. S. Sefat, Z. Wang, R. J. Birgeneau, and S. D. Wilson, *Phys. Rev. Lett.* **108**, 087001 (2012).
- [20] C. Dhital, T. Hogan, Z. Yamani, R. J. Birgeneau, W. Tian, M. Matsuda, A. S. Sefat, Z. Wang, and S. D. Wilson, *Phys. Rev. B* **89**, 214404 (2014).
- [21] Y. Song, S. V. Carr, X. Y. Lu, C. L. Zhang, Z. C. Sims, N. F. Luttrell, S. X. Chi, Y. Zhao, J. W. Lynn, and P. C. Dai, *Phys. Rev. B* **87**, 184511 (2013).
- [22] D. W. Tam, Y. Song, H. Man, S. C. Cheung, Z. Yin, X. Lu, W. Wang, B. A. Frandsen, L. Liu, Z. Gong, T. U. Ito, Y. Cai, M. N. Wilson, S. Guo, K. Koshiishi, W. Tian, B. Hitti, A. Ivanov, Y. Zhao, J. W. Lynn *et al.*, *Phys. Rev. B* **95**, 060505(R) (2017).
- [23] X. Lu, J. T. Park, R. Zhang, H. Luo, A. H. Nevidomskyy, Q. Si, and P. C. Dai, *Science* **345**, 657 (2014).
- [24] X. Lu, D. D. Scherer, D. W. Tam, W. Zhang, R. Zhang, H. Luo, L. W. Harriger, H. C. Walker, D. T. Adroja, B. M. Andersen, and P. C. Dai, *Phys. Rev. Lett.* **121**, 067002 (2018).
- [25] M. Yi, D. Lu, J.-H. Chu, J. G. Analytis, A. P. Sorini, A. F. Kemper, B. Moritz, S.-K. Mo, R. G. Moore, M. Hashimoto, W.-S. Lee, Z. Hussain, T. P. Devereaux, I. R. Fisher, and Z.-X. Shen, *Proc. Natl. Acad. Sci. USA* **108**, 6878 (2011).
- [26] Y. Zhang, F. Chen, C. He, B. Zhou, B. P. Xie, C. Fang, W. F. Tsai, X. H. Chen, H. Hayashi, J. Jiang, H. Iwasawa, K. Shimada, H. Namatame, M. Taniguchi, J. P. Hu, and D. L. Feng, *Phys. Rev. B* **83**, 054510 (2011).
- [27] V. Brouet, M. F. Jensen, P.-H. Lin, A. Taleb-Ibrahimi, P. Le Fèvre, F. Bertran, C.-H. Lin, W. Ku, A. Forget, and D. Colson, *Phys. Rev. B* **86**, 075123 (2012).
- [28] M. Yi, Y. Zhang, Z.-X. Shen, and D. H. Lu, *npj Quantum Mater.* **2**, 57 (2017).
- [29] H. Pfau, C. R. Rotundu, J. C. Palmstrom, S. D. Chen, M. Hashimoto, D. Lu, A. F. Kemper, I. R. Fisher, and Z.-X. Shen, *Phys. Rev. B* **99**, 035118 (2019).
- [30] M. D. Watson, P. Dudin, Luke C. Rhodes, D. V. Evtushinsky, H. Iwasawa, S. Aswartham, S. Wurmehl, B. Büchner, M. Hoesch, and T. K. Kim, *npj Quantum Mater.* **4**, 36 (2019).
- [31] X. Lu, Orbital selective neutron spin resonance in underdoped $\text{Ba}(\text{Fe}_{0.952}\text{Co}_{0.048})_2\text{As}_2$, STFC ISIS Neutron and Muon Source, [10.5286/ISIS.E.90683198](https://doi.org/10.5286/ISIS.E.90683198) (2018).
- [32] Y. Song, X. Lu, D. L. Abernathy, D. W. Tam, J. L. Niedziela, W. Tian, H. Luo, Q. Si, and P. Dai, *Phys. Rev. B* **92**, 180504(R) (2015).
- [33] T. Chen, Y. Chen, A. Kreisel, X. Lu, A. Schneidewind, Y. Qiu, J. T. Park, T. G. Perring, J. R. Stewart, H. Cao, R. Zhang, Y. Li, Y. Rong, Y. Wei, B. M. Andersen, P. J. Hirschfeld, C. Broholm, and P. Dai, *Nat. Mater.* **18**, 709 (2019).
- [34] C. Zhang, R. Yu, Y. Su, Y. Song, M. Wang, G. Tan, T. Egami, J. A. Fernandez-Baca, E. Faulhaber, Q. Si, and P. C. Dai, *Phys. Rev. Lett.* **111**, 207002 (2013).
- [35] P. Steffens, C. H. Lee, N. Qureshi, K. Kihou, A. Iyo, H. Eisaki, and M. Braden, *Phys. Rev. Lett.* **110**, 137001 (2013).
- [36] F. Waßer, C. H. Lee, K. Kihou, P. Steffens, K. Schmalzl, N. Qureshi, and M. Braden, *Sci. Rep.* **7**, 10307 (2017).
- [37] C. Zhang, W. Lv, G. Tan, Y. Song, S. V. Carr, S. Chi, M. Matsuda, A. D. Christianson, J. A. Fernandez-Baca, L. W. Harriger, and P. Dai, *Phys. Rev. B* **93**, 174522 (2016).
- [38] W. Wang, J. T. Park, R. Yu, Y. Li, Y. Song, Z. Zhang, A. Ivanov, J. Kulda, and P. Dai, *Phys. Rev. B* **95**, 094519 (2017).
- [39] C. Fang, H. Yao, W.-F. Tsai, J. P. Hu, and S. A. Kivelson, *Phys. Rev. B* **77**, 224509 (2008).
- [40] C. Xu, M. Müller, and S. Sachdev, *Phys. Rev. B* **78**, 020501(R) (2008).
- [41] R. M. Fernandes, A. V. Chubukov, and J. Schmalian, *Nat. Phys.* **10**, 97 (2014).
- [42] R. M. Fernandes, E. Abrahams, and J. Schmalian, *Phys. Rev. Lett.* **107**, 217002 (2011).
- [43] R. M. Fernandes and J. Schmalian, *Supercond. Sci. Technol.* **25**, 084005 (2012).
- [44] S. Liang, A. Moreo, and E. Dagotto, *Phys. Rev. Lett.* **111**, 047004 (2013).
- [45] Q. Si, R. Yu, and E. Abrahams, *Nat. Rev. Mater.* **1**, 16017 (2016).
- [46] M. A. Metlitski, D. F. Mross, S. Sachdev, and T. Senthil, *Phys. Rev. B* **91**, 115111 (2015).
- [47] S. Lederer, Y. Schattner, E. Berg, and S. A. Kivelson, *Phys. Rev. Lett.* **114**, 097001 (2015).
- [48] M. G. Kim, G. S. Tucker, D. K. Pratt, S. Ran, A. Thaler, A. D. Christianson, K. Marty, S. Calder, A. Podlesnyak, S. L. Bud'ko, P. C. Canfield, A. Kreyssig, A. I. Goldman, and R. J. McQueeney, *Phys. Rev. Lett.* **110**, 177002 (2013).
- [49] R. Zhang, W. Wang, T. A. Maier, M. Wang, M. B. Stone, S. Chi, B. Winn, and P. Dai, *Phys. Rev. B* **98**, 060502(R) (2018).
- [50] P. J. Hirschfeld, M. M. Korshunov, and I. I. Mazin, *Rep. Prog. Phys.* **74**, 124508 (2011).

- [51] A. Chubukov, *Annu. Rev. Condens. Phys.* **3**, 57 (2012).
- [52] P. C. Dai, J. P. Hu, and E. Dagotto, *Nat. Phys.* **8**, 709 (2012).
- [53] J. Zhang, R. Sknepnek, and J. Schmalian, *Phys. Rev. B* **82**, 134527 (2010).
- [54] P. J. Hirschfeld, *C. R. Phys.* **17**, 197 (2016).
- [55] M. G. Kim, R. M. Fernandes, A. Kreyssig, J. W. Kim, A. Thaler, S. L. Bud'ko, P. C. Canfield, R. J. McQueeney, J. Schmalian, and A. I. Goldman, *Phys. Rev. B* **83**, 134522 (2011).
- [56] L. W. Harriger, H. Q. Luo, M. S. Liu, C. Frost, J. P. Hu, M. R. Norman, and P. Dai, *Phys. Rev. B* **84**, 054544 (2011).
- [57] A. L. Wysocki, K. D. Belashchenko, and V. P. Antropov, *Nat. Phys.* **7**, 485 (2011).
- [58] D. Stanek, O. P. Sushkov, and G. S. Uhrig, *Phys. Rev. B* **84**, 064505 (2011).
- [59] R. Yu, Z. Wang, P. Goswami, A. H. Nevidomskyy, Q. Si, and E. Abrahams, *Phys. Rev. B* **86**, 085148 (2012).
- [60] W. Lv, A. Moreo, and E. Dagotto, *Phys. Rev. B* **89**, 104510 (2014).
- [61] R. Yu, J. X. Zhu, and Q. Si, *Phys. Rev. B* **89**, 024509 (2014).
- [62] A. E. Böhmer and A. Kreisel, *J. Phys.: Condens. Matter* **30**, 023001 (2018).
- [63] P. O. Sprau, A. Kostin, A. Kreisel, A. E. Böhmer, V. Taufour, P. C. Canfield, S. Mukherjee, P. J. Hirschfeld, B. M. Andersen, and J. C. Séamus Davis, *Science* **357**, 75 (2017).