



This is a repository copy of *Improving outcomes for older people in the emergency department : a review of reviews*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/167355/>

Version: Accepted Version

Article:

Preston, L. orcid.org/0000-0001-7477-4517, van Oppen, J.D., Conroy, S.P. et al. (3 more authors) (2021) *Improving outcomes for older people in the emergency department : a review of reviews*. *Emergency Medicine Journal*, 38 (12). pp. 882-888. ISSN 1472-0205

<https://doi.org/10.1136/emered-2020-209514>

© 2020 The Author(s). This is an author-produced version of a paper subsequently published in *Emergency Medicine Journal (EMJ)*. Reuse of this manuscript version (excluding any databases, tables, diagrams, photographs and other images or illustrative material included where a another copyright owner is identified) is permitted strictly pursuant to the terms of the Creative Commons Attribution-NonCommercial 4.0 International (<https://creativecommons.org/licenses/by-nc/4.0/>).

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial (CC BY-NC) licence. This licence allows you to remix, tweak, and build upon this work non-commercially, and any new works must also acknowledge the authors and be non-commercial. You don't have to license any derivative works on the same terms. More information and the full terms of the licence here:
<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Title

Improving outcomes for older people in the Emergency Department. A review of reviews

Authors

Louise Preston, James D van Oppen, Simon Conroy, Suzanne Ablard, Helen Buckley Woods and Suzanne Mason.

Corresponding Author

Dr Louise Preston, School of Health and Related Research, University of Sheffield, Western Bank, Sheffield, S1 4DA.
+441142220847, +441142220749

Other Authors

Dr James D van Oppen, Department of Health Sciences, University of Leicester, Leicester, United Kingdom

Professor Simon Conroy, Department of Health Sciences, University of Leicester, Leicester, United Kingdom

Ms Suzanne Ablard, School of Health and Related Research, University of Sheffield, Sheffield, United Kingdom

Dr Helen Buckley Woods, Information School, University of Sheffield, Sheffield, United Kingdom
Professor Suzanne Mason, School of Health and Related Research, University of Sheffield, Sheffield, United Kingdom

Grant

This research was funded by the National Institute for Health Research, Health Services and Delivery Research - 17/05/96, £931,653.

Word Count

2997 words (excluding title page, abstract, references, figures and tables)

Author Contributions

SC and SM conceptualised the study, SC, SM and LP obtained research funding for the study. LP, SC, SM, JvO, SA and HW developed the protocol. HW ran the literature searches. LP, SA, SC and JvO undertook screening of the included studies, data extraction and quality assessment. LP drafted the manuscript and all authors contributed substantially to its development and revision.

Keywords

emergency care systems, emergency departments, geriatrics

Disclaimer

"The views expressed are those of the author(s) and not necessarily of the NIHR or the Department of Health and Social Care"

Conflicts of Interest

None declared

Abstract

Background

There has been a recognised trend of increasing use of Emergency and Urgent Care (EUC) and Emergency Departments (EDs) by older people, which is marked by a substantial evidence base reporting interventions for this population and guidance from key organisations. Despite this, outcomes for this population remain suboptimal. A plethora of reviews in this area provides challenges for clinicians and commissioners in determining which interventions and models of care

best meet people's needs. The aim of this review was to identify effective ED interventions which have been reported for older people, and to provide a clear summary of the myriad reviews and numerous intervention types in this area.

Methods

A review of reviews, reporting interventions for older people, either initiated or wholly delivered within the ED.

Results

A total of 15 review articles describing 83 primary studies met our content and reporting standards criteria. The majority (n=13) were systematic reviews (four using meta-analysis.) Across the reviews, 26 different outcomes were reported with inconsistency. Follow up duration varied within and across the reviews. Based on how authors had reported results, evidence clusters were developed:

- (1) Staff-focused reviews
- (2) Discharge intervention reviews
- (3) Population-focused reviews
- (4) Intervention component review

Conclusions

The evidence base describing interventions is weak due to inconsistent reporting, differing emphasis placed on the key characteristics of primary studies (staff, location, outcome) by review authors and varying quality of reviews. No individual interventions have been found to be more promising, but interventions initiated in the ED and continued into other settings have tended to result in more favourable patient and health service outcomes. Despite many interventions reported within the reviews being holistic and patient focused, outcomes measured were largely service focused.

Introduction

Background

Use of Emergency and Urgent Care (EUC) and Emergency Departments (EDs) by older people is increasing. This has been variously attributed to the complex physical, social and mental health comorbidities that older people often live with, changes in the healthcare options available to patients, professional opinions on appropriate treatment, and the capacity of individual care systems. Caring for those older people living with frailty presents an urgent national and international public health issue. Despite guidance developed by organisations such as the British Geriatrics Society, the Royal College of Emergency Medicine, the American College of Emergency Physicians, and the International Federation of Emergency Medicine, fast-flowing EUC systems are yet to fully integrate person-centred case management designed to best meet the needs of older people (1). Existing reviews report a large body of evidence describing interventions for this population. However, there is a need to identify consistent messages around proposed approaches to older people's care in the ED, to ensure that care is sensitive, effective and efficient, encompassing individuals' clinical and wider social needs. This study aimed to review the evidence for ED interventions for older people and the characteristics of that evidence base, in particular the overlap in primary study coverage between reviews, the outcomes reported within reviews, and the consistency of intervention reporting. The study aimed to identify whether the literature demonstrated any evidence of intervention effectiveness and to identify which ED interventions best meet the needs of older people.

Methods

A review of reviews ('overview') of systematic and non-systematic reviews was undertaken, including both qualitative and quantitative studies. This method was chosen due to the number of existing reviews in this field (2, 3). Reviews of reviews offer benefits in that they "enable broader evidence synthesis questions to be addressed...in a faster timeframe" (4). In October 2018, the

protocol was published on PROSPERO (the international prospective register of systematic reviews) (CRD42018111461). No ethical approval was required for this review. The review was conducted and reported according to PRISMA standards (5).

Inclusion criteria

Reviews were appraised for eligibility against pre-defined inclusion criteria. Criteria for reporting standards were based on the Cochrane Handbook definition of a systematic review and criteria developed by Brunton (6). Reviews which met all of the inclusion criteria and three or more reporting standards criteria were included. These criteria are included in Supplementary Material 3.

Search approach

A comprehensive database search used existing strategies (7) combining terms for Emergency and Urgent Care (EUC) and for older people, limited by publication type (reviews), language (English Language studies only) and date (2000-2018). The search strategy for Medline (OVID SP) was developed by an information specialist and is reproduced in Supplementary Material. This was adapted for other databases: Embase (OVID SP), CINAHL (EBSCO), Cochrane Library (Wiley Interscience), Web of Science Core Collection (Clarivate), SCOPUS (Elsevier), and AgeINFO (<http://www.cpa.org.uk/>). Further review sources were searched using an adapted database strategy: Joanna Briggs Institute (<https://journals.lww.com/jbisrir/Pages/default.aspx>), the Campbell Collaboration (<https://campbellcollaboration.org/>), Epistemonikos (<https://www.epistemonikos.org/>) and PROSPERO (<https://www.crd.york.ac.uk/PROSPERO/>). In addition, we undertook forward and backward searching of included reviews using reference lists and Google Scholar for citation identification. Topic experts were consulted to identify missing reviews. A search alert was set up to identify additional reviews published following the database searches.

Study selection

References were managed in Endnote Version 8. Duplicates were removed prior to screening for inclusion at title and abstract level. This was undertaken by one first reviewer (JvO or LP), with 50% from each first reviewer also screened by a second reviewer (SA). All remaining potentially eligible reviews were double screened at full text by LP and SC. Reasons for excluding reviews were recorded. Figure 1 presents a PRISMA-SR flowchart of searching and study selection.

Data extraction and quality assessment

A data extraction sheet was designed in Microsoft Excel by LP and iteratively refined following piloting by SA and JvO. Data were single extracted by one reviewer (LP, SA or JvO). LP subsequently checked all extractions and a random sample of 10% were also checked by SA. Data were extracted on review type, review methods, description of included studies, all reported outcomes (including whether they had been synthesised or reported as individual studies) and a headline message or conclusion. We used the AMSTAR2 checklist (Supplementary material 1) to assess the quality of reviews. AMSTAR2 allows the appraisal of reviews that include non-randomised studies of interventions, in addition to randomised controlled trials (8). The findings from our quality assessment are reported narratively.

Overlap within reviews

A citation matrix was drawn up (9). This matrix assessed overlap in the evidence base by mapping each included review against all cited primary studies.

Data synthesis

Extracted data were summarised and presented in tables with a narrative synthesis. Due to the heterogeneity between reviews, no further statistical synthesis was undertaken.

PPI

This review of reviews formed part of a larger study of emergency care for older people. The wider project team included two lay collaborators, who advised on the research including this review of reviews.

Results

Overview

A total of 806 articles were retrieved from the database searches . From these, 15 eligible reviews were identified, published 2005-2019. These 15 reviews reported 83 unique primary studies (published 1994-2018). Of these 83 studies, 25 were included in more than one review, with the most frequently cited primary study included in 11 reviews (10). The review characteristics are presented in Table 1. Quality assessment is summarised in Supplementary Material 2.




Figure 1 PRISMA-SR flowchart

Table 1 Review characteristics

Author (date)	Topic	Review methods	Evidence synthesis method	Study types included in the review	Number of primary studies included in the review	Evidence cluster
Conroy (2011) (11)	Rapid-access nurse-led/geriatrician supported assessment with comprehensive geriatric assessment for patients post emergency department.	Systematic review	Quantitative synthesis including meta-analysis	RCTs	5	Discharge intervention reviews
Fan (2015) (12)	Strategies for older people in the emergency department	Literature review	Narrative synthesis of quantitative data	Any experimental or observational study.	20	Intervention component reviews
Fealy (2009) (13)	Nursing assessment and referral interventions	Systematic review	Narrative synthesis of quantitative data	Clinical trials, before-and-after designs, and descriptive-evaluative studies.	11	Staff focused reviews
Graf (2011) (14)	Comprehensive Geriatric Assessment interventions delivered by nurses in the Emergency Department	Systematic review	Narrative synthesis of quantitative data	RCTs or matched controlled trials (cohort, case control, case matched and cross sectional).	8	Staff focused reviews
Hastings (2005) (15)	Discharge interventions (all staff).	Systematic review	Narrative synthesis of quantitative data	RCTs, non-randomised CTs and observational studies.	14	Discharge intervention reviews
Hughes (2019)	Interventions according to strategy type, single or multi strategy, intervention components.	Systematic review	Narrative synthesis and meta-analysis	Randomised or quasi-experimental study types.	15	Intervention component reviews

Author (date)	Topic	Review methods	Evidence synthesis method	Study types included in the review	Number of primary studies included in the review	Evidence cluster
(16)						
Jay (2017) (17)	Consultant led comprehensive geriatric assessment.	Systematic review	Narrative synthesis of quantitative data	Randomised control trials, non-randomised controlled trials and observational studies.	5	Staff focused reviews
Karam (2015) (18)	Nurse/social worker / geriatrician led integrated (discharge) assessment interventions	Systematic review	Narrative synthesis of quantitative data	Studies with a comparison group.	9	Discharge intervention reviews
Lowthian (2015) (19)	Emergency Department to Community Transition Strategies	Systematic review and meta-analysis	Quantitative synthesis of quantitative data including meta-analysis	RCT and other quantitative studies.	11	Discharge intervention reviews
Malik (2018) (20)	Nurse-led emergency department based comprehensive geriatric assessment	Systematic review	Quantitative synthesis of quantitative data including meta-analysis	Quantitative research consisting of randomized control trials (RCTs), multicentre and observational studies.	9	Staff focused reviews

Author (date)	Topic	Review methods	Evidence synthesis method	Study types included in the review	Number of primary studies included in the review	Evidence cluster
McCusker (2006) (21)	Comprehensive geriatric assessment	Systematic review	Narrative synthesis of quantitative data	RCTs, non-randomised trials, before and after studies and cross sectional studies.	8	Discharge intervention reviews
Parke (2011) (22)	Prevalence and identification of cognitive impairment in emergency department (comprehensive geriatric assessment, training).	Scoping review	Narrative synthesis of quantitative data	Systematic review; meta-analysis; clinical trial; cohort study; evaluation study.	15	Population focused reviews
Pearce (2011) (23)	Nursing interventions in the emergency department to enhance older peoples comfort.	Systematic review	Narrative synthesis of quantitative data and meta synthesis of qualitative data	Quantitative research study designs as well as narrative opinion and text.	2	Staff focused reviews
Schnitker (2013) (24)	Screening, preventing, and managing of cognitive impairment	Systematic literature review	Narrative synthesis of quantitative data	"Research based literature".	12	Population focused reviews
Sinha (2011) (25)	Emergency Department based case management models	Systematic review	Narrative synthesis of quantitative data	RCT, non randomised trials, observational studies and program descriptions.	20	Intervention component reviews

Population definitions

Most reviews defined older people as being aged over 65, although some (13) and (21) included papers with populations aged 60 and older. Some reviews did not report a specific age, but rather reported interventions for participants who were ‘older’ or ‘elderly’. The majority of reviews reported ED care for a general population of older people, who were not stratified by condition or severity. However, Lowthian *et al.* (19) included a population of ‘high risk’ participants; this may indicate that there was some prior screening of patients before they were included in the intervention.

Outcome measures

Table 2 lists all outcomes reported in the 15 included reviews, organised according to Parker et al (26). There was inconsistency in the reporting of outcomes between reviews. Some reviews synthesised papers by outcome measures. There was a clear preference to measure outcomes in terms of service delivery metrics as opposed to patient centred outcomes. Other reviews did not synthesise outcomes across included studies but reported these narratively on an article-by-article basis. There was a high level of variability in the length of patient follow up from 0 days to 18 months. To some extent this depended on whether the intervention was wholly delivered in the ED or continued into other settings.

Table 2 Outcomes

Clinical outcomes	Cognition/cognitive decline Comfort Functional decline (Instrumental) Activities of Daily Living Medication adherence Mortality Use of advance directives
Other psychosocial outcomes	Quality of life Wellbeing
Operational outcomes	ED cost per patient ED length of stay ED readmission/return visit ED use ED utilisation ED visits per patient Hospital days (ED and inpatient) Inpatient admission Inpatient length of stay
Destinational outcomes	Care/nursing home admissions Community service referral rates Home care referral rates
Other outcomes	Carer satisfaction ED care provider satisfaction Patient satisfaction with care received Patient satisfaction with information received Primary care provider satisfaction

Intervention classification

Reviews of ED interventions were organised into four evidence clusters: discharge intervention reviews (11, 15, 18, 19, 21), staff-focused reviews (13, 14, 17, 20, 23), population focused reviews (22, 24), and intervention component reviews (12, 16, 24). Only 5 of the 15 reviews reported interventions delivered wholly within the ED – the remainder were continued into other settings.

Table 3 – Intervention cluster characteristics

Cluster	Reviews included	Range of primary studies included	Total number of primary studies included	Number of primary studies appearing in more than one review	Primary studies publication dates
Discharge intervention reviews	(11, 15, 18, 19, 21)	5-14	25	9	1996-2013
Staff-focused reviews	(13, 14, 17, 20, 23)	2-11	15	9	1996-2015
Population-focused reviews	(22, 24)	12-15	27	4	1994-2011
Intervention component reviews	(12, 16, 25)	15-20	38	13	1994-2018

* A subset of papers in the following reviews were reported (12, 14, 18, 21, 24).

Discharge intervention reviews

The reported discharge interventions included post-discharge follow up of patients by ED or community-based care professionals, although these were often reported incompletely.

Conroy et al (11) reported interventions delivered within 72 hours of ED attendance. These were Comprehensive Geriatric Assessment (CGA) interventions delivered either by nurses or geriatricians and were targeted at older people with frailty. The review by Hastings et al (15) looked at evidence for interventions to improve outcomes for older people discharged from the ED. Fourteen of the studies reported by Hastings et al (15) were of interventions either initiated or concluded in the ED. A wide variety of interventions were reported, from CGA to single screening and assessment interventions, delivered by single practitioners or multidisciplinary teams. Karam et al (18) limited inclusion criteria to interventions delivered within the ED and including CGA and other intervention types. Lowthian et al (19) reported on discharge interventions in the form of Community Transition Strategies from the ED. All of these strategies included geriatric assessment, but this was undertaken by a variety of professional groups including nurses, allied health professionals, and health visitors. Follow up interventions either consisted of referral to community services or direct linkages including telephone/GP follow up. Nine of the 18 primary studies included in the review of interventions to reduce ED visits by McCusker et al (21) were delivered in the ED; all had an ED and post discharge component.

Outcomes were reported using meta-analysis (11) and (19) and narrative synthesis (15, 18, 21). Conroy et al (11) found no clear evidence of benefit for CGA discharge interventions across all outcomes included in the review. Hastings (14) reported at the level of individual studies only across a wide variety of outcomes. Karam et al (18) developed themes for intervention types (referral, follow up, integrated model of care) and identification of study participants (risk screening or no risk screening). They found that the most effective interventions extended beyond referral and used a clinical risk prediction tool to identify those who would most benefit from the intervention. In the review by Lowthian et al (19), four of the nine studies were included in a meta-analysis, which found no benefit of interventions in terms of ED re-attendance, mortality and emergency hospitalisation. Individual studies were effective in reducing ED reattendance and nursing home admissions – Lowthian et al (19) attributed this potentially to the methods of telephone follow up of discharged patients. The review by McCusker (21) found that there was limited evidence of benefit of discharge interventions (two studies of borderline statistical significance) on ED visits and there was evidence of short term only increase in ED visits as a result of the intervention.

Summary – Discharge interventions varied in their components but tended to employ improved linkages between the ED and the community, either through direct linkage or referral interventions. CGA was frequently used and involved a variety of professional groups. There was limited evidence for the effectiveness of these interventions – two meta-analyses found no benefit to these interventions, and narrative synthesis reported an increase in ED readmissions in the short term among patients who had received these interventions.

Staff-focused reviews

Interventions were generally delivered by ED physicians, geriatricians working within the ED, and nurses with or without an advanced role. There was also evidence of wider MDT led interventions in a number of reviews (14, 15, 17-19, 21, 22, 25), where professionals included research assistants, occupational therapists, discharge co-ordinators, social workers, physiotherapists, and health visitors. Study outcomes were reported narratively and using meta-analysis. There was moderate but inconsistent agreement across the studies for the effectiveness of nurse led interventions.

Fealy et al (13) described 11 nurse-led interventions which included assessment, post discharge referral, patient education, and follow up. Five studies reported reduction in service use and three studies reported functional improvements. Three studies found no effect. Findings were contradictory – there was evidence of reduced service use in the ED leading to increases in primary care service use. The suggested characteristics of effective interventions included pre-intervention screening and better links with home care.

Graf et al (14) described eight nurse-led CGA interventions which included follow up. They reported that nurse-led CGA was effective in improving functional outcomes. There was varying evidence on ED readmissions (both reduced and increased admissions) and nursing home admissions. Three studies found no effect, attributed partly to study design limitations.

Malik et al (20) reported three different types of nurse intervention: assessment using risk screening, CGA, and nurse-led case and discharge management. This meta-analysis of nine studies found that nursing interventions did not have a significant statistical impact on any of four outcomes (hospitalisation, readmissions, length of hospital stay and ED revisits). This study did not examine functional decline. The researchers contrasted these findings with previous reviews (13, 15, 18) which had demonstrated reduced service use as a result of these interventions, and had also reported that ED risk screening led to reduced hospitalisation and nursing home admissions. These inconsistencies are attributed to methodological weaknesses in study designs, supporting an agenda for additional research on interventions that extend from the ED to the community.

Pearce et al (23) identified only two studies which evaluated patient focussed outcomes. The interventions were related to physical equipment supplied by nurses. Findings indicated that both warming blankets and seating position had a positive impact on patient comfort and wellbeing. The researchers noted the paucity of research around patient centred outcomes such as nutrition, hydration and communication.

Jay (17) reported reduced admissions rates (ranging between 2.6% and 9.7%). The evidence for length of stay and readmission rates was mixed. A number of their included studies also reported changes in admissions rates for the control groups, indicating that CGA may have altered culture and practices around the risks of admission versus discharge.

Summary – There was conflicting evidence around the benefits of nurse-led interventions for older people in the ED. Included reviews reported reduced service use and reduced functional decline, in contrast to evidence of increased service use as a result of interventions. The strongest evidence, in the form of meta-analysis, found no effect from nurse-led interventions. There was evidence of lowered admission rates following geriatrician led CGA interventions. There were common methodological limitations reported across studies.

Population-focused reviews

Schnitker et al (22) and Sinha et al (24) reported evidence for identification and management programmes which specifically targeted older people with cognitive impairment. Schnitker et al (22) also reported staffing interventions (team and individual changes to service delivery and staff training). Neither review reported patient or health service outcomes. Both reviews described intervention characteristics that report positive outcomes, but not the outcomes themselves. Both reviews summarised that interventions were poorly represented or described within the ED literature. There was more evidence from acute care settings, although transferability of these interventions to the ED is not well understood.

Summary – There was limited evidence for population-focused interventions. The reporting of evidence made any comparison between reviews challenging. It was not possible to summarise ED interventions for older people with cognitive impairment.

Intervention component reviews

Three reviews reported on the core components of successful interventions and their outcomes. Fan (12), Hughes (16), and Sinha (25) considered the key components or elements of effective interventions in addition to the overall effectiveness:

- Core operational components of interventions and the role of these components in the success of interventions (25)
- Key elements of effective interventions (12)
- Intervention components and intervention strategies adopted (16)

In terms of intervention effectiveness, the case management interventions reported by Sinha were reported as having positive effects (not statistically significant) on satisfaction levels, ED reattendances, admission rates (immediate and longer term), and nursing home admissions. Negative results included a small but significant negative effect on ED reattendances (25) and higher ED use (12). There was a statistically significant outcome of lowering ED use or length of stay in five of 20 studies (12). Hughes et al (16) found a small positive effect of ED interventions on functional status.

Table 4 reports components that were associated with interventions found in reviews to be effective. There was considerable overlap between the three reviews, indicated by shading.

Table 4 Effective components of interventions

Fan et al (12)	Hughes et al (25)	Sinha et al (16)
Multidisciplinary team and gerontological expertise	Inter-professional and capacity building work practices	
Integrated social and medical care		Multi strategy interventions
Risk screening and geriatric assessment	High risk screening. Focussed geriatric assessment	Assessment
Care planning and management		Case management
Discharge planning and referral coordination	Initiation of care and disposition planning in the ED	‘Bridge’ interventions (contact before and after discharge)
Follow up and regular group visits	Post ED discharge follow up with patients	Referral plus follow up

Fan et al (12)	Hughes et al (25)	Sinha et al (16)
Evidence based practice model	Establishment of evaluation and monitoring processes	
	Nursing clinical delivery involvement or leadership	

Summary – There was considerable agreement across three reviews for the components of successful interventions. Effective interventions: integrated strategies for social and medical care involvement; included screening and assessment; initiated care in the ED and bridged this with follow up; monitored and evidenced successful practices. Care quality indicators tended to focus on care processes rather than structures or outcomes and were generally lacking in evidence and limited in testing.

Discussion

This review of reviews summarised evidence on interventions to improve outcomes for older people (including those with frailty related conditions) attending an Emergency Department. Overall, the evidence base was inconsistent. Across the reviews there was incomplete reporting of interventions – a feature of reviews in which data lose details through abstraction from primary studies. In addition, there was high variability in the standards to which reviews were conducted and reported. Our findings are limited to each review authors’ interpretation of primary evidence. Some reviews reported primary studies by intervention type and others by their outcome, and this limits the potential for further synthesis of data. The evidence was broadly US focussed and relatively old in terms of the studies included in the reviews. Summaries commonly featured calls for more primary research using rigorous evaluation methods, and also acknowledged the challenges of researching a vulnerable population in a fast moving and high pressured environment.

The evidence for CGA and related multidisciplinary interventions has been widely studied, but inconsistent reporting makes definitive conclusions difficult. Despite this, there was some evidence for effectiveness. In particular, geriatrician-led CGA appeared to have stronger effect on reducing admission rates than nurse-led interventions. Following CGA, however, ED reattendance rates may be unchanged or even increased, particularly in the short-term. This may reflect the evidence for continuous rather than brief interventions: holistic, person-centred management plans take time to implement and to yield benefit, and there may be a short-term incidence of rebound problems for people while they adjust to change. Studies with a longer follow up period may be required to understand whether this is the case.

There was a lack of evidence appraising targeted interventions for older people with cognitive impairment attending EDs. There was widespread evidence of holistic interventions being undertaken, including CGA. However, despite being a holistic and person-centred intervention, the effectiveness of CGA tended to be measured with service-related metrics (such as mortality and admissions) as opposed to patient centred metrics (such as pain and quality of life). Future research and quality improvement innovations should ensure that patients are consulted on the outcomes of importance.

Successful interventions integrated social and medical care, included screening and assessment, were initiated in the ED and bridged to other settings with follow up, and monitored and evidenced successful practices. This has far reaching implications for service delivery and reconfiguration. There is a need for robust, multicentre controlled studies (e.g. cluster RCT) that examine CGA based interventions in the ED, focusing upon patient centred outcomes.

Key Messages Box

What is already known on this subject?	What this study adds?
ED care for older people can be complex. Those people living with frailty have poorer health outcomes.	Description and appraisal of healthcare interventions is inconsistent and therefore difficult to synthesise.
Caring for increasing numbers of older ED attendees is a critical health service issue.	No individual intervention was found to be more beneficial for older people with emergency care needs.
Numerous interventions have been trialled within Emergency and Urgent Care.	Interventions initiated in the ED and continued into other settings tended to result in improved outcomes.
	Most studies reported service metrics rather than person-centred outcomes.

References

- van Oppen JD, Keillor L, Mitchell Á, Coats TJ, Conroy SP. What older people want from emergency care: a systematic review. *Emerg Med J.* 2019;36(12):754-61.
- Pollock A, Campbell P, Brunton G, Hunt H, Estcourt L. Selecting and implementing overview methods: implications from five exemplar overviews. *Syst Rev.* 2017;6(1):145.
- Hunt H, Pollock A, Campbell P, Estcourt L, Brunton G. An introduction to overviews of reviews: planning a relevant research question and objective for an overview. *Syst Rev.* 2018;7(1):39.
- McKenzie JE, Brennan SE. Overviews of systematic reviews: great promise, greater challenge. *Syst.* 2017;6:185.
- Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JPA, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *Bmj.* 2009;339:b2700.
- Pollock M, Fernandes RM, Becker LA, Featherstone R, Hartling L. What guidance is available for researchers conducting overviews of reviews of healthcare interventions? A scoping review and qualitative metasummary. *Syst Rev.* 2016;5(1):190.
- Preston L, Chambers D, Campbell F, Cantrell A, Turner J, Goyder E. What evidence is there for the identification and management of frail older people in the emergency department? A systematic mapping review *Health Services and Delivery Research.* 2018.
- Banzi R, Cinquini M, Gonzalez-Lorenzo M, Pecoraro V, Capobussi M, Minozzi S. Quality assessment versus risk of bias in systematic reviews: AMSTAR and ROBIS had similar reliability but differed in their construct and applicability. *Journal of clinical epidemiology.* 2018;99:24-32.
- Pieper D, Antoine SL, Mathes T, Neugebauer EA, Eikermann M. Systematic review finds overlapping reviews were not mentioned in every other overview. *Journal of clinical epidemiology.* 2014;67(4):368-75.
- Caplan GA, Williams AJ, Daly B, Abraham K. A Randomized, Controlled Trial of Comprehensive Geriatric Assessment and Multidisciplinary Intervention After Discharge of Elderly from the Emergency Department--The DEED II Study. *J Am Geriatr Soc.* 2004;52(9):1417-23.
- Conroy SP, Stevens T, Parker SG, Gladman JR. A systematic review of comprehensive geriatric assessment to improve outcomes for frail older people being rapidly discharged from acute hospital: 'interface geriatrics'. *Age Ageing.* 2011;40(4):436-43.

12. Fan L, Lukin W, Zhao J, Sun J, Hou XY. Interventions targeting the elderly population to reduce emergency department utilisation: a literature review. *Emerg Med J*. 2015;32(9):738-43.
13. Fealy G, McCarron M, O'Neill D, McCallion P, Clarke M, Small V, et al. Effectiveness of gerontologically informed nursing assessment and referral interventions for older persons attending the emergency department: systematic review. *J Adv Nurs*. 2009;65(5):934-5.
14. Graf CE, Zekry D, Giannelli S, Michel JP, Chevalley T. Efficiency and applicability of comprehensive geriatric assessment in the emergency department: a systematic review. *Aging Clin Exp Res*. 2011;23(4):244-54.
15. Hastings SN, Heflin MT. A systematic review of interventions to improve outcomes for elders discharged from the emergency department. *Acad Emerg Med*. 2005;12(10):978-86.
16. Hughes JM, Freiermuth CE, Shepherd-Banigan M, Ragsdale L, Eucker SA, Goldstein K, et al. Emergency Department Interventions for Older Adults: A Systematic Review. *J Am Geriatr Soc*. 2019;Early View.
17. Jay S, Whittaker P, McIntosh J, Hadden N. Can consultant geriatrician led comprehensive geriatric assessment in the emergency department reduce hospital admission rates? A systematic review. *Age Ageing*. 2017;46(3):366-72.
18. Karam G, Radden Z, Berall LE, Cheng C, Gruneir A. Efficacy of emergency department-based interventions designed to reduce repeat visits and other adverse outcomes for older patients after discharge: A systematic review. *Geriatr Gerontol Int*. 2015;15(9):1107-17.
19. Lowthian JA, McGinnes RA, Brand CA, Barker AL, Cameron PA. Discharging older patients from the emergency department effectively: a systematic review and meta-analysis. *Age Ageing*. 2015;44(5):761-70.
20. Malik M, Moore Z, Patton D, O'Connor T, Nugent LE. The impact of geriatric focused nurse assessment and intervention in the emergency department: A systematic review. *Int Emerg Nurs*. 2018;37:52-60.
21. McCusker J, Verdon J. Do geriatric interventions reduce emergency department visits? A systematic review. *J Gerontol Ser A-Biol Sci Med Sci*. 2006;61(1):53-62.
22. Parke B, Beath A, Slater L, Clarke AM. Contextual factors influencing success or failure of emergency department interventions for cognitively impaired older people: a scoping and integrative review. *J Adv Nurs*. 2011;67(7):1426-48.
23. Pearce S, Rogers-Clark C, Doolan JM. A comprehensive systematic review of age-friendly nursing interventions in the management of older people in emergency departments. *JBI Libr Syst Rev*. 2011;9(20):679-726.
24. Schnitker L, Martin-Khan M, Beattie E, Gray L. What is the evidence to guide best practice for the management of older people with cognitive impairment presenting to emergency departments? A systematic review. *Adv*. 2013;35(2):154-69.
25. Sinha SK, Bessman ES, Flomenbaum N, Leff B. A systematic review and qualitative analysis to inform the development of a new emergency department-based geriatric case management model. *Ann Emerg Med*. 2011;57(6):672-82.
26. Parker SG, Du X, Bardsley MJ, Goodfellow J, Cooper RG, Cleary R, et al. Measuring outcomes in care of the elderly. *Journal of the Royal College of Physicians of London*. 1994;28(5):428-33.