



This is a repository copy of *Postprandial glycaemic responses in pre- and post-menopausal women*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/167552/>

Version: Published Version

Article:

Sayegh, M., Drew, J.E. and Ranawana, V. orcid.org/0000-0003-2461-7841 (2019)
Postprandial glycaemic responses in pre- and post-menopausal women. *Proceedings of the Nutrition Society*, 78 (OCE2). E47. ISSN 0029-6651

<https://doi.org/10.1017/S0029665119000739>

© 2019 The Authors. Reproduced in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Postprandial glycaemic responses in pre- and post-menopausal women

M. Sayegh, J.E. Drew and V. Ranawana

The Rowett Institute, University of Aberdeen, Foresterhill, Aberdeen, AB25 2ZD.

Postprandial hyperglycaemia increases diabetic and cardiovascular disease risk in healthy individuals⁽¹⁾. Co-ingesting vegetables with high carbohydrate meals can attenuate postprandial glycaemic responses (GR)⁽²⁾. This study aims to further investigate the effect of age and menstrual status on postprandial GR in pre- and post-menopausal women.

Eighteen, pre- (n = 7) and post-menopausal (n = 11) women (BMI 25.2 – 30.4 kg/m²) have been recruited to The impact of Vegetables on acute glycaemia and Glycaemia induced CVD risk in women: metabolic effects and Inter-individual variations study (VegGI). Postprandial GR following rice (75 g of available carbohydrate) (R), rice with bok choy (150g) (RB) or spinach (150g) (RS) was measured every 5 minutes for 4 hours using a Continuous Glucose Monitoring system (Medtronic Ltd). Anthropometric, cholesterol (Cholestech LDX, Alere) and HbA1c (Afinion AS100, Alere) measures were made in a fasted state. Oestradiol, progesterone, follicular stimulating hormone (FSH) and testosterone were measured (ADVIA Centaur). Principal component analysis (PCA) (Umetrics SIMCA-P + 12.0) was performed on GR data. Total GR was expressed as the incremental area under the curve (iAUC) and calculated geometrically. A linear mixed model was used to analyse the impact of R, RB, RS on GR using age and diet as factors. Independent samples t-test was used for further data comparisons (ver.24.0; SPSS Inc).

The post-menopausal group was significantly older with significantly higher HbA1c, systolic BP, total cholesterol, FSH levels and lower oestradiol levels than pre-menopausal (P < 0.05). PCA (PC1 and PC2) accounting for approximately 82% of the variation indicated age and hormonal status as factors contributing to postprandial GR rather than meal consumption. No significant difference was observed in postprandial GR with consumption of vegetables with the high carbohydrate meal (R) in either group. Inter-individual variation in postprandial GR was observed in both groups, with greatest variation observed in the post-menopausal group. Mean postprandial GR was significantly higher in the post-menopausal group following consumption of RB (P < 0.045) (Fig 1A). A linear mixed model revealed a significant impact of meals on postprandial GR associated with age (P = 0.010). The post-menopausal group had significantly higher postprandial GR following RB at 40 – 90 minutes (Fig 1B).




Fig. 1. [A] iAUC distribution for postprandial GR in pre- (R, RB n = 7, RS n = 5) and post-menopausal (R n = 11, RB n = 10, RS n = 11) individuals and mean values \pm SE of groups. [B] Mean change \pm SE in postprandial GR following consumption of RB using mixed linear model. *P < 0.05; [‡]P \leq 0.001

Preliminary analysis of data suggests a difference in glucose metabolism regulation between pre- and post-menopausal women, but no significant reduction in postprandial GR was associated with co-ingesting vegetables with a high carbohydrate meal. There was an indication that postprandial GR varied more in the post-menopausal group. Recruitment is ongoing to increase group size and further assess factors influencing GR responses in pre- and post-menopausal women.

This work was supported by Scottish Government’s Rural and Environment Science and Analytical Services Division

1. Woerle HJ, Neumann C, Zschau S *et al.* (2007) *Diabetes Res Clin Pract* 77, 280–285.
2. Kondo S, Suzuki A, Kurokawa M *et al.* (2016) *Biomed Rep* 5, 553–558.