



This is a repository copy of *Synthesis of boronic ester  $\gamma$ -lactam building blocks*.

White Rose Research Online URL for this paper:  
<https://eprints.whiterose.ac.uk/169069/>

Version: Published Version

---

**Article:**

Rodgers, G., Wilson, E.J., Robertson, C.C. [orcid.org/0000-0002-1419-9618](https://orcid.org/0000-0002-1419-9618) et al. (2 more authors) (2021) Synthesis of boronic ester  $\gamma$ -lactam building blocks. *Advanced Synthesis & Catalysis*, 363 (9). pp. 2392-2395. ISSN 1615-4150

<https://doi.org/10.1002/adsc.202001339>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:  
<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# Synthesis of Boronic Ester $\gamma$ -Lactam Building Blocks

George Rodgers,<sup>a</sup> Ellen J. Wilson,<sup>a</sup> Craig C. Robertson,<sup>a</sup> Daniel J. Cox,<sup>b</sup> and Benjamin M. Partridge<sup>a,\*</sup>

<sup>a</sup> Department of Chemistry, University of Sheffield, Dainton Building, Sheffield, S3 7HF, United Kingdom  
E-mail: b.m.partridge@sheffield.ac.uk

<sup>b</sup> Redbrick Molecular Ltd., 217 Portobello, Sheffield, S1 4DP, United Kingdom

Manuscript received: October 29, 2020; Revised manuscript received: December 3, 2020;

Version of record online: ■■■, ■■■■



Supporting information for this article is available on the WWW under <https://doi.org/10.1002/adsc.202001339>

© 2020 The Authors. Advanced Synthesis & Catalysis Published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

**Abstract:** Saturated heterocycles are found widely in biologically active compounds such as medicinal drugs and agrochemicals. However, boronic acid-derived building blocks for these structures have limited availability, particularly in comparison to heteroaromatic boronic acids. We report the preparation of boronic ester  $\gamma$ -lactams through a Cu-catalysed conjugate borylation-cyclisation protocol. Using a chiral catalyst, this can be performed in high enantioselectivity. Exploration of the further transformations of these reagents suggest that the boronic esters have much potential as chemical building blocks.


**Keywords:** boron; lactams; copper; building blocks; cross-coupling.

Boronic acids are valuable chemical building blocks, which are used in a broad range of disciplines from medicinal chemistry to materials science.<sup>[1]</sup> Heteroaromatic boronic acids, and their derivatives, are particularly useful for the preparation of biologically active molecules through reactions such as the Suzuki-Miyaura cross-coupling<sup>[2]</sup> and the Chan-Lam reaction.<sup>[3]</sup> This is further enabled by the wide commercial availability and ease of preparation of a broad range of heteroaromatic boronic acids and derivatives. Conversely, the corresponding boronic acid-containing saturated heterocycles have limited availability, despite the prevalence of such heterocycles in pharmaceutical drugs.<sup>[4]</sup> To illustrate this, at the time of publication Merck Sigma-Aldrich sold only 4


pyrrolidine- or piperidine-based boronic acids and derivatives.<sup>[5]</sup> There are a growing number of transformations of alkylboron reagents.<sup>[6]</sup> However, in order for such methods to be taken up widely by both academia and industry, access to a broader range of alkylboron building blocks is required.

As part of an ongoing research programme into the chemistry of alkylboronic esters,<sup>[7]</sup> we envisaged that beta-boronic ester  $\gamma$ -lactams could be prepared through conjugate borylation<sup>[8]</sup> of an *E*-amino enoate (scheme 1). Borylation would break the unsaturation of the enoate, allowing the pendent amine to form the lactam. Previous conjugate borylation-cyclisation strategies have instead involved C–C bond formation through reaction of an *in situ* formed enolate with a pendent electrophile.<sup>[9]</sup> This work complements recent reports from Lautens and co-workers, who have developed a borylation-acylation strategy of styrene-containing carbamoyl chlorides to form lactams.<sup>[10]</sup>


We started this investigation by subjecting enoate **1a** to a B<sub>2</sub>pin<sub>2</sub> in the presence of CuI as a catalyst (scheme 2). It was found that borylation proceeded smoothly to give boronic ester **2a** in good yield. Interestingly, the reaction was complete within 1 h, whereas the borylation of ethyl cinammate under the same reaction conditions requires > 12 h reaction time. Boronic ester **2a** did not lactamise spontaneously, presumably due to the relatively low nucleophilicity of the aniline nitrogen. Attempts to induce cyclisation under mildly basic conditions were successful, but were accompanied by a small amount of a degradation product from protodeboration of the boronic ester. Instead, treatment of boronic ester **2a** with acetic acid gave the corresponding lactam **3a**, without need for further purification. While the process can be tele-




Previous work including: Fernandez, Lam, Lautens, Xu



Lautens and co-workers:



**Scheme 1.** Examples of borylation-cyclisation strategies to form saturated carbo- and heterocyclic boronic esters.





**Scheme 2.** Borylation and cyclisation of enoate **1a**.

scoped, purification after conjugate borylation was found to be more successful.

We next subjected our protocol to the formation of a range of boronic ester-containing lactams (scheme 3). The method is tolerant to functional groups including esters, nitriles and aryl bromides. Lactamisation of anilines substituted with electron withdrawing groups required longer reaction times (**3d**, **3e**, **3g**). Both steps of the protocol could be performed on multigram scale.<sup>[11]</sup>

When *N*-benzyl enoate **1h** was subjected to conjugate borylation, the linear boronic ester underwent cyclisation *in situ*. This is presumably due to the higher nucleophilicity of alkyl amines versus anilines. While the boronic ester was formed in good yield, attempts at purification led to significant decomposition of the product. Instead, to give an indication of the




**Scheme 3.** Scope of boronic ester lactams. Yields are of isolated material unless otherwise stated. a) 24 h reaction time for step 2. b) 9 h reaction time for step 2. c) 5 h reaction time for step 2. d) isolated as a 1:1 mixture with pinacol. e) oxidation of the crude material using H<sub>2</sub>O<sub>2</sub>/NaOH was carried out. See the supporting information for full details.


efficiency of the reaction, oxidation of the boronic ester was performed to give alcohol **4**.

Access to enantiomerically enriched boronic ester was achieved using a catalyst derived from CuCl and (*R,S*)-Josiphos as a ligand<sup>[12]</sup> to give (*S*)-**2a** in 56% yield (Scheme 4). Cyclisation followed by oxidation of the C–B bond gave alcohol (*S*)-**5** in 90% e.e.. The absolute configuration of (*S*)-**5** was confirmed through X-ray crystallography.<sup>[11,13]</sup>

To demonstrate the utility of the boronic ester lactams, we explored their further reactivity (Scheme 5). Oxidation of the boronic ester to alcohol



**Scheme 4.** Enantioselective borylation-cyclisation and oxidation. e.e. determined by HPLC analysis of (*S*)-**5**.



**Scheme 5.** Transformations of boronic ester lactams. a)  $\text{NaBO}_3 \cdot \text{H}_2\text{O}$ , THF/ $\text{H}_2\text{O}$ . b)  $\text{AgNO}_3$ , Selectfluor, TFA,  $\text{CH}_2\text{Cl}_2/\text{H}_2\text{O}$ ,  $60^\circ\text{C}$ . c)  $\text{KHF}_2$ , MeOH. d) PhBr, cataCXium A Pd G3 (5 mol%),  $\text{Cs}_2\text{CO}_3$ , PhMe/ $\text{H}_2\text{O}$ ,  $110^\circ\text{C}$ . e) i) furan, nBuLi, THF,  $-78^\circ\text{C}$  ii) **3c**, iii) NBS, THF,  $-78^\circ\text{C}$ -RT. f) i) vinyl MgBr, THF,  $-78^\circ\text{C}$ , ii)  $\text{I}_2$ , MeOH,  $-78^\circ\text{C}$ , iii) NaOMe, MeOH,  $-78^\circ\text{C}$ -RT.

**6**, using  $\text{NaBO}_3 \cdot \text{H}_2\text{O}$ , and fluoride **7**, using  $\text{AgNO}_3$  and Selectfluor,<sup>[14]</sup> proceeded smoothly. Zweifel olefination and Aggarwal arylation<sup>[15]</sup> were both successful, despite the presence of enolisable protons from the lactam. Suzuki-Miyaura cross coupling could be carried out using the corresponding trifluoroborate salt **8**, albeit in moderate yield.<sup>[16]</sup>

In summary, we have developed a protocol for the preparation of  $\gamma$ -lactams containing an alkylboronic ester handle. These boronic esters have much potential as heterocyclic chemical building blocks, demonstrated by a range of further transformations of the boronic ester moiety. Further methods to produce boronic ester heterocycles are underdevelopment and will be reported in due course.

## Experimental Section

### General Procedure for the Preparation of $\gamma$ -lactam Boronic Esters (**3**)

An oven dried Schlenk flask was charged with  $\text{B}_2\text{pin}_2$  (0.305 g, 1.20 mmol), CuI (4.0 mg, 0.020 mmol),  $\text{K}_2\text{CO}_3$  (0.235 g, 1.70 mmol), and evacuated and backfilled with Ar. Anhydrous THF (2 mL) was added, and the mixture was stirred for 10 mins. A solution of the enoate **1** (1.00 mmol) in anhydrous THF (1.3 ml) and methanol (0.08 ml, 2 mmol) was added, and the mixture was stirred for 1 h. The mixture was passed through a plug of celite, and the solution was concentrated *in vacuo*. The crude material was purified by either chromatography or recrystallisation to give boronic ester **2**.

Boronic ester **2** (0.5 mmol) was stirred in AcOH (1 ml, 0.5 M) at  $50^\circ\text{C}$  for 4 h. The mixture was quenched with saturated aqueous  $\text{Na}_2\text{CO}_3$  (10.0 ml), and extracted with  $\text{CH}_2\text{Cl}_2$  ( $3 \times 15.0$  ml). The combined organic phases were dried ( $\text{MgSO}_4$ ), filtered, and concentrated *in vacuo* to give lactam **3**.

## Acknowledgements

This work was supported by EPSRC (EP/T009292/1) and the University of Sheffield. We thank Redbrick Molecular Ltd. for a CASE studentship for GR. We acknowledge Dr Antonio Romero-Arenas for performing the HPLC analysis.

## References

- [1] D. G. Hall, Editor, *Boronic Acids, Volume 1: Preparation and Applications in Organic Synthesis, Medicine and Materials*, Second Completely Revised Edition, Wiley-VCH Verlag GmbH & Co. KGaA, **2011**.
- [2] a) J. Magano, J. R. Dunetz, *Chem. Rev.* **2011**, *111*, 2177–2250; b) C. Torborg, M. Beller, *Adv. Synth. Catal.* **2009**, *351*, 3027–3043.
- [3] a) J. X. Qiao, P. Y. S. Lam, *Synthesis* **2011**, 829–856; b) M. J. West, J. W. B. Fyfe, J. C. Vantourout, A. J. B. Watson, *Chem. Rev.* **2019**, *119*, 12491–12523.
- [4] E. Vitaku, D. T. Smith, J. T. Njardarson, *J. Med. Chem.* **2014**, *57*, 10257–10274.
- [5] <https://www.sigmaaldrich.com/united-kingdom.html> (accessed 29/10/20).
- [6] For reviews, see: a) C. Sandford, V. K. Aggarwal, *Chem. Commun.* **2017**, *53*, 5481–5494; b) R. J. Armstrong, V. K. Aggarwal, *Synthesis* **2017**, *49*, 3323–3336; c) D. Leonori, V. K. Aggarwal, *Angew. Chem. Int. Ed.* **2015**, *54*, 1082–1096; *Angew. Chem.* **2015**, *127*, 1096–1111.
- [7] a) J. D. Grayson, B. M. Partridge, *ACS Catal.* **2019**, *9*, 4296–4301; b) F. M. Dennis, C. C. Robertson, B. M. Partridge, *Synthesis* **2020**, *52*, 1903–1914.
- [8] J.-B. Chen, A. Whiting, *Synthesis* **2018**, *50*, 3843–3861.
- [9] a) Y.-P. Bi, H.-M. Wang, H.-Y. Qu, X.-C. Liang, Y. Zhou, X.-Y. Li, D. Xu, M.-H. Shen, H.-D. Xu, *Org. Biomol. Chem.* **2019**, *17*, 1542–1546; b) J. Sendra, R. Manzano, E. Reyes, J. L. Vicario, E. Fernández, *Angew. Chem. Int. Ed.* **2020**, *59*, 2100–2104; c) Y.-J. Zuo, X.-T. Chang, Z.-M. Hao, C.-M. Zhong, *Org. Biomol. Chem.* **2017**, *15*, 6323–6327; d) E. M. Larin, J. Loup, I. Polishchuk, R. J. Ross, A. Whyte, M. Lautens, *Chem. Sci.* **2020**, *11*, 5716–5723; e) A. R. Burns, J. Solana González, H. W. Lam, *Angew. Chem. Int. Ed.* **2012**, *51*, 10827–10831; *Angew. Chem.* **2012**, *124*, 10985–10989.
- [10] a) A. Torelli, A. Whyte, I. Polishchuk, J. Bajohr, M. Lautens, *Org. Lett.* **2020**, *22*, 7915–7919; b) A. Whyte, K. I. Burton, J. Zhang, M. Lautens, *Angew. Chem. Int. Ed.* **2018**, *57*, 13927–13930.
- [11] For full details, see the supporting information.
- [12] J.-E. Lee, J. Yun, *Angew. Chem. Int. Ed.* **2008**, *47*, 145–147; *Angew. Chem.* **2008**, *120*, 151–153.

- [13] CCDC 2041142 ((S)-5) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.
- [14] Z. Li, Z. Wang, L. Zhu, X. Tan, C. Li, *J. Am. Chem. Soc.* **2014**, *136*, 16439–16443.
- [15] A. Bonet, M. Odachowski, D. Leonori, S. Essafi, V. K. Aggarwal, *Nat. Chem.* **2014**, *6*, 584–589.
- [16] a) L. Li, S. Zhao, A. Joshi-Pangu, M. Diane, M. R. Biscoe, *J. Am. Chem. Soc.* **2014**, *136*, 14027–14030; b) S. Zhao, T. Gensch, B. Murray, Z. L. Niemeyer, M. S. Sigman, M. R. Biscoe, *Science* **2018**, *362*, 670–674.
-


## COMMUNICATIONS

---

### Synthesis of Boronic Ester $\gamma$ -Lactam Building Blocks

*Adv. Synth. Catal.* **2020**, 362, 1–5

 G. Rodgers, E. J. Wilson, C. C. Robertson, D. J. Cox, B. M. Partridge\*



---