



**UNIVERSITY OF LEEDS**

This is a repository copy of *EULAR recommendations for the reporting of ultrasound studies in rheumatic and musculoskeletal diseases (RMDs)*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/169502/>

Version: Accepted Version

---

**Article:**

Costantino, F, Carmona, L, Boers, M et al. (20 more authors) (2021) EULAR recommendations for the reporting of ultrasound studies in rheumatic and musculoskeletal diseases (RMDs). *Annals of the Rheumatic Diseases*. ISSN 0003-4967

<https://doi.org/10.1136/annrheumdis-2020-219816>

---

© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ. This manuscript version is made available under the CC BY-NC 4.0 license <https://creativecommons.org/licenses/by-nc/4.0/>

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial (CC BY-NC) licence. This licence allows you to remix, tweak, and build upon this work non-commercially, and any new works must also acknowledge the authors and be non-commercial. You don't have to license any derivative works on the same terms. More information and the full terms of the licence here:  
<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

**EULAR reporting guidelines for ultrasound studies in rheumatic and musculoskeletal diseases**

Félicie Costantino, MD, PhD<sup>1,2</sup>, Loreto Carmona<sup>3</sup>, MD, PhD, Maarten Boers<sup>4</sup>, MD, PhD, Marina Backhaus<sup>5</sup>, MD\*, Peter V. Balint<sup>6</sup>, MD, PhD\*, Hilde B. Hammer<sup>7</sup>, MD, PhD\*, George A.W. Bruyn<sup>8</sup>, MD, PhD\*, Robin Christensen<sup>9</sup>, MSc, PhD\*, Philip G Conaghan<sup>10\*</sup>, MBBS, PhD, FRACP, FRCP, Ricardo J. O. Ferreira, RN, PhD<sup>11,12\*</sup>, Juan Luis Garrido-Castro<sup>13\*</sup>, Francis Guillemin<sup>14</sup>, MD, PhD\*, Désirée van der Heijde MD, PhD<sup>15\*</sup>, Annamaria Iagnocco<sup>16</sup>, MD\*, Marion Kortekaas<sup>15,17</sup>, MD, PhD\*, Robert Landewe<sup>18</sup>, MD, PhD\*, Peter Mandl<sup>19</sup>, MD, PhD\*, Esperanza Naredo<sup>20</sup>, MD\*, Wolfgang A. Schmidt<sup>21</sup>, MD\*, Lene Terslev<sup>22</sup>, MD, PhD\*, Caroline B Terwee<sup>23</sup>, PhD\*, Ralf Thiele<sup>24</sup>, MD\* and Maria Antonietta D'Agostino, MD, PhD<sup>1,2</sup>

**\*Alphabetical order; \*these authors have contributed equally to the project**

1 – Université Paris-Saclay, UVSQ, Inserm U1173, Infection et inflammation, Laboratory of Excellence INFLAMEX, 78180, Montigny-Le-Bretonneux, France.

2 – Rheumatology Department, AP-HP, Ambroise Paré Hospital, 92100, Boulogne-Billancourt, France.

3 – Instituto de Salud Musculoesquelética (InMusc), Madrid, Spain.

4 – Department of Epidemiology&Biostatistics, and Amsterdam Rheumatology and Immunology Center, Amsterdam University Medical Centers, Vrije Universiteit, Amsterdam, the Netherlands.

5 – Department of Internal Medicine - Rheumatology and Clinical Immunology, Park-Klinik Weissensee, Berlin, Germany

6 – 3rd Rheumatology Department, National Institute of Rheumatology and Physiotherapy, Budapest, Hungary.

7 – Dept of Rheumatology, Diakonhjemmet Hospital, and University of Oslo, Oslo, Norway.

8 – MC Groep Hospitals, Lelystad, the Netherlands.

9 – Musculoskeletal Statistics Unit, the Parker Institute, Bispebjerg and Frederiksberg Hospital, University of Copenhagen & Research Unit of Rheumatology, Department of Clinical Research, University of Southern Denmark, Odense University Hospital, Denmark.

10 – Leeds Institute for Rheumatic and Musculoskeletal Medicine, University of Leeds, and NIHR Leeds Biomedical Research Centre, Leeds, UK.

11 – Department of Rheumatology, Centro Hospitalar e Universitário de Coimbra, Coimbra, Portugal.

12 – Health Sciences Research Unit: Nursing (UICISA:E), Nursing School of Coimbra (ESEnfC), Coimbra, Portugal.

13 – Spanish Federation of Spondyloarthritis Associations (CEADE), Madrid, Spain.

14 – Université de Lorraine, EA 4360 APEMAC, Nancy, France.

15 – Department of Rheumatology, Leiden University Medical Center, Leiden, the Netherlands.

16 – Academic Rheumatology Centre, Università degli Studi di Torino, Turin, Italy.

17 – Department of Rheumatology, Flevoziekenhuis, Almere, the Netherlands.

18 – Amsterdam University Medical Center & Zuyderland Medical Center Heerlen, the Netherlands.

19 – Department of Rheumatology, Medical University of Vienna, Vienna, Austria.

20 – Department of Rheumatology, Bone and Joint Research Unit. Hospital Universitario Fundación Jiménez Díaz, IIS Fundación Jiménez Díaz, and Universidad Autónoma de Madrid. Madrid, Spain.

21 – Medical Centre for Rheumatology Berlin-Buch, Immanuel Krankenhaus Berlin, Germany.

22 – Center for Rheumatology and spine Diseases, Rigshospitalet, Copenhagen university, Copenhagen, Denmark.

23 – Amsterdam UMC, Vrije Universiteit Amsterdam, Department of Epidemiology and Biostatistics, Amsterdam, the Netherlands.

24 – Allergy/Immunology and Rheumatology Division, University of Rochester, Rochester, NY, USA.

Correspondence to:

Maria Antonietta D'Agostino,

Rheumatology Department

Hôpital Ambroise Paré, 9 avenue Charles de Gaulle, 92100, Boulogne, France

Ph: 33-(0)149 095 674; Fax: 33-(0)149 095 865

E-mail: maria-antonietta.dagostino@aphp.fr

**ABSTRACT** [*Word count 248/250*]

**Objectives:** To produce European League Against Rheumatism (EULAR) reporting guidelines for ultrasound studies in rheumatic and musculoskeletal diseases (RMDs).

**Methods:** Based on literature reviews and expert opinion (through Delphi surveys), a taskforce of 23 members (12 experts in ultrasound in RMDs, 9 in methodology and biostatistics together with a patient research partner and a health-care professional) developed a checklist of items to be reported in every RMDs study using ultrasound. This checklist was further refined by involving a panel of 79 external experts (musculoskeletal imaging experts, methodologists, journal editors), who evaluated its comprehensibility, feasibility and comprehensiveness. Agreement on each proposed item was assessed with an 11-point Likert scale, grading from 0 (total disagreement) to 10 (full agreement).

**Results:** Two face-to-face meetings, as well as two Delphi rounds of voting, resulted in a final checklist of 23 items, including a glossary of terminology. Twenty-one of these were considered “mandatory” items to be reported in every ultrasound study (such as blinding, development of scoring systems, definition of target pathologies) and 2 “optional” to be reported only if applicable, such as possible confounding factors (i.e. alcohol intake, ambient conditions) or experience of the sonographers.

**Conclusions:** A EULAR taskforce developed a checklist to ensure transparent and comprehensive reporting of aspects concerning research and procedures that need to be present in studies using ultrasound in RMDs. This checklist, if widely adopted by authors and editors, will greatly improve the interpretability of study development and results, including the assessment of validity, generalisability and applicability.

**Keywords:** reporting guidelines, checklist, musculoskeletal ultrasound, rheumatic and musculoskeletal diseases, EULAR

*Word count [2,958/3,000]*

Ultrasound is an imaging technique widely used in patients with rheumatic and musculoskeletal diseases (RMDs) to detect signs of inflammation and destructive changes.[1] Despite an increased use in clinical practice facilitated by major technical advances in the resolution of soft tissue contrast (B-mode or grey-scale (GS)) and of vascular perfusion (Doppler techniques), a relatively long learning curve [2] and, until recently, the absence of agreed scoring systems, have hampered its utilisation for research.[3,4]

The European League Against Rheumatism (EULAR) and the Outcome Measures in Rheumatology (OMERACT) Ultrasound Working Group have actively worked towards the standardization of the technique by developing educational programmes and by performing several studies evaluating its reliability, validity, and feasibility.[5–8] These initiatives have underlined that factors such as nomenclature, definitions of ultrasound-detected pathologies, scoring systems and technical issues with the ultrasound equipment may affect the validity and generalizability of these results. These aspects, along with critical design characteristics, such as reproducibility, blinding, patient selection, and clearly defined purposes of the ultrasound evaluation, are often sub-optimally reported in current ultrasound studies. [5,6,9,10]


Complete and accurate reporting is necessary to detect potential biases in the study (internal validity) and to assess the generalisability and applicability of the results (external validity). Over the last 20 years many guidelines have been developed to improve the quality of reporting of research articles, including those for randomized controlled trials (RCT) (CONSORT) [11] and diagnostic accuracy studies (STARD). [12,13] EULAR has also contributed by developing guidelines for reporting registers and clinical trial extension studies.[14,15] We are not aware of guidelines focused on how to report characteristics of imaging measurement tools such as the equipment characteristics, procedures or scoring, which can influence the validity and generalisability of study results. Therefore, a EULAR taskforce was convened to propose guidelines for the reporting of such aspects in ultrasound studies in RMDs.

## **METHODS**

The convenor (MADA), lead methodologist (LC) and project fellow (FC) led a multidisciplinary taskforce in accordance with the EULAR Standardised Operating Procedures (SOPs).[16] The taskforce included 23 members from 11 European countries and from the United States, and was composed as follows: 12 experts in ultrasound in RMDs, 7 in methodology, 2 in biostatistics, 1 patient research partner and 1 health-care professional in rheumatology. Three of the 23 members were members of EMEUNET and 13 of them were also part of an editorial board.

The taskforce employed a stepwise process summarised in **Figure 1**, including two face-to-face meetings and several Delphi rounds. First the lead methodologist, convenor and fellow searched for evidence of quality of reporting of ultrasound studies in RMDs. The choice was made to focus on an extensively studied topic, i.e. ultrasound assessment of synovitis in rheumatoid arthritis. In PubMed Clinical Queries a broad search was performed; 80 studies were randomly selected and divided in four categories: diagnosis, aetiology, prognosis, and therapy. The articles were summarised in table format to highlight objective, design, technical data, measures and outcomes (Supplementary file 1). These tables were sent to each member of the taskforce prior to the first face-to-face meeting, with the request to identify possible sources of bias and error and the absence of information considered important for the generalisability of the results. During the first face-to-face meeting, the members of the taskforce discussed the results and the unmet requirements in the selected literature, agreed on the format of presentation of the project (checklist or statement document), and elaborated a first list of items to be included. Other objectives of this meeting were the definition of a target audience and the need for systematic reviews. After the meeting, a number of focussed literature reviews addressed specific issues; a summary of their results, along with the total list of items, were subsequently sent to the taskforce members. Relevance and comprehensibility of each proposed item was tested in a Delphi exercise, first by the taskforce members (excluding the convenor, lead methodologist, and fellow), then by a panel of external experts chosen from the fields of musculoskeletal imaging, epidemiology and methodology, including journal editors. External experts were also asked if no key aspects were

missed (comprehensiveness). During the second face-to-face meeting, the optimal format of the checklist document was established. Inclusion of each item was either supported by empirical evidence, when available, or by consensus within the task-force, that the information requested by the item was methodologically important to assess in a study, as recommended by the Enhancing the QUALITY and Transparency Of health Research (EQUATOR) “guidance for developers of health research reporting guidelines” [17]. In the same way it was agreed not to include a level of evidence for each proposed item. The external experts were then invited to apply the checklist to a selection of ultrasound articles and to comment on its feasibility and comprehensibility; this resulted in minor modifications to the items. Finally, an online Delphi survey was performed among the taskforce experts to obtain their level of agreement with each final item, including each term of an accompanying glossary, included to define the checklist terminology. Agreement was assessed with a Likert scale, grading from 0 (total disagreement) to 10 (full agreement). Consensus was defined as a mean agreement  $\geq 7$  and with at least 8 responders (2/3 of participants) having an agreement  $\geq 7$ .



**Figure 1.** Process flow chart of the project.

## RESULTS

**Figure 1** shows the flowchart of the project. During the first face-to-face meeting, a preliminary checklist of 43 items was established, and three scoping reviews were requested on factors potentially influencing the ultrasound evaluation and therefore the generalisability of the results: a) contextual factors (e.g. smoking, temperature), b) machine quality (e.g. device, settings) and acquisition methods (e.g. joint or transducer position) (Supplementary file 2).

A first Delphi exercise helped to decide which of the 43 items should be considered as “mandatory” (always reported in every ultrasound study) or “optional” (reported only according to specific study designs). After two voting rounds, several items were rephrased, deleted or combined, resulting in a checklist of 17 “mandatory” and 8 “optional” items (**Figure 1**).

This new checklist was distributed among 218 external panellists (external Delphi exercise): 123 experts in MSK imaging, 67 in epidemiology, 7 in methodology as well as 21 journal editors. Seventy-nine of them (36%) participated. The external experts rated the initiative as very important (96%), the checklist comprehensive (95%), and all items were considered clear by the majority of them (median: 96%, range: 86-96%). Additional suggestions were made to clarify some terminology. The results were discussed during the second face-to-face meeting, where the format of the checklist was agreed. Each item was verified to ensure comprehensibility, and a preliminary glossary including 12 terms was prepared. After this process the checklist included 23 items (21 “mandatory” and 2 “optional”), organised into 13 categories. This version and the glossary were distributed to the 79 external experts who had participated in the previous evaluation. Twenty-nine (37%) of them agreed to test the new checklist on additional selected articles and to comment on the comprehensibility and comprehensiveness of the items and the glossary, as well as on the feasibility of applying the checklist. The median time needed to assess the articles for reporting the checklist items was 30 minutes (range 10 to 240). Comprehensibility was assessed as good with a median of 8 (range 1-10); and additional suggestions on the glossary terminology were made. The convenor and fellow incorporated all suggestions, and the final product (checklist with accompanying recommendation guidelines and

glossary) was submitted to the taskforce members. Two rounds of an additional Delphi exercise (the second) were needed to obtain a final agreement on each item of the checklist and each term of the glossary.

The final checklist, composed of 23 items (21 “mandatory” and 2 “optional”) and organised into 13 categories, along with the level of agreement for each item, is reported in **Table 1**; the accompanying glossary is presented in **Table 2**, and the recommendation guidelines on how to use and interpret each item in the following paragraphs.

### **Target audience and when to apply the checklist**

The target audience comprises health or scientific researchers reporting or assessing observational and interventional studies using ultrasound in RMDs: i.e. rheumatologists, radiologists, and health care professionals using ultrasound, manuscript reviewers, grant applicants and reviewers, journal editors. Each mandatory item of the checklist should be considered as essential to be reported in every ultrasound study regardless of the purpose of the study. Such a report will allow proper appraisal of the validity and applicability of the results. The checklist is meant to be applied whenever ultrasound is used (investigation of measurement properties, diagnostic or prognostic accuracy and therapeutic studies). It is focused specifically on ultrasound issues and is not intended to be totally comprehensive for all study design issues, nor intended to replace other existing reporting guidelines (e.g. for RCT, observational diagnostic studies, etc.).

### **General items**

The first six items on “Objective”, “Design”, “Participants”, and “Blinding” as well as items 20 (“Statistical analysis”), and 21 (“Disclosures”) are not specifically related to ultrasound and some of them have already been included in other reporting checklists. However, the taskforce members felt it was essential to include them in this checklist to emphasize their importance. For example, the objective of the ultrasound measurement in a study might be different from the main objective of the study, and then such difference should be clearly described. Another example is the blinding of the ultrasound evaluation. Blinding is of utmost importance, especially in diagnostic or therapeutic studies,



since the lack of patient or operator blinding can influence the results.[18,19] Item 20 (“Statistical analysis”) refers mainly to the way in which the analysis of ultrasound variables should be performed, for example the importance to clearly state whether analyses are performed at patient or at joint/site level.

### **Ultrasound features**

Item 7 refers to the ultrasound definition of the pathological lesions under study. It is crucial to be able to check for consistency between what authors say they want to measure (for example, erosions [target domain] as a measure of structural damage [broad domain of interest]) and what was really measured with ultrasound. The concepts of broad and target domains are explained in **Table 2**. In such cases, reporting the domain components (i.e. elementary lesions) really measured by ultrasound will help to evaluate consistency. A precise definition of ultrasound elementary lesions used in the study helps to check if ultrasound is able to measure what it is supposed to measure (domain match). Here we used the terminology proposed by OMERACT for the development of imaging outcome measurement instruments,[20] and in particular of ultrasound.[5] In recent years, the OMERACT ultrasound working group, frequently in collaboration with EULAR, has undertaken considerable efforts to develop and improve definitions of ultrasound elementary lesions for a defined pathology (e.g. synovitis, enthesitis, bone erosion).[5,8]

### **Scanning/acquisition procedures**

Several sources of variability may affect the reliability of ultrasound measurements and generalizability of the study results. These include the quality of the equipment, positioning of patient and transducer, and training of the examiner. One of the scoping reviews (Supplementary file 2) studied whether acquisition methods (i.e., joint or transducer position and dynamic acquisitions) affected the reliability and accuracy of ultrasound. All retrieved studies confirmed the importance of a standardised joint position for the reliability and generalisability of the results; this applies to all anatomical sites (e.g. knee, wrist, Achilles tendon, etc.) and all target pathologies under study (e.g. synovitis, joint effusion, etc.),[21–26]. In addition, appropriate transducer manipulation is needed to avoid artefacts.[23–25]

For example, transducer pressure may cause the synovial hypertrophy or Doppler signal to disappear or be reduced.[27] The 2001 EULAR guideline for performing ultrasound in rheumatology (updated in 2017) addresses both joint (and patient) positioning as well as transducer use.[28,29] Authors are invited to refer to these latest guidelines in their studies. Items 8 and 9 describe what details should be provided about the scanning and acquisition procedures to assess compliance with these guidelines.

### **Ultrasound scoring systems**

Items 10, 11 and 12 focus on a clear description of ultrasound scoring systems, especially if a newly developed scoring system is used. Special attention should be paid to the documentation of the development of the scoring system. As ultrasound is frequently considered the most operator-dependent imaging technique, intra- and inter-rater reliability is an important concern and a strong argument for standardisation. For new scoring systems, results of intra- and inter-sonographer/reader reliability studies should be reported. For existing scoring systems, reference to previous reliability studies should be given, as well as the results of reliability assessments among the sonographers/readers in the context of the study.

### **Sonographers / readers**

Depending on the setting, the person who performs the ultrasound acquisition of the images (sonographer) can be a health-care professional or a medical doctor (radiologist or rheumatologist). The images can be interpreted at the time of acquisition or later, by the same or another person. Choices made here affect ultrasound scores and generalisability, so details on who performs the acquisition and the interpretation are a mandatory reporting requirement. Item 14 on the experience of sonographer(s) and/or reader(s) is optional, mainly because no consensus exists on how to report such experience. EULAR and American College of Rheumatology suggest a competency assessment in ultrasound to improve the quality of the examination.[30,31]

### **Equipment**

Technical characteristics of the imaging device (item 15), ultrasound modalities and settings used (item 16) may affect the intra- and inter-sonographer/reader reliability and generalisability of the results. A

second scoping review (Supplementary file 2) addressed this question, i.e. whether the ultrasound device (model, age, acquisition software, transducers and settings) affect the reliability or accuracy of the ultrasound examination. We found no study investigating the influence of device age or software on ultrasound results. However, seven studies assessed the influence of the machine (e.g. ultrasound device, transducer frequency, settings) on ultrasound results whatever the anatomical site studied, the ultrasound modality used (i.e. Doppler, GS), or the target pathology under study (e.g. joint effusion, synovitis, erosion).[32–38] Three studies used a phantom to compare the ultrasound devices.[34,36,37] Five of the seven studies showed differences in the performance between machines and therefore an influence on the study results.[35,36] However, in these studies, the relevance and the magnitude of such differences were reported, but no sensitivity analysis was conducted.

### **Images, pictures and figures**

Since ultrasound is a tomographic technique, the appearance of the structures may change following the orientation and position of the transducer. Standardised images should always be presented (item 17) so that the reader can easily recognise the anatomical structures, as well as the target pathology and elementary lesions described in the study. The use of drawings can facilitate the interpretation of the images for readers not experienced in ultrasound. Images should never contain patient information and should be accompanied by clear legends and points of reference.

### **Contextual factors**

Item 18 deals with the feasibility of ultrasound, in particular the time necessary for the evaluation, which depends on the number of sites (or joints) examined and the number of ultrasound examinations performed over the duration of the study. Although these aspects are highly important for the acceptability of the technique, the taskforce members felt that time spent should be reported only if relevant to the study question.

Item 19 refers to additional potential sources of variability: ambient and patient conditions. This item was made optional because the third scoping review (Supplementary file 2) failed to find strong evidence of the influence of these factors on the ultrasound results. It reviewed the effect of three

ambient conditions (room temperature, atmospheric pressure, time of the day) and five patient conditions (exercise, skin temperature, smoking, alcohol consumption and caffeine) on ultrasound measurements. There was a potential influence of time of the day on Doppler signal evaluation (with contradictory effects) [39,40] and on GS results;[41] and potential influences on Doppler signal following the application of cold (ice and cold water),[42–44] and after physical exercise.[45–47]

### **DISCUSSION**

This EULAR taskforce developed a checklist to ensure transparent and comprehensive reporting of ultrasound research and procedures aspects, which may affect the interpretation and generalisability of the results. The checklist consists of 23 items (21 “mandatory” and 2 “optional”), organised into 13 categories. Its organisation allows authors to choose the order and format for presenting information, depending on their preferences and on journal style. Content validity of the checklist was confirmed by a panel of external experts, who considered each item of the checklist an essential reporting point, crucial to make an informed judgment on the quality of the scientific report. Moreover, all items were considered comprehensible and the checklist as a whole was considered to comprehensively cover all relevant reporting issues.

Along with sufficient content validity of this checklist, additional strengths include the development process that followed EULAR SOPs for a stepwise consensual approach [16] and the guidance from the EQUATOR Network.[17] Also, the panel members reflected a wide range of expertise and stakeholders. In addition, agreement on comprehensiveness and comprehensibility of the checklist was obtained in the first round of voting for all items of the checklist and all definitions of the glossary.

A possible limitation of this project is the fact that the face-to-face meetings comprised mostly Europeans, with only one colleague from United States, and only one patient. We partially overcame this in the external Delphi panels, including more international experts, including several radiologists. The checklist was purposefully focused and is complementary to other existing guidelines, depending on the study design.[59] It has not been developed as a tool to assess the quality of published research; however, it can certainly serve as a basis to develop such a tool, and its use may improve the quality of studies, as seen with other guidelines.[48,49] We hope this checklist will be widely adopted by authors and editors, which, in turn, will greatly improve the interpretability, reproducibility, and generalisability of study results.

**Acknowledgments**

The authors want to thank all contributing experts who participated in the online survey (see supplementary file 3 for the full list). PGC is supported in part by the UK National Institute for Health Research (NIHR) Leeds Biomedical Research Centre. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health. RC and the Parker Institute, Bispebjerg and Frederiksberg Hospital) is supported by a core grant from the Oak Foundation (OCAY-18-774-OFIL).

**Table 1. Reporting checklist for studies using Ultrasound in Rheumatic and Musculoskeletal Diseases.**

| Topic                                   | No.   | Item to report  | Agreement <sup>†</sup><br>(mean ± SD) |
|---|---|---|---------------------------------------|
| <b>Objective</b>                        | 1   | Objective of the ultrasound measurement in the study<br>( <i>e.g., description, prediction, diagnosis, validation...</i> )  | 9.9 ± 0.3                             |
|   | 2   | Study design<br>( <i>e.g., cross-sectional, case-control, cohort, randomised clinical trial, ...</i> )  | 9.9 ± 0.3                             |
| <b>Design</b>                           | 3   | Prospective or retrospective data collection*   | 9.7 ± 0.7                             |
|   | 4   | Informed consent procedure (written, oral)  | 9.2 ± 1.0                             |
| <b>Participants</b>                     | 5   | Source, selection criteria, and sampling of the participants<br>(including controls where appropriate)  | 9.9 ± 0.3                             |
|   | 6   | Procedures for blinding of sonographers and participants  | 9.3 ± 1.0                             |
| <b>Blinding</b>                         | 7   | a. Broad domain* of interest<br>( <i>e.g. inflammation or structural damage</i> )   | 8.9 ± 1.4                             |
|   |   | b. Target domain* with corresponding theoretical ultrasound definition(s)*<br>( <i>e.g. synovitis: synovial hypertrophy plus increased synovial blood flow</i> )                                |                                       |
|   |   | c. Domain components (i.e. elementary lesions)*<br>with corresponding operational definitions* ( <i>e.g. synovial hypertrophy: increased thickness of synovium with hypoechoic appearance</i> ) |                                       |
| <b>Ultrasound features</b>              | 8   | a. Anatomical region(s)* or structure(s)* that were studied   | 9.2 ± 1.2                             |
|   |   | b. Rationale for choosing these anatomical region(s)/structure(s)   |                                       |
|   | 9   | a. Patient position ( <i>e.g. prone, supine...</i> )  | 9.6 ± 0.8                             |
|   |   | b. Anatomical region position ( <i>e.g. neutral...</i> )  |                                       |
|   |   | c. Surfaces scanned ( <i>e.g. volar, dorsal</i> )   |                                       |
|   | d. Transducer position ( <i>e.g. transverse, longitudinal</i> )   |   |                                       |
|   | e. Whether the examination was dynamic*   |   |                                       |
| <b>Scanning/ acquisition procedures</b> | 10  | Scoring system used:  | 9.6 ± 0.5                             |
|   |   | a. Type ( <i>e.g. quantitative, semi-quantitative, binary</i> )<br>b. Level: ( <i>e.g. patient level, joint/anatomical region level</i> )   |                                       |
| <b>Ultrasound scoring system</b>        | 11  | For existing scoring systems:   | 9.3 ± 1.2                             |
|   |   | a. References or results of previous validity and reliability studies   |                                       |
|   |   | b. Score range (minimum-maximum), and meaning of the score<br>( <i>e.g. higher is .....</i> )   |                                       |
|   |   | c. Rationale for any thresholds or cut-offs   |                                       |
|   |   | d. Training session details if performed  |                                       |
|   |   | e. The reliability* of the scoring system in the hands<br>of the study sonographers/readers   |                                       |
|   | 12  | For new scoring systems:  | 9.4 ± 0.8                             |
|   |   | a. Rationale for developing a new scoring system  |                                       |
|   |   | b. Detailed description of the scoring system   |                                       |
|   |   | c. Reliability assessment:  |                                       |
|   |   | i. Type of reliability: inter-reader, other   |                                       |
|   |   | ii. Training session if performed   |                                       |
|   | iii. The reliability of the scoring system as applied<br>by the study sonographers/readers  |   |                                       |
|   | iv. Whether reliability was assessed on static images,<br>video-clips or real-time examination of patients  |   |                                       |
|   | v. Sample size of the reliability study   |   |                                       |
|   | vi. Reliability results ( <i>e.g. kappa or ICC with 95% confidence interval<br/>and type of kappa or ICC, prevalence of observed lesions,<br/>smallest detectable change, standard error of measurement</i> ) |   |                                       |

## Reporting Ultrasound Studies in RMDs

| Topic   | No. Item to report  | Agreement†<br>(mean ± SD) |
|---|---|---------------------------|
| <b>Sonographer(s)*/<br/>reader(s)*</b>        | 13 a. Whether acquisition and reading were performed at the same time<br>b. Whether acquisition and reading were performed by the same person<br>c. Number of sonographer or reader<br>d. In longitudinal studies, whether the same sonographer scanned the same patient at each assessment | 9.6 ± 0.6                 |
|   | 14 <i>Optional: Information about the experience of sonographer(s) and reader(s) (e.g. numbers of scanned performed, certification, qualification...)</i>   | 8.7 ± 2.3                 |
| <b>Equipment</b>                              | 15 a. Brand and model of the ultrasound device<br>b. Type and model of the transducer<br>c. Whether the ultrasound device (or software) was changed during the study  | 9.1 ± 1.3                 |
|   | 16 Ultrasound modalities* and settings<br>a. Grey scale<br>b. Doppler<br>c. Other   | 9.7 ± 0.9                 |
| <b>Images<br/>(pictures and<br/>drawings)</b> | 17 For images included into the manuscript, verify that:<br>a. Information identifying patient is deleted<br>b. Essential targets in the image(s) are clearly labelled<br>c. Images match the content of the manuscript<br>d. Quality of the images is adequate                             | 9.3 ± 1.6                 |
| <b>Contextual factors</b>                     | 18 Duration of ultrasound examination when relevant for the study question  | 8.8 ± 1.6                 |
|   | 19 <i>Optional:</i><br>a. <i>Whether ambient conditions (e.g. temperature, time of day) were kept stable during the study</i><br>b. <i>Potential confounding factors (e.g. exercise, alcohol, caffeine, smoking)</i>  | 7.3 ± 2.8                 |
| <b>Statistical analysis</b>                   | 20 a. Existence of a pre-specified statistical analysis plan and specification of post-hoc analyses<br>b. Analyses performed<br>c. Whether the analyses were performed at patient or at joint/region level<br>d. Extent of missing data<br>e. Handling of missing data                      | 9.3 ± 1.2                 |
| <b>Disclosures</b>                            | 21 Potential conflicts of interest including those related to ultrasound  | 9.2 ± 1.4                 |

\* Items are explained in detail in the glossary (Table 2).



Table 2. Glossary

| Item            | Terminology                                       | Definition  | Agreement†<br>(mean ± SD) |
|-----------------|---|---|---------------------------|
| <b>3</b>        | Prospective data collection                       | Data collection which starts before the outcome has occurred.   | 9.2 ± 1.7                 |
|                 | Retrospective data collection                     | Data collection which starts after outcome status has been determined, and refers to information up to that moment.   | 9.4 ± 1.2                 |
| <b>7</b>        | Broad domain                                      | A pathological (or pathophysiological) manifestation we are interested in assessing/measuring. For example, current broad domains in rheumatic and musculoskeletal diseases measured by ultrasound are “inflammation” and “structural damage”.  | 9.7 ± 0.8                 |
|                 | Target domain                                     | Further specification of the broad domain we are interested in assessing/measuring with ultrasound. For example, synovitis, enthesitis, erosion.  | 9.7 ± 0.6                 |
|                 | Theoretical definition of target domain           | The ultrasound definition of the target domain we want to measure, made up of domain components.  | 8.9 ± 1.7                 |
|                 | Target domain component                           | An individual characteristic of the target domain that can be measured. All domain components together constitute the theoretical definition. For example, synovial hypertrophy and synovial hyperaemia are the domain components that can be measured, and together define the target domain of synovitis as assessed by ultrasound.   | 9.1 ± 1.1                 |
|                 | Operational definition of target domain component | The ultrasound definition of a target domain component (i.e., the “signal” that can be detected by ultrasound). For example, synovial hypertrophy is defined as hypoechoic thickening of the synovium; and synovial hyperaemia as increased Doppler signal within the synovial hypertrophy.   | 8.9 ± 2.2                 |
| <b>8</b>        | Anatomical region                                 | The region of the body which is the focus of the ultrasound examination; it may include more than one related structure. For example, muscles, nerves and arteries, or joint cavity and tendons.  | 9.9 ± 0.2                 |
|                 | Anatomical structure                              | Isolated tissue(s) or organ(s) which is/are examined by ultrasound. For example, synovium, bone, tendon, muscle.  | 9.6 ± 1.3                 |
| <b>9</b>        | Dynamic examination                               | Procedure in which the transducer is moved along the anatomical region under study; or the anatomical region is moved during the ultrasound examination, for example through muscle contraction or tendon movement.   | 9.4 ± 1.3                 |
| <b>11,12</b>    | Reliability                                       | The degree to which the measurement is free from measurement error.   | 9.1 ± 1.4                 |
| <b>11,12,13</b> | Reader  | The person who is reading (i.e. interpreting) the ultrasound images or the video-clips of the examination. This interpretation may take place at the same time as the acquisition of the ultrasound images/video-clips, or later. In the latter case, the reader may be the same person who performed the ultrasound examination and the acquisition of the images/video-clips or a different person. | 9.9 ± 0.3                 |
|                 | Sonographer                                       | The person performing the ultrasound evaluation. Usually the sonographer is a health professional with the appropriate skills to perform an ultrasound examination in RMDs.   | 9.7 ± 0.9                 |
| <b>16</b>       | Ultrasound modalities                             | The ultrasound technique(s) used, i.e. grey-scale mode (or B mode), Doppler (colour, power, pulse), elastography, contrast-enhanced ultrasound, etc.  | 9.9 ± 0.2                 |

## References

- 1 Mandl P, Ciechomska A, Terslev L, *et al.* Implementation and role of modern musculoskeletal imaging in rheumatological practice in member countries of EULAR. *RMD Open* 2019;**5**:e000950. doi:10.1136/rmdopen-2019-000950
- 2 Scheel AK, Schmidt WA, Hermann K-GA, *et al.* Interobserver reliability of rheumatologists performing musculoskeletal ultrasonography: results from a EULAR “Train the trainers” course. *Ann Rheum Dis* 2005;**64**:1043–9. doi:10.1136/ard.2004.030387
- 3 Joshua F. Ultrasound applications for the practicing rheumatologist. *Best Practice & Research Clinical Rheumatology* 2012;**26**:853–67. doi:10.1016/j.berh.2012.10.002
- 4 Holsbeeck M van. Fury over sound. *Arthritis Care & Research* 2004;**51**:877–80. doi:10.1002/art.20835
- 5 Terslev L, Naredo E, Keen HI, *et al.* The OMERACT Stepwise Approach to Select and Develop Imaging Outcome Measurement Instruments: The Musculoskeletal Ultrasound Example. *J Rheumatol* 2019;**46**:1394–400. doi:10.3899/jrheum.181158
- 6 Joshua F, Lassere M, Bruyn GA, *et al.* Summary findings of a systematic review of the ultrasound assessment of synovitis. *J Rheumatol* 2007;**34**:839–47.
- 7 Wakefield RJ, Balint PV, Szkudlarek M, *et al.* Musculoskeletal ultrasound including definitions for ultrasonographic pathology. *J Rheumatol* 2005;**32**:2485–7.
- 8 Bruyn GA, Iagnocco A, Naredo E, *et al.* OMERACT Definitions for Ultrasonographic Pathologies and Elementary Lesions of Rheumatic Disorders 15 Years On. *J Rheumatol* 2019;**46**:1388–93. doi:10.3899/jrheum.181095
- 9 Mandl P, Naredo E, Wakefield RJ, *et al.* A systematic literature review analysis of ultrasound joint count and scoring systems to assess synovitis in rheumatoid arthritis according to the OMERACT filter. *J Rheumatol* 2011;**38**:2055–62. doi:10.3899/jrheum.110424
- 10 Lazarou I, D’Agostino M-A, Naredo E, *et al.* Ultrasound-guided synovial biopsy: a systematic review according to the OMERACT filter and recommendations for minimal reporting standards in clinical studies. *Rheumatology (Oxford)* 2015;**54**:1867–75. doi:10.1093/rheumatology/kev128
- 11 Schulz KF, Altman DG, Moher D, *et al.* CONSORT 2010 statement: updated guidelines for reporting parallel group randomized trials. *Ann Intern Med* 2010;**152**:726–32. doi:10.7326/0003-4819-152-11-201006010-00232
- 12 Bossuyt PM, Reitsma JB, Bruns DE, *et al.* Towards complete and accurate reporting of studies of diagnostic accuracy: The STARD Initiative. *Ann Intern Med* 2003;**138**:40–4. doi:10.7326/0003-4819-138-1-200301070-00010
- 13 Bossuyt PM, Reitsma JB, Bruns DE, *et al.* STARD 2015: an updated list of essential items for reporting diagnostic accuracy studies. *BMJ* 2015;**351**:h5527.

- 14 Dixon WG, Carmona L, Finckh A, *et al.* EULAR points to consider when establishing, analysing and reporting safety data of biologics registers in rheumatology. *Ann Rheum Dis* 2010;**69**:1596–602. doi:10.1136/ard.2009.125526
- 15 Buch MH, Silva-Fernandez L, Carmona L, *et al.* Development of EULAR recommendations for the reporting of clinical trial extension studies in rheumatology. *Ann Rheum Dis* 2015;**74**:963–9. doi:10.1136/annrheumdis-2013-204948
- 16 Heijde D van der, Aletaha D, Carmona L, *et al.* 2014 Update of the EULAR standardised operating procedures for EULAR-endorsed recommendations. *Annals of the Rheumatic Diseases* 2015;**74**:8–13. doi:10.1136/annrheumdis-2014-206350
- 17 Moher D, Schulz KF, Simera I, *et al.* Guidance for developers of health research reporting guidelines. *PLoS Med* 2010;**7**:e1000217. doi:10.1371/journal.pmed.1000217
- 18 Hróbjartsson A, Emanuelsson F, Skou Thomsen AS, *et al.* Bias due to lack of patient blinding in clinical trials. A systematic review of trials randomizing patients to blind and nonblind sub-studies. *Int J Epidemiol* 2014;**43**:1272–83. doi:10.1093/ije/dyu115
- 19 Hróbjartsson A, Thomsen ASS, Emanuelsson F, *et al.* Observer bias in randomized clinical trials with measurement scale outcomes: a systematic review of trials with both blinded and nonblinded assessors. *CMAJ* 2013;**185**:E201–211. doi:10.1503/cmaj.120744
- 20 Boers M, Beaton DE, Shea BJ, *et al.* OMERACT Filter 2.1: Elaboration of the Conceptual Framework for Outcome Measurement in Health Intervention Studies. *J Rheumatol* 2019;**46**:1021–7. doi:10.3899/jrheum.181096
- 21 Zappia M, Cuomo G, Martino MT, *et al.* The effect of foot position on Power Doppler Ultrasound grading of Achilles enthesitis. *Rheumatol Int* 2016;**36**:871–4. doi:10.1007/s00296-016-3461-z
- 22 Zayat AS, Freeston JE, Conaghan PG, *et al.* Does joint position affect US findings in inflammatory arthritis? *Rheumatology (Oxford)* 2012;**51**:921–5. doi:10.1093/rheumatology/ker383
- 23 Terslev L, D’Agostino MA, Brossard M, *et al.* Which knee and probe position determines the final diagnosis of knee inflammation by ultrasound? Results from a European multicenter study. *Ultraschall Med* 2012;**33**:E173–8. doi:10.1055/s-0031-1281973
- 24 Mandl P, Brossard M, Aegerter P, *et al.* Ultrasound evaluation of fluid in knee recesses at varying degrees of flexion. *Arthritis Care Res (Hoboken)* 2012;**64**:773–9. doi:10.1002/acr.21598
- 25 Hong BY, Lim SH, Cho YR, *et al.* Detection of knee effusion by ultrasonography. *Am J Phys Med Rehabil* 2010;**89**:715–21. doi:10.1097/PHM.0b013e3181e29c55
- 26 Koenig MJ, Torp-Pedersen ST, Christensen R, *et al.* Effect of knee position on ultrasound Doppler findings in patients with patellar tendon hyperaemia (jumper’s knee). *Ultraschall Med* 2007;**28**:479–83. doi:10.1055/s-2007-962865

- 27 Joshua F, de Carle R, Rayment M, *et al.* Power Doppler “blanching” after the application of transducer pressure. *Australas Radiol* 2005;**49**:218–21. doi:10.1111/j.1440-1673.2005.01435.x
- 28 Möller I, Janta I, Backhaus M, *et al.* The 2017 EULAR standardised procedures for ultrasound imaging in rheumatology. *Ann Rheum Dis* 2017;**76**:1974–9. doi:10.1136/annrheumdis-2017-211585
- 29 Backhaus M, Burmester G-R, Gerber T, *et al.* Guidelines for musculoskeletal ultrasound in rheumatology. *Annals of the Rheumatic Diseases* 2001;**60**:641–9. doi:10.1136/ard.60.7.641
- 30 US-2022. <https://esor.eular.org/enrol/index.php?id=159> (accessed 3 Apr 2020).
- 31 RhMSUS Certification. <https://www.rheumatology.org/Learning-Center/RhMSUS-Certification> (accessed 3 Apr 2020).
- 32 Torp-Pedersen S, Christensen R, Szkudlarek M, *et al.* Power and color Doppler ultrasound settings for inflammatory flow: impact on scoring of disease activity in patients with rheumatoid arthritis. *Arthritis & Rheumatology (Hoboken, NJ)* 2015;**67**:386–95. doi:10.1002/art.38940
- 33 Brulhart L, Ziswiler H-R, Tamborrini G, *et al.* The importance of sonographer experience and machine quality with regards to the role of musculoskeletal ultrasound in routine care of rheumatoid arthritis patients. *Clin Exp Rheumatol* 2015;**33**:98–101.
- 34 Ten Cate DF, Luime JJ, Swen N, *et al.* Role of ultrasonography in diagnosing early rheumatoid arthritis and remission of rheumatoid arthritis--a systematic review of the literature. *Arthritis Res Ther* 2013;**15**:R4. doi:10.1186/ar4132
- 35 Peterlein C-D, Fuchs-Winkelmann S, Schüttler K-F, *et al.* Does probe frequency influence diagnostic accuracy in newborn hip ultrasound? *Ultrasound Med Biol* 2012;**38**:1116–20. doi:10.1016/j.ultrasmedbio.2012.02.033
- 36 Koski JM, Alasaarela E, Soini I, *et al.* Ability of ultrasound imaging to detect erosions in a bone phantom model. *Ann Rheum Dis* 2010;**69**:1618–22. doi:10.1136/ard.2009.126680
- 37 Koski JM, Saarakkala S, Helle M, *et al.* Assessing the intra- and inter-reader reliability of dynamic ultrasound images in power Doppler ultrasonography. *Ann Rheum Dis* 2006;**65**:1658–60. doi:10.1136/ard.2005.051250
- 38 Albrecht K, Grob K, Lange U, *et al.* Reliability of different Doppler ultrasound quantification methods and devices in the assessment of therapeutic response in arthritis. *Rheumatology (Oxford)* 2008;**47**:1521–6. doi:10.1093/rheumatology/ken318
- 39 Lazaar H, Lhoste-Trouilloud A, Pereira B, *et al.* Does rheumatoid synovitis activity vary during the day? Evaluation with color doppler sonography. *BMC Musculoskelet Disord* 2017;**18**:98. doi:10.1186/s12891-017-1450-3
- 40 Semerano L, Gutierrez M, Falgarone G, *et al.* Diurnal variation of power Doppler in metacarpophalangeal joints of patients with rheumatoid arthritis: a preliminary study. *Ann Rheum Dis* 2011;**70**:1699–700. doi:10.1136/ard.2010.146761

- 41 Kilic G, Kilic E, Akgul O, *et al.* Ultrasonographic assessment of diurnal variation in the femoral condylar cartilage thickness in healthy young adults. *Am J Phys Med Rehabil* 2015;**94**:297–303. doi:10.1097/PHM.0000000000000179
- 42 Guillot X, Tordi N, Prati C, *et al.* Cryotherapy decreases synovial Doppler activity and pain in knee arthritis: A randomized-controlled trial. *Joint Bone Spine* 2017;**84**:477–83. doi:10.1016/j.jbspin.2016.09.004
- 43 Ellegaard K, Torp-Pedersen S, Henriksen M, *et al.* Influence of recent exercise and skin temperature on ultrasound Doppler measurements in patients with rheumatoid arthritis--an intervention study. *Rheumatology (Oxford)* 2009;**48**:1520–3. doi:10.1093/rheumatology/kep294
- 44 Strunk J, Strube K, Müller-Ladner U, *et al.* Three dimensional power Doppler ultrasonography confirms early reduction of synovial perfusion after intra-articular steroid injection. *Ann Rheum Dis* 2006;**65**:411–2. doi:10.1136/ard.2005.041210
- 45 Malliaras P, Chan O, Simran G, *et al.* Doppler ultrasound signal in Achilles tendinopathy reduces immediately after activity. *Int J Sports Med* 2012;**33**:480–4. doi:10.1055/s-0032-1304636
- 46 Boesen MI, Koenig MJ, Torp-Pedersen S, *et al.* Tendinopathy and Doppler activity: the vascular response of the Achilles tendon to exercise. *Scand J Med Sci Sports* 2006;**16**:463–9. doi:10.1111/j.1600-0838.2005.00512.x
- 47 Cook JL, Kiss ZS, Ptasznik R, *et al.* Is vascularity more evident after exercise? Implications for tendon imaging. *AJR Am J Roentgenol* 2005;**185**:1138–40. doi:10.2214/AJR.04.1205
- 48 Plint AC, Moher D, Morrison A, *et al.* Does the CONSORT checklist improve the quality of reports of randomised controlled trials? A systematic review. *Med J Aust* 2006;**185**:263–7.
- 49 Smidt N, Rutjes AWS, van der Windt D a. WM, *et al.* The quality of diagnostic accuracy studies since the STARD statement: has it improved? *Neurology* 2006;**67**:792–7. doi:10.1212/01.wnl.0000238386.41398.30