

Psychotic-Like Experiences in Adolescence Occurring in Combination or Isolation: Associations with Schizophrenia Risk Factors

Alastair G. Cardno, MB.ChB., Ph.D., Saskia Selzam, Ph.D., Daniel Freeman, Ph.D., Angelica Ronald, Ph.D.

Objectives: Individual adolescent psychotic-like experiences (PLEs) are associated with schizophrenia risk factors. As DSM-5 schizophrenia requires the co-occurrence of at least two psychotic symptoms, we investigated whether co-occurring adolescent PLEs have stronger associations with schizophrenia risk factors, lower quality of life and functioning, and have higher heritability, than individual PLEs.

Methods: Participants were 9646 16-year-old twins from the longitudinal Twins Early Development Study. We investigated co-occurrence of high questionnaire scores for three PLE combinations: (1) paranoia and hallucinations; (2) paranoia or hallucinations, and cognitive disorganization; and (3) paranoia or hallucinations, and negative symptoms, and their associations with 11 schizophrenia-relevant variables by regression analysis and structural equation twin modeling.

Results: Against expectation, none of the co-occurring PLEs had the nominally strongest associations significantly more often than individual PLEs. Co-occurring PLEs had

the strongest associations with bullying victimization, cannabis use and lower life satisfaction, but individual PLEs had the strongest associations with cognitive function variables. Obstetric complications were most associated with negative symptoms. Secondary analysis revealed that co-occurrence of cognitive disorganization and negative symptoms had the nominally strongest associations with most schizophrenia-relevant variables overall and relatively high heritability (67%).

Conclusions: Focusing on co-occurrence enhances some individual PLE associations but obscures others. The combination of subjective cognitive disorganization plus observed negative symptoms showed a broad range of enhanced associations with schizophrenia-relevant variables. Future research could investigate associations with other risk factors and the ability of this PLE combination to predict onset of schizophrenia.

Psych Res Clin Pract. 2021; 3:67–75; doi: 10.1176/appi.prcp.20200010

There is considerable interest in the relationships between psychotic-like experiences (PLEs) in the general population and schizophrenia (1, 2), particularly PLEs occurring during adolescence, just prior to the main period of onset of schizophrenia (3–5). As these PLEs are unaffected by the consequences of having a clinical disorder, such as the effects of antipsychotic medication, they may give new insights into the etiology and conceptualization of schizophrenia, and potentially inform intervention and prevention approaches.

The term PLE covers a wide range of phenomena occurring in the general population, from mild feelings of suspiciousness or unusual perceptions to positive psychotic symptoms such as delusions and hallucinations (3, 6). Broad definitions also include cognitive

HIGHLIGHTS

- Focusing on co-occurring psychotic-like experiences (PLEs) in adolescence enhances some individual PLE associations but may obscure others
- While victimization, cannabis use, and lower life satisfaction have broad associations with PLE combinations, cognitive variables are most associated with cognitive disorganization and negative symptoms, and obstetric complications are most associated with negative symptoms
- The combination of subjective cognitive disorganization and observed negative symptoms has a relatively broad range of enhanced associations with schizophrenia-relevant variables and relatively high heritability

disorganization and negative symptoms (3). PLEs are also known as psychotic experiences or psychotic experiences and negative symptoms (7), and have some similarities to the concept of schizotypy (8).

Individual PLEs are associated with a range of schizophrenia risk factors (9–14). DSM-5 schizophrenia requires the co-occurrence of at least two psychotic symptoms. Additionally, schizophrenia is associated with lower quality of life and functioning (15, 16), and has higher twin heritability (~80%) (17, 18) than individual general population PLEs (15%–59%) (14) as well as higher single nucleotide polymorphism heritability (7, 19). This raises the question of whether co-occurring PLEs have stronger associations with schizophrenia-relevant variables than their individual component PLEs.

Previous studies have found that, compared to one form of PLE occurring alone, co-occurrence of delusional and hallucinatory PLEs is associated with poorer functioning (20) and greater risk of psychotic disorder (21, 22), family history of psychosis (23), anxiety symptoms, negative symptoms, childhood trauma, persistence of PLEs, and clinical need (24). Co-occurrence of positive PLEs and negative symptoms is associated with poorer functioning (25) and greater risk of schizophrenia (2).

We aimed to investigate, for the first time specifically in adolescents, and broadening the range of prospectively assessed risk factors compared to previous studies, whether co-occurring PLEs have stronger associations with (a) schizophrenia risk factors, (b) lower quality of life and functioning, and (c) have higher twin heritability than individual PLEs.

METHODS

Participants

Participants were members of the Longitudinal Experiences And Perceptions (LEAP) study (3), that is part of the Twins Early Development Study (TEDS), a general population sample of monozygotic (MZ) and dizygotic (DZ) twins born in England and Wales in 1994–1996 and assessed longitudinally across childhood and adolescence (26). TEDS and LEAP have full ethical approval, and written informed consent was obtained after the procedures had been fully explained at each point of contact.

In total 10,874 families from TEDS were invited to take part in LEAP. Parent reports for 5076 (46.7%) families and twin reports for 5059 (46.5%) pairs were obtained (3). Adolescents involved in the LEAP project had a mean age of 16.3 years (SD 0.68) (3). Individuals were excluded if they did not provide consent at first contact (when TEDS was started) or for this study, had severe perinatal complications, or a severe medical disorder. After exclusions, the sample comprised 9646 individual twins with data on multiple PLEs.

Psychotic-Like Experiences

PLEs were assessed using the Specific Psychotic Experiences Questionnaire (SPEQ) at age 16 years (3). SPEQ includes self-report subscales for paranoia, hallucinations, and cognitive disorganization, and a parent-rated subscale for negative symptoms. Each subscale includes multiple items relating to the relevant concept (see supplementary methods for further information).

The primary analysis focused on the following PLE combinations: (1) paranoia and hallucinations; (2) paranoia or hallucinations, and cognitive disorganization; and (3) paranoia or hallucinations, and negative symptoms.

We defined PLEs as binary variables, being counted as present if their score was over the threshold closest to the top 15%, and co-occurring phenotypes as over this threshold for multiple PLEs, for example, for both paranoia and hallucinations. This gave at least 500 individual twins in each PLE group, and in previous analysis of this sample the heritabilities at this threshold were not significantly different from those at more extreme thresholds (14).

Schizophrenia Risk Factors

Schizophrenia risk factors were as follows: family history (27) measured as presence/absence of schizophrenia in a first or second degree relative (14) and validated with the schizophrenia polygenic risk score; older paternal age (28); ethnic minority status (29); obstetric complications (30); slower developmental milestones (31) measured with total vocabulary at age 2 years; lower premorbid IQ (32) measured as general cognitive ability (g) at age 12 years; bullying victimization (33) at age 12 years (12); and cannabis use (34) by age 16 years (13). Further information is presented in Table 1 and supplementary methods.

Quality of Life and Functioning Variables

Quality of life and functioning variables were life satisfaction measured at age 16 years and General Certificate of Secondary Education (GCSE) attainment at age 16 years, respectively. (See Table 1 and supplementary methods.)

Analysis

For schizophrenia risk factors, we used logistic regression analysis with the PLE groups as dependent variables. For example, for analysis of paranoia (P) and hallucinations (H) there were four exclusive groups: twins who scored high on neither P nor H (baseline group); those who scored high on P only; those who scored high on H only; and those who scored high on both P and H (P+H). The neither P nor H baseline group was compared with the P only group, the H only group, and the P+H group. The risk factor was the independent variable (with adjustment for birth order, age, and sex). We also adjusted for socioeconomic status at first contact whenever there were significant associations. We included all twins and adjusted for clustering within twin pairs using a generalized estimating equations approach.

TABLE 2. Demographic variables for PLE groups

PLE Group ^a	Demographic variable						
	No. individual twins		Sex ^b	Age in years		SES at first contact ^c	
	N	%	% Male	Mean	SD	Mean	SD
Paranoia (P) and hallucinations (H) (N = 9646)							
Neither P nor H	7217	74.8	45.6	16.3	0.68	0.25	0.99
P and not H	814	8.4	42.1	16.4	0.67	0.22	1.02
H and not P	986	10.2	41.2	16.3	0.67	0.11	0.96
Both P and H	629	6.5	38.5	16.3	0.72	0.15	1.00
Cognitive disorganization (CD) and paranoia or hallucinations (P/H) (N = 9629)							
Neither CD nor P/H	6633	68.9	46.9	16.3	0.68	0.27	0.98
CD and not P/H	572	5.9	31.1	16.3	0.68	0.11	0.99
P/H and not CD	1672	17.4	45.3	16.3	0.68	0.20	0.99
Both CD and P/H	752	7.8	30.7	16.3	0.68	0.07	0.99
Negative symptoms (NS) and paranoia or hallucinations (P/H) (N = 9579)							
Neither NS nor P/H	6147	64.2	44.3	16.3	0.68	0.29	0.98
NS and not P/H	1018	10.6	53.7	16.3	0.68	0.03	0.98
P/H and not NS	1844	19.3	40.7	16.3	0.67	0.25	0.98
Both NS and P/H	570	6.0	41.1	16.3	0.70	-0.13	0.98

Abbreviations: PLE, psychotic-like experience; SES, family socioeconomic status as a standardized composite score based on parental qualifications and employment - higher score means higher socioeconomic status.

^aPLE was defined as present if scoring in the top ~15% of the sample.

^bSex distributions differed significantly for all three PLE groupings: P and H ($\chi^2 = 19.20$, df 3, $p < 0.001$); CD and P/H ($\chi^2 = 115.29$, df 3, $p < 0.001$); NS and P/H ($\chi^2 = 48.59$, df 3, $p < 0.001$).

^cSES distributions differed significantly for all three PLE groupings: P and H ($F = 6.93$, df 9136, $p < 0.001$); CD and P/H ($F = 12.68$, df 9121, $p < 0.001$); NS and P/H ($F = 42.76$, df 9074, $p < 0.001$).

hallucinations; (2) paranoia or hallucinations, and cognitive disorganization; and (3) paranoia or hallucinations, and negative symptoms).

Analysis Software

We used SPSS version 24 (<https://www.ibm.com/products/spss-statistics>) for the analysis of risk factors, quality of life and functioning variables, and calculation of probandwise concordances. We used OpenMx (<https://openmx.ssri.psu.edu/>) for calculating tetrachoric correlations and twin modeling analyses.

RESULTS

Descriptive statistics are shown in Table 2. Paranoia, hallucinations, and cognitive disorganization were more common in females, while negative symptoms without paranoia or hallucinations was more common in males, consistent with previous reports (3). Family socioeconomic status at first contact was lower in individuals with negative symptoms, and to a lesser extent also in individuals with cognitive disorganization, and hallucinations (All p -values < 0.001 : Table 2).

Primary Analysis

Results are summarized in Table 3 and detailed in Tables S1.01–S3.29 of the online supplement. In none of the three primary analyses did the co-occurring PLE group have the strongest association with significantly more variables than its individual component PLEs overall (last row of Table 3: p -values 0.16–0.39).

Three variables had broad associations (bullying victimization, cannabis use, and lower life satisfaction), being significantly associated with almost every co-occurring and individual PLE group in all three analyses (the one exception was no significant association between cannabis use and negative symptoms only). For victimization, and life satisfaction, the co-occurring PLE group had the significantly strongest association in all three analyses, and for cannabis use, the co-occurring PLE group had the significantly strongest association in two out of the three analyses. Effect sizes were greatest for cannabis use, with ORs of 2.61 to 3.15 for co-occurring PLE groups.

Family history of schizophrenia had its nominally strongest association with the co-occurring PLE group in all three analyses. The association was significant for cognitive disorganization plus paranoia or hallucinations, and for negative symptoms plus paranoia or hallucinations. The family history variable was validated by showing significant association with the schizophrenia polygenic risk score (PRS; $n = 3951$, OR = 1.73 (95% CI 1.16 to 2.57), $p = 0.007$). We also performed a supplementary analysis of associations between PLE groups and schizophrenia PRS, but there were no statistically significant associations (Tables S4.01–S4.03 in the online supplement). This was probably due to insufficient sample size, as PLEs have shown significant associations in larger analyses that included the current sample (7, 11).

The three variables relating to cognitive function (vocabulary, general cognitive ability, and GCSE score) had their nominally strongest association with an individual

heritability was nominally highest for cognitive disorganization co-occurring with paranoia or hallucinations, while cognitive disorganization without paranoia or hallucinations had zero heritability due to unexpectedly higher DZ than MZ concordance. This is more likely to be a chance finding than to be scientifically meaningful because plausible mechanisms for the DZ>MZ pattern are hard to envisage. In the negative symptoms analysis, all three PLE groups had significant heritability, and negative symptoms without paranoia or hallucinations was nominally highest.

Secondary Analysis

As cognitive disorganization and negative symptoms groups were both associated with all three of the cognitive function variables, we speculated that co-occurrence of these two PLEs might have stronger associations with cognitive variables than either occurring alone. We therefore conducted a secondary analysis of cognitive disorganization and negative symptoms occurring together or individually.

The summary results are given in Table S5.01 and detailed in Tables S5.02–S5.33 of the online supplement. Co-occurring cognitive disorganization and negative symptoms had the nominally strongest association with two of the cognitive variables (general cognitive ability, and GCSE score—significantly strongest for GCSE score), while cognitive disorganization without negative symptoms was nominally strongest for vocabulary. Overall, the co-occurring cognitive disorganization and negative symptoms group had the nominally strongest association with 7 of the 10 variables where at least one PLE group was significantly associated (Table S5.01 in the online supplement: proportion 0.70, $p=0.014$).

The co-occurring cognitive disorganization and negative symptoms group had the nominally highest heritability from all of the PLE analyses (0.67 or 67%), although CIs overlapped with the cognitive disorganization only and negative symptoms only groups. The heritability estimate remained similar when based on residualized PLEs after regressing out the effects of sex (0.65, 95%CI 0.23 to 0.77), in male (0.64, 95%CI 0.36 to 0.83) and female (0.67, 95%CI 0.18 to 0.79) twin pairs, and when including opposite-sex DZ twin pairs (0.65, 95%CI 0.41 to 0.76).

In order to investigate the effects of applying a higher PLE threshold, we conducted the following analyses using a 10% cut-off. For all PLE combinations, we investigated associations with two prospectively assessed risk factors that showed contrasting patterns of association, general cognitive ability, and victimization. Additionally, for paranoia or hallucinations, and negative symptoms we investigated associations with obstetric complications, and for cognitive disorganization and negative symptoms we investigated twin heritability. The results remained very similar (Tables S6.01–S6.15 in the online supplement). The same associations were statistically significant at 15% and 10% cut-offs. Among these, the only change in order of nominal effect

sizes was that at the 10% cut-off negative symptoms without paranoia or hallucinations had a marginally stronger association with obstetric complications than negative symptoms plus paranoia or hallucinations (OR 3.04, 95%CI 1.76 to 5.26 vs. 2.87, 95%CI 1.30 to 6.33), reinforcing the previous impression that obstetric complications were most associated with negative symptoms. Also, for most associations at the 10% cut-off the largest effect size was marginally greater than at the 15% cut-off.

DISCUSSION

Three variables—bullying victimization, cannabis use, and lower life satisfaction—had broad associations with most PLE groups and their strongest associations with co-occurring PLEs, as predicted. These variables might be particularly related to clinical service use, through reduced quality of life or as proximal precipitants of clinical episodes. They may thus contribute to effects seen in adult samples where, relative to individual PLEs, co-occurring PLEs are associated with increased risk of clinical need (24), schizophrenia (2), and psychotic disorder (21, 22), and other clinically relevant variables, including poorer functioning (20, 25), anxiety symptoms (24) and persistence of PLEs (24). Also, consistent with the current study, co-occurring PLEs have previously been associated with childhood trauma (24): in that study any lifetime trauma when interviewed at 14–24 years, and in this study with bullying victimization assessed at age 12 years.

Co-occurring PLEs have previously been associated with more frequent family history of psychosis (23). In the current study, family history of schizophrenia in a first or second degree relative had its nominally strongest association with the co-occurring PLE group in all three primary comparisons.

The three variables relating to cognitive function had their strongest association with an individual PLE in all three primary analyses. This may have been due to paranoia not being associated with reduced cognitive function (and having a trend in the opposite direction in some analyses) leading to associations for co-occurring PLE groups that included paranoia being attenuated. To our knowledge, previous studies of co-occurring PLEs have not investigated cognitive functioning prospectively. However, consistent with our findings, the Avon Longitudinal Study of Parents and Children (ALSPAC) study found only weak evidence for cognitive deficits in adolescence for a psychotic experiences group defined by the presence of hallucination- or delusion-like experiences (36).

Obstetric complications have previously been studied in relation to positive PLEs with mixed results (37, 38), while we found them to be most associated with negative symptoms. Ethnic minority status has been associated with positive PLEs (39). Unfortunately our findings for PLE combinations were inconsistent, and further research is needed in samples that are more ethnically diverse. We

found that paternal age was not significantly associated with any PLE group. The ALSPAC study found a non-significant trend towards an association between positive PLEs at age 12 years and older paternal age (40). As some studies of schizophrenia have found the association only with the oldest fathers (28), and there was a trend consistent with this in the current study, albeit with too small numbers for formal analysis, it is possible that an association with PLEs might be evident in larger samples.

In the secondary analysis of cognitive disorganization and negative symptoms, the group where these two PLEs co-occurred had the strongest association with schizophrenia-relevant variables most frequently compared with their individual component PLEs, that is, cognitive disorganization alone and negative symptoms alone. It was also notable that this PLE combination had the highest heritability out of all the PLE groups analyzed (67%). To our knowledge this PLE combination has not been investigated in previous co-occurrence studies. A negative/disorganized PLE group has been investigated in a past study, but it was predominantly composed of negative symptoms and the component PLEs were not disaggregated (25).

Are there potential implications of these PLE results for clinical diagnoses that are based on the co-occurrence of symptoms? Where processes underlying PLEs and symptoms are similar, a diagnostic approach might enhance associations with risk factors where most individual symptoms are associated, but could attenuate associations with risk factors where only one type of symptom, for example, negative symptoms, is associated. However, this depends on factors including the relationships between PLEs and symptoms and the processes that underlie the co-occurrence (22–24). Further insights may be gained from ongoing research, including longitudinal studies of PLEs and subsequent clinical disorders and their symptom profiles, where relevant risk factors are assessed.

Limitations

The variables used covered a broader range of prospectively measured relevant variables than previously investigated, but it is possible that the results could have differed if other schizophrenia-relevant variables were used that were not available in this sample, for example, measures of motor development, other forms of childhood trauma, or other aspects of impairment in life roles. Also results could have differed if alternative questionnaires or interview measures were employed, although the SPEQ is based on well-established measures that provide reliable and valid PLE subscales.

We defined PLEs in terms of dichotomized scores, that may have reduced power compared to quantitative variables, but gave relative clarity for defining PLE groups and flexibility in allowing both “and” and “or” combinations of PLEs.

Correlations between PLEs and schizophrenia-relevant variables could be elevated when they had the same rater, and lowered when there were different raters, and the power to detect associations may vary between risk factors because of differences in their prevalence, but the analysis focused on comparing associations for different PLE combinations with each risk factor individually, and then conducting an overall test based on how frequently the co-occurring PLE group had the strongest association.

We were able to investigate more schizophrenia risk factors than measures of quality of life and functioning. Future research in older participants could extend this to include assessment of, for example, further education, employment, and social relationships.

As part of the development of the PLE measures, twin and singleton comparisons were made and showed no notable differences (3)—twin-related effects are unlikely but cannot be excluded.

These findings require independent replication, especially as many are reported here for the first time.

CONCLUSIONS

First, we found the patterns of association between PLE combinations and schizophrenia-relevant variables to vary considerably. Focusing on co-occurrence enhances some individual PLE associations but obscures others. Second, in terms of which variables are most associated with co-occurring PLEs, bullying victimization, cannabis use, and lower life satisfaction had the strongest associations with co-occurring PLEs compared with PLEs in isolation. These variables may be particularly relevant to the association between co-occurring PLEs and increased risk of clinical disorder and service use. Third, obstetric complications were most associated with negative symptoms, a finding that suggests opportunities for future research into the underlying neurodevelopmental pathways to psychosis vulnerability. Fourth, we found that co-occurring subjective cognitive disorganization and observed negative symptoms had the at-least nominally strongest associations across a broad range of schizophrenia-relevant variables and relatively high heritability. Future research could investigate associations between this PLE combination and other schizophrenia risk factors, and also cumulative exposure to risk factors, and assess how well it predicts onset of schizophrenia.

AUTHOR AND ARTICLE INFORMATION

Division of Psychological and Social Medicine, Faculty of Medicine and Health, University of Leeds, Leeds (Cardno); Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London (Selzam); Oxford Cognitive Approaches to Psychosis (O-CAP), Department of Psychiatry, University of Oxford, and Oxford Health NHS Foundation Trust, Oxford (Freeman); Genes Environment Lifespan (GEL) Laboratory, Centre for Brain and Cognitive Development, Department of Psychological Sciences, Birkbeck, University of London, London (Ronald).

Send correspondence to Dr. Cardno (a.g.cardno@leeds.ac.uk).

We are very grateful to the participants of TEDS for making this research possible, to Robert Plomin and Andrew McMillan for the TEDS collaboration, and to Frühling Rijdsdijk for the twin modeling script and comments. The TEDS study of psychotic-like experiences was funded by Medical Research Council grant G1100559 to AR and a Wellcome Trust ISSF grant to AR. TEDS is funded by Medical Research Council grant MR/M021475/1 to Robert Plomin. High performance computing facilities were funded with capital equipment grants from the Guy's and St Thomas' Charity (TR130505) and Maudsley Charity (980). Alastair G. Cardno has received research funding from the Qatar National Research Fund. Saskia Selzam reports no financial relationships with commercial interests. Daniel Freeman is a co-founder, and shareholder in Oxford VR, a University of Oxford spin-out company. Daniel Freeman has received grant funding from the National Institute for Health Research, Medical Research Council, and Wellcome Trust. Angelica Ronald has received fees for consultancy work for the National Childbirth Trust. Angelica Ronald receives an annual honorarium as joint editor of the *Journal of Child Psychology and Psychiatry*. Angelica Ronald has received research funding from the Wellcome Trust.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *Psychiatric Research and Clinical Practice* published by Wiley Periodicals LLC. on behalf of the American Psychiatric Association.

Received April 1, 2020; revised October 27, 2020; accepted October 28, 2020.

REFERENCES

- Fisher HL, Caspi A, Poulton R, et al: Specificity of childhood psychotic symptoms for predicting schizophrenia by 38 years of age: a birth cohort study. *Psychol Med*. 2013; 43:2077–2086
- Werbelloff N, Dohrenwend BP, Yoffe R, et al: The association between negative symptoms, psychotic experiences and later schizophrenia: a population-based longitudinal study. *PLoS One*. 2015; 10:e0119852
- Ronald A, Sieradzka D, Cardno AG, et al: Characterization of psychotic experiences in adolescence using the specific psychotic experiences questionnaire: findings from a study of 5000 16-year-old twins. *Schizophr Bull*. 2014; 40:868–877
- Wigman JT, Vollebergh WA, Raaijmakers QA, et al: The structure of the extended psychosis phenotype in early adolescence—a cross-sample replication. *Schizophr Bull*. 2011; 37:850–860
- Yung AR, Nelson B, Baker K, et al: Psychotic-like experiences in a community sample of adolescents: implications for the continuum model of psychosis and prediction of schizophrenia. *Aust N Z J Psychiatry*. 2009; 43:118–128
- van Os J, Linscott RJ, Myin-Germeys I, et al: A systematic review and meta-analysis of the psychosis continuum: evidence for a psychosis proneness-persistence-impairment model of psychotic disorder. *Psychol Med*. 2009; 39:179–195
- Ronald A, Pain O: A systematic review of genome-wide research on psychotic experiences and negative symptom traits: new revelations and implications for psychiatry. *Hum Mol Genet*. 2018; 27:R136–R152
- Mason O, Linney Y, Claridge G: Short scales for measuring schizotypy. *Schizophr Res*. 2005; 78:293–296
- Kelleher I, Cannon M: Psychotic-like experiences in the general population: characterizing a high-risk group for psychosis. *Psychol Med*. 2011; 41:1–6
- Linscott RJ, van Os J: An updated and conservative systematic review and meta-analysis of epidemiological evidence on psychotic experiences in children and adults: on the pathway from proneness to persistence to dimensional expression across mental disorders. *Psychol Med*. 2013; 43:1133–1149
- Pain O, Dudbridge F, Cardno AG, et al: Genome-wide analysis of adolescent psychotic-like experiences shows genetic overlap with psychiatric disorders. *Am J Med Genet B Neuropsychiatr Genet*. 2018; 177:416–425
- Shakoor S, McGuire P, Cardno AG, et al: A shared genetic propensity underlies experiences of bullying victimization in late childhood and self-rated paranoid thinking in adolescence. *Schizophr Bull*. 2015; 41:754–763
- Shakoor S, Zavos HM, McGuire P, et al: Psychotic experiences are linked to cannabis use in adolescents in the community because of common underlying environmental risk factors. *Psychiatry Res*. 2015; 227:144–151
- Zavos HM, Freeman D, Haworth CM, et al: Consistent etiology of severe, frequent psychotic experiences and milder, less frequent manifestations: a twin study of specific psychotic experiences in adolescence. *JAMA Psychiatry*. 2014; 71:1049–1057
- Kahn RS, Sommer IE, Murray RM, et al: Schizophrenia. *Nat Rev Dis Primers*. 2015; 1:15067
- Tandon R, Keshavan MS, Nasrallah HA: Schizophrenia, “Just the Facts”: what we know in 2008 part 1: overview. *Schizophr Res*. 2008; 100:4–19
- Cardno AG, Gottesman II: Twin studies of schizophrenia: from bow-and-arrow concordances to star wars Mx and functional genomics. *Am J Med Genet*. 2000; 97:12–17
- Sullivan PF, Kendler KS, Neale MC: Schizophrenia as a complex trait: evidence from a meta-analysis of twin studies. *Arch Gen Psychiatry*. 2003; 60:1187–1192
- Legge SE, Jones HJ, Kendall KM, et al: Association of genetic liability to psychotic experiences with neuropsychotic disorders and traits. *JAMA Psychiatry*. 2019; 76:1256–1265
- Nuevo R, Van Os J, Arango C, et al: Evidence for the early clinical relevance of hallucinatory-delusional states in the general population. *Acta Psychiatr Scand*. 2013; 127:482–493
- Kirli U, Binbay T, Drukker M, et al: DSM outcomes of psychotic experiences and associated risk factors: 6-year follow-up study in a community-based sample. *Psychol Med*. 2019; 49:1346–1356
- Krabbendam L, Myin-Germeys I, Hanssen M, et al: Hallucinatory experiences and onset of psychotic disorder: evidence that the risk is mediated by delusion formation. *Acta Psychiatr Scand*. 2004; 110:264–272
- Smeets F, Lataster T, van Winkel R, et al: Testing the hypothesis that psychotic illness begins when subthreshold hallucinations combine with delusional ideation. *Acta Psychiatr Scand*. 2013; 127:34–47
- Smeets F, Lataster T, Dominguez MD, et al: Evidence that onset of psychosis in the population reflects early hallucinatory experiences that through environmental risks and affective dysregulation become complicated by delusions. *Schizophr Bull*. 2012; 38:531–542
- Dominguez MD, Saka MC, Lieb R, et al: Early expression of negative/disorganized symptoms predicting psychotic experiences and subsequent clinical psychosis: a 10-year study. *Am J Psychiatry*. 2010; 167:1075–1082
- Haworth CM, Davis OS, Plomin R: Twins Early Development Study (TEDS): a genetically sensitive investigation of cognitive and behavioral development from childhood to young adulthood. *Twin Res Hum Genet*. 2013; 16:117–125
- Lichtenstein P, Yip BH, Bjork C, et al: Common genetic determinants of schizophrenia and bipolar disorder in Swedish families: a population-based study. *Lancet*. 2009; 373:234–239
- de Kluiver H, Buizer-Voskamp JE, Dolan CV, et al: Paternal age and psychiatric disorders: a review. *Am J Med Genet B Neuropsychiatr Genet*. 2017; 174:202–213

29. Kirkbride JB, Hameed Y, Ioannidis K, et al: Ethnic minority status, age-at-immigration and psychosis risk in rural environments: evidence from the SEPEA study. *Schizophr Bull.* 2017; 43:1251–1261
30. Cannon M, Jones PB, Murray RM: Obstetric complications and schizophrenia: historical and meta-analytic review. *Am J Psychiatry.* 2002; 159:1080–1092
31. Jones P, Rodgers B, Murray R, et al: Child development risk factors for adult schizophrenia in the British 1946 birth cohort. *Lancet.* 1994; 344:1398–1402
32. Woodberry KA, Giuliano AJ, Seidman LJ: Premorbid IQ in schizophrenia: a meta-analytic review. *Am J Psychiatry.* 2008; 165:579–587
33. Varese F, Smeets F, Drukker M, et al: Childhood adversities increase the risk of psychosis: a meta-analysis of patient-control, prospective- and cross-sectional cohort studies. *Schizophr Bull.* 2012; 38:661–671
34. Marconi A, Di Forti M, Lewis CM, et al: Meta-analysis of the association between the level of cannabis use and risk of psychosis. *Schizophr Bull.* 2016; 42:1262–1269
35. Rijdsdijk FV, Sham PC: Analytic approaches to twin data using structural equation models. *Brief Bioinform.* 2002; 3:119–133
36. Mollon J, David AS, Zammit S, et al: Course of cognitive development from infancy to early adulthood in the psychosis spectrum. *JAMA Psychiatry.* 2018; 75:270–279
37. Spauwen J, Krabbendam L, Lieb R, et al: Early maternal stress and health behaviours and offspring expression of psychosis in adolescence. *Acta Psychiatr Scand.* 2004; 110:356–364
38. Zammit S, Odd D, Horwood J, et al: Investigating whether adverse prenatal and perinatal events are associated with non-clinical psychotic symptoms at age 12 years in the ALSPAC birth cohort. *Psychol Med.* 2009; 39:1457–1467
39. Leane E, Dealberto MJ, Luck D, et al: Ethnic minority position and migrant status as risk factors for psychotic symptoms in the general population: a meta-analysis. *Psychol Med.* 2019; 49:545–558
40. Zammit S, Horwood J, Thompson A, et al: Investigating if psychosis-like symptoms (PLIKS) are associated with family history of schizophrenia or paternal age in the ALSPAC birth cohort. *Schizophr Res.* 2008; 104:279–286