



UNIVERSITY OF LEEDS

This is a repository copy of *On the formal theory of pseudomonads and pseudodistributive laws*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/170302/>

Version: Accepted Version

Article:

Gambino, N orcid.org/0000-0002-4257-3590 and Lobbia, G orcid.org/0000-0002-2732-0317 (2021) On the formal theory of pseudomonads and pseudodistributive laws. *Theory and Applications of Categories*, 37 (2). pp. 14-56. ISSN 1201-561X

© Nicola Gambino and Gabriele Lobbia, 2021. Permission to copy for private use granted. This is an author produced version of a paper published in *Theory and Applications of Categories*. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

ON THE FORMAL THEORY OF PSEUDOMONADS AND PSEUDODISTRIBUTIVE LAWS

NICOLA GAMBINO AND GABRIELE LOBBIA

ABSTRACT. We contribute to the formal theory of pseudomonads, *i.e.* the analogue for pseudomonads of the formal theory of monads. In particular, we solve a problem posed by Lack by proving that, for every Gray-category \mathcal{K} , there is a Gray-category $\mathbf{Psm}(\mathcal{K})$ of pseudomonads, pseudomonad morphisms, pseudomonad transformations and pseudomonad modifications in \mathcal{K} . We then establish a triequivalence between $\mathbf{Psm}(\mathcal{K})$ and the Gray-category of pseudomonads introduced by Marmolejo and give a simpler proof of the equivalence between pseudodistributive laws and liftings of pseudomonads to 2-categories of pseudoalgebras.

INTRODUCTION

Context and motivation. Monads are one of the fundamental notions of category theory [22, Chapter VI]. For example, they provide a homogeneous approach to the study of categories of sets equipped with algebraic structure, such as groups and monoids [1]. Furthermore, Beck’s theorem on distributive laws between monads [2] describes concisely the structure that is necessary and sufficient in order to combine two algebraic structures, so that the operations of one distribute over those of the other. For example, the monads for groups and for monoids can be combined via a distributive law to define the monad for rings. Subsequently, the formal theory of monads, originally introduced by Street [28] and later developed further by Lack and Street [21], has offered an elegant and mathematically efficient account of the theory of monads, starting from the observation that the notion of a monad can be defined within any 2-category (so that the usual notion is recovered by considering the 2-category of categories, functors and natural transformations). Among many other results, it provides a characterisation of the existence of categories of Eilenberg-Moore algebras as a completeness property and, importantly for our purposes, a simple account of Beck’s theorem on distributive laws.

In recent years, motivation from pure mathematics, *e.g.* in the theory of operads [9–12], and theoretical computer science, *e.g.* in the study of variable binding [4, 6, 31], led to significant interest in pseudomonads [19, 23, 25, 32], which are the counterparts of monads in 2-dimensional category theory, obtained by requiring the axioms for a monad to hold only up to coherent isomorphism rather than strictly [3]. Here, one of the key issues has been the proof of a counterpart of Beck’s theorem on distributive laws, which requires a satisfactory axiomatisation of the notion of a pseudodistributive law [5, 24, 26, 29, 30], building on early work of Kelly [16] on semi-strict distributive laws. This is a difficult question because such a notion necessarily involves complex coherence conditions.

Date: January 16th, 2021.

Given how the formal theory of monads offers a simple proof of Beck’s theorem on distributive laws, it seems natural to attack this problem by developing a formal theory of pseudomonads. In order to do this, however, one needs to face the challenge that, just as the formal theory of monads is formulated within 2-dimensional category theory [18], the formal theory of pseudomonads is formulated within 3-dimensional category theory [14, 15], which is notoriously hard. In this setting, it is convenient to work with Gray-categories, *i.e.* semistrict tricategories [14, Section 4.8], which are easier to handle than tricategories, but sufficiently general for many purposes, since every tricategory is triequivalent to a Gray-category [14, Theorem 8.1].

In spite of significant advances in the creation of a formal theory of pseudomonads in the works cited above, there are still fundamental questions to be addressed. In particular, there is not yet a direct counterpart of the 2-category $\mathbf{Mnd}(\mathcal{K})$ of monads, monad morphisms and monad transformations in a 2-category \mathcal{K} , which is the starting point of the formal theory of monads [28]. Filling this gap would involve the definition, for a Gray-category \mathcal{K} , of a 3-dimensional category $\mathbf{Psm}(\mathcal{K})$ having pseudomonads in \mathcal{K} as 0-cells, pseudomonad morphisms as 1-cells, and appropriately defined pseudomonad transformations and pseudomonad modifications as 2-cells and 3-cells, respectively. This issue was raised by Lack in [19, Section 6], who suspected that defining $\mathbf{Psm}(\mathcal{K})$ in this way would give rise only to a tricategory, not a Gray-category, and hence require lengthy verifications of the coherence conditions. For this reason, Lack preferred to define a Gray-category of pseudomonads in \mathcal{K} using the description of pseudomonads in \mathcal{K} as suitable lax functors and developing parts of the theory using enriched category theory.

Yet another approach was taken earlier by Marmolejo in [24], who introduced, for a Gray-category \mathcal{K} , a Gray-category that we denote here $\mathbf{Lift}(\mathcal{K})$ to avoid confusion, that has pseudomonads in \mathcal{K} as 0-cells and liftings of 1-cells, 2-cells and 3-cells of \mathcal{K} to 2-categories of pseudoalgebras as 1-cells, 2-cells and 3-cells, respectively. He then used $\mathbf{Lift}(\mathcal{K})$ to introduce the notion of a lifting of a pseudomonad and of a pseudodistributive law, proving the fundamental result that pseudodistributive laws are equivalent liftings of pseudomonads are equivalent to [24, Theorems 6.2, 9.3 and 10.2], thus obtaining an analogue of Beck’s result on distributive laws. Here, Marmolejo defined pseudodistributive laws explicitly, giving nine coherence conditions for them [24]. Later, Marmolejo and Wood [26] showed not only that an additional tenth coherence condition, introduced by Tanaka [29], can be derived from Marmolejo’s conditions, but also that one of the original nine conditions introduced by Marmolejo is derivable from the others, thus reducing the number of coherence axioms for a pseudodistributive law to eight.

Main results. The aim of this paper is take some further steps in the development of the formal theory of pseudomonads. In particular, our main contributions are the following:

- Theorem 2.5, which answers the question raised in [19] by showing that for every Gray-category \mathcal{K} , there is a Gray-category $\mathbf{Psm}(\mathcal{K})$ of pseudomonads, pseudomonad morphisms, pseudomonad transformations and pseudomonad modifications in \mathcal{K} ;
- Theorem 3.4, the analogue of a fundamental result of the formal theory of monads, asserting that $\mathbf{Psm}(\mathcal{K})$ is equivalent, in a suitable 3-categorical sense, to the Gray-category $\mathbf{Lift}(\mathcal{K})$;

- Proposition 4.4, recording that an object of $\mathbf{Psm}(\mathbf{Psm}(\mathcal{K}))$ is the same thing as a pseudodistributive law in \mathcal{K} ;
- a new, simpler proof of Marmolejo’s theorem equivalence between pseudodistributive laws and liftings of pseudomonads to 2-categories of pseudoalgebras, given as the proof of Theorem 4.5.

Theorem 2.5 supports the definition of a pseudodistributive law of [24, 26], since it allows us to show that a pseudodistributive law is the same thing as a pseudomonad in $\mathbf{Psm}(\mathcal{K})$ (Proposition 4.4), as one would expect by analogy with the situation in the formal theory of monads. Thanks to this observation, we provide an interpretation of the complex coherence conditions for a pseudodistributive law in terms of the simpler ones, namely those for a pseudomonad morphism, a pseudomonad transformation and a pseudomonad modification (see Table 2 for details). This point of view allows us to give a principled presentation of the conditions for pseudodistributive laws of [24, 29], included in Appendix A, which hopefully provides a useful reference for future work in this area. For the convenience of readers, we also describe how our formulation relates to the ones of Marmolejo and of Tanaka (see Table 1).

Theorem 3.4, which establishes the equivalence between $\mathbf{Psm}(\mathcal{K})$ and $\mathbf{Lift}(\mathcal{K})$, does not seem to be part of the literature (in part because its very statement requires the introduction of the 3-dimensional category $\mathbf{Psm}(\mathcal{K})$, which is defined here for the first time), but extends existing results. In particular, the equivalence between pseudomonad morphisms and liftings of morphisms to categories of pseudoalgebras is proved in [26]. Related results appear also in [29], but with important differences. First, the work carried out therein is developed for the particular tricategory $\mathbf{2-Cat}_{\text{psd}}$ of 2-categories, pseudofunctors, pseudonatural transformations and modifications, rather than for a general tricategory or Gray-category. While that is an important example (*cf.* Remark 4.7), restricting to a particular tricategory does not allow us to exploit the various dualities that are essential to derive results in the formal theory. Secondly, the results obtained therein focus on hom-2-categories of pseudomonad endomorphisms, *i.e.* of the form $\mathbf{Psm}(\mathcal{K})((X, S), (X, S))$, rather than on general hom-2-categories of pseudomonad morphisms.

Our proof Theorem 4.5, which asserts the equivalence between pseudodistributive laws and liftings of pseudomonads to 2-categories of pseudoalgebras established in [24], follows naturally combining Theorem 2.5 and Theorem 3.4. More specifically, combining our identification of pseudodistributive laws with pseudomonads in $\mathbf{Psm}(\mathcal{K})$ of Proposition 4.4 with the fact that a pseudomonad in $\mathbf{Lift}(\mathcal{K})$ is a lifting of a pseudomonad T to the 2-categories of pseudoalgebras of another pseudomonad S , we obtain the desired equivalence between a pseudodistributive law of S over T and a lifting of T to pseudo- S -algebras. This proof is simpler than that in [24] since it takes a modular, more abstract, approach to the verification of the coherence conditions and avoids completely the notion of a composite of pseudomonads with compatible structure.

As the proofs of our main results involve lengthy, subtle calculations with pasting diagrams, we tried to strike a reasonable compromise between rigour and conciseness by giving what we hope are the key diagrams of the proofs, and describing the additional steps in the text. When in doubt, we preferred to err on the side of rigour, since one of our initial goals was to answer the question raised in [19] about whether $\mathbf{Psm}(\mathcal{K})$ is

a Gray-category or not. We hope that this did not make the paper too long. For the convenience of the readers, some of the diagrams are confined to the Appendices.

Outline of the paper. Section 1 provides background on Gray-categories, pseudomonads and 2-categories of pseudoalgebras. Section 2 defines the Gray-category $\mathbf{Psm}(\mathcal{K})$. We prove the equivalence of $\mathbf{Psm}(\mathcal{K})$ and $\mathbf{Lift}(\mathcal{K})$ in Section 3. We conclude the paper in Section 4 by discussing pseudodistributive laws.

Acknowledgements. We are very grateful to Steve Lack for patiently discussing his work on pseudomonads with us, especially during the 2019 International Conference on Category Theory at the University of Edinburgh. We wish to thank also John Bourke, Emily Riehl and Nick Gurski for fruitful conversations, Martin Hyland for encouraging us to consider carefully the directions of 3-cells, and the anonymous referee for helpful comments. Nicola Gambino gratefully acknowledges the support of EPSRC under grant EP/M01729X/1, of the US Air Force Office for Scientific Research under agreement FA8655-13-1-3038 and the hospitality of the School of Mathematics of the University of Manchester while on study leave from the University of Leeds in 2018/19. Gabriele Lobbia gratefully acknowledges the support of an EPSRC PhD Scholarship.

1. PRELIMINARIES

Gray-categories. We begin by reviewing the notion of a Gray-category and fixing some notation. A Gray-category can be defined very succinctly in terms of enriched category theory (see Remark 1.4). For our purposes, however, it is useful to give an explicit definition, which we recall from [24, Section 2] in Definition 1.1 below. The explicit definition makes it easier to see that Gray-categories are special tricategories [14, Proposition 3.1] in which the only non-strict operation is horizontal composition of 2-cells [14, Section 5.2]. Throughout this paper, for a Gray-category \mathcal{K} , we use X, Y, Z, \dots to denote its 0-cells, $F: X \rightarrow Y, G: Y \rightarrow Z, \dots$ for its 1-cells, $f: F \rightarrow F', g: G \rightarrow G' \dots$ for 2-cells, and $\alpha: f \rightarrow f', \beta: g \rightarrow g' \dots$ for 3-cells.

When stating the definition of a Gray-category below, we make use of the notion of a cubical functor from [14], which we unfold in Remark 1.2.

Definition 1.1. A Gray-category \mathcal{K} consists of the the data in (G1)-(G4), subject to axioms (G5) and (G6), as given below.

- (G1) A class of objects \mathcal{K}_0 . We call the elements of \mathcal{K}_0 the 0-cells of \mathcal{K} .
- (G2) For every $X, Y \in \mathcal{K}_0$, a 2-category $\mathcal{K}(X, Y)$. We refer to the n -cells of these 2-categories as the $(n + 1)$ -cells of \mathcal{K} .
- (G3) For every $X, Y, Z \in \mathcal{K}_0$, a cubical functor

$$\mathcal{K}(Y, Z) \times \mathcal{K}(X, Y) \rightarrow \mathcal{K}(X, Z),$$

whose action on $F: X \rightarrow Y$ and $G: Y \rightarrow Z$ is written $GF: X \rightarrow Z$, and whose action on $f: F \rightarrow F'$ and $g: G \rightarrow G'$ gives rise to an invertible 3-cell

$$\begin{array}{ccc} GF & \xrightarrow{Gf} & GF' \\ gF \downarrow & \Downarrow g_f & \downarrow gF' \\ G'F & \xrightarrow{G'f} & G'F' \end{array}$$

called the *interchange maps* of \mathcal{K} .

(G4) For any $X \in \mathcal{K}_0$, a 1-cell $1_X: X \rightarrow X$. We call these the identity 1-cells of \mathcal{K} .

(G5) For every

$$F \begin{array}{c} \xrightarrow{f} \\ \Downarrow \alpha \\ \xrightarrow{f'} \end{array} F' \quad G \begin{array}{c} \xrightarrow{g} \\ \Downarrow \beta \\ \xrightarrow{g'} \end{array} G' \quad K \begin{array}{c} \xrightarrow{k} \\ \Downarrow \gamma \\ \xrightarrow{k'} \end{array} K'$$

in $\mathcal{K}(X, Y)$, $\mathcal{K}(Y, Z)$ and $\mathcal{K}(Z, W)$, respectively,

$$\begin{aligned} (KG)F &= K(GF), \\ (KG)f &= K(Gf), \quad (Kg)F = K(gF), \quad (kG)F = k(GF), \\ (KG)\alpha &= K(G\alpha), \quad (K\beta)F = K(\beta F), \quad (\gamma G)F = \gamma(GF), \\ (Kg)_f &= K(g_f), \quad (kG)_f = k_{Gf}, \quad (k_g)F = k_{(gF)}. \end{aligned}$$

(G6) For every X , the 2-functors

$$1_X(-): \mathcal{K}(X, Y) \rightarrow \mathcal{K}(X, Y), \quad (-)1_X: \mathcal{K}(X, Y) \rightarrow \mathcal{K}(X, Y)$$

defined by composition with $1_X: X \rightarrow X$, are identities.

Remark 1.2. Asserting that composition in a Gray-category \mathcal{K} is a cubical functor means that the properties in (i)-(v) below hold, for every $F, F', F'': X \rightarrow Y$, $G, G', G'': Y \rightarrow Z$ and

$$F \begin{array}{c} \xrightarrow{f} \\ \Downarrow \phi \\ \xrightarrow{f'} \end{array} F' \xrightarrow{f''} F'', \quad G \begin{array}{c} \xrightarrow{g} \\ \Downarrow \psi \\ \xrightarrow{g'} \end{array} G' \xrightarrow{g''} G''.$$

(i) Composition with 1-cells on either side,

$$(-)F: \mathcal{K}(Y, Z) \rightarrow \mathcal{K}(X, Z) \quad G(-): \mathcal{K}(X, Y) \rightarrow \mathcal{K}(X, Z),$$

is a strict 2-functor.

(ii) Composition with 2-cells,

$$(-)f: (-)F \rightarrow (-)F', \quad g(-): G(-) \rightarrow G'(-),$$

is a pseudo-natural transformation.

(iii) Composition with 3-cells,

$$(-)\varphi: (-)f \rightarrow (-)f', \quad \psi(-): g(-) \rightarrow g'(-),$$

is a modification,

(iv) The following coherence equations hold:

$$(1.1) \quad \begin{array}{ccc} & \begin{array}{ccc} & \xrightarrow{Gf} & \\ & \Downarrow G\varphi & \\ & \xrightarrow{Gf'} & \end{array} & \\ \begin{array}{ccc} g'F & \xleftarrow{\gamma_F} & gF \\ & \searrow g_f & \\ & & gF' \end{array} & \xrightarrow{gF'} & \begin{array}{ccc} & \xrightarrow{gF'} & \\ & \Downarrow G'\varphi & \\ & \xrightarrow{G'f'} & \end{array} \\ G'F & \xrightarrow{G'f'} & G'F' \end{array} = \begin{array}{ccc} GF & \xrightarrow{G'f} & GF' \\ & \searrow g'_f & \swarrow g_{F'} \\ & & \begin{array}{ccc} & \xrightarrow{Gf} & \\ & \Downarrow G'\varphi & \\ & \xrightarrow{G'f'} & \end{array} \\ g'F & \xrightarrow{gF'} & gF' \end{array}$$

$$(1.2) \quad \begin{array}{ccc} GF & \xrightarrow{Gf} & GF' \\ \downarrow gF & \Downarrow g_f & \downarrow gF' \\ G'F & \xrightarrow{G'f} & G'F' \\ \downarrow g''F & \Downarrow g''_f & \downarrow g''F' \\ G''F' & \xrightarrow{G''f} & G''F' \end{array} = \begin{array}{ccc} GF & \xrightarrow{Gf} & GF' \\ \downarrow (g'g)F & \Downarrow (g''g)_f & \downarrow (g''g)F' \\ G''F' & \xrightarrow{G''f} & G''F' \end{array}$$

$$(1.3) \quad \begin{array}{ccccc} GF & \xrightarrow{Gf} & GF' & \xrightarrow{Gf''} & GF'' \\ \downarrow gF & \Downarrow g_f & \downarrow gF' & \Downarrow g_{f''} & \downarrow gF'' \\ G'F & \xrightarrow{G'f} & G'F' & \xrightarrow{G'f''} & G'F'' \end{array} = \begin{array}{ccc} GF & \xrightarrow{G(f''f)} & GF'' \\ \downarrow gF & \Downarrow g_{(f''f)} & \downarrow gF'' \\ G'F & \xrightarrow{G'(f''f)} & G'F'' \end{array}$$

(v) The interchange map g_f is the identity 3-cell when either f or g is the identity.

Remark 1.3. When working with a Gray-category, we sometimes write $G \circ F$ instead of GF for cubical composition of 1-cells. For 2-cells, we write $g' \cdot g$ (or $g'g$) for the vertical composition and $g \circ f$ for cubical composition. For 3-cells, we write $\beta \circ \alpha$ for cubical composition, $\alpha' * \alpha$ for vertical composition in $\mathcal{K}(X, Y)$ and $\bar{\alpha} \cdot \alpha$ for horizontal composition in $\mathcal{K}(X, Y)[F', F'']$.

Remark 1.4. We write **Gray** for the category of 2-categories and 2-functors. For 2-categories X and Y , let $[X, Y]$ be the 2-category of 2-functors from X to Y , pseudonatural transformations, and modifications [18]. This definition equips the category **Gray** with the structure of a closed category [8]. The closed structure of **Gray** is part of symmetric monoidal structure, whose the tensor product is known as the *Gray tensor product* [14, Section 4.8]. We will write $X \otimes Y$ for the Gray tensor product of 2-categories X

and Y . A Gray-category can then be defined equivalently as a **Gray**-enriched category [14, Section 5.1]. Since **Gray** is a monoidal closed category, it is enriched over itself. Therefore, it can be viewed as a Gray-category, as we will do from now on. More explicitly, **Gray** is the Gray-category having 2-categories as 0-cells, 2-functors as 1-cells, pseudonatural transformations as 2-cells, and modifications as 3-cells.

The notions of a *Gray-functor* and of a *Gray-natural transformation* are instances of the general notions of enriched functor and enriched natural transformation [17, Section 1.2]. We will use the terminology of *Gray-modification* and *Gray-perturbation* to denote the strict counterparts of the corresponding tricategorical notions [14, Section 3.3].

When working with the Yoneda embedding for Gray-categories, which is just an instance of the Yoneda embedding for enriched categories [17, Section 2.4], we often identify an object $X \in \mathcal{K}$ with the representable Gray-functor $\mathcal{K}(-, X): \mathcal{K}^{op} \rightarrow \mathbf{Gray}$ associated to it. Analogous conventions will be used also for the n -cells of \mathcal{K} , where $n = 1, 2, 3$. For further information on Gray-categories and tricategories, we invite the reader to refer to [13–15, 20].

Pseudomonads and their pseudoalgebras. Let \mathcal{K} be a Gray-category, to be considered fixed for the rest of this section. We recall the definition of a pseudomonad.

Definition 1.5. Let $X \in \mathcal{K}$. A *pseudomonad* on X in \mathcal{K} consists of:

- a 1-cell $S : X \rightarrow X$ in \mathcal{K} ;
- two 2-cells $m : S^2 \rightarrow S$ and $s : 1_X \rightarrow S$ in \mathcal{K} ;
- three invertible 3-cells in \mathcal{K} of the following form:

$$\begin{array}{ccc}
 S^3 & \xrightarrow{Sm} & S^2 \\
 mS \downarrow & & \Downarrow \mu \\
 S^2 & \xrightarrow{m} & S
 \end{array}
 \qquad
 \begin{array}{ccccc}
 S & \xrightarrow{Ss} & S^2 & \xleftarrow{sS} & S \\
 & \searrow \lambda & \downarrow m & \swarrow \rho & \\
 & & S & & \\
 & \nearrow 1_s & & \nwarrow 1_s &
 \end{array}$$

satisfying the coherence axioms in (1.4) and (1.5) below:

$$(1.4) \quad
 \begin{array}{ccccc}
 S^4 & \xrightarrow{S^2m} & S^3 & & \\
 mS^2 \downarrow & \searrow SmS & \Downarrow S\mu & \searrow Sm & \\
 S^3 & \xrightarrow{Sm} & S^2 & & \\
 mS \downarrow & \searrow mS & \Downarrow \mu & \searrow m & \\
 S^2 & \xrightarrow{m} & S & &
 \end{array}
 =
 \begin{array}{ccccc}
 S^4 & \xrightarrow{S^2m} & S^3 & & \\
 mS^2 \downarrow & \searrow m_m & \downarrow mS & \searrow Sm & \\
 S^3 & \xrightarrow{Sm} & S^2 & & \\
 mS \downarrow & \searrow \mu & \downarrow m & \searrow m & \\
 S^2 & \xrightarrow{m} & S & &
 \end{array}$$

$$(1.5) \quad \begin{array}{ccc} S^2 & \xrightarrow{1_{S^2}} & S^2 \\ \downarrow SsS & \searrow & \downarrow S\rho \\ S^3 & \xrightarrow{Sm} & S^2 \\ \downarrow \lambda S & \downarrow Sm & \downarrow \mu \\ S^2 & \xrightarrow{m} & S \end{array} \quad = \quad S^2 \xrightarrow{m} S$$

For brevity, we will refer to an object $X \in \mathcal{K}$ and a pseudomonad $(S, m, s, \mu, \lambda, \rho)$ on X simply as a pseudomonad in \mathcal{K} and write simply (X, S) to denote it.

Note that the notion of a pseudomonad is self-dual, in the sense that a pseudomonad in \mathcal{K} is the same thing as a pseudomonad in \mathcal{K}^{op} , where \mathcal{K}^{op} is the Gray-category obtained from \mathcal{K} by reversing the direction of the 1-cells, but not that of the 2-cells and 3-cells. As in the formal theory of monads, this is important to obtain results by duality.

Let (X, S) be a pseudomonad in \mathcal{K} . For $I \in \mathcal{K}$, there is a 2-category $\text{Ps-}S\text{-Alg}(I)$ of I -indexed pseudo- S -algebras, pseudoalgebra morphisms, and pseudoalgebra 2-cells, whose definitions we recall below. An I -indexed pseudoalgebra for S consists of a 1-cell $A: I \rightarrow X$, called the underlying 1-cell of the pseudoalgebra, a 2-cell $a: SA \rightarrow A$, called the *structure map* of the pseudoalgebra, and invertible 3-cells

$$\begin{array}{ccc} S^2 A & \xrightarrow{Sa} & SA \\ m_A \downarrow & \Downarrow \bar{a} & \downarrow a \\ SA & \xrightarrow{a} & A \end{array}, \quad \begin{array}{ccc} A & \xrightarrow{s_A} & SA \\ \searrow 1_A & \xrightarrow{\bar{a}} & \downarrow a \\ & & A \end{array},$$

called the *associativity* and *unit* of the pseudoalgebra, satisfying the coherence axioms (1.6) and (1.7) stated below.

$$(1.6) \quad \begin{array}{ccc} S^3 A & \xrightarrow{S^2 a} & S^2 A \\ \downarrow m_{SA} & \searrow Sm_A & \searrow S\bar{a} \\ S^2 A & \xrightarrow{Sa} & SA \\ \downarrow m_A & \Downarrow \alpha_A & \downarrow \bar{a} \\ SA & \xrightarrow{a} & A \end{array} = \begin{array}{ccc} S^3 A & \xrightarrow{S^2 a} & S^2 A \\ \downarrow m_{SA} & \downarrow m_a & \downarrow m_A \\ S^2 A & \xrightarrow{Sa} & SA \\ \downarrow m_A & \Downarrow \bar{a} & \downarrow a \\ SA & \xrightarrow{a} & A \end{array}.$$

$$(1.7) \quad \begin{array}{ccc} SA & \xrightarrow{1_{SA}} & SA \\ \downarrow s_{SA} & \searrow \tilde{a} & \downarrow a \\ S^2A & \xrightarrow{Sa} & SA \\ \downarrow \lambda_A & \downarrow m_A & \downarrow \tilde{a} \\ SA & \xrightarrow{a} & A \end{array} \quad = \quad SA \xrightarrow{a} A.$$

As usual, we refer to a pseudoalgebra by the name of its underlying 1-cell, leaving the rest of its data implicit. Similar conventions will be implicitly assumed for other kinds of structures.

Proposition 1.6 (Marmolejo). *Let (X, S) be a pseudomonad in \mathcal{K} , $I \in \mathcal{K}$ and A an I -indexed pseudoalgebra for S . Then, the coherence condition*

$$(1.8) \quad \begin{array}{ccc} SA & \xrightarrow{a} & A \\ \downarrow s_{SA} & \searrow s_a & \downarrow s_A \\ S^2A & \xrightarrow{Sa} & SA \\ \downarrow m_A & \searrow \tilde{a} & \downarrow a \\ SA & \xrightarrow{a} & A \end{array} \quad \overset{1_{SA}}{=} \quad \begin{array}{ccc} SA & \xrightarrow{1_{SA}} & SA \\ \downarrow s_{SA} & \searrow \rho_A & \downarrow m_A \\ S^2A & \xrightarrow{m_A} & SA \\ \downarrow m_A & \searrow a & \downarrow a \\ SA & \xrightarrow{a} & A \end{array}$$

is derivable.

Proof. See [23, Lemma 9.1]. □

Given pseudoalgebras A and B , a *pseudoalgebra morphism* $f : A \rightarrow B$ consists of a 2-cell $f : A \rightarrow B$ and an invertible 3-cell

$$\begin{array}{ccc} SA & \xrightarrow{Sf} & SB \\ \downarrow a & \Downarrow \bar{f} & \downarrow b \\ A & \xrightarrow{f} & B \end{array}$$

satisfying the coherence conditions (1.9) and (1.10) stated below.

$$(1.9) \quad \begin{array}{ccc} S^2A & \xrightarrow{S^2f} & S^2B \\ m_A \downarrow & \searrow^{S_a} & \searrow^{S\bar{f}} \\ SA & \xrightarrow{Sf} & SB \\ \bar{a} \Downarrow & \searrow^a & \searrow^b \\ A & \xrightarrow{f} & B \end{array} = \begin{array}{ccc} S^2A & \xrightarrow{S^2f} & SB \\ m_A \downarrow & \searrow^{m_f} & \searrow^{m_B} \\ SA & \xrightarrow{Sf} & SB \\ a \Downarrow & \searrow^a & \searrow^b \\ A & \xrightarrow{f} & B \end{array}$$

$$(1.10) \quad \begin{array}{ccc} A & & \\ s_A \downarrow & \searrow^{1_A} & \\ SA & \xrightarrow{\bar{a}} & A \\ a \downarrow & & \\ A & \xrightarrow{f} & B \end{array} = \begin{array}{ccc} A & \xrightarrow{f} & B \\ s_A \downarrow & \searrow^{s_f} & \searrow^{s_B} \\ SA & \xrightarrow{Sf} & SB \\ a \downarrow & \searrow^a & \searrow^b \\ A & \xrightarrow{f} & B \end{array}$$

Given pseudoalgebra morphisms $f : A \rightarrow B$ and $g : A \rightarrow B$, a *pseudoalgebra 2-cell* consists of a 3-cell $\alpha : f \rightarrow g$ satisfying the coherence condition (1.11).

$$(1.11) \quad \begin{array}{ccc} SA & \xrightarrow{Sf} & SB \\ \bar{a} \Downarrow & \searrow^{S\alpha} & \searrow^{Sg} \\ A & \xrightarrow{f} & B \\ \bar{g} \Downarrow & & \end{array} = \begin{array}{ccc} SA & \xrightarrow{Sf} & SB \\ a \downarrow & \searrow^{\bar{f}} & \searrow^b \\ A & \xrightarrow{f} & B \\ \alpha \Downarrow & & \end{array}$$

There is a forgetful 2-functor $U_I : \text{Ps-}S\text{-Alg}(I) \rightarrow \mathcal{K}(I, X)$, defined by mapping a pseudo- S -algebra to its underlying 1-cell, which has a left pseudoadjoint, defined by mapping a 1-cell $A : I \rightarrow X$ to the free pseudoalgebra on it, given by the composite 1-cell $SA : I \rightarrow X$. Attentive readers will have observed that the directions of the structural 3-cells μ and λ for a pseudomonad as in Definition 1.5 match those of the 3-cells necessary to make SA into a pseudoalgebra.

The function mapping an object $I \in \mathcal{K}$ to the 2-category $\text{Ps-}S\text{-Alg}(I)$ extends to a Gray-functor $\text{Ps-}S\text{-Alg} : \mathcal{K}^{op} \rightarrow \mathbf{Gray}$. We also have a Gray-transformation

$$(1.12) \quad U : \text{Ps-}S\text{-Alg} \rightarrow X,$$

with components given by the forgetful 2-functors $U_I : \text{Ps-}S\text{-Alg}(I) \rightarrow \mathcal{K}(I, X)$, for $I \in \mathcal{K}$. Note the use of our convention on the Yoneda lemma in (1.12). Note that the structure

of pseudo- S -algebra on a 1-cell $A: I \rightarrow X$ can be viewed as a left S -action on A , associative and unital up to coherent isomorphism. For this reason, we sometimes refer to pseudoalgebras as *left pseudomodules*. This terminology is convenient when we discuss dualities in Section 4.

2. THE GRAY-CATEGORY OF PSEUDOMONADS

The aim of this section is to introduce the 3-dimensional category $\mathbf{Psm}(\mathcal{K})$ of pseudomonads in a Gray-category \mathcal{K} and prove that it is a Gray-category. In order to do so, we review the notion of a pseudomonad morphism from [26] and introduce the notions of a pseudomonad transformation and modification. Again, we fix a Gray-category \mathcal{K} . When working with two pseudomonads (X, S) and (Y, T) , we use m and s for the multiplication and unit of S , n and t for the multiplication and unit of T , but we use the same letters μ, λ, ρ for the structural 3-cells of both monads to simplify notation, as the context makes it always clear to which we are referring.

Definition 2.1. Let (X, S) and (Y, T) be pseudomonads in \mathcal{K} . A *pseudomonad morphism* $(F, \phi): (X, S) \rightarrow (Y, T)$ consists of a 1-cell $F: X \rightarrow Y$, a 2-cell $\phi: TF \rightarrow FS$ and two invertible 3-cells

$$\begin{array}{ccc}
 T^2F & \xrightarrow{T\phi} & TFS \\
 \downarrow nF & & \downarrow \phi S \\
 & \Downarrow \bar{\phi} & FS^2 \\
 & & \downarrow Fm \\
 TF & \xrightarrow{\phi} & FS
 \end{array}
 \qquad
 \begin{array}{ccc}
 F & \xrightarrow{tF} & TF \\
 \downarrow Fs & \Downarrow \tilde{\phi} & \downarrow \phi \\
 & & FS
 \end{array}$$

These data are required to satisfy the coherence axioms in (2.1) and (2.2).

$$(2.1) \quad \begin{array}{ccccc} T^3F & \xrightarrow{T^2\phi} & T^2FS & \xrightarrow{T\phi S} & \\ \downarrow nTF & \searrow TnF & \downarrow T\bar{\phi} & TFS^2 & \xrightarrow{TFm} \\ T^2F & \xrightarrow{\downarrow \mu F} & T^2F & \xrightarrow{T\phi} & TFS = \\ & \searrow nF & \downarrow nF & \downarrow \bar{\phi} & FS^2 \\ & & TF & \xrightarrow{h} & FS \end{array}$$

$$\begin{array}{ccccccc} T^3F & \xrightarrow{T^2\phi} & T^2FS & & & & \\ \downarrow nTF & & \downarrow n\phi & & & & \\ T^2F & \xrightarrow{T\phi} & TFS & & & & \\ & \searrow nF & & & & & \\ & & TF & \xrightarrow{\phi} & FS & & \\ & & & & \downarrow \phi S & & \\ & & & & FS^2 & & \\ & & & & \downarrow Fm & & \\ & & & & FS & & \\ & & & & \downarrow \phi S & & \\ & & & & FS^2 & & \\ & & & & \downarrow Fm & & \\ & & & & FS & & \\ & & & & \downarrow \phi S & & \\ & & & & FS^2 & & \\ & & & & \downarrow Fm & & \\ & & & & FS & & \\ & & & & \downarrow \phi S & & \\ & & & & FS^2 & & \\ & & & & \downarrow Fm & & \\ & & & & FS & & \end{array}$$

$$(2.2) \quad \begin{array}{ccc} TF & \xrightarrow{TFs} & TFS \\ \downarrow 1_{TF} & \searrow TtF & \downarrow T\bar{\phi} \\ TF & \xrightarrow{\downarrow \lambda H} & T^2F & \xrightarrow{T\phi} & TFS \\ & \searrow nF & \downarrow nF & \downarrow \bar{\phi} & FS^2 \\ & & TF & \xrightarrow{\phi} & FS \end{array} = \begin{array}{ccc} TF & \xrightarrow{TFs} & TFS \\ \downarrow \phi & & \downarrow \phi S \\ FS & \xrightarrow{FSs} & TFS \\ & \searrow 1_{FS} & \downarrow \phi S \\ & & FS^2 \\ & & \downarrow Fm \\ & & FS \end{array}$$

Proposition 2.2 (Marmolejo and Wood). *Let $(F, \phi) : (X, S) \rightarrow (Y, T)$ be a pseudomonad morphism. The coherence condition*

$$\begin{array}{ccc}
 TF & \xrightarrow{\phi} & FS \\
 \downarrow tTF & \Downarrow t\phi & \downarrow tFS \\
 T^2F & \xrightarrow{T\phi} & TFS \\
 \downarrow nF & \Downarrow \bar{\phi} & \downarrow \phi S \\
 TF & \xrightarrow{\phi} & FS
 \end{array}
 \quad
 \begin{array}{ccc}
 TF & \xrightarrow{\phi} & FS \\
 \downarrow tTF & \Downarrow \rho_{TF} & \downarrow \rho_{TF} \\
 T^2F & \xrightarrow{\phi} & TFS \\
 \downarrow nF & \Downarrow \bar{\phi} & \downarrow \phi S \\
 TF & \xrightarrow{\phi} & FS
 \end{array}$$

is derivable.

Proof. See [26, Theorem 2.3]. □

Definition 2.3. Let $(F, \phi), (F', \phi') : (X, S) \rightarrow (Y, T)$ be pseudomonad morphisms. A pseudomonad transformation $(p, \bar{p}) : (F, \phi) \rightarrow (F', \phi')$ consists of a 2-cell $p : F \rightarrow F'$ and an invertible 3-cell

$$\begin{array}{ccc}
 TF & \xrightarrow{Tp} & TF' \\
 \downarrow \phi & \Downarrow \bar{p} & \downarrow \phi' \\
 FS & \xrightarrow{pS} & F'S
 \end{array}$$

satisfying the coherence conditions in (2.3) and (2.4) below:

$$(2.3) \quad
 \begin{array}{ccc}
 T^2F & \xrightarrow{T^2p} & T^2F' \\
 \downarrow nF & \searrow T\phi & \downarrow T\bar{p} \\
 TF & \xrightarrow{\phi} & FS \\
 \downarrow \phi & \Downarrow \bar{\phi} & \downarrow \phi S \\
 FS & \xrightarrow{pS} & F'S
 \end{array}
 \quad
 \begin{array}{ccc}
 T^2F & \xrightarrow{T^2p} & T^2F' \\
 \downarrow nF & \searrow T\phi & \downarrow n_p \\
 TF & \xrightarrow{Tp} & TF' \\
 \downarrow \phi & \Downarrow \bar{p} & \downarrow \phi' \\
 FS & \xrightarrow{pS} & F'S
 \end{array}$$

$$(2.4) \quad \begin{array}{ccc} F & \xrightarrow{p} & F' \\ tF \downarrow & \searrow^{Fs} & \downarrow^{F's} \\ TF & & TF' \\ \phi \searrow & \Downarrow^{\tilde{\phi}} & \Downarrow^{p_s^{-1}} \\ FS & \xrightarrow{pS} & F'S \end{array} = \begin{array}{ccc} F & \xrightarrow{p} & F' \\ tF \downarrow & \Downarrow^{t_p} & \downarrow^{tF'} \\ TF & \xrightarrow{Tp} & TF' \\ \phi \searrow & \Downarrow^{\tilde{p}} & \Downarrow^{\tilde{\phi}'} \\ FS & \xrightarrow{pS} & F'S \end{array}$$

Definition 2.4. Let $(p, \tilde{p}), (p', \tilde{p}') : (F, \phi) \rightarrow (F', \phi')$ be pseudomonad transformations. A *pseudomonad modification* $\alpha : (p, \tilde{p}) \rightarrow (p', \tilde{p}')$ is a 3-cell $\alpha : p \rightarrow p'$ satisfying the coherence condition

$$(2.5) \quad \begin{array}{ccc} TF & \xrightarrow{Tp} & TF' \\ \phi \downarrow & \Downarrow^{T\alpha} & \downarrow^{\phi'} \\ FS & \xrightarrow{pS} & F'S \end{array} = \begin{array}{ccc} TF & \xrightarrow{Tp} & TF' \\ \phi \downarrow & \Downarrow^{\tilde{p}} & \downarrow^{\phi'} \\ FS & \xrightarrow{pS} & F'S \end{array}$$

The following is our first main result, which solves the problem raised in [19, Section 6].

Theorem 2.5. *Let \mathcal{K} be a Gray-category. Then there is a Gray-category $\mathbf{Psm}(\mathcal{K})$, called the Gray-category of pseudomonads in \mathcal{K} , having pseudomonads in \mathcal{K} as 0-cells, pseudomonad morphisms as 1-cells, pseudomonad transformations as 2-cells, and pseudomonad modifications as 3-cells.*

The rest of this section is devoted to the proof of Theorem 2.5, which will be obtained by combining Lemmas 2.6, 2.8, 2.9 and 2.10 below. We begin by giving the definition of the hom-2-categories of $\mathbf{Psm}(\mathcal{K})$.

Lemma 2.6. *Let (X, S) and (Y, T) be two pseudomonads in \mathcal{K} . Then there is a 2-category $\mathbf{Psm}(\mathcal{K})((X, S), (Y, T))$ having pseudomonad morphisms from (X, S) to (Y, T) as 0-cells, pseudomonad transformations as 1-cells and pseudomonad modifications as 2-cells.*

Proof. First of all, for any pair of composable 1-cells $(p_0, \tilde{p}_0) : (F_0, \phi_0) \rightarrow (F_1, \phi_1)$ and $(p_1, \tilde{p}_1) : (F_1, \phi_1) \rightarrow (F_2, \phi_2)$ we define their composition as $(p_1 p_0, \tilde{p}_1 \tilde{p}_0)$ where $\tilde{p}_1 \tilde{p}_0$ is defined as the pasting of

$$\begin{array}{ccccc}
TF_0 & \xrightarrow{Tp_0} & TF_1 & \xrightarrow{Tp_1} & TF_2 \\
\downarrow \phi_0 & & \Downarrow \tilde{p}_0 & \downarrow \phi_1 & \Downarrow \tilde{p}_1 & \downarrow \phi_2 \\
F_0S & \xrightarrow{p_0S} & F_1S & \xrightarrow{p_1S} & F_2S.
\end{array}$$

We want to show that composition is strictly associative. So let us consider three composable 1-cells

$$(F_0, \phi_0) \xrightarrow{(p_0, \tilde{p}_0)} (F_1, \phi_1) \xrightarrow{(p_1, \tilde{p}_1)} (F_2, \phi_2) \xrightarrow{(p_2, \tilde{p}_2)} (F_3, \phi_3)$$

By definition, the two possible composites are

$$\begin{aligned}
(p_2, \tilde{p}_2) \cdot ((p_1, \tilde{p}_1) \cdot (p_0, \tilde{p}_0)) &= (p_2(p_1p_0), \widetilde{p_2(p_1p_0)}), \\
((p_2, \tilde{p}_2) \cdot (p_1, \tilde{p}_1)) \cdot (p_0, \tilde{p}_0) &= ((p_2p_1)p_0, \widetilde{(p_2p_1)p_0}).
\end{aligned}$$

We want to show that these are equal. Since \mathcal{K} is a Gray-category, $p_2(p_1p_0) = (p_2p_1)p_0$. Moreover, $\widetilde{p_2(p_1p_0)} = \widetilde{(p_2p_1)p_0}$ since they are both the pasting of

$$\begin{array}{ccccccc}
TF_0 & \xrightarrow{Tp_0} & TF_1 & \xrightarrow{Tp_1} & TF_2 & \xrightarrow{Tp_2} & TF_3 \\
\downarrow \phi_0 & & \Downarrow \tilde{p}_0 & \downarrow \phi_1 & \Downarrow \tilde{p}_1 & \downarrow \phi_2 & \Downarrow \tilde{p}_2 & \downarrow \phi_3 \\
F_0S & \xrightarrow{p_0S} & F_1S & \xrightarrow{p_1S} & F_2S & \xrightarrow{p_2S} & F_3S.
\end{array}$$

It remains to define the identity 1-cells of $\mathbf{Psm}(\mathcal{K})((X, S), (Y, T))$. For a pseudomonad morphism $(F, \phi): (X, S) \rightarrow (Y, T)$, we define the identity on it to be

$$(1_F, 1_\phi): (F, \phi) \rightarrow (F, \phi).$$

This is allowed since $T1_F = 1_{TF}$ and $1_FS = 1_{FS}$. These can be shown to be a strict identities, using that \mathcal{K} is a Gray-category and in particular Axiom (G6). \square

We proceed by defining the composition of 1-cells in $\mathbf{Psm}(\mathcal{K})$ and proving that is *strictly* associative, as required to have a Gray-category. Since a pseudomonad morphism is a tuple of the form $(F, \phi, \bar{\phi}, \tilde{\phi})$, where F is a 1-cell, ϕ is a 2-cell while $\bar{\phi}$ and $\tilde{\phi}$ are 3-cells, we will need to check equalities at three levels. The key level of the verification is that of 2-cells. Indeed, strict associativity at the level of 1-cells will follow easily from the strict associativity of composition of 1-cells in \mathcal{K} . The key issue are the equalities at the level of 2-cells, since 2-cells could be isomorphic (by means of an invertible 3-cell), but not equal. Instead, equalities of 3-cells will be quite straightforward. In fact, the required equations for 3-cells either hold strictly or they fail completely, since there are no 4-cells that could make these equations hold only up to isomorphism.

In the following, for a pseudomonad morphism $\underline{F} = (F, \phi, \bar{\phi}, \tilde{\phi})$, we define

$$\underline{F}^- := \bar{\phi}, \quad \underline{F}^\sim := \tilde{\phi}.$$

Let $(F, \phi) : (X, S) \rightarrow (Y, T)$ and $(G, \psi) : (Y, T) \rightarrow (Z, Q)$ be two pseudomonad morphisms. We define their composition as

$$(2.6) \quad (G, \psi, \underline{G}^-, \underline{G}^-) \circ (F, \phi, \underline{F}^-, \underline{F}^-) := (GF, G\phi \cdot \psi F, \underline{G} \circ \underline{F}^-, \underline{G} \circ \underline{F}^-)$$

where the invertible 3-cells are defined by the following pasting diagrams:

$$\begin{array}{ccccc} & Q^2GF & \xrightarrow{Q\psi F} & QGTF & \xrightarrow{QG\phi} & QGFS \\ & \downarrow & & \downarrow \psi TF & \downarrow \psi_\phi \Downarrow & \downarrow \psi FS \\ & & & GT^2F & \xrightarrow{GT\phi} & GTFS \\ \underline{G} \circ \underline{F}^- := & \downarrow m_{QGF} & & \downarrow \bar{\psi} F \Downarrow & & \downarrow G\phi S \\ & & & GnF & \downarrow G\bar{\phi} \Downarrow & GFS^2 \\ & & & & & \downarrow GFm \\ & QGF & \xrightarrow{\psi F} & GTF & \xrightarrow{G\phi} & GFS \\ & & & & & \\ & GF & \xrightarrow{GFs} & & & GFS \\ & \downarrow qGF & \searrow GtF & \downarrow G\tilde{\phi} \Downarrow & \nearrow G\phi & \\ & & & GTF & & \\ & & & \downarrow \tilde{\psi} F \Downarrow & & \\ & & & QGF & \nearrow \psi F & \end{array}$$

The proof that this definition gives a pseudomonad morphism is in Appendix B.

Remark 2.7. We did not consider any parenthesis in the diagrams above thanks to axiom (G5) for a Gray-category. Moreover since $Q(-)$ is a strict 2-functor we have $Q(G\phi \cdot \psi F) = QG\phi \cdot Q\psi F$ (and similarly for other compositions in the diagrams).

Lemma 2.8. *The composition of pseudomonad morphisms defined in (2.6) is strictly associative.*

Proof. From now on, let us consider three pseudomonad morphisms in \mathcal{K} :

$$(X, S) \xrightarrow{(F, \phi)} (Y, T) \xrightarrow{(G, \psi)} (Z, Q) \xrightarrow{(H, \xi)} (V, R)$$

In order to prove this statement we have to prove that the equation for associativity holds for the respective 1-, 2- and 3-cell components. For 1-cells, since \mathcal{K} is a Gray-category, then $H(GF) = (HG)F$.

For 2-cells, the idea is to reduce both composites to $HG\phi \cdot H\psi F \cdot \xi GF$. On the one hand,

$$\begin{aligned} H(G\phi \cdot \psi F) \cdot \xi GF &= [H(G\phi) \cdot H(\psi F)] \cdot \xi GF && \text{(because } H(-) \text{ is strict)} \\ &= [HG\phi \cdot H\psi F] \cdot \xi GF && \text{(by (G5))} \\ &= HG\phi \cdot H\psi F \cdot \xi GF && \text{(since } \mathcal{K}(X, V) \text{ is a 2-category).} \end{aligned}$$

On the other hand,

$$\begin{aligned} HG\phi \cdot (H\psi \cdot \xi G)F &= HG\phi \cdot [(H\psi)F \cdot (\xi G)F] && \text{(because } (-)F \text{ is strict)} \\ &= HG\phi \cdot [H\psi F \cdot \xi GF] && \text{(by (G5))} \\ &= HG\phi \cdot H\psi F \cdot \xi GF && \text{(since } \mathcal{K}(X, V) \text{ is a 2-category).} \end{aligned}$$

For 3-cells, to prove that $(\underline{H}(\underline{GF}))^\sim = ((\underline{HG})\underline{F})^\sim$ we just need to notice that, using the fact that $H(-)$ and $(-)^F$ are strict 2-functors, both of them are pasting of:

$$\begin{array}{ccc} HGF & \xrightarrow{HG\phi s} & HGFS \\ & \searrow^{HGtF} & \downarrow^{HG\phi} \\ & & HGTf \\ & \searrow^{HqGF} & \downarrow^{H\psi F} \\ & & HQGF \\ & \searrow^{rHGF} & \downarrow^{\xi GF} \\ & & RHGF \end{array}$$

We get the required equality by the pasting theorem for 2-categories [27]. Finally, let us prove the equality on the other 3-cell component. By definition,

$$\begin{array}{ccccc} R^2 HGF & \longrightarrow & RHQGF & \longrightarrow & RHGFS \\ \downarrow & & \downarrow & \searrow^{\xi(G\phi \cdot \psi F)} & \downarrow \\ \underline{H}(\underline{GF})^- = & & \downarrow^{\xi GF} & HQ^2 GF & \longrightarrow & HQGFS \\ & & & \downarrow^{H(\underline{GF}^-)} & & \downarrow \\ RHGF & \longrightarrow & HQGF & \longrightarrow & HGFS. \end{array}$$

Using the definition of \underline{GF}^- and (1.3), the right-hand side pasting becomes

$$\begin{array}{ccccccc}
R^2HGF & \longrightarrow & RHQGF & \longrightarrow & RHGTF & \longrightarrow & RHGFS \\
\downarrow & & \downarrow & & \Downarrow \xi_{\psi F} & & \Downarrow \xi_{G\phi} \\
& & HQ^2GF & \longrightarrow & HGTFS & \longrightarrow & HQGFS \\
& & \Downarrow \bar{\xi}_{GF} & & \downarrow & & \Downarrow H(\psi_\phi) \\
& & & & H\bar{\psi}F \Downarrow & & HGT^2F \longrightarrow HGTFS \\
& & & & \downarrow & & \Downarrow HG\bar{\phi} \\
RHGF & \longrightarrow & HQGF & \longrightarrow & HGTF & \longrightarrow & HGFS.
\end{array}$$

Let us notice that, by (G5), $H(\psi_\phi) = H\psi_\phi$, $\xi_{\psi F} = (\xi_\psi)F$ and $\xi_{G\phi} = \xi G_\phi$. Moreover, using the definition of \underline{HG}^- and (1.2), the diagram above is equal to

$$\begin{array}{ccccc}
R^2HGF & \longrightarrow & RHGTF & \longrightarrow & RHGFS \\
\downarrow & & \downarrow & & \Downarrow (H\psi \cdot \xi G)_\phi \\
& & HGT^2FS & \longrightarrow & HGTFS \\
& & \Downarrow (HG^-)F & & \Downarrow HG\bar{\phi} \\
RHGF & \longrightarrow & HGTF & \longrightarrow & HGFS,
\end{array}$$

which is exactly the definition of $\underline{H}(\underline{GF})^-$. \square

For brevity, we sometimes write $P_{\mathcal{K}}$ instead of $\mathbf{Psm}(\mathcal{K})$, so for any pair of pseudomonads (X, S) and (Y, T) the 2-category of pseudomonads morphisms from (X, S) to (Y, T) can be written as $P_{\mathcal{K}}((X, S), (Y, T))$.

Lemma 2.9. *The definition of composition of pseudomonad morphisms extends to a cubical functor*

$$- \circ - : P_{\mathcal{K}}((Y, T), (Z, Q)) \times P_{\mathcal{K}}((X, S), (Y, T)) \longrightarrow P_{\mathcal{K}}((X, S), (Z, Q))$$

for $(X, S), (Y, T), (Z, Q) \in \mathbf{Psm}(\mathcal{K})$.

Proof. This is a just a long verification, but we spell it out in some detail. By the definition of a cubical functor, for $(F, \phi) : (X, S) \rightarrow (Y, T)$ and $(G, \psi) : (Y, T) \rightarrow (Z, Q)$ in $\mathbf{Psm}(\mathcal{K})$, we need to define strict 2-functors

$$(2.7) \quad F_\phi : P_{\mathcal{K}}((Y, T), (Z, Q)) \rightarrow P_{\mathcal{K}}((X, S), (Z, Q)),$$

$$(2.8) \quad G_\psi : P_{\mathcal{K}}((X, S), (Y, T)) \rightarrow P_{\mathcal{K}}((X, S), (Z, Q))$$

such that

$$(2.9) \quad F_\phi((G, \psi)) = G_\psi((F, \phi)) = (G, \psi) \circ (F, \phi),$$

plus, for 2-cells $(p, \tilde{p}) : (F, \phi) \rightarrow (F', \phi')$ and $(q, \tilde{q}) : (G, \psi) \rightarrow (G', \psi')$, an invertible 3-cell in $\mathbf{Psm}(\mathcal{K})$

$$(2.10) \quad \begin{array}{ccc} (G, \psi) \circ (F, \phi) & \xrightarrow{(G, \psi) \circ (p, \tilde{p})} & (G, \psi) \circ (F', \phi') \\ \downarrow (q, \tilde{q}) \circ (F, \phi) & \Downarrow \Sigma_{(p, \tilde{p}), (q, \tilde{q})} & \downarrow (q, \tilde{q}) \circ (F', \phi') \\ (G', \psi') \circ (F, \phi) & \xrightarrow{(G', \psi') \circ (p, \tilde{p})} & (G', \psi') \circ (F', \phi') \end{array}$$

satisfying axioms (1.1), (1.2) and (1.3).

We begin by defining F_ϕ in (2.7). Its action on objects is determined by (2.9). For its action on 1-cells, we send $(q, \tilde{q}) : (G, \psi) \rightarrow (G', \psi')$ to the pseudomonad modification $(qF, \tilde{q}F) : (GF, G\phi \cdot \psi F) \rightarrow (G'F, G'\phi \cdot \psi'F)$, where $\tilde{q}F$ is defined as the following pasting:

$$\begin{array}{ccc} QGF & \xrightarrow{QqF} & QG'F \\ \psi F \downarrow & \Downarrow \tilde{q}F & \downarrow \psi'F \\ GTF & \xrightarrow{qTF} & G'TF \\ G\phi \downarrow & \Downarrow q_\phi^{-1} & \downarrow G'\phi \\ GFS & \xrightarrow{qFS} & G'FS. \end{array}$$

The action of F_ϕ on 3-cells $\beta : (q, \tilde{q}) \rightarrow (q', \tilde{q}')$ is defined by letting $\beta \circ (F, \phi) := \beta F$ in \mathcal{K} . The proof that this is a pseudomonad modification, and therefore a 3-cells in $\mathbf{Psm}(\mathcal{K})$, is in Appendix B.

We now show that F_ϕ is a 2-functor. For this, we use extensively the notation of Remark 1.3 to avoid writing some diagrams. To prove that composition is preserved strictly, we show that

$$(2.11) \quad F_\phi((q', \tilde{q}') \cdot (q, \tilde{q})) = F_\phi(q', \tilde{q}') \cdot F_\phi(q, \tilde{q})$$

for any

$$(G, \psi) \xrightarrow{(q, \tilde{q})} (G', \psi') \xrightarrow{(q', \tilde{q}')} (G'', \psi'')$$

in $P_{\mathcal{K}}((Y, T), (Z, Q))$. The composition $(q', \tilde{q}') \cdot (q, \tilde{q})$ is defined as $(q'q, \tilde{q}'\tilde{q})$ where $\tilde{q}'\tilde{q}$ is defined as the pasting of

$$\begin{array}{ccccc}
QG & \longrightarrow & QG' & \longrightarrow & QG' \\
\downarrow & & \Downarrow \tilde{q} & & \downarrow \\
GT & \longrightarrow & G'T & \longrightarrow & G'T.
\end{array}$$

Using the equation (1.3) we can see that the 3-cells components of $F_\phi((q', \tilde{q}') \cdot (q, \tilde{q}))$ and $F_\phi(q', \tilde{q}') \cdot F_\phi(q, \tilde{q})$ are two pasting of the diagram below

$$\begin{array}{ccccc}
QG & \longrightarrow & QG' & \longrightarrow & QG' \\
\downarrow & & \Downarrow \tilde{q} & & \downarrow \\
GT & \longrightarrow & G'T & \longrightarrow & G'T \\
\downarrow & & \Downarrow q_\phi^{-1} & & \downarrow \\
GF & \longrightarrow & G'F & \longrightarrow & G'F.
\end{array}$$

Moreover, $(q' \cdot q)F = q'F \cdot qF$ since $(-)F$ is a strict 2-functor (since \mathcal{K} is a Gray category). Hence, the required equality in (2.11) holds. Let us also verify that F_ϕ preserves identities strictly. Recall from Lemma 2.6 that $1_{(G, \psi)} := (1_G, 1_\psi)$ in $P_{\mathcal{K}}((Y, T), (Z, Q))$. Therefore,

$$F_\phi(1_G, 1_\psi) = (1_{GF}, \widetilde{1_{GF}})$$

and moreover

$$\begin{aligned}
(1_{GF}, \widetilde{1_{GF}}) &= (1_{GF}, ((1_G)_\phi \cdot 1_{\psi F}) * (1_{G\phi} \cdot 1_{\psi F})) && \text{(by definition of } F_\phi) \\
&= (1_{GF}, (1_{G\phi} \cdot 1_{\psi F}) * (1_{G\phi} \cdot 1_{\psi F})) && \text{(by Remark 1.2)} \\
&= (1_{GF}, (1_{G\phi \cdot \psi F}) * (1_{G\phi \cdot \psi F})) && \text{(since } \cdot \text{ preserves identities)} \\
&= (1_{GF}, 1_{G\phi \cdot \psi F}) && \text{(by (G2))} \\
&= 1_{(GF, G\phi \cdot \psi F)} \\
&= 1_{F_\phi(G, \psi)},
\end{aligned}$$

as required.

We now define the 2-functor G_ϕ of (2.8). Again, its action on objects is determined by (2.9). On morphisms, it sends $(p, \tilde{p}): (F, \phi) \rightarrow (F', \phi')$ to

$$(Gp, \widetilde{Gp}): (GF, G\phi \cdot \psi F) \rightarrow (GF', G\phi' \cdot \psi F'),$$

where \widetilde{Gp} is defined as the following pasting:

$$\begin{array}{ccc}
QGF & \xrightarrow{QGp} & QGF' \\
\downarrow \psi F & \Downarrow \psi_p & \downarrow \psi F' \\
GTF & \xrightarrow{GTp} & GTF' \\
\downarrow G\phi & \Downarrow G\tilde{p} & \downarrow G\phi' \\
GFS & \xrightarrow{GpS} & GF'S.
\end{array}$$

On 3-cells $\alpha : (p, \tilde{p}) \rightarrow (p', \tilde{p}')$ we let $(G, \phi) \circ \alpha := G\alpha$, which is a 3-cell in $\mathbf{Psm}(\mathcal{K})$ by a similar argument to the one used for F_ϕ . The proof that this is a 2-functor is completely analogous to the one for F_ϕ and hence omitted.

To conclude the proof, we need to define the 3-cell $\Sigma_{(p, \tilde{p}), (q, \tilde{q})}$ in (2.10). We take this to be q_p , which is shown to be a pseudomonad modification in Appendix B. The required axioms for $\Sigma_{(p, \tilde{p}), (q, \tilde{q})}$, as in (1.1), (1.2) and (1.3), hold as they are instances of the ones for q_p for \mathcal{K} . \square

Lemma 2.10. *The cubical functor providing composition in $\mathbf{Psm}(\mathcal{K})$ satisfies the coherence conditions of Axiom (G5).*

Proof. The first one is just Lemma 2.8. Since the definitions on 3-cells coincide with the ones in \mathcal{K} , all the equations regarding them hold directly. Therefore, we only need to prove the ones for 2-cells. Let us consider the following diagram in $\mathbf{Psm}(\mathcal{K})$:

$$\begin{array}{ccccc}
& (F, \phi) & & (G, \psi) & & (H, \xi) \\
(X, S) & \xrightarrow{\quad} & (Y, T) & \xrightarrow{\quad} & (Z, Q) & \xrightarrow{\quad} & (V, R) \\
& \Downarrow (p, \tilde{p}) & & \Downarrow (q, \tilde{q}) & & \Downarrow (r, \tilde{r}) \\
& (F', \phi') & & (G', \psi') & & (H', \xi')
\end{array}$$

We need to prove:

- (i) $(H_\xi \circ G_\psi)(p, \tilde{p}) = H_\xi(G_\psi(p, \tilde{p}))$,
- (ii) $(H_\xi(q, \tilde{q}))F_\phi = H_\xi((q, \tilde{q})F_\phi)$,
- (iii) $((r, \tilde{r})G_\psi)F_\phi = (r, \tilde{r})(G_\psi \circ F_\phi)$.

At the 2-cells level we have $(HG)p = H(Gp)$ because \mathcal{K} is a Gray-category. The same happens in (ii) and (iii) so we will just prove that the associated 3-cells are equal in each case. Let us start with part (i). On the one hand, by definition,

$$\widetilde{(HG)p} = \widetilde{HGp},$$

and therefore

$$\begin{array}{ccc}
RHGF & \longrightarrow & RHGF' \\
\downarrow & & \downarrow \\
HQGF & \longrightarrow & HQGF' \\
\downarrow & & \downarrow \\
HGFS & \longrightarrow & HGF'S
\end{array}
=
\begin{array}{ccc}
RHGF & \longrightarrow & RHGF' \\
\downarrow & \Downarrow_{\xi_{G_p}} & \downarrow \\
HQGF & \longrightarrow & HQGF' \\
\downarrow & \Downarrow_{H\psi_p} & \downarrow \\
HGTF & \longrightarrow & HGTF' \\
\downarrow & \Downarrow_{HG\tilde{p}} & \downarrow \\
HGFS & \longrightarrow & HGF'S.
\end{array}$$

On the other hand $H_\xi(G_\psi(p, \tilde{p})) = (HG, H\psi \cdot \xi G)(p, \tilde{p})$ so the associated 3-cell is, using (1.2),

$$\begin{array}{ccc}
RHGF & \longrightarrow & RHGF' \\
\downarrow & & \downarrow \\
HGTF & \longrightarrow & HGTF' \\
\downarrow & & \downarrow \\
HGFS & \longrightarrow & HGF'S
\end{array}
=
\begin{array}{ccc}
RHGF & \longrightarrow & RHGF' \\
\downarrow & \Downarrow_{\xi_{G_p}} & \downarrow \\
HQGF & \longrightarrow & HQGF' \\
\downarrow & \Downarrow_{H\psi_p} & \downarrow \\
HGTF & \longrightarrow & HGTF' \\
\downarrow & \Downarrow_{HG\tilde{p}} & \downarrow \\
HGFS & \longrightarrow & HGF'S.
\end{array}$$

But $\xi_{G_p} = \xi_{G_p}$, since \mathcal{K} is a Gray-category, and so the required equality holds.

For part (ii), by definition, the 3-cell component of $H_\xi(q, \tilde{q})F_\phi$ is:

$$\begin{array}{ccccc}
& & RHGF & \longrightarrow & RHGF' \\
& & \downarrow & & \downarrow \\
RHGF & \longrightarrow & RHGF' & & \\
\downarrow & & \Downarrow_{\widetilde{H}qF} & & \downarrow \\
HGTF & \longrightarrow & HG'TF & = & \\
\downarrow & & \Downarrow_{Hq_\phi^{-1}} & & \downarrow \\
HGFS & \longrightarrow & HG'FS & & \\
& & \downarrow & & \downarrow \\
& & RHGF & \longrightarrow & RHGF' \\
& & \downarrow & & \downarrow \\
& & HQGF & \longrightarrow & HQG'F \\
& & \downarrow & & \downarrow \\
& & \Downarrow_{H\bar{q}F} & & \\
& & HGTF & \longrightarrow & HG'TF \\
& & \downarrow & & \downarrow \\
& & \Downarrow_{Hq_\phi^{-1}} & & \\
& & HGFS & \longrightarrow & HG'FS
\end{array}$$

Finally, part (iii) is completely analogous to the first one using the inverse of 1.3 instead of 1.2. \square

The combination of Lemmas 2.6, 2.8, 2.9 and 2.10 proves Theorem 2.5.

3. LIFTINGS TO PSEUDOALGEBRAS

We now recall from [24, Section 7] and [19, Section 6] the definition of the Gray-category $\mathbf{Lift}(\mathcal{K})$ of pseudomonads in \mathcal{K} and liftings to pseudoalgebras. In [24] this was written as $\mathbf{Psm}(\mathcal{K})$, but we prefer to use that notation for the Gray-category introduced in Section 2, since it seems the natural generalization of the 2-category of monads defined by Street in [28]. We will then show that $\mathbf{Lift}(\mathcal{K})$ is equivalent to $\mathbf{Psm}(\mathcal{K})$, which will be used in Section 4 for our results on pseudodistributive laws.

The 0-cells of $\mathbf{Lift}(\mathcal{K})$ are pseudomonads (X, S) in \mathcal{K} . For 0-cells (X, S) and (Y, T) , a 1-cell $(F, \hat{F}) : (X, S) \rightarrow (Y, T)$ consists of a 1-cell $F : X \rightarrow Y$ in \mathcal{K} and a Gray-transformation $\hat{F} : \text{Ps-}S\text{-Alg} \rightarrow \text{Ps-}T\text{-Alg}$ making the following diagram commute

$$\begin{array}{ccc}
\text{Ps-}S\text{-Alg} & \xrightarrow{\hat{F}} & \text{Ps-}T\text{-Alg} \\
U \downarrow & & \downarrow U \\
X & \xrightarrow{F} & Y
\end{array}$$

where, using implicitly the Yoneda lemma for Gray-categories, we write X and Y instead of $\mathcal{K}(X, -)$ and $\mathcal{K}(Y, -)$. We refer to \hat{F} as a *lifting* of F to pseudoalgebras. Analogous terminology will be used for the 2- and 3-cells introduced below.

Lemma 3.1. *Let $(F, \phi) : (X, S) \rightarrow (Y, T)$ be a pseudomonad morphism. Then, there exists a lifting $\hat{F} : \text{Ps-}S\text{-Alg} \rightarrow \text{Ps-}T\text{-Alg}$ of $F : X \rightarrow Y$.*

Proof. Let us consider a fixed $I \in \mathcal{K}$. First, let us observe that if A is an I -indexed pseudo- S -algebra, then FA is naturally an I -indexed pseudo- T -algebra, with structure map given by the composite

$$TFA \xrightarrow{\phi_A} FSA \xrightarrow{Fa} FA$$

and associativity and unit 3-cells provided by the pasting diagrams

$$\begin{array}{ccc}
T^2FA & \xrightarrow{T\phi_A} & TFSa & \xrightarrow{TFa} & TFA \\
\downarrow n_{(FA)} & & \downarrow \phi_{SA} & \Downarrow \phi_a & \downarrow \phi_A \\
& & \downarrow \bar{\phi}_A & & \downarrow FSA \\
& & F S^2 A & \xrightarrow{FSa} & FSA \\
& & \downarrow Fm_A & \Downarrow F\bar{a} & \downarrow Fa \\
TFA & \xrightarrow{\phi_A} & FSA & \xrightarrow{Fa} & FA,
\end{array}
\qquad
\begin{array}{ccc}
FA & \xrightarrow{s_{FA}} & TFA \\
\downarrow F_{SA} & \xrightarrow{\tilde{\phi}_A} & \downarrow \phi_A \\
& & FSA \\
\downarrow 1_{FA} & \xrightarrow{F\tilde{a}} & \downarrow Fa \\
& & FA.
\end{array}$$

The coherence condition (1.6) for FA follows by an application of the coherence condition (2.1) for F and the coherence condition (1.6) for A . The coherence condition (1.7) for FA follows by applying the coherence condition (2.2) for F and the coherence condition (1.7) for A . Secondly, we observe that if $f : A \rightarrow B$ is a pseudo- S -algebra morphism, then $Ff : FA \rightarrow FB$ is naturally a pseudo- T -algebra morphism, as we have the following pasting diagram:

$$\begin{array}{ccc}
TFA & \xrightarrow{TFf} & TFB \\
\downarrow \phi_A & \Downarrow \phi_f & \downarrow \phi_B \\
FSA & \xrightarrow{FSf} & FSB \\
\downarrow Fa & \Downarrow F\bar{f} & \downarrow Fb \\
FA & \xrightarrow{Ff} & FB.
\end{array}$$

The coherence conditions (1.9) and (1.10) follow immediately by the axioms for a Gray-category. Finally, if $\alpha : f \rightarrow g$ is a pseudo- S -algebra 2-cell, the required pseudo- T -algebra 2-cell is given by $F\alpha : Ff \rightarrow Fg$. We have thus defined the components of a Gray-natural transformation $\hat{F} : \text{Ps-}S\text{-Alg} \rightarrow \text{Ps-}T\text{-Alg}$, which is clearly a lifting of $F : X \rightarrow Y$. \square

Given 1-cells $(F, \hat{F}) : (X, S) \rightarrow (Y, T)$ and $(F', \hat{F}') : (X, S) \rightarrow (Y, T)$, a 2-cell $(p, \hat{p}) : (F, \hat{F}) \rightarrow (F', \hat{F}')$ in $\mathbf{Lift}(\mathcal{K})$ consists of a 2-cell $p : F \rightarrow F'$ in \mathcal{K} and a Gray-modification $\hat{p} : \hat{F} \rightarrow \hat{F}'$ such that the following diagram commutes

$$\begin{array}{ccc}
U\hat{F} & \xrightarrow{U\hat{p}} & U\hat{F}' \\
\parallel & & \parallel \\
FU & \xrightarrow{pU} & F'U.
\end{array}$$

The vertical arrows are the identities, which hold by the assumption that \hat{F} and \hat{F}' are liftings of F and F' , respectively.

Lemma 3.2. *Let $(p, \tilde{p}) : (F, \phi) \rightarrow (F', \phi')$ be a pseudomonad transformation. Then, there exists a lifting $\hat{p} : \hat{F} \rightarrow \hat{F}'$ of $p : F \rightarrow F'$, where \hat{F} and \hat{F}' are the liftings of F and F' associated to the pseudomonad morphisms (F, ϕ) and (F', ϕ') , respectively, defined as in Lemma 3.1.*

Proof. Let $I \in \mathcal{K}$. We need to define a pseudonatural transformation $\hat{p} : \hat{F}_I \rightarrow \hat{F}'_I$. We define the component of \hat{p} associated to an I -indexed pseudo- S -algebra A to be the I -indexed pseudo- T -algebra morphism given by $p_A : FA \rightarrow F'A$ and the 2-cell

$$\begin{array}{ccc} TFA & \xrightarrow{Tp_A} & TF'A \\ \phi_A \downarrow & \Downarrow \bar{p}_A & \downarrow \phi'_A \\ FSA & \xrightarrow{p_{SA}} & F'SA \\ Fa \downarrow & \Downarrow p_a^{-1} & \downarrow F'a \\ FA & \xrightarrow{p_A} & F'A. \end{array}$$

To prove the condition (1.9) for the pseudoalgebra morphism p_A , we apply the axioms for a Gray-category and then condition (2.3) for the pseudomonad transformation p . To establish condition (1.10), it is sufficient to apply the coherence condition (2.4) for the pseudomonad transformation p , and then the axioms for a Gray-category. By definition, \hat{p} is a lifting of p as required. \square

Finally, for 2-cells (p, \hat{p}) and (q, \hat{q}) , a 3-cell $\alpha : (p, \hat{p}) \rightarrow (q, \hat{q})$ consists of a 3-cell and $\alpha : p \rightarrow q$ and a Gray-perturbation $\hat{\alpha} : \hat{p} \rightarrow \hat{q}$ making the following diagram commute

$$\begin{array}{ccc} U\hat{p} & \xrightarrow{U\hat{\alpha}} & U\hat{q} \\ \parallel & & \parallel \\ pU & \xrightarrow{\alpha U} & qU. \end{array}$$

As before, the vertical arrows are the identities that are part of the assumption that \hat{p} and \hat{q} are liftings of p and q , respectively. Composition and identities of $\mathbf{Lift}(\mathcal{K})$ are defined in the evident way, using those of \mathcal{K} and \mathbf{Gray} .

Lemma 3.3. *Given a pseudomonad modification $\alpha : (p, \tilde{p}) \rightarrow (q, \tilde{q})$ we can define a lifting $\hat{\alpha} : \hat{p} \rightarrow \hat{q}$ of α as the Gray-perturbation whose components are, for a pseudo- S -algebra A , the 3-cells $\alpha_A : p_A \rightarrow q_A$.*

Proof. It suffices to check that, these 3-cells are a pseudo- T -algebra 2-cells. To prove this, apply the axioms for a Gray-category and the coherence axiom (2.5). \square

We use these results to define a Gray-functor

$$\Phi : \mathbf{Psm}(\mathcal{K}) \rightarrow \mathbf{Lift}(\mathcal{K}).$$

On objects, Φ acts as the identity. For two pseudomonads (X, S) and (Y, T) in \mathcal{K} , the hom-2-functors

$$\Phi_{(X,S), (Y,T)} : \mathbf{Psm}(\mathcal{K})((X, S), (Y, T)) \longrightarrow \mathbf{Lift}(\mathcal{K})((X, S), (Y, T))$$

are defined sending a pseudomonad morphism, pseudomonad transformation and pseudomonad modification to the associated liftings, using Lemmas 3.1, 3.2 and 3.3, respectively. Here, the Gray-functoriality of Φ is standard verification, which we omit for

brevity, limiting ourselves to highlight that this includes checking that Φ preserves composition strictly, *i.e.* that the lifting associated to the composite of two pseudomonad morphisms is equal (rather than just equivalent by invertible 2-cells) to the composite of the liftings obtained from the pseudomonad morphisms. Theorem 3.4 states that the construction of $\mathbf{Psm}(\mathcal{K})$ given in Section 2 is equivalent to the one by Marmolejo in [24].

Theorem 3.4. *The Gray-functor $\Phi: \mathbf{Psm}(\mathcal{K}) \rightarrow \mathbf{Lift}(\mathcal{K})$ is a triequivalence.*

Proof. Since Φ is clearly bijective on objects, it suffices to prove that locally it is a biequivalence. Let us begin by considering a lifting $\hat{F}: \mathbf{Ps}\text{-}S\text{-Alg} \rightarrow \mathbf{Ps}\text{-}T\text{-Alg}$ of a 1-cell $F: X \rightarrow Y$. By the definition of a lifting, the following diagram of 2-categories and 2-functors commutes:

$$(3.1) \quad \begin{array}{ccc} \mathbf{Ps}\text{-}S\text{-Alg}(X) & \xrightarrow{\hat{F}_X} & \mathbf{Ps}\text{-}T\text{-Alg}(X) \\ U_X \downarrow & & \downarrow U_X \\ \mathcal{K}(X, X) & \xrightarrow{\mathcal{K}(X, F)} & \mathcal{K}(X, Y). \end{array}$$

Let us now observe that $S: X \rightarrow X$ can be regarded as an X -indexed pseudo- S -algebra, with structure map given by the 2-cell $m: S^2 \rightarrow S$. By the commutativity of the diagram (3.1), this pseudo- S -algebra is mapped by the 2-functor \hat{F}_X into a pseudo- T -algebra with underlying 1-cell $FS: X \rightarrow Y$, with structure map a 2-cell of the form $\phi_0: TFS \rightarrow FS$, and invertible 3-cells fitting in the diagrams

$$\begin{array}{ccc} T^2FS & \xrightarrow{T\phi_0} & TFS \\ n_{FS} \downarrow & \Downarrow \bar{\phi}_0 & \downarrow \phi_0 \\ TFS & \xrightarrow{h'} & FS \end{array} \quad \begin{array}{ccc} FS & \xrightarrow{t_{FS}} & TFS \\ \searrow 1_{FS} & \xRightarrow{\tilde{\phi}_0} & \downarrow h' \\ & & FS. \end{array}$$

The desired pseudomonad morphism $(F, \phi): (X, S) \rightarrow (Y, T)$ is then obtained by letting $\phi: TF \rightarrow FS$ be the composite

$$TF \xrightarrow{TFs} TFS \xrightarrow{\phi_0} FS.$$

The appropriate 3-cells are provided by the following pasting diagrams

$$\begin{array}{ccc} T^2F & \xrightarrow{T^2Fs} & T^2FS & \xrightarrow{T\phi_0} & TFS & \xrightarrow{TFsS} & TFS^2 \\ n_F \downarrow & \Downarrow n_{Fs} & \downarrow n_{FS} & \Downarrow \bar{\phi}_0 & \downarrow \phi_0 & \Downarrow \gamma & \downarrow \phi_0 S \\ TF & \xrightarrow{TFs} & TFS & \xrightarrow{\phi_0} & FS & & FS^2 \\ & & & & & & \downarrow Fm \\ & & & & & & FS \end{array} \quad \begin{array}{ccc} F & \xrightarrow{tF} & TF \\ F_s \downarrow & \xRightarrow{t_{Fs}} & \downarrow TF_s \\ FS & \xrightarrow{tFS} & TFS \\ \searrow 1_{FS} & \xRightarrow{\tilde{\phi}_0} & \downarrow \phi_0 \\ & & FS, \end{array}$$

where γ is the inverse to the 2-cell obtained from the following pasting of invertible 2-cells:

$$\begin{array}{ccc}
TFS & & \\
\downarrow TFsS & \searrow^{1_{TFS}} & \\
TFS^2 & \xrightarrow{TFm} & TFS \\
\downarrow \phi_0 S & \Downarrow F\alpha & \downarrow \phi_0 \\
FS^2 & \xrightarrow{Fm} & FS.
\end{array}$$

Let us now consider a lifting $(p, \hat{p}) : (F, \hat{F}) \rightarrow (F', \hat{F}')$ of a 2-cell $p : F \rightarrow F'$. We can define a pseudomonad transformation $p : (F, \phi) \rightarrow (F', \phi')$ by considering the following pasting diagram:

$$\begin{array}{ccc}
TF & \xrightarrow{Tp} & TF' \\
\downarrow TFs & \Downarrow Tp_u^{-1} & \downarrow TF's \\
TFS & \xrightarrow{TpS} & TF'S \\
\downarrow \phi_o & \Downarrow p\bar{S} & \downarrow \phi'_o \\
FS & \xrightarrow{pS} & F'S,
\end{array}$$

in which the bottom 3-cell is part of the structure making $pS : FS \rightarrow F'S$ into a pseudoalgebra morphism. Finally, if $(\alpha, \hat{\alpha}) : (p, \hat{p}) \rightarrow (q, \hat{q})$ is a lifting of a 3-cell $\alpha : p \rightarrow q$, then $\alpha : p \rightarrow q$ is a pseudomonad modification. These definitions determine a 2-functor

$$\Psi_{(X,S),(Y,T)} : \mathbf{Lift}(\mathcal{K})((X, S), (Y, T)) \longrightarrow \mathbf{Psm}(\mathcal{K})((X, S), (Y, T))$$

which provides the required quasi-inverse to $\Phi_{(X,S),(Y,T)}$. We omit the construction of the required invertible pseudonatural transformations $\eta : 1 \rightarrow \Psi\Phi$ and $\varepsilon : \Phi\Psi \rightarrow 1$, since this is not difficult. \square

4. PSEUDODISTRIBUTIVE LAWS

Definition 4.1. Let (X, S) and (X, T) be pseudomonads in \mathcal{K} . A *pseudodistributive law* of T over S consists of a 2-cell $d : ST \rightarrow TS$ and invertible 3-cells

$$\begin{array}{ccc}
S^2T \xrightarrow{Sd} STS & & T \xrightarrow{sT} ST \\
\downarrow mT & \Downarrow \bar{m} & \downarrow d \\
ST \xrightarrow{d} TS & & TS
\end{array}$$

$$\begin{array}{ccc}
ST^2 & \xrightarrow{Sn} & ST \\
dT \downarrow & & \downarrow d \\
TST & \Downarrow \bar{n} & \\
Td \downarrow & & \\
T^2S & \xrightarrow{nS} & TS
\end{array}
\qquad
\begin{array}{ccc}
& & ST \\
& \nearrow St & \downarrow d \\
S & \xrightarrow{tS} & TS \\
& & \Downarrow \bar{t}
\end{array}$$

satisfying the coherence conditions (C1)-(C8) stated in Appendix A.

Remark 4.2. For the convenience of the reader, Table 1 describes the correspondence between the presentation of the coherence conditions for pseudodistributive laws here and in [24, 29]. In the table, each row lists different formulations of the same axiom.

Appendix A	Marmolejo [24]	Tanaka [29]
(C1)	(coh 4)	(T6)
(C2)	(coh 2)	(T2)
(C3)	(coh 5)	(T9)
(C4)	(coh 3)	(T8)
(C5)	(coh 1)	(T1)
(C6)	(coh 6)	(T10)
(C7)	(coh 9)	(T7)
(C8)	(coh 7)	(T5)
(C9)	-	(T3)
(C10)	(coh 8)	(T4)

TABLE 1. Comparison of coherence conditions.

Remark 4.3. Our development in Section 3 allows us to give a clear explanation for the coherence conditions for pseudodistributive laws, summarised in Table 2.

Axiom	Coherence condition
(C1) and (C2)	$(T, d) : (X, S) \rightarrow (X, S)$ is a pseudomonad morphism
(C3) and (C4)	$(n, \bar{n}) : (T, d)^2 \rightarrow (T, d)$ is a pseudomonad transformation
(C5) and (C6)	$(t, \bar{t}) : (X, 1_X) \rightarrow (T, d)$ is a pseudomonad transformation
(C7)	α is pseudomonad modification
(C8)	ρ is a pseudomonad modification
(C9)	λ is a pseudomonad modification

TABLE 2. Coherence axioms for pseudodistributive laws.

The coherence axioms, (C9) and (C10) of Appendix A have been shown to be derivable from the others in [26, Theorem 2.3 and Proposition 4.2]. Indeed, axiom (C9) is a particular case of a provable coherence condition for a pseudomonad morphism and follows from (C1) and (C2) (*cf.* Proposition 2.2). By duality, one can see that axiom (C10)

is a particular case of a provable coherence condition for a pseudomonad op-morphism and follows from (C7) and (C8).

The explanation of the axioms for a pseudodistributive law in Remark 4.3 proves the following straightforward, but important, proposition.

Proposition 4.4. *The objects of $\mathbf{Psm}(\mathbf{Psm}(\mathcal{K}))$ are exactly pseudodistributive laws in \mathcal{K} .*

Proof. An object of $\mathbf{Psm}(\mathbf{Psm}(\mathcal{K}))$ consists of an object (X, S) of $\mathbf{Psm}(\mathcal{K})$, i.e. a pseudomonad in \mathcal{K} , together with a pseudomonad $(T, d): (X, S) \rightarrow (X, S)$ on it in $\mathbf{Psm}(\mathcal{K})$, which is exactly a pseudodistributive law by Remark 4.3. \square

We can now give a new simple proof of Marmolejo's fundamental result asserting the equivalence between a pseudodistributive law of a pseudomonad T over a pseudomonad S and a lifting of the pseudomonad T to the 2-categories of pseudoalgebras for S [24].

Theorem 4.5. *Let \mathcal{K} be a Gray-category, (X, S) and (X, T) be pseudomonads in \mathcal{K} . A pseudodistributive law $d: ST \rightarrow TS$ is equivalent to a lifting of T to pseudo- S -algebras.*

Proof. By Theorem 2.5, $\mathbf{Psm}(\mathcal{K})$ is a Gray-category and therefore we can consider the Gray-category $\mathbf{Psm}(\mathbf{Psm}(\mathcal{K}))$.

Next, observe that that $\mathbf{Psm}(-)$ preserves triequivalences between Gray-categories, i.e. given a triequivalence of Gray-categories $\Phi: \mathcal{K} \rightarrow \mathcal{K}'$, then it is possible to define a triequivalence $\mathbf{Psm}(\Phi): \mathbf{Psm}(\mathcal{K}) \rightarrow \mathbf{Psm}(\mathcal{K}')$. The construction of $\mathbf{Psm}(\Phi)$ is evident, and the verification that it is a triequivalence is a long, but routine, calculation. For example, to prove essential surjectivity, we need to show that for every pseudomonad (X', T') in \mathcal{K}' , there is a pseudomonad (X, T) in \mathcal{K} that is mapped by $\mathbf{Psm}(\Phi)$ to a pseudomonad that is biequivalent to (X', T') in $\mathbf{Psm}(\mathcal{K}')$. For this, one defines (X, T) using the essential surjectivity of Φ , carefully inserting coherence isomorphisms that are part of the given triequivalence where appropriate.

Applying this fact to the triequivalence of Theorem 3.4, we get a triequivalence:

$$\mathbf{Psm}(\mathbf{Psm}(\mathcal{K})) \simeq \mathbf{Psm}(\mathbf{Lift}(\mathcal{K})).$$

An object on the left hand side is exactly a pseudodistributive law by Proposition 4.4. Similarly, an object on the right hand side consists exactly of a pseudomonad (X, S) in \mathcal{K} and a pseudomonad $T: X \rightarrow X$ with a lifting $\hat{T}: \text{Ps-}S\text{-Alg} \rightarrow \text{Ps-}S\text{-Alg}$. \square

We conclude the paper by outlining how duality can be applied as in [28, Section 4] to obtain an equivalence between pseudodistributive laws and extensions to Kleisli objects. Fix a Gray-category \mathcal{K} and let (X, T) be a pseudomonad in it. By definition, a *right pseudo- T -module* in \mathcal{K} is a left T -module in \mathcal{K}^{op} . We then have a Gray-functor

$$(4.1) \quad \text{Mod}_T: \mathcal{K}^{op} \rightarrow \mathbf{Gray}.$$

Assuming the evident definition of a lifting to 2-categories of right pseudomodules, we have the following corollary of Theorem 4.5.

Corollary 4.6. *Let (X, S) and (X, T) be pseudomonads in \mathcal{K} . A pseudodistributive law $d: ST \rightarrow TS$ is equivalent to a lifting of S to right pseudo- T -modules.* \square

The equivalence of Corollary 4.6 becomes more familiar under the assumption that \mathcal{K} has Kleisli objects. Recall that a *Kleisli object* for a pseudomonad (X, T) in \mathcal{K} is an 0-cell $X_T \in \mathcal{K}$ and a right pseudo- T -module $J_T: X \rightarrow X_T$, which is universal in the sense that the 2-functor

$$\mathcal{K}(X_T, I) \rightarrow \text{Mod}_T(I),$$

induced by composition with J_T , is an equivalence of 2-categories, thus making the Gray-functor in (4.1) representable. Now, a pseudodistributive law $d: ST \rightarrow TS$ is equivalent to a lifting of S to right pseudo- T -modules, as in

$$\begin{array}{ccc} \text{Mod}_T & \xrightarrow{\hat{S}} & \text{Mod}_T \\ \downarrow U & & \downarrow U \\ \mathcal{K}(-, X) & \xrightarrow{S \circ -} & \mathcal{K}(-, X). \end{array}$$

This, in turn, is equivalent to

$$\begin{array}{ccc} X_T & \xrightarrow{\hat{S}} & X_T \\ \uparrow J_T & & \uparrow J_T \\ X & \xrightarrow{S} & X, \end{array}$$

which describes an extension of S to the Kleisli object of T .

Remark 4.7. We conclude the paper by briefly discussing the question of whether the Gray-category **Gray** has Kleisli objects. Given a 2-category X and pseudomonad $T: X \rightarrow X$, there are two reasonable options to be the Kleisli object for T , mirroring the one-dimensional situation. In both cases, the objects are the same objects as those of X , but they have different hom-categories of morphisms. The first option is to define the hom-category between two objects $x, y \in X$ to be $X(x, Ty)$. With this definition we only get a bicategory, not a 2-category, and so we step outside **Gray**. The second option (which we will call X_T), is to take the hom-category of morphisms between x and y to consist of pseudoalgebra morphisms from Tx to Ty (considered as free algebras). This is a 2-category and so one could try to show that it is a Kleisli object for **Gray**. In order to do this, one should prove that, for any 2-category I , there is an equivalence as in (4.1). However, the construction taking a I -indexed right pseudo- T -module to a 2-functor $X_T \rightarrow I$ is only a pseudofunctor and not a strict 2-functor, thus leading again outside **Gray**. The reason for this is that we need to use the pseudonaturality of the module action λ and other coherence isomorphisms. Because of this, it seems that **Gray** does not have Kleisli objects. We suspect that, once it is defined what it means for a tricategory to have Kleisli objects, it should be possible to show that the tricategory $\mathbf{2-Cat}_{\text{psd}}$ of 2-categories, pseudofunctors, pseudonatural transformations and modifications has Kleisli objects. The same should hold also for **Bicat**, the tricategory

of bicategories, pseudofunctors, pseudonatural transformation and modifications. We leave the investigation of these problems to future research.

APPENDIX A. COHERENCE CONDITIONS FOR PSEUDODISTRIBUTIVE LAWS

We limit ourselves to drawing the boundaries of these diagrams and explain in text which 3-cells should be inserted in them, except for the 3-cells coming from the structure of a Gray-category of \mathcal{K} .

(C1)

=

In (C1), the left-hand side pasting is obtained using $S\bar{m}$, \bar{m} , and the associativity 3-cell of the pseudomonad S ; the right-hand side pasting is obtained using the associativity 3-cell of the pseudomonad S and \bar{m} .

$$(C2) \quad \begin{array}{ccc} ST & \xrightarrow{ST_s} & STS \\ \downarrow S_s T & \searrow S_d & \downarrow dS \\ S^2 T & \xrightarrow{S_d} & STS \\ \downarrow mT & & \downarrow dS \\ ST & \xrightarrow{d} & TS \\ \uparrow 1_{ST} & & \downarrow Tm \\ & & TS^2 \end{array} = \begin{array}{ccc} ST & \xrightarrow{ST_s} & STS \\ \downarrow d & & \downarrow dS \\ TS & \xrightarrow{TS_s} & TS^2 \\ \downarrow Id & & \downarrow Tm \\ & & TS \end{array}$$

In (C2), the left-hand side pasting is obtained using $S\bar{s}$, \bar{m} , and the left unit 3-cell of the pseudomonad S ; the right-hand side pasting is obtained using the left unit 3-cell of the pseudomonad S .

$$(C3) \quad \begin{array}{ccc} S^2 & \xrightarrow{S^2 t} & S^2 T \\ \downarrow m & \searrow stS & \downarrow dS \\ S & \xrightarrow{tS} & TS \\ \downarrow tS & & \downarrow Tm \\ & & TS^2 \end{array} = \begin{array}{ccc} S^2 & \xrightarrow{S^2 t} & S^2 T \\ \downarrow m & \searrow mT & \downarrow dS \\ S & \xrightarrow{St} & ST \\ \downarrow tS & & \downarrow Tm \\ & & TS \end{array}$$

For (C3), the left-hand side pasting is obtained using $S\bar{t}$, $\bar{t}S$; the right-hand side is obtained using \bar{m} and \bar{m} .

$$(C4) \quad \begin{array}{ccc} 1_X & \xrightarrow{t} & T \\ \downarrow s & \searrow sT & \downarrow Ts \\ S & \xrightarrow{St} & ST \\ \downarrow tS & \searrow d & \downarrow \\ & & TS \end{array} = \begin{array}{ccc} 1_X & \xrightarrow{t} & T \\ \downarrow s & & \downarrow Ts \\ S & \xrightarrow{tS} & TS \end{array}$$

For (C4), the left-hand side pasting is obtained using \bar{s} and \bar{t} ; the right-hand side is obtained from pseudonaturality of t .

$$\begin{array}{c}
 \begin{array}{c}
 (C5) \quad S^2T^2 \xrightarrow{S^2n} S^2T \\
 \downarrow mT^2 \quad \searrow SdT \\
 STST \\
 \downarrow dST \quad \searrow STd \\
 TS^2T \quad \xrightarrow{SnS} \quad STS \\
 \downarrow TSd \quad \searrow dTS \\
 TSTS \\
 \downarrow TmT \quad \searrow TdS \\
 ST^2 \quad \xrightarrow{nS^2} \quad TS^2 \\
 \downarrow dT \quad \searrow T^2m \\
 TST \quad \xrightarrow{nS} \quad TS \\
 \downarrow Td \\
 T^2S
 \end{array}
 & = &
 \begin{array}{c}
 S^2T^2 \xrightarrow{S^2n} S^2T \\
 \downarrow mT^2 \quad \searrow SdT \\
 ST^2 \quad \xrightarrow{Sn} \quad ST \\
 \downarrow dT \quad \searrow d \\
 TST \quad \xrightarrow{Td} \quad TS \\
 \downarrow Td \\
 T^2S \quad \xrightarrow{nS} \quad TS \\
 \downarrow Tm \\
 TS^2 \\
 \downarrow dS \\
 STS
 \end{array}
 \end{array}$$

For (C5), the left-hand side pasting is obtained using $S\bar{n}$, $\bar{n}S$ and $\bar{m}T$; the right-hand side pasting is obtained using \bar{m} and \bar{n} .

(C6)

In (C6), the left-hand side pasting is obtained using \bar{s} and \bar{n} ; the right-hand side pasting is obtained using $\bar{s}T$.

(C7)

For (C7), the left-hand side pasting is obtained using the associativity 3-cell of the pseudomonad T , \bar{n} and $\bar{n}T$; the right-hand side pasting is obtained using $T\bar{n}$, \bar{n} and the associativity 3-cell of the pseudomonad T .

(C8)

The diagram (C8) consists of two parts separated by an equals sign. The left part is a commutative diagram with nodes ST , ST^2 , TST , TS , and T^2S . Arrows include 1_{ST} (curved top), StT (top-left), S_n (top-right), d (vertical down from ST and TS), tST (diagonal down-left), dT (vertical down from ST^2), Td (vertical down from TST), tTS (bottom-left), and nS (bottom-right). The right part is a simpler commutative diagram with nodes ST , TS , and T^2S . Arrows include 1_{TS} (curved top), tTS (bottom-left), and nS (bottom-right).

For (C8), the left-hand side pasting is obtained using the right unit 3-cell of the pseudomonad T , \bar{n} , $\bar{t}S$; the right-hand side pasting is the right unit 3-cell of the pseudomonad T .

(C9)

The diagram (C9) consists of two parts separated by an equals sign. The left part is a commutative diagram with nodes ST , TS , S^2T , STS , TS^2 , and ST . Arrows include d (top), sST (left), sTS (middle-left), TsS (middle-right), 1_{TS} (curved top-right), Sd (bottom-left), dS (bottom-middle), Tm (bottom-right), and mT (diagonal down-left). The right part is a simpler commutative diagram with nodes ST , S^2T , and ST . Arrows include sST (left), 1_{ST} (curved top-right), mT (diagonal down-left), and d (bottom).

For (C9), the left-hand side pasting is obtained using the right unit 3-cell of the pseudomonad S , $\bar{s}S$ and \bar{m} ; the right-hand side pasting is obtained using the right unit 3-cell of the pseudomonad S .

(C10)

For (C10), the left-hand side pasting uses the left unit 3-cell of the pseudomonad T . The right-hand side pasting is obtained using \bar{n} and the left unit 3-cell of the pseudomonad T .

APPENDIX B. SOME TECHNICAL PROOFS

Coherence diagrams for $(GF, G\phi \cdot \psi F)$. We show only equation in (2.2). Using the coherence diagram (2.2) for (G, ψ) and for (F, ϕ) , it suffices to prove that the following two diagrams are equal:

$$\begin{array}{ccc}
QGF & \xrightarrow{QGF_s} & QGFS \\
\downarrow \psi_F & \Downarrow \psi_{F_s} & \downarrow \psi_{FS} \\
GTF & \xrightarrow{GTF_s} & GTFS \\
\downarrow GT_tF & \Downarrow GT_{\tilde{\phi}} & \downarrow G\phi_S \\
& GT^2F \xrightarrow{GT_\phi} & GTFS \\
\downarrow 1_{GTF} & \Downarrow G\eta_{TF} & \downarrow G\phi_S \\
& GnF \Downarrow G\bar{\phi} & GFS^2 \\
& \downarrow GnF & \downarrow GF_m \\
GTF & \xrightarrow{G\psi} & GFS.
\end{array}$$

This equality holds using (1.1) and (1.2).

F_ϕ is well-defined. Given a pseudomonad transformation $(q, \bar{q}) : (G, \psi) \rightarrow (G', \psi')$ in $P_{\mathcal{K}}((Y, T), (Z, Q))$ we want to show that (qF, \overline{qF}) is a pseudomonad transformation as well. We will show just equation (2.3), since (2.4) can be proved similarly. The required equality follows from equation (2.3) for q and the equation below, which can be proved using (1.1) twice.

$$\begin{array}{ccc}
QGTF & \xrightarrow{QqTF} & QG'TF \\
\downarrow \psi TF & \searrow QG\phi & \searrow Qq\phi^{-1} \quad \searrow QG'\phi \\
& & QGFS \xrightarrow{QqFS} QG'FS \\
& \searrow \psi\phi \Downarrow & \downarrow \psi FS \quad \Downarrow \bar{q}FS \quad \downarrow \psi'FS \\
GT^2F & \xrightarrow{GT\phi} & GTFS \xrightarrow{qTFS} G'TFS \\
& \searrow GnF & \downarrow G\phi S \quad \Downarrow q\phi S^{-1} \quad \downarrow G'\phi S \\
& & GFS^2 \xrightarrow{qFS^2} G'FS^2 \\
& \searrow G\bar{\phi} \Downarrow & \downarrow GFm \quad \Downarrow qFm^{-1} \quad \downarrow G'Fm \\
GTF & \xrightarrow{G\phi} & GFS \xrightarrow{qFS} G'FS
\end{array}
=
\begin{array}{ccc}
QG'TF & \xrightarrow{QqTF} & QG'TF \\
\downarrow \psi TF & \searrow \bar{q}TF \Downarrow & \downarrow \psi'TF \quad \searrow QG'\phi \\
& & GT^2F \xrightarrow{qT^2F} G'T^2F \\
& \searrow GnF & \downarrow qnF \quad \Downarrow qnF \quad \downarrow G'nF \\
& & GTF \xrightarrow{qTF} G'TF \\
& \searrow G\phi & \searrow q\phi^{-1} \Downarrow \quad \searrow G'\phi \\
& & GFS \xrightarrow{qFS} G'FS
\end{array}$$

Let

$$(q, \bar{q}), (q', \bar{q}') : (G, \psi) \rightarrow (G', \psi')$$

be pseudomonads transformations in $P_{\mathcal{K}}((Y, T), (Z, Q))$. Given a pseudomonad modification $\beta : (q, \bar{q}) \rightarrow (q', \bar{q}')$ we want to show that βF is a pseudomonad modification from $(qF, \bar{q}F)$ to $(q'F, \bar{q}'F)$. So we need to show the following equation

$$\begin{array}{ccc}
 \begin{array}{ccc}
 QGF & \xrightarrow{QqF} & QG'F \\
 \downarrow \psi F & \Downarrow Q\beta F & \downarrow \psi' F \\
 QGF & \xrightarrow{Qq'F} & QG'F \\
 \downarrow \psi F & \Downarrow \bar{q}'F & \downarrow \psi' F \\
 GTF & \xrightarrow{q'TF} & G'TF \\
 \downarrow G\phi & \Downarrow q_\phi^{-1} & \downarrow G'\phi \\
 GFS & \xrightarrow{q'FS} & G'FS
 \end{array} & = &
 \begin{array}{ccc}
 QGF & \xrightarrow{QqF} & QG'F \\
 \downarrow \psi F & \Downarrow \bar{q}F & \downarrow \psi' F \\
 GTF & \xrightarrow{qTF} & G'TF \\
 \downarrow G\phi & \Downarrow q_\phi^{-1} & \downarrow G'\phi \\
 GFS & \xrightarrow{qFS} & G'FS \\
 \downarrow G\phi & \Downarrow \beta FS & \downarrow G'\phi \\
 GFS & \xrightarrow{q'FS} & G'FS
 \end{array}
 \end{array}$$

This can be shown to hold using the coherence axiom for β and (1.3).

Coherence for q_p . Given $(p, \bar{p}) : (F, \phi) \rightarrow (F', \phi')$ and $(q, \bar{q}) : (G, \psi) \rightarrow (G', \psi')$ two 2-cells in $P_{\mathcal{K}}$ we want to prove that $q_p : G'p \cdot qF \rightarrow qF' \cdot Gp$ is a 3-cell in $P_{\mathcal{K}}$. First of all, it is useful to note that

$$\begin{array}{ccc}
 \begin{array}{ccc}
 GTF & \xrightarrow{qTF} & G'TF \\
 \downarrow G\phi & \Downarrow q_\phi^{-1} & \downarrow G'\phi \\
 GTF & \xrightarrow{qTF} & G'TF \\
 \downarrow G\phi & \Downarrow q_\phi^{-1} & \downarrow G'\phi \\
 GFS & \xrightarrow{qFS} & G'FS \\
 \downarrow G\phi & \Downarrow q_p S^{-1} & \downarrow G'\phi \\
 GFS & \xrightarrow{qFS} & G'FS
 \end{array} & = &
 \begin{array}{ccc}
 GTF & \xrightarrow{qTF} & G'TF \\
 \downarrow G\phi & \Downarrow q_{Tp}^{-1} = (qT)_p^{-1} & \downarrow G'\phi \\
 GTF & \xrightarrow{qTF} & G'TF \\
 \downarrow G\phi & \Downarrow q_{Tp}^{-1} = (qT)_p^{-1} & \downarrow G'\phi \\
 GFS & \xrightarrow{qFS} & G'FS \\
 \downarrow G\phi & \Downarrow q_{\phi'}^{-1} & \downarrow G'\phi \\
 GFS & \xrightarrow{qFS} & G'FS
 \end{array}
 \end{array}$$

This equality is true since both diagrams are equal to the following one, using (1.3) for the one on the left-hand side and (1.1) for the one on the right-hand side.

$$\begin{array}{ccc}
GTF & \xrightarrow{qTF} & G'TF \\
\downarrow G(pS \cdot \phi) & \swarrow q_{(pS \cdot \phi)}^{-1} & \downarrow G'(\phi' \cdot Tp) \\
GF'S & \xrightarrow{g'F} & G'F'S
\end{array}$$

$\begin{array}{c} \left(\begin{array}{c} \leftarrow \\ \leftarrow \\ \leftarrow \\ \leftarrow \\ \leftarrow \end{array} \right) \\ \leftarrow \\ \leftarrow \\ \leftarrow \\ \leftarrow \\ \leftarrow \end{array}$

The proof can be concluded using the invertibility of the 3-cells involved and (1.1).

REFERENCES

- [1] M. Barr and C. Wells. *Toposes, triples, and theories*. Springer, 1985.
- [2] J. Beck. Distributive laws. In *Seminar on Triples and Categorical Homology Theory (ETH, Zürich, 1966/67)*, pages 119–140. Springer, 1969.
- [3] M. Bunge. Coherent extensions and relational algebras. *Transactions of the American Mathematical Society*, 197:355–390, 1974.
- [4] G. L. Cattani and G. Winskel. Profunctors, open maps, and bisimulation. *Mathematical Structures in Computer Science*, 15(3):553–614, 2005.
- [5] E. Cheng, M. Hyland, and J. Power. pseudodistributive laws. *Electronic Notes in Theoretical Computer Science*, 83, 2003.
- [6] P.-L. Curien. Operads, clones, and distributive laws. Proc. of the International Conference on Operads and Universal Algebra, Nankai Series in Pure, Applied Mathematics and Theoretical Physics, Vol. 9, World Scientific, Singapore, 25-50 (2012).
- [7] M. Dostál. Two-dimensional universal algebra. PhD Thesis, Czech Technical University in Prague, 2018. <https://math.feld.cvut.cz/dostamat/research/phd-thesis.pdf>.
- [8] S. Eilenberg and G. M. Kelly. Closed categories. In *Proceedings of La Jolla Conference on Categorical Algebra*, pages 421–562. Springer-Verlag, 1966.
- [9] M. Fiore, N. Gambino, M. Hyland, and G. Winskel. The cartesian closed bicategory of generalised species of structures. *Journal of the London Mathematical Society*, 77(2):203–220, 2008.
- [10] M. Fiore, N. Gambino, M. Hyland, and G. Winskel. Relative pseudomonads, Kleisli bicategories, and substitution monoidal structures. *Selecta Mathematica New Series*, 24: 2791, 2018.
- [11] N. Gambino and A. Joyal. On operads, bimodules and analytic functors. *Memoirs of the American Mathematical Society*, 289(1184), (v) + 110pp, 2017.
- [12] R. Garner. Polycategories via pseudodistributive laws. *Advances in Mathematics*, 218(3):781–827, 2008.
- [13] R. Garner and N. Gurski. The low-dimensional structures formed by tricategories. *Mathematical Proceedings of the Cambridge Philosophical Society*, 146(3):551–589, 2009.
- [14] R. Gordon, A. J. Power, and R. Street. Coherence for tricategories. *Memoirs of the American Mathematical Society*, 117(558), 1995.
- [15] N. Gurski. *Coherence in Three-Dimensional Category Theory*. Cambridge University Press, 2013.
- [16] G. M. Kelly. Coherence theorems for lax algebras and for distributive laws. In G. M. Kelly and R. H. Street, editors, *Category Seminar (Proc. Sem., Sydney, 1972/1973)*, volume 420 of *Lecture Notes in Mathematics*, pages 281–375. Springer, 1974.
- [17] G. M. Kelly. *Basic Concepts of Enriched Category Theory*. Cambridge University Press, 1982.
- [18] G. M. Kelly and R.H. Street. Review of the elements of 2-categories. In G. M. Kelly and R. H. Street, editors, *Category Seminar (Proc. Sem. Sydney 1972/1973)*, volume 420 of *Lecture Notes in Mathematics*, pages 75–103. Springer, 1974.
- [19] S. Lack. A coherent approach to pseudomonads. *Advances in Mathematics*, 152:179–202, 2000.
- [20] S. Lack. Bicats is not triequivalent to Gray. *Theory and Applications of Categories*, 18(1):1–3, 2007.

- [21] S. Lack and R. Street. The formal theory of monads. II. *Journal of Pure and Applied Algebra*, 175(1-3):243–265, 2002.
- [22] S. Mac Lane. *Categories for the working mathematician*. 2nd ed., Springer, 1998.
- [23] F. Marmolejo. Doctrines whose structure forms a fully faithful adjoint string. *Theory and applications of categories*, 3(2):24–44, 1997.
- [24] F. Marmolejo. Distributive laws for pseudomonads. *Theory and Applications of Categories*, 5(5):91–147, 1999.
- [25] F. Marmolejo. Distributive laws for pseudomonads II. *Journal of Pure and Applied Algebra*, 195(1-2):169–182, 2004.
- [26] F. Marmolejo and R. J. Wood. Coherence for pseudodistributive laws revisited. *Theory and Applications of Categories*, 20(6):74–84, 2008.
- [27] A. J. Power. A 2-Categorical Pasting Theorem. *Journal of Algebra*, volume 129, pages 439-445, 1990.
- [28] R. Street. The formal theory of monads. *Journal of Pure and Applied Algebra*, 2(2):149–168, 1972.
- [29] M. Tanaka. *Pseudodistributive laws and a unified framework for variable binding*. PhD thesis, Laboratory for the Foundations of Computer Science, School of Informatics, University of Edinburgh, 2004. Available as LFCS Technical Report ECS-LFCS-04-438.
- [30] M. Tanaka and J. Power. Pseudodistributive laws and axiomatics for variable binding. *Higher-order and Symbolic Computation*, 19(2-3):305–337, 2006.
- [31] M. Tanaka and J. Power. A unified category-theoretic semantics for binding signatures in substructural logics. *Journal of Logic and Computation*, 16:5–25, 2006.
- [32] C. Walker. Distributive laws via admissibility. *Applied Categorical Structures*, 27(6) 567–617, 2019.