

Strengthening International Research in Long-Term Care: Recommended Common Data Elements to Support Clinical Staff Training

Gerontology & Geriatric Medicine
Volume 7: 1–17
© The Author(s) 2021
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: [10.1177/233721421999312](https://doi.org/10.1177/233721421999312)
journals.sagepub.com/home/ggm



Charlene H. Chu, RN, GNC(c), PhD^{1,2} , Katherine S. McGilton, RN, PhD, FAAN, FCAHS^{1,2} , Karen Spilsbury, RN, PhD³, Kim N. Le, RN, HBSc, BScN¹ , Veronique Boscart, RN, PhD^{2,4} , Annica Backman, RN, PhD⁵, Anette Fagertun, PhD⁶, Reena Devi, PhD³, and Franziska Zúñiga, RN, PhD⁷ 

Abstract

The purpose of this study is to develop candidate common data element (CDE) items related to clinical staff training in long-term care (LTC) homes that can be used to enable international comparative research. This paper is part of the WE-THRIVE (Worldwide Elements to Harmonize Research in Long-Term Care Living Environments) group's initiative which aims to improve international academic collaboration. We followed best practices to develop CDEs by conducting a literature review of clinical staff (i.e., Regulated Nurses, Health Care Aides) training measures, and convening a subgroup of WE-THRIVE experts to review the literature review results to develop suitable CDEs. The international expert panel discussed and critically reflected on the current knowledge gaps from the literature review results. The panel proposed three candidate CDEs which focused on the *presence of* and the *measurement of* training. These three proposed CDEs seek to facilitate international research as well as assist in policy and decision-making regarding LTC homes worldwide. This study is a critical first step to develop candidate CDE items to measure staff training internationally. Further work is required to get feedback from other researchers about the proposed CDEs, and assess the feasibility of these CDEs in high and low resourced settings.

Keywords

common data elements, training, nursing home, long-term care, measurement

Manuscript received: February 5, 2021; **final revision received:** February 5, 2021; **accepted:** February 9, 2021.

As the global population ages, the spotlight is on long-term care (LTC) homes (e.g., care homes, nursing homes) to meet the needs of older adults with multiple complex and chronic conditions (Beard et al., 2016). Staff training is a widely accepted method to build staff capacity with broad-reaching benefits for residents (Alzheimers Disease International, 2013; Fujisawa & Colombo, 2009; Hussein & Manthorpe, 2005; Spector et al., 2016). Staff training involves programs (e.g., in-services and workshops) offered by the workplace that impart a set of specific skills or knowledge to improve employee performance in certain areas and settings (Davis & Lundstrom, 2011; Saponaro & Baughman, 2009). Such training augments and enhances individuals' knowledge, understanding, behaviors, skills, values,

¹Lawrence S. Bloomberg Faculty of Nursing, University of Toronto, Toronto, ON, Canada

²KITE—Toronto Rehabilitation Institute, University Health Network, Toronto, ON, Canada

³University of Leeds, West Yorkshire, UK

⁴Conestoga College Institute of Technology and Advanced Learning, Kitchener, ON, Canada

⁵Umeå University, Umeå, Sweden

⁶Western Norway University of Applied Sciences, Bergen, Vastland, Norway

⁷Nursing Science, Department Public Health, University of Basel, Basel, Switzerland

Corresponding Author:

Charlene H. Chu, Lawrence S. Bloomberg Faculty of Nursing, University of Toronto, 155 College Street, Unit 130, Toronto, ON M5T 1P8, Canada.

Email: charlene.chu@utoronto.ca



and beliefs (Davis & Lundstrom, 2011). Training, alongside other professional development opportunities, can also improve societal perceptions of careers in LTC homes and therefore attract and retain more people to the sector (Fujisawa & Colombo, 2009; Hussein & Manthorpe, 2005).

LTC homes are defined as facilities certified within their countries to provide continuous nursing care and personal support to residents that include assistance with their activities of daily living and complex health conditions (National Institute on Ageing, 2019). This care is often provided by informal caregivers (e.g., family members) and clinical staff. The current paper will focus on clinical staff defined as staff who are formally trained and employed part- or full-time by LTC homes (e.g., Registered Nurses, Registered Practical Nurses, Licensed Practical Nurses, and Health Care Aides/Personal Support Workers/Nursing Aides) to provide direct patient care. As clinical staff provide direct patient care, inadequate workplace training can negatively impact staff's attitudes and skills, the quality of care provided and worsen health outcomes for older people residing in LTC (Cooper et al., 2016). While preparatory education may encourage individuals to potentially seek employment in LTC, this paper is focused on supporting the clinical staff in LTC by means of on-the-job training which may be associated with staff retention, and the ability to enhance the quality of care and quality of life for residents in LTC (Caspar et al., 2016; Fujisawa & Colombo, 2009; Kane, 2003; Spector et al., 2016).

Staff Training in LTC: Challenges for International Research in LTC Homes

The provision of staff training in LTC is met with logistical and methodological challenges; first, previous research indicates that the variability of the LTC workforce, such as their professional backgrounds, cultural and sociodemographic characteristics, and educational levels, coupled with high rates of LTC staff turnover of an aging workforce in LTC, complicates the process to provide training (Fujisawa & Colombo, 2009; Hussein & Manthorpe, 2005). Second, while staff training is often prompted by the care needs of the residents in the LTC home, current training often fails to scaffold on current knowledge and inadequately addresses the knowledge, skills and competencies required (Stone & Harahan, 2010). Third, the OECD (Fujisawa & Colombo, 2009) has reported significant within and between country differences in the availability of the content, duration and emphasis on on-the-job training versus theoretical learning in training programs for LTC staff. Lastly, from a resource perspective, individual LTC homes' access to expertise, technology, and time will also influence the number and quality of staff training opportunities.

The lack of a minimum international standard for LTC training is prohibitive to accurately compare country experiences, curricula, and requirements (Fujisawa &

Colombo, 2009). Methodologically, the resultant heterogeneity with respect to the training details like frequency, content, modality of delivery (e.g., online or in-person), and evaluation, prevents the aggregation of data to build cross-national evidence about effective training in LTC. Most training initiatives have been in experimental stages in small-scale and/or did not include an evaluation component (Aylward et al., 2003; Beeber et al., 2010; Hussein & Manthorpe, 2005; Kuske et al., 2007). These previous literature reviews referred to the evidence on staff training from a variety of settings and countries to suggest that the effectiveness of training and its long-term impact on resident care remains unclear. In sum, gaps in evidence due to methodological weaknesses, differences in training content, training duration, organizational structures in LTC, and evaluation method makes it difficult to determine what common data elements (CDEs) are important to effective staff training from a cross-national perspective (Beeber et al., 2010; Hussein & Manthorpe, 2008; Kuske et al., 2007). There is a need to create a measurement infrastructure to better understand the current landscape of clinical staff training in LTC homes located in high and low resourced LTC settings.

International LTC research is a key lever to improve international development, policy, and practice, and more broadly enable the LTC workforce to prepare for the increasingly complex care needs of residents (Corazzini et al., 2019; World Health Organization [WHO], 2011). To this end, the “Worldwide Elements To Harmonize Research In LTC Living Environments” (WE-THRIVE) international group was formed to aggregate cross-country data to support the decision making in LTC (Corazzini et al., 2019; Lepore & Corazzini, 2019). WE-THRIVE aims to identify, develop, and evaluate the effectiveness of ecologically viable CDEs to further mutual understanding and facilitate international research in four overarching domains: LTC context; workforce and staffing; person-centered care; and care outcomes (Corazzini et al., 2019). This paper is a part of a broader body of literature examining the central concepts related to “workforce and staffing” domain (Zúñiga et al., 2019, McGilton et al., 2020). An international approach with CDEs to measure staff training in LTC will facilitate comparative research focused on the role of training in supporting staff and the workforce, and contribute to an international measurement infrastructure (Lepore & Corazzini, 2019).

The purpose of this study is to develop feasible candidate CDE items related to clinical staff training in LTC that can be used to enable international comparative research. This study is a critical first step to develop a small number of candidate CDE items that are ecologically viable to measure staff training in high to low income countries.

Methods

Following the similar approaches of recently published WE-THRIVE work (McGiltn et al., 2020), we followed the Best Practices for Identifying Common Data Elements

identified by Redeker et al. (2015) which involved: (1) completed a literature review of staff training outcome measures; and (2) convening a group of WE-THRIVE experts to review the literature review results to develop suitable CDEs. This is an iterative process by which the expert panel is convened, and the CDEs are developed based on the evidence over the course of multiple subgroup discussions (Redeker et al., 2015). The process of developing and proposing CDEs is ideally transparent, inclusive, and involves identifying and developing CDEs by national and international experts in the field (Redeker et al., 2015). The discussion points from the expert panel will be described in the results.

Literature Review

A WE-THRIVE subcommittee group focused on the “work staff and staffing” domain advised on the broad research questions and identified search terms to include in the search strategy. A literature review was completed to gain a broad understanding and describe the state of the literature of the “on-the-job training” provided to clinical staff in LTCs around the globe, followed by utilizing expert consensus based on the findings of the literature review. Arksey and O’Malley’s (2005) scoping review framework was used: (i) identifying the research questions, (ii) identifying relevant studies, (iii) study selection, (iv) extracting and charting the data, and (v) summarizing and reporting the results. The expert panel wanted to understand what the evidence to inform the development of CDEs were and asked the following questions: what are the characteristics of training provided to clinical staff in LTC home settings? What measurement tools are used to evaluate the training? Is staff training effective in influencing staff and resident outcomes?

Search strategy. An information specialist conducted a search in July 2019 in four electronic data sources: Medline, Embase, CINAHL, and HAPI. A secondary search was conducted on Google Scholar and only the first 200 results were screened. These databases were selected to cover a broad range of disciplines from various academic sources. The search was limited to peer-reviewed publications that were full-text manuscripts and published in the last 20 years in English.

The search query consisted of terms and their synonyms suggested by the expert panel and the information specialist. Clinical staff include health care workers and professionals like nurses (“Registered Nurses; Regulated Nurses; Registered Practical Nurses; Licensed Practical Nurses), aides (“Personal Support Workers, nursing aids, nursing assistants, care aids”), training (“education”), paired with terms “nursing homes” (“LTC homes; home for the aged”), and measurement (“questionnaire; survey”). Inclusion criteria included studies that quantitatively evaluated on-the-job training on any topic for clinical staff providing direct resident care in LTC home settings. Studies focusing on pre-service education were

excluded because WE-THRIVE is focused on building the capacity of the workforce in LTC settings. Citations were imported into Covidence, a web-based systematic review software to remove duplicates, and to conduct the title and abstract relevance screening.

Three research assistants (RAs) with nursing backgrounds alongside the PI (CC) conducted the screening. First, the title and abstract of citations were independently screened by two reviewers, and double checked. Studies that did not meet the eligibility criteria described above were excluded. Reviewers discussed any uncertainties to resolve conflicts related to study inclusion. Next, full-text review of the studies that were deemed relevant was completed. Articles were obtained through University institutional access. For articles that could not be accessed, the corresponding author was contacted to obtain the article. An extraction form in Excel was created based on the discussions and questions from the expert panel and included: author name, publication year, journal name, manuscript type, sample size, demographics of the sample, number of homes participating, research questions or aims, format, duration, and frequency of training, and forms of evaluation of the training, research design, attributes of staff training (training topic, summary of content, type and number of trainers, evaluation measures, follow-up duration), tools used to measure the training, how training is documented, impact of training on any nurse or resident outcomes, source of training funding, and study limitations. Outcomes related to training were included to inform a baseline understanding of the effect of training programs.

Expert panel. The formation of the expert panel has been described elsewhere (Corazzini et al., 2019; Zúñiga et al., 2019). We followed the best practices for identifying and developing CDEs (Redeker et al., 2015), and group processes were established within WE-THRIVE including forming a subgroup focused on the “workforce and staffing” domain comprised of colleague experts from Canada, USA, Switzerland, Sweden, and the United Kingdom. The subgroup held regular quarterly meetings throughout 2019 and 2020 to discuss the staff training literature review search parameters which informed the extraction form described above. Subsequently, the results of the review were discussed with the intention of determining the basic data elements related to staff training that would be applicable and inclusive of low- and middle-income countries (LMICs).

Results

The search generated 356 manuscripts for screening, and after duplicates were removed, 282 manuscripts were left to be screened. Only 47 met the eligibility criteria for full-text screening based on the abstracts. During the full screen, articles that were not relevant were




Figure 1. PRISMA flow diagram for article selection.

removed (Figure 1 shows the PRISMA flow diagram). Twenty-nine studies were included in the scoping review. Table 1 outlines the study characteristics such as study design, number of sites, analysis methods, and effectiveness of training on outcomes. All the included studies were from high-income countries. Most studies occurred in a single LTC home, or in multiple LTC homes in the same geographical region. The total sample size across all 29 studies included 4,933 staff participants (range = 4–1,076). Nursing Assistants (NAs) were the most common type of learners in 20 of the studies (68.97%) followed by Registered Nurses (RNs) who were involved in 13 of the studies (44.83%).

Characteristics of the Staff Training

Table 2 presents the training content, components, formats, duration, funding source, evaluation measures, and instructor. The most commonly taught topic involved dementia care ($n=16$) including topics such as caring for residents with dementia through QoL and person-centered care (PCC) training (Ballard et al., 2018; Kemeny et al., 2006), managing apathy and depression in residents with dementia (Leone et al., 2013; Rosen et al., 2002), communication and behavior management with residents with dementia (Burgio et al., 2001, 2002; Magai et al., 2002), nutrition in dementia residents (Chang et al., 2006), restraint use on dementia residents (Huizing et al., 2006), self-efficacy training to prevent staff burnout related to caring for residents with dementia (Mackenzie & Peragine, 2003), pain practices and beliefs in dementia care (Ghandehari et al., 2013) and engaging in advanced care planning (ACP) related conversations with dementia residents (Wils et al., 2017). Two studies each provided training about oral

care (Janssens et al., 2016; Williams et al., 2016), nutrition (Crogan & Evans, 2001; Greene et al., 2018), and depression and mental health (McAiney et al., 2007; Smith et al., 2013). Remaining training topics were: communication styles with residents (Soderlund et al., 2014), ergonomics (Peterson et al., 2004), pain management (Jones et al., 2004), staff hand hygiene (Huang & Wu, 2008), PCC (Bökberg et al., 2019), restorative care (Resnick et al., 2009), and behavioral modification (Arco & Du Toit, 2006).

Training program duration ranged from a single 2-hour session to a full year's intervention and follow-up. The most common format was didactic in-person educational sessions in the LTC home. The most common type of trainer were the authors of the studies, with a majority holding clinical psychology titles and/or nurse researchers.

Measurement Tools Used in Staff Training Studies

Table 2 presents all the measurement tools included in the review. Overall, studies used validated, non-validated, or a combination, to measure the effectiveness of training with a lack of consistency of measurement tools between the studies. For example, each of the 16 studies that provided dementia training used a unique combination of tools to measure resident outcomes, and while the majority of studies used validated scales including dementia severity (e.g., Clinical Dementia Rating [Morris, 1997]), behaviors (e.g., Cohen-Mansfield Agitation Inventory [Cohen-Mansfield, 1997], Neuropsychiatric Inventory–Nursing Home (Wood et al., 2000), and depression (Cornell Scale for Depression in Dementia [Alexopoulos et al., 1988]), a small number of studies used self-developed tools to measure staff knowledge and attitudes related to dementia staff training (Chang et al., 2006; Kemeny et al., 2006; Rosen et al., 2002). The heterogeneity of tools is observed across the training topics listed in Table 2.

Effectiveness of Training on Staff and Resident Outcomes

Twenty-eight of the 29 included studies reported staff outcomes (see Table 1). Twenty-five had favorable staff outcomes and three showed no statistical difference (Bökberg et al., 2019; Crogan & Evans, 2001; Huizing et al., 2006). Staff knowledge was the most frequently measured staff outcome that showed statistically significant improvement after the training intervention ($n=15$), other studies reported statistically significant improvements in staff confidence and competence after the training intervention ($n=5$), staff attitudes after training ($n=3$), and staff communication skills after training ($n=3$).

Thirteen studies reported on a wide range of resident outcomes (listed in Table 1) related to training and each

Table I. Summary of Literature Review Findings.

Authors (country)	Type	NHs (#)	Learner	N (Female %)	Aim	Topic	Pre and post measurement	Analysis	Staff outcomes	Patient outcomes
Arce and Du Toit (2006) (Australia)	Mixed Methods; Multiple baseline	1	NAs	Staff: 4 (100) Resident: 1 (100)	To examine effects of on-the-job feedback after staff training and to verify prior findings that competence was maintained after on-the-job feedback was ceased	Behavior modification	Observations Mon-Fri for 26 weeks. Baseline: observations up to 90 minutes. Over course of study: decreased to <60 minutes	Sum of resident behavior and consequences. Social validation questionnaire	Increased ability to cope with resident behaviors; increased competency	Increase in QoL ($p = .0042$), improvement in agitation ($p = .0076$), and neuropsychiatric symptoms ($p < .001$) in residents with moderately severe dementia. No reduction in antipsychotic usage, global deterioration, unmet needs, pain, or mood
Ballard et al. (2018) (U.K.)	RCT	69	Care staff	Residents: 553 (70.7)	To evaluate the efficacy of a PCC and psychosocial intervention on QoL, agitation, and antipsychotic use in people with dementia living in NHs and to determine its cost	Resident QoL and behavior	Resident QoL and Baseline and at 9 months	ANCOVA	Increase in positive care interactions ($p = .003$)	Increase in QoL ($p = .0042$), improvement in agitation ($p = .0076$), and neuropsychiatric symptoms ($p < .001$) in residents with moderately severe dementia. No reduction in antipsychotic usage, global deterioration, unmet needs, pain, or mood
Böckberg et al. (2019) (Sweden)	Pre- and post-test experimental design	20	Assistant nurse, RNs, OTs, PTs, Frontline managers	Staff: 365 (94.79)	To assess whether an educational intervention can effect staff perception of providing PCC for palliative persons in NHs	PCC	T=0 and 3 months after intervention (9 months from T=0)	Within-group comparisons, Wilcoxon signed rank test, subgroup analyses within the intervention group. Pearson's χ^2 test or Fisher's exact test. Mann-Whitney U-test, Kruskal-Wallis test, and a univariate logistic regression analysis	No improvement on any of the subscales and measures	—
Burgio et al. (2001) (U.S.A)	Mixed methods experimental design	5	NAs	Staff: 64 (84.38) Residents: 67 (74.63)	To examine communication skills training and use of memory books in improving communication between NAs and residents	Communication	Baseline, after intervention	ANOVA; ANCOVAs	Increased knowledge ($p < .05$); improved communication skills ($p < .05$), increased amount of positive statements to resident ($p < .05$); increased verbal interaction with residents ($p = .02$); no change in amount of time spent with resident; maintenance at 2 months f/u	Increased positive interactions ($p = .01$); increased independence in self-care ($p = .04$); no change in coherent verbal interactions
Burgio et al. (2002) (U.S.A)	RCT	2	NAs	Staff: 85 (9.56) Residents: 79 (6.00)	To evaluate a behavior management skills training program for improving NA behavioral skill performance and any resulting effects on residents' behaviors	Behavior management	Baseline, immediately after and 3 and 6 months after intervention	Baseline, immediately Cronbach's α , ANOVA, ANCOVA	Decreased knowledge ($p < .001$); decreased ineffective behavioral management techniques ($p < .05$), not maintained at f/u; improvement in six out of seven measured communication skills ($p < .05$), maintained at f/u ($p < .05$)	Decrease in agitation ($p < .05$) maintained at f/u
Chang et al. (2006) (Taiwan)	Quasi-experimental	2	NAs	Staff: 67 (97.01) Residents: 36 (N/A)	To observe the feeding behaviors of NAs after implementation of a feeding skills training program	Feeding skills	Immediately before and after training program, and 4 weeks later	Cronbach's α , ANOVA, ANCOVA	Increased knowledge ($p < .001$). No change in attitude or perceived behavior control. Increased intention frequency ($p < .05$) although no change in intention belief	—
Crogan and Evans (2001) (U.S.A)	Pre- and post-test experimental design	1	NAs	Staff: 20 (N/A)	To measure NAs' knowledge of nutritional care	Nutritional care	Before and after training	Performance observation	No statistical difference between pre and posttest scores. No improvement in essential principles of care. Improvement in 10 problematic areas while problems persisted in all other areas.	—
Dassel et al. (2020) (U.S.A)	Pre- and post-test experimental design	20	RNs, NAs, OTs, informal caregivers, students, admin, faculty	Staff: 94 (8.19)	To improve the care of residents with ADRD through community-based education for interprofessional team members	Dementia	Before and after training	Paired t-tests, Cronbach's alpha	Increased knowledge ($p < .01$); satisfaction with relevance and applicability of training to practice	—

(continued)

Table 1. (continued)

Authors (country)	Type	NHs (#)	Learner	N (Female %)	Aim	Topic	Pre and post measurement	Analysis	Staff outcomes	Patient outcomes
Ghandehari et al. (2013) (Canada)	Mixed methods experimental	Two large health regions	Nurses, special care aides	Staff: 13! (N/A)	To examine a pain assessment/management PE program aimed at improving staff beliefs, attitudes, and overall knowledge	Pain assessment and management	2 weeks before PE, after three educational sessions and 2 weeks after completion of PE	χ^2 tests, ANOVA, Tukey's honestly significant difference (HSD) tests, QSR N6ivo, thematic analysis comprised the framework for examining the data	Increased knowledge ($p < .001$) and improved pain beliefs ($p < .009$)	—
Greens et al. (2018) (U.K.)	Pre- and post-test experimental design	2	HCA's, RNs, catering or domestic staff, activity co-ordinators, PTs, OTs, and management	Staff: 16! (N/A)	To design, deliver and evaluate a hydration training session for care home staff that developed their knowledge and skills	Hydration	Before and after training	Wilcoxon signed-rank test of evaluation form. Qualitative data (field notes)	Increased knowledge re: dehydration ($p = .009$), —	Reduced infection rate ($p < .001$)
Huang and Wu (2008) (Taiwan)	Pre- and post-test experimental	3	NAs	Staff: 40 (100)	To test a hand-hygiene intervention for NAs in LTC on outcomes for NAs (knowledge, behavior) and residents (infection rate)	Hand hygiene	Before (pre-test), 1 month after (post-test I), and 3 months after (post-test II)	Descriptive statistics (means, SDs, frequencies, and percentages), paired t-test, McNemar test, χ^2 -test, and logistic regression test	NAs with more years of education were more likely to improve their knowledge. Increased rates of hand hygiene ($p < .001$).	Reduced infection rate ($p < .001$)
Huijzing et al. (2006) (Netherlands)	Cluster RCT	1	Nurses	Residents: 144 (71.33)	To investigate an educational intervention and its effect on the use of physical restraints in psycho-geriatric NH residents	Physical restraints	Baseline and 1 month post-intervention	Frequency tables, means, χ^2 test, t-test, Fisher's exact test, McNemar tests, and gain scores, logistic regression analysis	No change in use, intensity, number or types of restraints used. No change in time of day when restraint used.	Decreased risk of restraint use. Increased depression. Decreased cognitive status
Janssens et al. (2016) (Belgium)	Cluster RCT	12	Nurses and NAs	Staff: 259 (95)	To explore the impact of an oral healthcare protocol, in addition to education, on nurses' and NAs' oral health-related knowledge and attitude	Oral health and hygiene	Baseline and 6 months after the start of the study	Bivariate analyses: nonparametric tests, Mann-Whitney U-test, Wilcoxon matched-pairs signed-rank test, linear mixed-model analyzes, post hoc power calculation, ANOVAs, GLMs, GEEs, χ^2 tests, observations	Increased knowledge ($p < .0001$), no change in attitudes; nurses demonstrated more knowledge although less favorable attitude versus NAs. Overall better attitude on psychogeriatric ward versus mixed ward	—
Jones et al. (2004) (U.S.A.)	Mixed methods experimental	12	25% RN, 26% LPN, and 49% PSW	Staff: 628 (N/A) Residents: 1,899 (N/A)	To develop a culturally competent intervention to improve NH pain practices, improve staff knowledge and attitudes about pain, improve pain practices in NHs and improve NH pain policies and procedures	Pain	Staff: before and after training Residents: quarterly (before, during and after)	Subtle increase in knowledge and attitudes. Decline in perceived barriers	Decreased reports of constant pain ($p < .001$). Improved pain reassessments ($p < .05$) No difference in overall pain reporting and acute moderate/severe pain.	
Kemeny et al. (2006) (U.S.A.)	Multiple baseline	1	NAs and nurses	Staff: 77 (92)	To describe experiential techniques used by Project RELATE in training for PCC, and NAs' and nurses' response	PCC	After each session and 2 months after	Likert scale; percent correct; effect size, d-scores	Favorable reactions to implement training in practice ($p < .05$); NAs reported more favorable reactions versus nurses. Increased knowledge (more so in nurses vs. NAs) ($p < .05$). Retention of knowledge 2 months later ($p < .05$); Increased confidence	

(continued)

Table 1. (continued)

Authors (country)	Type	NHs (#)	Learner	N (Female %)	Aim	Topic	Pre and post measurement	Analysis	Staff outcomes	Patient outcomes
Leone et al. (2013) (France)	Mixed methods experimental	16	Psychologists, physicians, practical nurses, agent of hospital service	Staff: 563 (N/A) Residents: 230 (79.5)	To evaluate the effectiveness of staff education for the management of apathy in older adults with dementia	Apathy	Baseline, at the end of the training program (week 4) and 3 months later (week 17)	Quantitative evaluation; change in ADL scores, NPI-NH, Katz ADL Scale and on the two observation scales, mean comparisons using t-test, χ^2 test; multiple linear regression analysis	Improved knowledge although not significant at 3 months f/u, INT group had increased knowledge ($p < .01$). Increased self-efficacy accomplishment ($p < .05$) not maintained at f/u	Improvements in emotional blunting ($p < .01$). More self-sufficient in "dressing" and "transferring" ($p < .05$) on Katz ADL Scores. Increase in affective and psychotic symptoms ($p < .01$). No change in number of drugs prescribed
Mackenzie and Peragine (2003) (Canada)	Quasi-experimental	1	RNs, RPNs, HCs, privately paid caregiver	Staff: 41 (92.68)	To describe the development and outcome of a stress and burnout relieving intervention by enhancing self-efficacy in managing challenging teams, residents, and family situations. Secondary purpose is to present a self-efficacy inventory to measure the effectiveness of the intervention	Managing stress and self-efficacy	Before, immediately after, and 3 months after	t-test and χ^2 analyzes; ANCOVAs	Similar knowledge between groups at post-test; at 3 months f/u, INT group had increased knowledge ($p < .01$). Increased self-efficacy accomplishment ($p < .05$) not maintained at f/u	—
Magai et al. (2002) (U.S.A.)	RCT	3	Nursing staff	Staff: 20 (100) Residents: 91 (93)	To assess a nonverbal sensitivity training program on the care provided to dementia patients and on staff caregiver well-being	Nonverbal communication	Baseline and 4 × 3 week intervals	Repeated measures ANOVAs, Wilks's lambda	Improved affective state; Improved BSI scores ($p < .01$)	Increased positive affect ($p < .05$), which converged with control groups at f/u
McAiney et al. (2007) (Canada)	Pre-and post-test experimental	439	RNs, RPNs, SVs, and other health disciplines	Staff: 1,076 (N/A)	To describe an education program for the management of mental health problems in LTC and the evaluation of its impact and sustainability	Mental health	Before start of program, and 6 weeks after	Frequencies, percentages, ranges, means, standard deviations, paired t-tests	Improved confidence ($p < .001$), understanding and — assessment of mental health problems ($p < .01$). Increased confidence and use of assessment tools ($p < .01$)	—
Peterson et al. (2002) (U.S.A.)	Pre-and post-test experimental	6	NAs, LPNs, RNs, SVs admins, music therapist, Porter	Staff: 72 (93.1)	To evaluate the effect of an educational course on dementia on staff knowledge, stress, and self-esteem	Dementia	Baseline, immediately after and 6 to 8 weeks after training	Pearson correlation, General Linear Model, Cramer's V	Increased knowledge in all groups, although only significant in those with prior training ($p = .004$; no significant change in stress or self-esteem scores	—
Peterson et al. (2004) (U.S.A.)	Pre-and post-test experimental	1	NAs, RNs, LPNs	Staff: 35 (N/A)	To develop an ergonomics training program for selected NAs at a state-run veterans' home to decrease musculoskeletal disorders	Ergonomics	3 months before training, at the end of the training program and 1 month after training	Two-paired student's t-test; ANOVA	Increased knowledge ($p < .001$). No change in level of stress for risk factors, pain or discomfort, or perceived general health	—
Resnick et al. (2009) (U.S.A.)	RCT	12	NAs	Staff: 523 (93)	To describe the implementation process of the educational component of the restorative care intervention, the outcomes and the effect on NA knowledge	Restorative care	Baseline and immediately after intervention	($SD = 2.7$, $F = 280.4$, $p < .05$)	Increased knowledge re: restorative care ($p < .05$)	—
Rosen et al. (2002) (U.S.A.)	RCT	3	RNs, LPNs and NAs	Staff: 279 (N/A)	To design and assess a curriculum of staff training on depression and dementia	Depression/ dementia	Before each training session and after final session	χ^2 test; Kolmogorov-Smirnov (K-S) test; Mann-Whitney (M-W) test; Wilcoxon rank sum test, ANOVA, posthoc pair-wise comparisons	Increased knowledge in all sites but significant in computer site ($p < .005$)	—

(continued)

Table I. (continued)

Authors (country)	Type	NHs (#)	Learner	N (Female %)	Aim	Topic	Pre and post measurement	Analysis	Patient Outcomes	
Scerri and Scerri Pre- and post-test experimental (2019) (Malta)	–	Nursing offices, deputy charge nurses, staff nurses, and enrolled nurses	Staff: 214 (68.2)	To investigate a dementia training program on nursing staff working in public nursing/residential homes on their knowledge, attitudes, and confidence	Dementia	Beginning of the first session and the end of the last session	Shapiro-Wilk test; independent sample t-test, ANOVA, post-hoc Tukey's test, Pearson correlation test, Pearson Chi-square, Cohen's d, stepwise regression analysis	Increased knowledge ($p < .001$), Improved staff attitude ($p = .001$), Increased confidence ($p = .017$). Training found to have a cumulative effect	–	
Smith et al. (2013) (U.S.A)	Mixed methods experimental	13	RNs and LPNs	Staff: 24 (N/A) Residents: 50 (76)	To describe a CD-based depression training program and its use and feasibility of nurses using it with older adults in their care, and to evaluate training-related outcomes among those residents	Depression	Baseline and at 8, 12, and 16 weeks	t-tests and χ^2 analyzes, linear mixed modeling, non-parametric Kruskal-Wallis test	No difference between groups in method of training, Improved knowledge, care and outcomes	Improved depression scores from baseline to flu ($p < .001$). Decreased pain in all groups ($p = .006$). flu tests suggests that only usual care group pain improved significantly ($p < .01$). No improvement in QoL or anxiety scores
Söderlund et al. (2014) (Sweden)	Mixed methods experimental	1	RNs, LPNs, NAs	Staff: 12 (100)	To explore nurses' experiences of attending a YM training program and to describe ratings of the work climate among the entire nursing staff	Communication style	Before and after intervention	Descriptive statistics	Difficulty changing communication style. Increased self-reflection. Increased positive interactions. Increased confidence. Improved work environment	–
Williams et al. (2016) (U.S.A)	Pre- and post-test experimental	2	CNAs, LPNs, RNs	Staff: 26 (83.4) Residents: 176 (N/A)	To facilitate the implementation of oral health protocols in NHs	Oral health	3 months before and 3 months after intervention.	Likert survey, Cronbach's α test, Wilcoxon signed rank test, McNemar's test	Increased feelings of responsibility on resident to make referral ($p = .02$), Increased confidence in performing oral assessments ($p = .009$) and identifying oral conditions that need referrals ($p = .03$)	Increased dental referrals ($p = .0018$)
Wils et al. (2017) (Belgium)	Mixed methods experimental	1	Nursing staff	Staff: 13 (84.62) Residents: 124 (72.58)	To assess the effect of an education program on the registration of care goals in a NH with dementia residents and to explore the views of staff on advance care planning	ACP	Baseline and 12 months	ANOVA, F-statistic	Increased communication regarding ACP with resident and appointed representative ($p < .02$). Increased care goal planning ($p = .05$)	Increased WIB values ($p < .001$). Increased social interactions in DCM ($p = .041$)
Yasuda and Sakakibara (2017) (Japan)	One group repeated measures	1	Care staff	Staff: 40 (N/A) Residents: 40 (77.5)	To determine how educational intervention for care staff can help to improve the status of residents with dementia	Dementia	Baseline, 1 month later and after intervention	Wilcoxon signed-rank test, DCM – data processing, ME value	–	–

Note. Abbreviations used based on order of appearance: NH = nursing home; NA = nursing assistant; RCT = randomized control trial; PCC = person centered care; QoL = quality of life; RN = registered nurse; OT = occupational therapist; PT = physical therapist; ADRD = Alzheimer's disease and related dementias; PE = pain education; HCA = healthcare aide; LTC = long term care; LPN = licensed practical nurse; PSW = personal support worker; AIC = Apathy Inventory–Clinician version; NPI-NH = neuropsychiatric inventory–nursing home version; ADL = activities of daily living; SW = social worker; CTN = control group; INT = intervention group; ACP = advanced care planning; DCM = dementia care mapping; ME value = mood and engagement value; WIB = well-being and ill-being.

Table 2. Summary of Training

Author	Format of training	Components of training	Quantitative measure	Psychometric properties	Duration of training (hour)	Content	Funder	Instructor
Arcu and Du Toit (2006)	Workshops	Training, observation, feedback	Social validation questionnaire, observational coding	Interobserver agreement during observational coding	2 hours/week for 3 weeks	Changes associated with aging and how institutional living and relations with staff affect behavior. Behavioral management principles were described. Behavior support plan	Australian Commonwealth Government	First author (clinical psychologist) and second author (postgraduate student)
Ballard et al. (2018)	Interview, lecture, experiential learning, and application in NH	Training, medication review, cost analysis	CDR, FAST, DEMQOL-Proxy, CANI, NPI-NH, CSDD, CSRI, Quality of Interactions Scale, Abey Pain Scale	Reliable and valid as per previous studies (refer to references for each instrument)	2 days or four half-days for 1 month. After, 6 hours/month for 4 months. Finally, 8 hours/month for 4 months	Person-centered activities and social interactions. Review of antipsychotic medications	National Institute of Health Research	Research therapist. Two lead care staff members (VHFIELD champions)
Böökberg et al. (2019)	Educational seminars	Training only	P-CAT, PCQ-S	This version of the P-CAT and PCQ-S is reported to be valid, reliable, and applicable for continued use	Five 2-hour seminars over 6 months	Key principles of palliative care and clinical practice guidelines (both based on the WHO definition of palliative care)	Swedish Research Council; the Vardal Foundation; the Gyllenhjälterna Kräpplup's Foundation; Medical Faculty, Lund University; the City of Lund; Faculty of Health and Life Sciences, Linnaeus University; the Greta and John Kock Foundation, and the Ribbingska Memorial Foundation	Palliative and geriatric RNs and researchers
Burgio et al. (2001)	Didactic training, role play, case studies, group discussions	Training, chart review, environment assessment	CNA Communication Skills Checklist-CABOS, MMSE, (FIM)—REACH Version	Validated tools supported by previous research	3 hours (additional hour for supervisors) over 1 week, followed by hands-on training for 4 weeks	Memory books and general communication skills; staff motivational system	The National Institute on Aging Foundation	Project manager (licensed clinical psychologist)
Burgio et al. (2002)	In-service classes, hands-on training, case studies, videos, group discussion, workbooks	Training, medical records review, family interview	MMSE, The Barthel Self-Care Rating Scale, CDR, CNAI, the Behavior Management Skills Checklist, computer-assisted behavioral observation systems	Validated tools supported by previous research	5 hours over three consecutive days followed by hands-on training for 2 weeks	Factors in environment that can affect resident behavior, communication skills and behavior management techniques	The National Institute of Nursing Research	Geropsychologist from the research staff
Chang et al. (2006)	In-service classes and hands-on training	Training only	Formal Caregivers' Knowledge of Feeding Dementia Patients Questionnaire, the Formal Caregivers' Attitude toward Feeding Dementia Patients Questionnaire, the Perceived Behavior Control Scale, the Intention Scale, the Formal Caregivers' Behaviors in Feeding Dementia Patients Observation Checklist	Comprehensive literature review, clinical experience, and observations were used to create the questionnaires. Reviewed by gerontological nursing expert.	4 hours over two consecutive days	Overview of dementia, common eating behaviors and protocol for managing feeding problems associated with dementia patients.	—	The PI (RN, PhD)
Crogan and Evans (2001)	Discussion, audiovisual presentations, experiential exercises, and role-playing	Training only	Multiple choice pre/post-test	—	—	Cues to feeding procedures and nutritional aspects of caring for resident; teaching plan, student note-taking guide, handouts, and a pocket guide	A skilled gerontological nurse	—
Dassel et al. (2020)	Audio-visual recorded presentations, case study, and supplemental information	Training only	Modified ADKS	Approx 3 hours online module	Overview of dementia, understanding behaviors and approach, effective communication	Health Resources and Services Administration	Faculty members and clinicians at the University of Utah	(continued)

Table 2. (continued)

Author	Format of training	Components of training	Quantitative measure	Psychometric properties	Duration of training (hour)	Content	Funder	Instructor
Ghandehari et al. (2013)	Focus group sessions; seminars with discussion, participation and critical thinking	Training; focus groups	PKBQ; Modified Pain Beliefs Questionnaire; session content knowledge test	The PKBQ has been previously found to be valid and internally consistent (Cronbach's alpha = 0.78). The Modified PBO has been previously found to have criterion validity, concurrent validity, and satisfactory reliability.	3 hours/weeks for 3 weeks (total 9 hours)	Assessing and managing pain in LTC based on empirical evidence (pharmacological and nonpharmacological); nutrition; physical functioning/physical activity; individual-centered care	Saskatchewan Health Research Foundation; Canadian Institutes of Health Research	Experts in pain management
Greene et al. (2018)	Emotional mapping, hydration quiz, case studies, hands-on-activity	Training; observations	Self-developed questionnaires	—	2 hours	Fluid preferences, s/s of dehydration + treatment, hydration principles	The National Institute for Health Research	Two staff from project team
Huang and Wu (2008)	In-service classes + hands-on training, motivating and giving feedback to NAs, engineering controls and placing reminders in the workplace	Training; infection rates	Hand-Hygiene Questionnaire, Behavior toward the Hand-Hygiene Observation Checklist	Comprehensive literature review, clinical experience and observations used to develop the instruments. Cronbach's alpha for Hand Hygiene Questionnaire= 0.76. 92% inter-rater reliability reached for the checklist. Cronbach's alpha for the checklist= 0.85	—	Purpose of intervention, overview of infection in NH residents, etiology and importance of hand hygiene, and timing and protocol for hand hygiene.	Chung Gung Medical Research Foundation	The two authors affiliated with Chang Gung University, and Zhong-Xian Hospital in Taiwan
Huijzing et al. (2006)	Small-scale meetings with an active learning environment and consultation with a nurse specialist	Training; restraint use observation	MDS CPS; MDS ADL Self-Performance Hierarchy, the DRS, the Social Engagement Scale, a mobility scale, accident registration form	CPS scale corresponds with the MMSE and the Test for Severe Impairment. The internal consistency of the mobility scale was high (Cronbach's α = 0.97). The reliability and validity of the scales were found to be sufficient from other studies found to be sufficient from other studies	Five meetings of 2 hours, followed by a 1.5 hour session, over 2 months	Philosophy of restraint-free care and techniques of individualized care; decision-making process towards restraint use, effects and consequences of restraint use; strategies to analyze risk behavior of residents and alternatives for restraints; discussion of real-life cases	MeanderGroup Zuid-Limburg, the Provincial Council for the Public Health (Limburg) and Maastricht University	Nurse specialized in restraints
Janssens et al. (2016)	Three educational stages; oral presentation/lecture; practical education; oral healthcare record keeping; supervised implementation; train-the-trainer concept	Training only	Oral healthcare questionnaire	Content and construct validity reviewed by experts in the field of gerontology.	Initial 1.5 hour presentation, followed by 2 hours lecture + 1 hour practical component, and finally a 1.5 hour theoretical + practical session	Theoretical and practical essentials of the guideline; summary of the guideline and all oral hygiene actions, such as tooth brushing	—	Project supervisor, at least two Ward Oral healthcare Organizers (nurses or NAs) per ward, a coordinating physician and optionally an occupational and/or speech therapist; the second author was also involved in supervised implementation; fourth author (dental hygienist) provided support
Jones et al. (2004)	Interactive educational sessions; case studies; videos	Training; pain assessments, medical record review	Staff pain surveys, modified Quick Pain Assessment	Internal consistency reliabilities and Cronbach's α reliability was adequate for the surveys	4 × 30 minutes sessions over 6 months	Pain problem and assessment; pharmacologic management; communication issues; pain case studies	Agency for Healthcare Research and Quality to the School of Nursing, University of Colorado Health Sciences Center	The first author
Kemeny et al. (2006)	Didactic sessions; coaching sessions; role play; simulations; debriefing	Training only	General reaction question scales, training evaluation	Face validity was reviewed by the researchers. Cronbach's alphas ranged from 0.89 to 0.97 for each instrument.	—	Dementia; PCC; communication skills	Budgett Butterworth Health Care Foundation	—

(continued)

Table 2. (continued)

Author	Format of training	Components of training	Quantitative measure	Psychometric properties	Duration of training (hour)	Content	Funder	Instructor
Leone et al. (2013)	Didactic session; interactive, hands-on teaching	Training only	Katz ADL Scale, NPI-NH, the Apathy Inventory—Clinician Observation Scale, an Individual Observation Scale	Katz ADL Scale: valid and predictive Norwegian version of NPI-NH found to be reliable and valid	Initially 2 hours, followed by 4 hours/weeks for a month	AD and BPSD; apathy and depression s/s and practical advice and methods to counteract; techniques for dealing with deficits in ADLs; nonpharmacological interventions	Fondation de Coopération Scientifique and the ARMEP association	Two psychologists
Mackenzie and Peragine (2003)	Didactic information and discussion; experiential role-play	Training only	The Inventory of Geriatric Nursing self-Efficacy, knowledge questionnaire, 22-item NB28, the Organizational Job Satisfaction Scale	Self-efficacy scale: Cronbach's alpha=0.96, average item-total correlation =0.83. Knowledge questionnaire: Internal consistency = 0.78.	4 × 2 hours modules (one session/week for 1 month)	Teamwork module; challenging behavior module; family module; review module	Morris Sivka Fellowship	—
Maggi et al. (2002)	Didactic/experiential sessions; videos, hands-on training	Training, caregiver interviews, behavior and facial expression assessment	BEHAVE-AD, CMAI, CDS, BSI, the Adult Developmental Interview	Previous literature supports the reliability and validity of the MBI as the gold standard for measuring The MBI is the gold standard for measuring burnout Internal consistency reliabilities for all measurement scales is high (alpha > 0.69)	10 × 1-hour sessions over 2 weeks	Issues of nonverbal communication and emotion expression: cognitive and behavioral aspects of dementia	Clinical psychologist	—
McAiney et al. (2007)	Peer mentoring and coaching; homework assignments; case-based; small group work; minimum lecture time	Training, long-term sustainability review	Pre/post-test questionnaires	Face and content validity of the measures ensured through reflecting material from the education program as well as receiving input from clinical experts. Cronbach's $\alpha=0.74$	18 hours over 3 days followed by 12 hours over 2 days	OA Physical and cognitive/mental health problems and behaviors; ADRD	MOHLTC	Clinicians and group facilitators
Peterson et al. (2002)	Didactic session, videos, role play, sensitization exercises, interactive discussion	Training only	Dementia Quiz, FCSI, Ownership subscale of the Reciprocal Empowerment Scale	Reliability of tools were high (alpha > 0.79) and face, concurrent, and predictive validity acceptable	6 hours class	Physiology of dementia, coping with challenging behaviors, performing ADLs	Emmett J. and Mary Martha Doer C. Center for Social Justice Education and Research in the School of Social Service at Saint Louis University	Professional educators with CNA or RN experience
Peterson et al. (2004)	Didactic session; floor supervision	Training, work environment review	Pre/post-test questionnaires	—	—	Top 20 perceived risk factors; correct ergonomic work practices; administrative strategies; use of engineering controls.	—	Research assistant
Resnick et al. (2009)	Educational sessions, group discussion, individual instruction, role play, case studies	Training only	The Theoretical Testing of Restorative Care Nursing	Evidence of test-retest reliability and validity based on prior use (Resnick et al., 2009)	30 minutes/weeks for 6 weeks	Intro to restorative care, resident motivation, specific skills associated with restorative care, documenting restorative care activities, overcoming challenges to implementation	Agency for Healthcare Research and Quality Grant	Advanced practice nurse
Rosen et al. (2002)	Computer-based interactive video training; lecture; individual, self-paced training	Training only	Satisfaction/relevance questionnaire, pre/post-training test	—	12 modules of 35 to 45 minutes (CS); 30 to 45 minutes/month for 6 months (LS)	Mental health in aging, depression, and dementia; changes associated with aging; behavior + nonpharmacological interventions; AD (and other dementias); behavior management: fundamentals of agitation and aggression	National Institute on Aging and National Institute of Mental Health	Computer (CS); advanced degree nurse (LS)
Scerri and Scerri (2019)	Didactic session; discussions	Training only	ADKS, DAS, Confidence in Dementia Scale	ADKS: has adequate reliability and content, predictive, concurrent, and convergent validity. DAS: adequate reliability and convergent validity compared to similar instruments	7 × 2 hours sessions (14 hours total)	Into to dementia care and services; activities for dementia OA; dementia-friendly design and assistive technologies; policy and development	European Social Fund	Local experts in dementia and old age mental health

(continued)

Table 2. (continued)

Author	Format of training	Components of training	Quantitative measure	Psychometric properties	Duration of training (hour)	Content	Funder	Instructor
Smith et al. (2013)	CD-based training; psychiatric nurse enhanced; digitized presentation; handouts; work place exercise; case-based learning	Training, feasibility review, chart review	Self-report measure, DTPE, PHQ-9, GAD-7, IPT	Established psychometric properties; tools commonly used in geriatric research	4-part training completed in 4 to 6 weeks	Depression and comorbid conditions; standardized rating scales; interventions; communication and teamwork	Wellmark Foundation of Iowa; Iowa Geriatric Education Center; American Psychiatric Nurses Foundation; Hartford Center of Geriatric Nursing Excellence	Computer (CD); psychiatric nurse (PDE)
Söderlund et al. (2014)	Group of 10 to 12 nurses; didactic session; practical training; videotape; written reflections; written test	Training, feedback	Creative climate questionnaire	Previously found to have good validity, reliability and internal consistency (Cronbach's Alpha = 0.709)	1-year program of 10 days of theoretical training/month; practical training 2-3×/week	Confirmatory; empathetic approach; verbal and non-verbal VM techniques for communication	Ersta Skondal University College; Swedish Order of St. John; Alzheimer Foundation; Dementia Association	External certified supervisor
Williams et al. (2016)	Chart review; didactic session; decision tree Didactic session; role play; workplace application; debriefing; videos	Training, chart review	Pre and post-training survey	Acceptable internal consistency for survey items (Cronbach's Alpha = 0.709)	—	Oral assessment technique; common oral conditions and abnormalities	PI (RDH, MSDH)	—
Wils et al. (2017)	Didactic sessions; group discussions	Training, debriefing, patient file chart review	Recording of ACP in the patient file	—	6×2 hours	Legal and ethical issues; communication skills	Law expert; one of the researchers	—
Yasuda and Sakakibara (2017)	Didactic sessions; group discussions	Training only	MMSE, Barthel Index, DCM	MMSE Barthel Index; commonly used tool DCM: reliability and content reported by Brooker and Sure (2006)	3 × 60-90 minutes	Dementia basics; PCC; communication and interactions; behaviors	—	—

Note. Abbreviations used based on order of appearance: CDR = clinical dementia rating; FAST = functional assessment staging tool; CMAI = Cohen-Mansfield agitation inventory; CSDD/CDS = cornell scale for depression in dementia; CANE = camberwell assessment of need for the elderly; CSRI = client service receipt inventory; QUS = quality of interaction scale; NIHRS = national institute of health research scale; WHELD = well-being and health for people with dementia; P-CAT = person centered care assessment tool; PCQ-S = person centered climate questionnaire; WHO = world health organization; CNA CSC = certified nursing assistant communications skills checklist; CABOS = computer-assisted behavioral observation system; MMSE = mini-mental state exam; FIM-REACH = functional independence measure-resources for enhancing Alzheimer's caregiver health version; NIA = national institute on aging; PI = principal investigator; ADKS = Alzheimer's disease knowledge scale; PKBQ = pain knowledge and beliefs questionnaire; PBQ = pain beliefs questionnaire; MDS = minimum data set; CPS = cognitive performance scale; DRS = depression rating scale; GOS = group observation scale; IOS = individual observation scale; BPSD = behavioral and psychological symptoms of dementia; AD = Alzheimer's disease; MBI = Maslach Burnout Inventory; BEHAVE-AD = behavioral pathology in Alzheimer's disease rating scale; BSI = brief symptom inventory; MOHLTC = ministry of health and long term care; FCSI = formal caregiver stress index; RA = research assistant; APN = advanced practice nurse; CS = computer site; LS = lecture site; DAS = dementia attitudes scale; OA = older adult; PHQ-9 = patient health questionnaire; 3MS = modified mini-mental state exam; PNE = psychiatric nurse enhanced; VM = validation method; RDH = registered dental hygienist; MSDH = masters of science in dental hygiene; NIR = national institute of nursing research; BMSC = behavior management skills checklist; SHRF = saskatchewan health research foundation; CIHR = Canadian institutes of health research; HRSA = health resources and services administration; SES = the social engagement scale; FCS = Fondation de Coopération Scientifique; RES = reciprocal empowerment scale; NIMH = national institute of mental health; NIA = national institute on aging; CODE = confidence in dementia scale; APNA = American psychiatric nurses foundation; IGEC = Iowa geriatric education center; NHCNE = National Hartford center of geriatric nursing excellence; CCQ = creative climate questionnaire; AHRO = agency for healthcare research and quality.

of these studies reported at least one positive resident outcome. In seven studies, the authors found that resident behavior was positively affected after implementing a training intervention: reduced responsive behaviors (Arco & Du Toit, 2006), increased positive affect (Magai et al., 2002), improved QoL ($p=.0042$), decreased agitation ($p=.0076$) and decreased overall neuropsychiatric symptoms ($p<.001$) (Ballard et al., 2018), and decreased agitation which was maintained at the 3- and 6-month follow-up ($p=.01$) (Burgio et al., 2002). Additionally, two studies showed that residents had increased independence in self-care activities ($p=.04$) (Burgio et al., 2001) and improved “dressing” and “transferring” scores on the Katz ADL instrument (Leone et al., 2013) compared to the control groups, after staff participated in a communications training intervention, and apathy training, respectively. Lastly, there were significant reductions in resident reports of constant pain ($p\leq .001$) and improvements in resident pain reassessments ($p<.05$) after staff were provided pain management training (Jones et al., 2004). In a separate study, decreased pain levels in residents were found ($p=.006$) after staff completed the depression training program (Smith et al., 2013).

Expert Panel Discussion

The literature review results were examined by the expert panel and the salient findings and gaps in knowledge identified by the panel were: (1) staff training was heterogeneous in almost all aspects including content, facilitation, duration, processes for monitoring effectiveness, and measurement tools; (2) the lack of any evidence of staff training in LMICs; (3) the absence of any international collaboration in training, and the local and regional scale of the training that was often limited to one home; (4) there was no macro-level measurement of staff training or indication of how staff training was documented that would allow aggregation of data between LTC homes; and (5) training appeared to be effective and positively influenced a broad range of resident and staff outcomes. Based on these findings, it was clear that there was no extant international measurement infrastructure and that the current literature was exclusive of LMICs; therefore, there was a need to create inclusive and accessible CDEs that could capture basic data internationally.

First, the expert panel acknowledged individual privileges and the discrepancies in resources and staffing as well as differences in system priorities between high, middle, and low-income countries. From this lens, the expert panel required that the proposed CDE candidates be feasible and applicable in a multitude of sociopolitical conditions in order to include data from LMICs. Further, we recognized that due to differences in languages, a CDE with nuanced or complex theoretical concepts may be lost in translation (e.g., self-efficacy as a primarily Western notion may not be easily

translated or understood in other cultures) or may not be considered relevant. The proposed CDEs needed to be single-items that were simple, clear, and have face validity in a wide variety of social contexts in order to have buy-in from understaffed or under-resourced settings.

Second, the expert panel discussed how staff training could be reflected in a small number of meaningful CDEs. For example, it was important that training be current in order to keep staff up-to-date with relevant practices. There was further expert consensus that fewer items were preferred to a higher number of CDEs to increase the probability of implementation in high and low-income countries given the complex characteristics of LTC settings. Third, the expert panel discussed the critical need to evaluate the staff training using a validated measure as the literature review suggested that training has positive staff and resident impacts, but several studies reported their own unstandardized instruments. The expert panel considered the quality of the measurement tool and included the aspect of using standardized measurement tools to ensure that the same information can be gathered, compared and aggregated about staff and residents. To this end, the expert panel determined that CDEs focused on the *presence* and the *measurement* of staff training will begin to fill a current international data gap. Lastly, the expert panel discussed that the permissible values should be binary “yes” or “no” rather than Likert scales or scales that use terms with room for interpretation (e.g., “sometimes,” “often”). The CDEs needed to be accessible, feasible, clear, and not resource-intensive to answer in order to increase their likelihood of implementation into LTC settings.

From the WE-THRIVE expert panel discussions described above, the members focused on the necessity of basic data elements, that is, the *presence* and *measurement* of staff training related to staff and resident outcomes in LTC as an initial step to conduct cross-country comparative research. Therefore, three candidate CDEs were suggested to lay the groundwork for this critical knowledge building. The suggested CDEs are: (1) “was there institutional training provided to staff in the last 12 months?” with a binary choice of “yes” or “no,” followed by (2) “was there evaluation of a staff-related outcome using a standardized measure?,” and (3) “was there evaluation of resident-related outcome using a standardized measure?” both with a “yes” or “no”. This critical knowledge will support LTC workforce needs and continue to contribute to evidence related to staff training, which may inform the development of policies for a sustainable and equitable LTC system which are needed (United Nations, Department of Economic and Social Affairs Population Division, 2019; WHO, 2017). It is important to note that these are proposed CDEs are an initial starting point, and subject to change according to the emerging literature and future ongoing consultations with the broader academic and LTC community.

Discussion

This work presents the process of developing CDEs related to staff training in LTC homes; a literature review about staff training was completed, and the results were presented and discussed by an international expert panel as part of WE-THRIVE's "Workforce and Staffing" subgroup. Best practices and a systematic method combining evidence in expert opinion were used to develop the candidate CDEs. The literature review results illuminated significant heterogeneity in staff training and use of measurement, a paucity of any international collaboration to evaluate staff training in LTC, as well as an absence of evidence about staff training from LMICs. The expert panel determined that a greater baseline understanding about whether training is provided and measured across different jurisdictions and countries is critical to understand inequities in professional development opportunities. A macro-level measurement infrastructure that can be applied to high- and low-income countries will begin to inform the development of LTC staff training benchmarks and quality standards aimed at ensuring research integrity, public accountability, and equitable allocation training.

Considerations from the expert panel about the quality of measurement tools used is consistent with previous staff training literature. Concerns with measurement have been longstanding in previous reviews of staff training programs in LTC (Aylward et al., 2003; Beeber et al., 2010; Kuske et al., 2007), and there is a consensus for a greater need for more rigorous evaluation of training with validated and standardized tools. Proposed CDEs that are explicit about the use of standardized tools may motivate more robust measurement and emulate the aspirational values of WE-THRIVE to build capacity and support cross-country research.

Significance and Next Steps

The WE-THRIVE initiative seeks to facilitate international research in order to support high quality of care for older adults and comparative research in LTC, such that findings between different researchers in diverse countries can be seamlessly pooled to create appropriate policies and interventions for LTC staff training (Corazzini et al., 2019). The findings of the current review can aid in the creation and development of appropriate LTC strategies and documentation related to staff training (Lepore & Corazzini, 2019). Accurately capturing such foundational data can benefit policymakers to create guidelines and benchmarks about staff training, and also help inform LTC home administrators about resource allocation. For example, if the proposed CDE data about staff training was collected and shared between regions, it could promote a greater understanding about the degree of capacity building and effects of staff training for staff and residents.

The iterative process and stakeholder engagement are critical elements to the successful development and use

of CDEs according to Redeker et al. (2015). The input of stakeholders in the development and use of CDEs promotes harmonization nationally and internationally (Choquet et al., 2015; Redeker et al., 2015). We plan on engaging in dialogue and discussions with the researchers and LTC stakeholders to get feedback about the proposed CDEs and the feasibility of these CDEs. Future directions include efforts to advance the development of CDEs through continued international research as part of WE-THRIVE. Further investigations are required to examine if the proposed CDEs are able to be translated across cultural, geographical and institutional barriers, as well as how the proposed CDEs will improve data quality and opportunities for comparison between international researchers.

Strengths and Weaknesses

One strength of this review is in the robust approach of expert consultation based on the diverse group of researchers from the WE-THRIVE initiative. Another strength is that we used a systematic approach when conducting the literature review that included a search generated by an information specialist spanning the last 20 years, and independent reviewers who completed the full-text screen and extractions. The search was limited to English only titles which may have excluded training programs from non-English speaking parts of the world. Additionally, publication bias could be a limitation as training that was not found to positively influence outcomes may not have been published. Lastly, although there are dozens of countries represented in the WE-THRIVE group, not all countries are presented in the panel, specifically countries in the Global South and Africa. The expert panel was composed of experts from high-income English-speaking countries. Moving forward, WE-THRIVE has active online campaigns to strongly encourage researchers in LTC from LMIC and non-English speaking countries to participate with the WE-THRIVE initiative and the workforce domain to ensure the selected LTC CDEs are relevant and applicable to these contexts.

Conclusion

In summary, this paper presented a systematic process by which an international expert panel proposed three CDEs as part of the WE-THRIVE CDE initiative to advance international research related to staff training in LTC. Proposing appropriate CDEs can bolster comparative research and create the evidence-base for policy and LTC decision-making. The three candidate CDEs proposed are accessible, feasible, and intended to be ecologically viable in LMICs in order to fill current knowledge gaps to evaluate staff training. Researchers must continue to be mindful that observed differences in training across countries are likely influenced by social policies and context specific to each jurisdiction, thus continuous assessments are required. These proposed

CDEs are positioned as the starting point for staff training, and are subject to change based on new evidence, feedback from LTC stakeholders, and the cultural, political, and institutional changes that may occur.

Acknowledgements

The authors would like to acknowledge the guidance and support of Drs Kirsten Corazzini and Michael Lepore, co-chairs of the WE-THRIVE consortium.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iDs

- Charlene H. Chu  <https://orcid.org/0000-0002-0333-7210>
 Katherine S. McGilton  <https://orcid.org/0000-0003-2470-9738>
 Kim N. Le  <https://orcid.org/0000-0002-3517-9878>
 Veronique Boscart  <https://orcid.org/0000-0002-7420-1978>
 Franziska Zúñiga  <https://orcid.org/0000-0002-8844-4903>

References

- Alexopoulos, G. S., Abrams, R. C., Young, R. C., & Shamoian, C. A. (1988). Cornell scale for depression in dementia. *Biological Psychiatry*, 23(3), 271–284.
- Alzheimers Disease International. (2013). *World Alzheimer report 2013 journey of caring: Analysis of long-term care for Dementia*. Author. <http://www.alz.co.uk/research/WorldAlzheimerReport2013.pdf>
- Arco, L., & Du Toit, E. (2006). Effects of adding on-the-job feedback to conventional analog staff training in a nursing home. *Behavior Modification*, 30(5), 713–735. <https://doi.org/10.1177/0145445505281058>
- Arksey, H., & O’Malley, L. (2005). Scoping studies: Towards a methodological framework. *International Journal of Social Research Methodology*, 8(1), 19–32.
- Aylward, S., Stolee, P., Keat, N., & Johncox, V. (2003). Effectiveness of continuing education in long-term care: A literature review. *The Gerontologist*, 43(2), 259–271. <https://doi.org/10.1093/geront/43.2.259>
- Ballard, C., Corbett, A., Orrell, M., Williams, G., Moniz-Cook, E., Romeo, R., Woods, B., Garrod, L., Testad, I., Woodward-Carlton, B., & Woodward-Carlton, B. (2018). Impact of person-centred care training and person-centred activities on quality of life, agitation, and antipsychotic use in people with dementia living in nursing homes: A cluster-randomised controlled trial. *PLoS Medicine*, 15(2), e1002500.
- Beard, J. R., Officer, A., de Carvalho, I. A., Sadana, R., Pot, A. M., Michel, J.-P., Lloyd-Sherlock, P., Epping-Jordan, J. E., Peeters, G. G., Mahanani, W. R., & Chatterji, S. (2016). The World report on ageing and health: A policy framework for healthy ageing. *Lancet (London, England)*, 387(10033), 2145–2154. [https://doi.org/10.1016/S0140-6736\(15\)00516-4](https://doi.org/10.1016/S0140-6736(15)00516-4)
- Beeber, A. S., Zimmerman, S., Fletcher, S., Mitchell, C. M., & Gould, E. (2010). Challenges and strategies for implementing and evaluating dementia care staff training in long-term care settings. *Alzheimer’s Care Today*, 11(1), 17–39. <https://doi.org/10.1097/ACQ.0b013e3181cd1a52>
- Bökberg, C., Behm, L., Wallerstedt, B., & Ahlström, G. (2019). Evaluation of person-centeredness in nursing homes after a palliative care intervention: Pre-and post-test experimental design. *BMC Palliative Care*, 18(1), 1–10.
- Brooker, D. J., & Sure, C. (2006). Dementia Care Mapping (DCM): Initial validation of DCM in UK field trials. *International Journal of Geriatric Psychiatry*, 21, 1018–1025.
- Burgio, L. D., Allen-Burge, R., Roth, D. L., Bourgeois, M. S., Dijkstra, K., Gerstle, J., & Bankester, L. (2001). Come talk with me: Improving communication between nursing assistants and nursing home residents during care routines. *The Gerontologist*, 41(4), 449–460.
- Burgio, L. D., Stevens, A., Burgio, K. L., Roth, D. L., Paul, P., & Gerstle, J. (2002). Teaching and maintaining behavior management skills in the nursing home. *The Gerontologist*, 42(4), 487–496.
- Casper, S., Cooke, H. A., Phinney, A., & Ratner, P. A. (2016, September 1). Practice change interventions in long-term care facilities: What works, and why? *Canadian Journal on Aging*, 35(3), 372–384. <https://doi.org/10.1017/S0714980816000374>
- Chang, C. C., Wykle, M. L., & Madigan, E. A. (2006). The effect of a feeding skills training program for nursing assistants who feed dementia patients in Taiwanese nursing homes. *Geriatric Nursing*, 27(4), 229–237.
- Choquet, R., Maaroufi, M., De Carrara, A., Messiaen, C., Luigi, E., & Landais, P. (2015). A methodology for a minimum data set for rare diseases to support national centers of excellence for healthcare and research. *Journal of the American Medical Informatics Association*, 22(1), 76–85. <https://doi.org/10.1136/amiajnl-2014-002794>
- Cohen-Mansfield, J. (1997). Conceptualization of agitation: Results based on the Cohen-Mansfield agitation inventory and the agitation behavior mapping instrument. *International Psychogeriatrics*, 8(s3), 309–315.
- Cooper, E., Spilsbury, K., McCaughan, D., Thompson, C., Butterworth, T., & Hanratty, B. (2016). Priorities for the professional development of registered nurses in nursing homes: A Delphi study. *Age and Ageing*, 46(1), 39–45. <https://doi.org/10.1093/ageing/awf160>
- Corazzini, K. N., Anderson, R. A., Bowers, B. J., Chu, C. H., Edvardsson, D., Fagertun, A., Gordon, A. L., Leung, A. Y., McGilton, K. S., Meyer, J. E., Lepore, M. J., & WE-THRIVE. (2019). Toward common data elements for international research in long-term care homes: Advancing person-centered care. *Journal of the American Medical Directors Association*, 20(5), 598–603. <https://doi.org/10.1016/j.jamda.2019.01.123>
- Crogan N. L., & Evans B. C. (2001). Nutrition education for nursing assistants: An important strategy to improve long-term care. *Journal of Continuing Education in Nursing*, 32(5), 216–218.
- Dassel, K., Butler, J., Telonis, J., & Edelman, L. (2020). Development and evaluation of Alzheimer’s Disease and Related Dementias (ADRD) best care practices in

- long-term care online training program. *Educational Gerontology*, 46(3), 150–157.
- Davis, E., & Lundstrom, K. (2011, July). Creating effective staff development committees: A case study. *New Library World*, 112(7/8), 334–346. <https://doi.org/10.1108/03074801111150468>
- Fujisawa, R., & Colombo, F. (2009). *The long-term care workforce: Overview and strategies to adapt supply to a growing demand, OECD health working papers*. OECD. https://www.oecd-ilibrary.org/social-issues-migration-health/the-long-term-care-workforce-overview-and-strategies-to-adapt-supply-to-a-growing-demand_225350638472
- Ghandehari, O. O., Hadjistavropoulos, T., Williams, J., Thorpe, L., Alfano, D. P., Dal Bello-Haas, V., Malloy, D. C., Martin, R. R., Rahaman, O., Zwakhalen, S. M. G., Carleton, R. N., Hunter, P. V., & Lix, L. M. (2013). A controlled investigation of continuing pain education for long-term care staff. *Pain Research and Management*, 18(1), 11–18.
- Greene, C., Canning, D., Wilson, J., Bak, A., Tingle, A., Tsiami, A., & Loveday, H. (2018). I-Hydrate training intervention for staff working in a care home setting: An observational study. *Nurse Education Today*, 68, 61–65.
- Huang, T.-T., & Wu, S.-C. (2008). Evaluation of a training programme on knowledge and compliance of nurse assistants' hand hygiene in nursing homes. *The Journal of Hospital Infection*, 68(2), 164–170.
- Huizing, A. R., Hamers, J. P. H., Gulpers, M. J. M., & Berger, M. P. F. (2006). Short-term effects of an educational intervention on physical restraint use: A cluster randomized trial. *BMC Geriatrics*, 6, 17.
- Hussein, S., & Manthorpe, J. (2005). An international review of the long-term care workforce: Policies and shortages. *Journal of Aging and Social Policy*, 17(4), 75–94. https://doi.org/10.1300/J031v17n04_05
- Janssens, B., De Visschere, L., van der Putten, G. J., de Lugt-Lustig, K., Schols, J. M., & Vanobbergen, J. (2016). Effect of an oral healthcare protocol in nursing homes on care staffs' knowledge and attitude towards oral health care: A cluster-randomised controlled trial. *Gerodontology*, 33(2), 275–286.
- Jones, K. R., Fink, R., Vojir, C., Pepper, G., Hutt, E., Clark, L., Scott, J., Martinez, R., Vincent, D., & Mellis, B. K. (2004). Translation research in long-term care: Improving pain management in nursing homes. *Worldviews on Evidence-Based Nursing*, 1, S13–S20.
- Kane, R. A. (2003). *Human resources for long-term care: Lessons from the United States*. World Health Organization Collection on Long-Term Care. https://www.who.int/chp/knowledge/publications/policy_issues_ltc.pdf#page=205
- Kemeny, B., Boettcher, I. F., DeShon, R. P., & Stevens, A. B. (2006). Using experiential techniques for staff development: Liking, learning, and doing. *Journal of Gerontological Nursing*, 32(8), 9–14.
- Kuske, B., Hanns, S., Luck, T., Angermeyer, M. C., Behrens, J., & Riedel-Heller, S. G. (2007, October). Nursing home staff training in dementia care: A systematic review of evaluated programs. *International Psychogeriatrics*, 17(5), 818–841. <https://doi.org/10.1017/S1041610206004352>
- Leone, E., Deudon, A., Bauchet, M., Laye, M., Bordone, N., Lee, J., Piano, J., Friedman, L., David, R., Delva, F., & Delva, F. (2013). Management of apathy in nursing homes using a teaching program for care staff: The STIM-EHPAD study. *International Journal of Geriatric Psychiatry*, 28(4), 383–392.
- Lepore, M., & Corazzini, K. (2019). Advancing international research on long-term care: Using adaptive leadership to build consensus on international measurement priorities and common data elements. *Gerontology and Geriatric Medicine*, 5, 233372141986472. <https://doi.org/10.1177/2333721419864727>
- Mackenzie, C. S., & Peragine, G. (2003). Measuring and enhancing self-efficacy among professional caregivers of individuals with dementia. *American Journal of Alzheimer's Disease & Other Dementias®*, 18(5), 291–299.
- Magai, C., Cohen, C. I., & Gomberg, D. (2002). Impact of training dementia caregivers in sensitivity to nonverbal emotion signals. *International Psychogeriatrics*, 14(1), 25–38.
- McAiney, C. A., Stolee, P., Hillier, L. M., Harris, D., Hamilton, P., Kessler, L., Madsen, V., & Le Clair, J.K. (2007). Evaluation of the sustained implementation of a mental health learning initiative in long-term care. *International Psychogeriatrics*, 19(5), 842–858.
- McGilton, K. S., Backman, A., Boscart, V., Chu, C., Gea Sánchez, M., Irwin, C., Meyer, J., Spilsbury, K., Zheng, N., & Zúñiga, F. (2020). Exploring a common data element for international research in long-term care homes: A measure for evaluating nursing supervisor effectiveness. *Gerontology and Geriatric Medicine*, 6(6), 233372142097981. <https://doi.org/10.1177/2333721420979812>
- Morris, J. C. (1997). Clinical dementia rating: a reliable and valid diagnostic and staging measure for dementia of the Alzheimer type. *International Psychogeriatrics*, 9(S1), 173–176.
- National Institute on Ageing. (2019). *Enabling the Future Provision of Long-Term Care in Canada*. Author.
- Peterson, D., Berg-Weger, M., McGillick, J., & Schwartz, L. (2002). Basic care I: The effect of dementia-specific training on certified nursing assistants and other staff. *American Journal of Alzheimer's Disease & Other Dementias®*, 17(3), 154–164.
- Peterson, E. L., McGlothlin, J. D., & Blue C. L. (2004). The development of an ergonomics training program to identify, evaluate, and control musculoskeletal disorders among nursing assistants at a state-run veterans' home. *Journal of Occupational and Environmental Hygiene*, 1(1), D10–D16.
- Redeker, N. S., Anderson, R., Bakken, S., Corwin, E., Docherty, S., Dorsey, S. G., Heitkemper, M., McCloskey, D. J., Moore, S., Pullen, C., & Grady, P. (2015). Advancing symptom science through use of common data elements. *Journal of Nursing Scholarship*, 47(5), 379–388. <https://doi.org/10.1111/jnu.12155>
- Resnick, B., Cayo, J., Galik, E., & Pretzer-Aboff, I. (2009). Implementation of the 6-week educational component in the Res-Care intervention: process and outcomes. *Journal of Continuing Education in Nursing*, 40(8), 353–360.
- Rosen, J., Mulsant, B. H., Kollar, M., Kastango, K. B., Mazumdar, S., & Fox, D. (2002). Interactive video training test. *Journal of the American Medical Directors Association*, 3, 291–296.
- Saponaro, M. Z., & Baughman, S. (2009). You came for the snacks, but what have you learned? Evaluation of a staff learning program at the University of Maryland Libraries.

- In E. Connor (Ed.), *An introduction to staff development in academic libraries* (pp. 215–233). Routledge.
- Scerri, A., & Scerri, C. (2019). Outcomes in knowledge, attitudes and confidence of nursing staff working in nursing and residential care homes following a dementia training programme. *Aging & Mental Health*, 23(8), 919–928.
- Smith, M., Stolder, M. E., Jaggers, B., Liu, M. F., & Haedtke, C. (2013). Depression Training in Nursing Homes: Lessons Learned from a Pilot Study. *Issues in Mental Health Nursing*, 34(2), 90–102.
- Soderlund, M., Norberg, A., & Hansebo, G. (2014). Validation method training: Nurses' experiences and ratings of work climate. *International Journal of Older People Nursing*, 9(1), 79–89.
- Spector, A., Revolta, C., & Orrell, M. (2016). The impact of staff training on staff outcomes in dementia care: A systematic review. *International Journal of Geriatric Psychiatry*, 31(11), 1172–1187. <https://doi.org/10.1002/gps.4488>
- Stone, R., & Harahan, M. F. (2010, January 2). Improving the long-term care workforce serving older adults. *Health Affairs*, 29(1), 109–115. <https://doi.org/10.1377/hlthaff.2009.0554>
- United Nations, Department of Economic and Social Affairs Population Division. (2019). *World population prospects 2019 highlights*. Author.
- Williams, K., Stolberg, R., O'Kelley-Wetmore, A., Porter, J., Jackson, S., & Van Son, C. (2016). Oral health education for nursing home staff. *Annals of Long Term Care*, 24(7), 19–24.
- Wils, M., Verbakel, J., & Lisaerde, J. (2017). Improving advance care planning in patients with dementia: The effect of training nurses to engage in ACP-related conversations. *Journal of Clinical Gerontology and Geriatrics*, 8(1), 17–20.
- Wood, S., Cummings, J. L., Hsu, M. A., Barclay, T., Wheatley, M. V., Yarema, K. T., & Schnelle, J. F. (2000). The use of the neuropsychiatric inventory in nursing home residents: characterization and measurement. *The American Journal of Geriatric Psychiatry*, 8(1), 75–83.
- World Health Organization. (2011). *Global health and aging*. Author. http://www.who.int/ageing/publications/global_health.pdf
- World Health Organization. (2017). *Global strategy and action plan on ageing and health*. Author. <http://apps.who.int/bookorders>
- Yasuda, M., & Sakakibara, H. (2017). Care staff training based on person-centered care and dementia care mapping, and its effects on the quality of life of nursing home residents with dementia. *Aging & Mental Health*, 21(9), 991–996.
- Zúñiga, F., Chu, C. H., Boscart, V., Fagertun, A., Gea-Sánchez, M., Meyer, J., Spilsbury, K., Devi, R., Haunch, K., Zheng, N., & McGilton, K. S. (2019). Recommended common data elements for international research in long-term care homes: Exploring the workforce and staffing concepts of staff retention and turnover. *Gerontology and Geriatric Medicine*, 5, 2333721419844344. <https://doi.org/10.1177/2333721419844344>