



UNIVERSITY OF LEEDS

This is a repository copy of *Causes of subcutaneous emphysema following dental procedures: a systematic review of cases 1993-2020*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/171424/>

Version: Supplemental Material

Article:

Jones, A, Stagnell, S, Renton, T et al. (2 more authors) (2021) Causes of subcutaneous emphysema following dental procedures: a systematic review of cases 1993-2020. *British Dental Journal*, 231. pp. 493-500. ISSN 0007-0610

<https://doi.org/10.1038/s41415-021-3564-0>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>




Figure 5. Most teeth affected were molar teeth (72%). Third molars were disproportionately represented, accounting for 52% of all affected molars.