

This is a repository copy of *COVID-19 lockdowns highlight a risk of increasing ozone pollution in European urban areas*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/171541/>

Version: Published Version

Article:

Grange, S. K., Lee, J. D. orcid.org/0000-0001-5397-2872, Drysdale, W. S. orcid.org/0000-0002-7114-7144 et al. (4 more authors) (2020) COVID-19 lockdowns highlight a risk of increasing ozone pollution in European urban areas. Atmospheric Chemistry and Physics Discussions. ISSN 1680-7367

<https://doi.org/10.5194/acp-2020-1171>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



COVID-19 lockdowns highlight a risk of increasing ozone pollution in European urban areas

Stuart K. Grange^{1,2}, James D. Lee², Will S. Drysdale², Alastair C. Lewis^{2,3}, Christoph Hueglin¹, Lukas Emmenegger¹, and David C. Carslaw^{2,4}

¹Empa, Swiss Federal Laboratories for Materials Science and Technology, Überlandstrasse 129, 8600 Dübendorf, Switzerland

²Wolfson Atmospheric Chemistry Laboratories, University of York, York, YO10 5DD, United Kingdom

³National Centre for Atmospheric Science, University of York, Heslington, York, YO10 5DD, United Kingdom

⁴Ricardo Energy & Environment, Harwell, Oxfordshire, OX11 0QR, United Kingdom

Correspondence: Stuart K. Grange (stuart.grange@empa.ch); David C. Carslaw (david.carslaw@york.ac.uk)

Abstract.

In March 2020, non-pharmaceutical interventions in the form of lockdowns were applied across Europe to urgently reduce the transmission of SARS-CoV-2, the virus which causes the COVID-19 disease. The near-complete shutdown of the European economy had widespread impacts on atmospheric composition, particularly for nitrogen dioxide (NO₂) and ozone (O₃). To investigate these changes, we analyze data from 246 ambient air pollution monitoring sites in 102 urban areas and 34 countries in Europe between February and July, 2020. Counterfactual, business as usual air quality time series are created using machine learning models to account for natural weather variability. Across Europe, we estimate that NO₂ concentrations were 34 and 32 % lower than expected for traffic and urban-background locations while O₃ was 30 and 21 % higher (in the same environments) at the point of maximum restriction on mobility. The European urban NO₂ experienced in the 2020 lockdown was equivalent to that which might be anticipated in 2028 based on average trends since 2010. Despite NO₂ concentrations decreasing by approximately a third, total oxidant (O_x) changed little, suggesting that the reductions of NO₂ were substituted by increases in O₃. The lockdown period demonstrated that the expected future reductions in NO₂ in European urban areas are likely to lead to a widespread increase in urban O₃ pollution unless additional mitigation measures are introduced.

1 Introduction

On December 31, 2019, a cluster of unexplained pneumonia cases in Wuhan, Hubei, China was reported to the World Health Organization (WHO) (World Health Organization (WHO), 2020a; Wu et al., 2020). Subsequent research in January, 2020 identified the disease to be caused by a previously unknown betacoronavirus (SARS-CoV-2), and the disease was given the name coronavirus disease 2019 (COVID-19) (Zhou et al., 2020; World Health Organization (WHO), 2020c). Due to rapid human-to-human transmission and the introduction of the virus to countries outside China, cases of COVID-19 were soon detected in all continents of the world, with the exception of Antarctica, and on March 11, the WHO declared a COVID-19 pandemic (World Health Organization (WHO), 2020b).



Europe was named the epicentre of the pandemic on March 13, and most European countries undertook unprecedented non-pharmaceutical interventions to reduce the transmission rate of SARS-CoV-2 in early or mid-March (BBC, 2020; Dehning et al., 2020; Remuzzi and Remuzzi, 2020). The exact nature and duration of the measures varied by country, but collectively they are often referred to as “lockdowns” (Ruktanonchai et al., 2020). The lockdowns generally resulted in the closure of all shops, schools, universities, and restaurants with the exception of supermarkets, pharmacies, and other services deemed essential. Working from home whenever possible was encouraged and some countries also controlled, or restricted travel, exercise, and leisure activities. All these measures created a situation where European economic activity was reduced to a bare minimum within a matter of days, and mobility of the European population was severely altered (Figure 1).




Figure 1. European mobility changes based on © Google’s mobility indices between February and July, 2020 (© Google, 2020).

30 The rapid reduction of economic activity had many positive environmental impacts with the improvement of air quality being widely reported, especially via striking satellite observations of column NO_2 (Liu et al., 2020; Patel et al., 2020; Venter et al., 2020). Reductions of CO_2 emissions have also been reported globally due to heavily curtailed economic activities (Le



35 Quéré et al., 2020; Forster et al., 2020). Many of the reports of improved air quality were preliminary, and further research was required to fully understand and quantify the improvements observed throughout Europe, particularly after accounting for meteorological factors (Grange et al., 2020; Carslaw, 2020; Lee et al., 2020; Wang et al., 2020).

The European lockdowns can be thought of and approached as an air quality ‘experiment’ where economic activity was curtailed to near-minimum levels. Questions can be asked from the data such as: what were the results, how do they compare to other planned interventions such as low emission or clean air zones, and whether the observations were inline with what would be expected? The rate and severity of the changes imposed on European populations due to the lockdowns is something that previously could only be investigated by atmospheric modeling. Therefore, the COVID-19 lockdowns have provided a unique ‘real-world modeling scenario’ which represents a plausible future with far fewer internal combustion engine vehicles in use across Europe.

40 Here, we report an analysis based on counterfactual business as usual scenarios using predictive machine learning models. This allows for robust comparisons of the observed concentrations of air pollutants with those which would have been expected without the lockdown measures. The primary objective of this study is to report the response of NO₂ and O₃ concentrations throughout European urban areas caused by mobility restrictions due to COVID-19 lockdown measures. A secondary objective is to outline the implications for European air quality management which the dramatic changes in population mobility exposed.

2 Materials and methods

2.1 Data

50 Up-to-date (UTD) hourly NO₂ and O₃ monitoring data were retrieved from the European Air Quality Portal (European Environment Agency, 2019) for the period between 2018 and 2020 for 102 urban areas in 33 European countries (Figure 2). For the 34th country, the United Kingdom, observations were directly retrieved from the countries’ individual (England, Wales, and Scotland) and national networks (Automatic Urban and Rural Network; AURN) (Department for Environment Food & Rural Affairs, 2020).

55 The 102 urban areas were chosen because they are the capital, a “principal”, or a particularly relevant city for the included European countries (Figure 2). In each urban area, at least one representative traffic site and at least one urban-background site were chosen (if available) to represent the area. Notably, UTD data are not validated, are subject to change, and will only be finalised (at the time of writing) in 11 months time (the deadline is September, 2021). However, the time series were screened for undesirable features such as calibration issues, frequent missing data, or long periods of no reported data. Time series with such obvious issues were not included in the analysis. Unfortunately, oxides of nitrogen (NO_x = NO₂ + NO) data were not available because most countries which participate in the UTD process do not report NO_x (or NO) since it is not a regulated, ambient pollutant in Europe (Grange, 2019). Additionally, total oxidant (O_x = NO₂ + O₃) was calculated (in ppb) and included in the analysis as a third variable.



© OpenStreetMap contributors. Distributed under a Creative Commons BY-SA License

Figure 2. The 102 European urban areas included in the data analysis.

Hourly surface-based meteorological data were downloaded from the Integrated Surface Database (ISD). For the 102 urban areas, these sites were generally airports (NOAA, 2016; Grange, 2020). A total of 246 air quality monitoring sites and 91 meteorological sites were included in the analysis. For details of the sites, see the tables available online³.

In the current work, we focus on changes in the concentrations of NO₂ and O₃ at urban-traffic and urban-background locations. NO₂ and O₃ in such locations are strongly influenced by local road vehicle emissions and not, for example, transboundary contributions, which would be the case for particulate matter (PM_{2.5} and PM₁₀). Furthermore, the concentrations of NO₂ and O₃ in urban areas are strongly influenced by local meteorological effects. Generally, traffic sites are located in close proximity to roads, and pollutant concentrations are forced by local vehicular emissions. The urban-background classification is more

³Temporary location: http://skgrange.github.io/www/data/data_analysis_links.html



varied, but can be thought as environments away from the immediate vicinity of roads and industrial facilities but are still located within an urban area.

2.2 Business as usual (BAU) modeling

75 A central issue when considering changes in atmospheric concentrations due to an intervention is whether the change is due to variations in meteorological conditions or emission source strength (Grange and Carslaw, 2019). This problem is widespread and affects time scales from hours to years. It is particularly important in ‘before-after’ studies where meteorological change, rather than changes in emission source strength, can easily dominate the variation in concentrations. This ambiguity can be somewhat reduced by averaging over several years to account for past inter-annual variability. However, this approach cannot
80 account for the significant impact that meteorology may have on a specific observation period.

In the current context of the changes in activities brought about by COVID-19 lockdowns, the changes are over a duration of several months and span a period from spring to summertime conditions. This period straddles important natural changes in meteorological conditions and atmospheric composition. For example, during February, 2020 the UK and much of western Europe experienced exceptionally high mean wind speeds due to storms Ciara, Dennis, and Jorge. Surface wind speed
85 records in Southern England suggest February, 2020 had the highest mean wind speed of any month for over 40 years. This demonstrates that the state of the atmospheric dispersion across Europe at the time of COVID-19 lockdowns was different than experienced in previous years. Similarly, urban-background concentrations of O_3 in the northern hemisphere tend to increase from the beginning of the year and peak in April, which will also influence NO_2 (Monks, 2000). These, and other factors suggest that considerable care is needed for the quantification of an intervention such as the COVID-19 lockdowns on surface
90 concentrations of primary and secondary pollutants.

To address the above issues, random forest models were trained to explain hourly mean NO_2 and O_3 concentrations using surface meteorological and time explanatory variables for each monitoring site (Breiman, 2001). The explanatory variables used were: wind direction, wind speed, air temperature, relative humidity, atmospheric pressure (if available in the ISD database), a trend term in the form of Unix date, a seasonal term in the form of Julian day, weekday, and hour of day. The following random
95 forest hyper-parameters were kept constant for all models: 300 trees, three variables to split at each node, and a minimal node size of five. The training period spanned just over two years and was between January 1, 2018 and February 14, 2020. The training-testing split percentage was 80 and 20 respectively. From February 14 to July 31, 2020, the models were used in predictive mode to predict pollutant concentrations based on the observed meteorological variables.

The models’ predictions can be thought of as business as usual (BAU) scenarios based on past behaviour of pollutant
100 concentrations and the weather which was experienced after February 14 at each monitoring site. Thus, the model represents a *counterfactual* which observed concentrations can be compared with (for example, see Figure 3).

February 14 to March 1, 2020 was considered a validation period where the models’ skill were checked for adequate performance. Summaries of the models’ performance based on the random forest model objects and predictions during the validation period in the form of R^2 are shown in Figure A1. From the start date of the lockdowns (the earliest was March 9 in Italy), the
105 application period began and gave estimates of BAU, *i.e.*, what concentrations would have been if the lockdown measures were




Figure 3. A NO_2 example where the observed concentrations clearly diverged from the business as usual (BAU) scenario for the Nice Promenade (France) traffic monitoring site between February and July, 2020.

not implemented. The modeling was conducted using the **rmweather** R package (Grange et al., 2018; Grange and Carslaw, 2019; Grange, 2018).

110 During the validation phase, a number of models showed bias in prediction, most notably, NO_2 was under-predicted at many locations. The under-prediction was on average, $-3.7 \mu\text{g m}^{-3}$ (95 % CI [-4.2, -3.3]). This under-prediction was most likely
115 caused by already-curtailed economic activity and reduced emissions throughout Europe at the very end of February and the beginning of March, *i.e.*, before the formal lockdowns were implemented. The beginning of 2020 was also mild in respect to ambient temperature and rather windy at most locations (discussed above) which may have resulted in some models under-predicting concentrations at this time of the year. For consistency and to create a reference point in time, the model predictions were corrected by calculating the model offset validation phase (February 14 to March 1) and subtracting this offset from the predictions. This ensured that the counterfactual predictions were calibrated at the start of the application phase and represented the changes in concentrations after March 1, 2020.



2.2.1 Change point analysis

To link NO₂ and O₃ concentration changes in March–April, 2020 to the lockdown restrictions placed on European populations, change point models were calculated. These change point models were conceptually simple – an intercept change was the expected *a priori* assumption. There were two motivations for these change point models. The first was to identify both the time, and magnitude of concentration response with an objective, data-driven approach rather than using a subjective and manual classifier. The second was to use such a technique to identify an atmospheric response after an intervention (an unplanned one in this case) which is a general goal of air quality data analysis.

The change point logic was implemented with the **mcp** R package with Bayesian inference (Lindeløv, 2020). To detect the change points, three Markov chains were run with 9000 iterations. The change point models tested the delta between the observed and counterfactual, however, the change-points were calibrated back to their pre-lockdown concentrations to conduct the (relative) percentage change calculations.

2.2.2 Presentation of results

When presenting the results of the analysis, most time series are displayed as seven-day rolling means. These rolling means act as a smoothing filter to make patterns clearer and remove the day-to-day variations generally seen in air quality monitoring data. Thirty-four countries were included in the analysis (Figure 2), but to avoid overwhelming plots and figures, a consistent set of six European countries (France, Germany, Italy, Spain, Switzerland, and the United Kingdom) were chosen to be displayed when discussing the counties' air quality patterns.

3 Results and discussion

3.1 Mean concentration changes

For all 34 European countries analysed, the observed concentrations of NO₂ were lower than those predicted by the counterfactual business as usual (BAU) scenarios between February 14 and July 31, 2020 (deltas (Δ) between the observed concentrations and predicted counterfactual shown in Figure 4). The reductions of NO₂ were greater in both an absolute and relative sense at the sites classified as either roadside or traffic environments compared to urban-background locations which can be explained by NO₂ being primarily a traffic-sourced pollutant (Grange et al., 2017). The impacts of vehicle-flow reductions during the lockdowns were more dramatic in the close proximity of roads when compared to more distant urban-background locations.

Mean O₃ concentrations increased at a similar magnitude to which NO₂ decreased throughout Europe between February and July, 2020 (Figure 4). Like NO₂, O₃ at roadside locations showed a greater divergence from the BAU predictions than urban-background sites. The near-mirror image of NO₂ and O₃ can be explained by the relationship between NO_x and O₃. The reduction of NO_x emissions and concentrations across Europe drove decreased O₃ destruction via the NO titration cycle during this period. In many countries, the 8-hour legal limit for O₃ of 120 $\mu\text{g m}^{-3} \text{ 8 h}^{-1}$ was breached during this time period.





Figure 4. Seven-day rolling means of the observed-predicted concentrations deltas for NO_2 and O_3 for all European sites analysed between February 14 and July 31, 2020.

Unlike NO_2 where concentrations remained below their BAU estimates until the end of the analysis period, O_3 concentrations returned to their expected values by the end of July, 2020.

3.2 Timing of changes

150 Figure 4 clearly indicates that concentrations in the first half of 2020 diverged from what was predicted by the counterfactual modelling. To objectively identify the date and magnitude of maximum divergence, change points were identified with a data-driven approach using Bayesian inference. The mean dates when NO_2 started to diverge at their greatest extent from the BAU scenarios along with national lockdown dates for six European countries are displayed in Figure 5. For the complete set of dates for all countries included in the analysis, see Table A1.

155 For NO_2 , the change points were between seven days before and seven days after the countries' lockdown date (excluding the outlier of Denmark). For O_3 , this range was greater, between -12 and 8 days. Italy was the first country in Figure 5 where change points were identified for NO_2 concentrations on March 13, 2020 and this was four days before Italy's nationwide lockdown date while Spain's NO_2 change point was the same as the country's lockdown date. Change points were often identified a day or two earlier than the lockdown date when the lockdown began on a Sunday or a Monday, for example, in



Vertical dashed lines are dates of nationwide lockdowns

Figure 5. Estimated timing of changes to NO_2 concentrations for six European countries between March and May, 2020. The distribution shown for each country is the dimensionless probability distribution of the estimated change-point in concentration. The country panels are ordered by nationwide lockdown date.



160 Germany. For almost every site included in the analysis, the change points for NO_2 were ones of decreases while those for O_3 were increases (as seen in Figure 4).

Figure 5 shows that some countries had very consistent changes in concentrations for the sites which were analysed, for example Spain. Changes in other countries were less consistent which may indicate regional differences within countries. The UK showed two peaks in density for the NO_2 change points which were separated by a week. This feature represents a two-
165 phase reduction in emissions because staggered lockdown measures were announced – the first was a set of recommendations for social distancing and not visiting restaurants and other social establishments (on March 16), while the second announcement (March 23) was one of a more strict lockdown.

Although the identified change point dates for NO_2 were broadly consistent with the various countries' lockdown dates, the change points for O_3 were not aligned as closely (Table A1). There was also no correlation between the magnitude of
170 NO_2 reduction and the time required for an O_3 change point to be identified. This suggests that O_3 's secondary generation processes did not immediately respond to reductions of ambient NO_x concentrations after lockdowns were imposed due to less NO titration. For this process to be identifiable, O_3 generation must occur, and this requires sunlight. Therefore, the lack of sunny conditions in some urban areas around the time of the NO_2 atmospheric response may have resulted in varying duration lags before changes in O_3 could be observed.

175 3.3 Concentration changes among different countries

At a European level, maximum divergence of NO_2 and O_3 from the counterfactual predictions was reached in late-March, 2020 (Figure 4). However, there was some diversity among European country NO_2 and O_3 divergence from their counterfactuals for the analysis periods (Figure 6). All countries analysed passed their maximum divergences for NO_2 and O_3 in late-April, and the shape of the recovery is of a “swoosh” with a sharp plunge away from the counterfactual around the date of the lockdown
180 implementations (Figure 6), but the rapid plunge was followed by a slower, and more gradual return to the BAU until the end of July. This pattern is very much reminiscent of the mobility changes shown in Figure 1.

Some countries experienced a smaller reduction in NO_2 than others. Germany and Switzerland for example, experienced lower NO_2 reductions when compared to France, Italy, and Spain. Some countries' greater reductions in ambient NO_2 concentrations could be explained by the level of “stringency” of the countries' lockdowns and resulting changes in mobility
185 (Hale et al., 2020; © Google, 2020). For example, Germany and Switzerland's measures were very strong recommendations with few legally enforceable restrictions on recreational or leisure activities, while France, Italy, and Spain had more stringent requirements where movement and travel were restricted and enforced in a much stronger manner. It is very likely that these different levels (or enforcement) of restrictions had implications for emissions of atmospheric pollutants. However, meteorological conditions, perhaps similar synoptic scale patterns likely played a role in the differences observed among the countries
190 too.

After late-April, concentrations moved towards their predicted counterfactual values and this continued to the end of the analysis period (Figure 6). Some European countries began to remove lockdown restrictions in the second half of April which increased traffic-sourced emissions, and this is consistent with the observations in Figure 4 and Figure 6. O_3 concentrations




Figure 6. Seven-day rolling means of the observed-predicted concentrations deltas for NO_2 , O_3 , and O_x for six selected countries in Europe between February 14 and July 31, 2020.

195 returned to approximately their BAU levels by the end of July, but NO_2 had yet to do so at the end of the analysis period, with the exception of Italy. This indicates that NO_x emissions (mostly traffic-sourced) had not yet reached their estimated BAU levels by the end of July across most of Europe after the country lockdowns were released.



3.4 Quantifying the changes in concentrations

The change point dates identified by Bayesian inference shown in Figure 5 and Table A1 were used to classify the time series as pre-lockdown, within lockdown, or post-lockdown periods. With this classification, concentrations were compared to calculate concentration deltas and percentage changes. At a European level, the mean NO₂ percentage changes for NO₂ at traffic and urban-background sites were -34 % (95 % CI [-36, -31]) and -32 % (95 % CI [-35, -29]) respectively (which equalled concentration reductions of -11 and -7 μg m⁻³). The European annual NO₂ standard is 40 μg m⁻³ y⁻¹, and the mean reduction of 11 μg m⁻³ is 27 % of the legal limit (European Commission, 2019). For O₃, the mean European percentage change for traffic and urban-background sites were estimated at 30 % (95 % CI [26, 35]) and 21 % (95 % CI [18, 24]), and the concentration changes were 12 and 9 μg m⁻³ respectively. The concentration deltas and percentage changes attributed to the European lockdown measures are listed by country and site type in Table 1.

To put these concentration changes into context, NO₂ and O₃ trend analysis between 2010 and 2019 for the 246 sites was conducted. Based on the sites which had a complete data record, the mean trends were -1.44 and -0.72 μg m⁻³ y⁻¹ for NO₂ at traffic and urban-background locations, while O₃ trends in the same environments were 0.2 and 0.49 μg m⁻³ y⁻¹. Therefore, at the roadside, the mean reduction of NO₂ across Europe due to the COVID-19 lockdown measures was equivalent to that of 7.6 years of continued concentration reduction, or equivalent to the anticipated European atmosphere in 2028 (Figure 7). O₃ however, increased at an equivalent of 17 years of the rate of change determined by trend analysis in urban-background locations.

The changes at traffic sites will strongly reflect the influence of changes in traffic activity in close proximity to each site for NO_x, NO₂ and O₃. Close to roads, the origins of NO₂ can be thought of as the combination of a background component, a component which is generated from the fast reaction between vehicular NO emissions and O₃, and directly emitted (primary) NO₂. The primary NO₂ contribution is known to have decreased in recent years from a peak around 2010. In London for example, the analysis of 35 traffic-influenced sites showed a reduction in the mean NO₂/NO_x vehicle emission ratio from around 25% in 2010 to about 15% in 2014, (Carslaw et al., 2016) while at a European level, the NO₂/NO_x emission ratio peaked at 16 % (also in 2010) (Grange et al., 2017). This decrease is believed to be driven by improvements in selective catalytic reduction control systems used on vehicles to reduce NO_x and also to the effect of ageing of diesel oxidation catalysts (Carslaw et al., 2019).

The decrease in primary NO₂ emissions over the past decade would have acted to reduce ambient NO₂ concentrations close to roads. Indeed, if the traffic reductions experienced across Europe through country-wide lockdowns had occurred closer to 2010, the reductions in road vehicle NO₂ emissions would have been much more important in affecting ambient concentrations than was experienced in early 2020.

The posterior draws (a type of model prediction) from the change point models show that in some countries, the reduction of traffic volumes during the COVID-19 lockdowns reduced NO₂ concentrations to those which are experienced at urban-background locations (United Kingdom shown in Figure 8). The roadside increment or enhancement of NO₂ was essentially eliminated by reducing traffic to the levels which were experienced while in the lockdown state. However, as discussed above



Table 1. Mean concentration deltas/differences and percentage changes of NO₂, O₃, and O_x for different countries and site types attributed to COVID-19 lockdown measures in March, 2020. Values which are missing indicates that there were not data and NC indicate no change point was identified.

Country	Site type	NO ₂		O ₃		O _x	
		Δ (μg m ⁻³)	% change	Δ (μg m ⁻³)	% change	Δ (ppb)	% change
Andorra	Traffic	–	–	–	–	–	–
Andorra	Urban-back.	-19.8	-59.7	16.1	43.0	-3.4	-9.8
Austria	Traffic	-7.6	-24.5	–	–	–	–
Austria	Urban-back.	-5.2	-23.1	11.3	19.5	4.3	11.2
Belgium	Traffic	-10.8	-45.3	5.0	10.5	-2.2	-6.5
Belgium	Urban-back.	-9.5	-38.4	8.9	19.2	2.4	6.5
Bosnia and Herzegovina	Traffic	–	–	–	–	–	–
Bosnia and Herzegovina	Urban-back.	-1.8	-11.9	1.4	15.0	-1.3	-3.4
Bulgaria	Traffic	-13.8	-29.5	14.0	29.6	0.9	2.2
Bulgaria	Urban-back.	-10.4	-34.2	13.9	33.6	3.0	8.4
Croatia	Traffic	-16.2	-42.3	–	–	–	–
Croatia	Urban-back.	-12.4	-43.9	21.5	34.1	4.4	9.6
Cyprus	Traffic	-15.3	-47.0	–	–	-2.8	-7.2
Cyprus	Urban-back.	-16.7	-59.7	6.1	10.9	-5.0	-11.8
Czechia	Traffic	NC	NC	–	–	–	–
Czechia	Urban-back.	NC	NC	–	–	–	–
Czechia	Urban-back.	–	–	9.0	18.3	4.9	13.8
Denmark	Traffic	-6.7	-28.0	15.7	31.7	3.9	9.8
Denmark	Urban-back.	-4.2	-49.0	7.6	12.3	3.1	8.4
Estonia	Traffic	-5.0	-35.2	0.7	1.3	-1.8	-5.2
Estonia	Urban-back.	-2.4	-29.2	6.4	10.7	-0.4	-1.2
Finland	Traffic	-9.4	-42.5	–	–	–	–
Finland	Urban-back.	-4.3	-34.1	–	–	–	–
France	Traffic	-20.3	-54.2	–	–	–	–
France	Urban-back.	-11.2	-44.1	13.9	35.0	-4.9	-12.1
Germany	Traffic	-10.5	-29.3	15.1	37.3	3.0	7.5
Germany	Urban-back.	-4.9	-21.6	8.8	16.6	3.5	9.1
Greece	Traffic	-12.3	-37.1	–	–	-1.1	-0.4
Greece	Traffic	–	–	NC	NC	–	–
Greece	Urban-back.	-9.5	-43.9	–	–	-3.8	-8.5
Greece	Urban-back.	–	–	NC	NC	–	–
Hungary	Traffic	NC	NC	–	–	–	–
Hungary	Urban-back.	NC	NC	–	–	–	–
Hungary	Urban-back.	–	–	5.0	15.7	-4.2	-11.4
Iceland	Traffic	-5.3	-33.7	–	–	–	–
Iceland	Urban-back.	-3.4	-23.5	–	–	–	–
Ireland	Traffic	–	–	–	–	–	–
Ireland	Urban-back.	-4.9	-33.6	–	–	-1.3	-3.5
Ireland	Urban-back.	–	–	NC	NC	–	–
Italy	Traffic	-17.3	-31.9	–	–	–	–
Italy	Urban-back.	-12.5	-32.7	3.8	14.1	-1.5	-2.2
Lithuania	Traffic	-7.0	-25.9	13.8	34.3	2.8	7.3
Lithuania	Urban-back.	-4.5	-21.0	–	–	–	–
Luxembourg	Traffic	-15.5	-53.2	–	–	–	–
Luxembourg	Urban-back.	-10.3	-47.0	9.6	17.0	-0.1	-0.3




Figure 7. Mean European roadside NO₂ trend with the reduction of NO₂ concentrations attributed to the COVID-19 lockdowns put in context.

and shown in Figure 8, O₃ increased in response to the reductions of NO₂ and O_x only altered very slightly. The same patterns in the United Kingdom were also experienced in other European countries such as France and Spain, but were not as clear for counties such as Switzerland and Germany.

3.5 O_x – NO₂ and O₃ repartitioning

235 Figure 4 and Figure 6 demonstrate that NO₂ concentrations and emissions decreased throughout Europe due to the COVID-19
lockdown measures, especially at the roadside. However, the reduction of NO₂ was accompanied by an increase of O₃ at
a similar magnitude and resulted in O_x showing little change despite the large reductions in traffic-sourced NO₂ (for example,
Figure 8).

240 Mean European changes in O_x were variable between the two site environments. At traffic sites, O_x decreased by -1 ppb
(-1.8 %; 95 % CI [-4, 0.7]) while in urban-background locations, O_x increased by 0.7 ppb (2.1 %; 95 % CI [-0.2, 4]). In the
case of the traffic sites, the modest decrease of O_x can be partially explained by decreased emissions of primary NO₂ (Grange
et al., 2017). However, in urban-background locations, O_x remained nearly constant. This is a very important observation



Table 1. Table 1 continued.

Country	Site type	NO ₂		O ₃		O _x	
		Δ (μg m ⁻³)	% change	Δ (μg m ⁻³)	% change	Δ (ppb)	% change
Malta	Traffic	-13.2	-38.7	10.0	15.4	-4.1	-8.1
Malta	Urban-back.	–	–	–	–	–	–
Netherlands	Traffic	-6.2	-28.3	–	–	1.3	3.5
Netherlands	Traffic	–	–	NC	NC	–	–
Netherlands	Urban-back.	-3.5	-21.2	–	–	4.1	11.2
Netherlands	Urban-back.	–	–	NC	NC	–	–
North Macedonia	Traffic	-8.6	-33.2	–	–	-1.9	-6.8
North Macedonia	Traffic	–	–	NC	NC	–	–
North Macedonia	Urban-back.	–	–	–	–	–	–
Norway	Traffic	-7.7	-30.0	–	–	–	–
Norway	Urban-back.	-2.8	-17.1	–	–	0.9	2.2
Norway	Urban-back.	–	–	NC	NC	–	–
Poland	Traffic	-11.7	-27.6	–	–	–	–
Poland	Urban-back.	-3.6	-12.7	7.1	15.1	2.1	5.5
Portugal	Traffic	-25.9	-53.8	20.2	46.8	-10.7	-24.6
Portugal	Urban-back.	-11.9	-40.5	13.8	26.8	4.7	12.1
Romania	Traffic	-5.8	-7.2	–	–	–	–
Romania	Urban-back.	-7.5	-26.3	13.0	39.9	-0.5	-0.5
Serbia	Traffic	–	–	–	–	–	–
Serbia	Urban-back.	-10.4	-56.4	15.6	44.9	-4.1	-12.6
Slovakia	Traffic	-6.8	-19.5	–	–	–	–
Slovakia	Urban-back.	–	–	–	–	–	–
Slovenia	Traffic	-9.6	-30.5	–	–	–	–
Slovenia	Urban-back.	-5.0	-18.9	20.9	55.7	8.2	26.1
Spain	Traffic	-22.8	-57.2	21.0	61.9	-1.5	-2.8
Spain	Urban-back.	-16.4	-55.7	15.9	37.5	-2.2	-5.4
Sweden	Traffic	-4.9	-17.0	–	–	–	–
Sweden	Urban-back.	-1.5	-12.5	6.5	12.2	0.6	2.0
Switzerland	Traffic	-5.5	-17.2	10.9	22.1	5.1	13.0
Switzerland	Urban-back.	-3.3	-10.1	11.7	21.7	5.2	14.4
United Kingdom	Traffic	-14.4	-50.8	14.4	45.8	-3.8	-8.3

for European air quality management. It suggests that the 34 % reduction of NO₂ concentrations was equalled by a similar absolute increase in O₃, which is clearly an undesirable outcome because of the deleterious effects of O₃ on population health, buildings, and vegetation.

The repartitioning of NO₂ to O₃ is of importance from a public health perspective. As Williams et al. (2014) argue, there are good reasons from an atmospheric chemistry perspective to consider NO₂ and O₃ together in epidemiological studies, rather than either of the two pollutants separately in single-pollutant models. Indeed, Williams et al. (2014) found that there were larger associations (on mortality) for mean 24 hour concentrations of O_x than for either O₃ or NO₂ individually. On this basis, the current analysis suggest that the health impacts may have been small because O_x concentrations changed little in urban environments. The analysis conducted here was exclusively concerned with daily mean O₃ concentrations, and does not




Figure 8. Posterior draws for NO₂, O₃, and O_x two-intercept change point models for the United Kingdom between March and May, 2020.

explore the subtleties associated with peak and/or increases in daily minima O₃ concentrations which are also important when considering the deleterious effects of O₃.

Efficacious management of O₃ has proven to be a challenge in Europe and in many other locations around the world (Sillman, 1999; Wang et al., 2017; Chang et al., 2017; Li et al., 2019). The struggle with O₃ control is partly due to the highly non-linear chemistry of O₃ production based on the precursors volatile organic compounds (VOCs) and NO_x. There are two regimes: NO_x-sensitive and VOC-sensitive – and the O_x analysis presented here strongly suggests that O₃ production is overwhelmingly VOC-sensitive across urban Europe. Therefore, if higher O₃ concentrations are to be avoided in the future where reductions in NO_x emissions of the scale seen in lockdown are likely, enhanced control of VOC emissions will be critical in the European urban environment. The prominence given to NO₂ as a pollutant following the dieselgate scandal of 2015 (Anenberg et al., 2017) has led to far more ambitious NO₂ emissions reductions policies in Europe than are currently in place for VOCs.

VOCs are only measured routinely in a few locations throughout Europe's urban areas, and represent a broad class of pollutant that are emitted from a wide range of sources. Whilst in the 1980s and 1990s VOC emissions were dominated by gasoline vehicle emissions (both tailpipe and evaporative) in more recent years their abundance has become increasingly influenced by non-transport sources such as natural gas leakage and wider solvent use (Lewis et al., 2020).

Data from the London Eltham site, the only suburban VOC monitoring site in the UK, indicates that for many VOCs lockdown did not lead to significant changes in overall emissions or atmospheric concentrations (Figure A3). A conclusion from this albeit anecdotal evidence would be that further reductions in only traffic-related VOC emissions would not likely generate the desired air quality improvements in O₃ and that reducing emissions from other sectors would be essential.



270 Although out of scope for this current work, an obvious avenue for future research is to further explore how individual
VOC concentrations responded during the lockdown periods in European urban areas in order to evaluate the proportion of
VOCs that still come from traffic. This, combined with chemical modelling on a species by species basis to fully assess O_3
production chemistry, would help direct where future VOC reduction strategies should be focused. An analysis such as this
would also strongly benefit from the access of NO_x data which, arguably, would be a better pollutant to analyse than NO_2
275 from an emissions perspective. We strongly encourage the institutions which are involved with reporting ambient air quality
data to the European Environment Agency to include NO_x alongside the legally required NO_2 observations for the air quality
community.

4 Conclusions

This work represents a classic air quality data analysis where atmospheric responses are linked to an intervention. In this case,
280 the intervention was an unplanned, likely unique, and extreme event with very different characteristics when compared to
typical interventions such as the introduction of new emission standards and low emission zones. Despite the extreme nature of
the COVID-19 lockdowns and their results being much more impactful on urban atmospheric composition than other policies
over a short time period, the analysis still demonstrates the difficulty of detecting “change upon change” for atmospheric
pollutants – especially for locations where concentrations are close to background. However, this analysis presents a robust
285 and portable framework for intervention analysis using a combination of machine learning-derived counterfactuals and change
point analysis to identify the timing and magnitude of an atmospheric response.

Analysis of the effect of the European COVID-19 lockdowns on NO_2 , O_3 , and O_x concentrations combining machine
learning derived BAU modelling and Bayesian change point models indicate that NO_2 concentrations reduced by 34 % at
roadside locations. However, the widespread reductions of NO_2 concentrations was accompanied by increases of O_3 at a
290 similar magnitude (30 %), and thus, O_x altered only very slightly due to the lockdowns when considering Europe as a whole.

This insight has important implications for the implementation of future air quality management policies. The COVID-19
lockdown conditions give a glimpse of a realistic, and indeed likely, future environment where NO_x emissions continue to
reduce at their current rate, primarily because of the increasing stringency of vehicular emission standards (Carslaw et al.,
2016; Grange et al., 2017). The future reduction of NO_x concentrations will likely result in repartitioning of O_x and the
295 increase of O_3 concentrations across most European urban areas. Although increases in European O_3 concentrations have been
acknowledged, the further rise should be pre-empted by the European air quality management community through increased
focus on VOC emission controls and the more holistic combined management of NO_2 , O_3 , and VOCs. This will allow for
continued improvements to air quality in a general sense, rather than focusing on reductions of individual pollutants.



300 *Code and data availability.* The data sources used in this work are described and are publicly accessible in a temporary data repository (http://skgrange.github.io/www/data/data_analysis_links.html). These data will be moved to a persistent data repository once the analysis is has been reviewed and finalised. Additional information about these data are available from the authors on reasonable request.

Author contributions. SKG and DCC conceived the research questions, conducted the analysis, and wrote the manuscript. JDL, WSD, and ACL contributed to the research design and writing of the manuscript. CH and LE reviewed and contributed to the manuscript writing.

Competing interests. The authors declare no competing interest.

305 *Acknowledgements.* SKG is supported by the Swiss Federal Office for the Environment (FOEN) and the Natural Environment Research Council (NERC) while holding associate status at the University of York.




Figure A1. Summaries of R^2 values from the random forest model objects and for predictions during the model validation period (February 14 to March 1, 2020) for the three predicted variables.




Figure A2. Oxford COVID-19 Government Response Tracker's (OxCGRT) stringency index of COVID-19 lockdown measures imposed by different countries' governments between February and July, 2020 (Hale et al., 2020).




Figure A3. Volatile organic compounds (VOCs) time series at London Eltham, an urban-background site in United Kingdom.



Table A1. Most commonly identified dates where observed and BAU modeled concentrations diverged in March, 2020. Dates which are missing indicates no change point was detected in March, 2020.

Country	Lockdown date	NO ₂ date	O ₃ date
Andorra	Fri., Mar. 13, 2020	Sat., Mar. 14, 2020	Thu., Mar. 19, 2020
Austria	Mon., Mar. 16, 2020	Thu., Mar. 19, 2020	Mon., Mar. 16, 2020
Belgium	Wed., Mar. 18, 2020	Sun., Mar. 15, 2020	Sat., Mar. 21, 2020
Bosnia and Herzegovina	Sat., Mar. 21, 2020	Thu., Mar. 19, 2020	Thu., Mar. 12, 2020
Bulgaria	Fri., Mar. 13, 2020	Wed., Mar. 11, 2020	Wed., Mar. 18, 2020
Croatia	Thu., Mar. 19, 2020	Fri., Mar. 20, 2020	Fri., Mar. 20, 2020
Cyprus	Sun., Mar. 15, 2020	Fri., Mar. 13, 2020	Thu., Mar. 19, 2020
Czechia	Mon., Mar. 16, 2020	-	Fri., Mar. 20, 2020
Denmark	Fri., Mar. 13, 2020	Fri., Mar. 27, 2020	Tue., Mar. 17, 2020
Estonia	Fri., Mar. 13, 2020	Mon., Mar. 16, 2020	Sat., Mar. 21, 2020
Finland	Mon., Mar. 16, 2020	Tue., Mar. 17, 2020	-
France	Tue., Mar. 17, 2020	Sat., Mar. 14, 2020	Wed., Mar. 11, 2020
Germany	Sun., Mar. 22, 2020	Sun., Mar. 22, 2020	Sat., Mar. 28, 2020
Greece	Mon., Mar. 16, 2020	Tue., Mar. 17, 2020	-
Hungary	Mon., Mar. 16, 2020	-	Sat., Mar. 14, 2020
Iceland	Mon., Mar. 16, 2020	Sat., Mar. 14, 2020	-
Ireland	Fri., Mar. 13, 2020	Thu., Mar. 19, 2020	-
Italy	Mon., Mar. 09, 2020	Fri., Mar. 13, 2020	Thu., Mar. 19, 2020
Lithuania	Mon., Mar. 16, 2020	Tue., Mar. 17, 2020	Wed., Mar. 11, 2020
Luxembourg	Mon., Mar. 16, 2020	Sat., Mar. 14, 2020	Fri., Mar. 20, 2020
Malta	Sun., Mar. 22, 2020	Sat., Mar. 14, 2020	Sun., Mar. 15, 2020
Netherlands	Mon., Mar. 16, 2020	Mon., Mar. 16, 2020	-
North Macedonia	Wed., Mar. 18, 2020	Fri., Mar. 13, 2020	-
Norway	Thu., Mar. 12, 2020	Tue., Mar. 17, 2020	-
Poland	Thu., Mar. 12, 2020	Tue., Mar. 17, 2020	Tue., Mar. 24, 2020
Portugal	Wed., Mar. 18, 2020	Sat., Mar. 14, 2020	Wed., Mar. 18, 2020
Romania	Mon., Mar. 16, 2020	Sat., Mar. 21, 2020	Tue., Mar. 17, 2020
Serbia	Sat., Mar. 21, 2020	Tue., Mar. 17, 2020	Mon., Mar. 16, 2020
Slovakia	Mon., Mar. 16, 2020	Sun., Mar. 22, 2020	-
Slovenia	Mon., Mar. 16, 2020	Thu., Mar. 12, 2020	Tue., Mar. 17, 2020
Spain	Sat., Mar. 14, 2020	Sat., Mar. 14, 2020	Sun., Mar. 15, 2020
Sweden	-	Wed., Mar. 18, 2020	Fri., Mar. 20, 2020
Switzerland	Tue., Mar. 17, 2020	Sun., Mar. 22, 2020	Thu., Mar. 26, 2020
United Kingdom	Mon., Mar. 23, 2020	Mon., Mar. 23, 2020	Thu., Mar. 26, 2020



References

- Anenberg, S. C., Miller, J., Minjares, R., Du, L., Henze, D. K., Lacey, F., Malley, C. S., Emberson, L., Franco, V., Klimont, Z., and Heyes, C.: Impacts and mitigation of excess diesel-related NO_x emissions in 11 major vehicle markets, *Nature*, 545, 467–471, <http://dx.doi.org/10.1038/nature22086>, 2017.
- BBC: Coronavirus: Europe now epicentre of the pandemic, says WHO, <https://www.bbc.com/news/world-europe-51876784>, 13 March 2020., 2020.
- Breiman, L.: Random Forests, *Machine Learning*, 45, 5–32, <https://doi.org/10.1023/A:1010933404324>, <http://dx.doi.org/10.1023/A:1010933404324>, 2001.
- 315 Carslaw, D.: Blog update on COVID-19 and changes in air pollution, <https://ee.ricardo.com/news/blog-update-on-covid-19-and-changes-in-air-pollution>, ricardo Energy & Environment. 07 May 2020, 2020.
- Carslaw, D. C., Murrells, T. P., Andersson, J., and Keenan, M.: Have vehicle emissions of primary NO₂ peaked?, *Faraday Discussions*, 189, 439–454, <https://doi.org/10.1039/C5FD00162E>, <http://dx.doi.org/10.1039/C5FD00162E>, 2016.
- Carslaw, D. C., Farren, N. J., Vaughan, A. R., Drysdale, W. S., Young, S., and Lee, J. D.: The diminishing importance of nitrogen dioxide emissions from road vehicle exhaust, *Atmospheric Environment: X*, p. 100002, <http://www.sciencedirect.com/science/article/pii/S2590162118300029>, 2019.
- 320 Chang, K.-L., Petropavlovskikh, I., Cooper, O. R., Schultz, M. G., and Wang, T.: Regional trend analysis of surface ozone observations from monitoring networks in eastern North America, Europe and East Asia, *Elementa: Science of the Anthropocene*, 5, 22pp, <https://doi.org/http://doi.org/10.1525/elementa.243>, <https://www.elementalscience.org/articles/10.1525/elementa.243/>, 2017.
- 325 © Google: COVID-19 Community Mobility Reports, <https://www.google.com/covid19/mobility/>, 2020.
- Dehning, J., Zierenberg, J., Spitzner, F. P., Wibral, M., Neto, J. P., Wilczek, M., and Priesemann, V.: Inferring change points in the spread of COVID-19 reveals the effectiveness of interventions, *Science*, p. eabb9789, <http://science.sciencemag.org/content/early/2020/05/14/science.abb9789.abstract>, 2020.
- Department for Environment Food & Rural Affairs: UK AIR Air Information Resource – Automatic Urban and Rural Network (AURN), <https://uk-air.defra.gov.uk/networks/network-info?view=aur>, 2020.
- 330 European Commission: Air Quality Standards, <https://ec.europa.eu/environment/air/quality/standards.htm>, under Directive 2008/50/EU. Last updated: 31/12/2019., 2019.
- European Environment Agency: Download of air quality data. Download service for E1a and E2a data, <http://discomap.eea.europa.eu/map/fme/AirQualityExport.htm>, 2019.
- 335 Forster, P. M., Forster, H. I., Evans, M. J., Gidden, M. J., Jones, C. D., Keller, C. A., Lamboll, R. D., Quéré, C. L., Rogelj, J., Rosen, D., Schleussner, C.-F., Richardson, T. B., Smith, C. J., and Turnock, S. T.: Current and future global climate impacts resulting from COVID-19, *Nature Climate Change*, <https://doi.org/10.1038/s41558-020-0883-0>, 2020.
- Grange, S. K.: **rmweather**: Tools to Conduct Meteorological Normalisation on Air Quality Data, <https://CRAN.R-project.org/package=rmweather>, r package version 0.1.2, 2018.
- 340 Grange, S. K.: Technical note: **saqgetr** R package, <https://doi.org/10.13140/RG.2.2.15533.44001>, <https://drive.google.com/file/d/1IgDODHqBHewCTKLdAAXRy7ml8ht6Ods/view>, 2019.
- Grange, S. K.: **isdR**: Import NOAA's Integrated Surface Database (ISD) Data, <https://github.com/skgrange/isdr>, r package, 2020.



- Grange, S. K. and Carslaw, D. C.: Using meteorological normalisation to detect interventions in air quality time series, *Science of the Total Environment*, 653, 578–588, <http://www.sciencedirect.com/science/article/pii/S004896971834244X>, 2019.
- 345 Grange, S. K., Lewis, A. C., Moller, S. J., and Carslaw, D. C.: Lower vehicular primary emissions of NO₂ in Europe than assumed in policy projections, *Nature Geoscience*, 10, 914–918, <https://doi.org/10.1038/s41561-017-0009-0>, 2017.
- Grange, S. K., Carslaw, D. C., Lewis, A. C., Boleti, E., and Hueglin, C.: Random forest meteorological normalisation models for Swiss PM₁₀ trend analysis, *Atmospheric Chemistry and Physics*, 18, 6223–6239, <https://doi.org/https://doi.org/10.5194/acp-18-6223-2018>, <https://www.atmos-chem-phys.net/18/6223/2018/>, 2018.
- 350 Grange, S. K., Hüglin, C., and Emmenegger, L.: Influence of COVID-19 lockdowns on Switzerland’s air quality, https://empa-interim.github.io/empa.interim/swiss_air_quality_and_covid_19.html, empas Laboratory for Air Pollution/Environmental Technology, 2020.
- Hale, T., Webster, S., Petherick, A., Phillips, T., and Kira, B.: Oxford Covid-19 Government Response Tracker (OxCGRT), <https://github.com/OxCGRT/covid-policy-tracker>, blavatnik School of Government, 2020.
- Le Quéré, C., Jackson, R. B., Jones, M. W., Smith, A. J. P., Abernethy, S., Andrew, R. M., De-Gol, A. J., Willis, D. R., Shan, Y., Canadell, J. G., Friedlingstein, P., Creutzig, F., and Peters, G. P.: Temporary reduction in daily global CO₂ emissions during the COVID-19 forced confinement, *Nature Climate Change*, <https://doi.org/10.1038/s41558-020-0797-x>, 2020.
- 355 Lee, J. D., Drysdale, W. S., Finch, D. P., Wilde, S. E., and Palmer, P. I.: UK surface NO₂ levels dropped by 42% during the COVID-19 lockdown: impact on surface O₃, *Atmospheric Chemistry and Physics Discussions*, 2020, 1–27, <https://acp.copernicus.org/preprints/acp-2020-838/>, 2020.
- 360 Lewis, A. C., Hopkins, J. R., Carslaw, D. C., Hamilton, J. F., Nelson, B. S., Stewart, G., Dornie, J., Passant, N., and Murrells, T.: An increasing role for solvent emissions and implications for future measurements of volatile organic compounds, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 378, 20190328, <https://doi.org/10.1098/rsta.2019.0328>, <https://doi.org/10.1098/rsta.2019.0328>, 2020.
- Li, K., Jacob, D. J., Liao, H., Zhu, J., Shah, V., Shen, L., Bates, K. H., Zhang, Q., and Zhai, S.: A two-pollutant strategy for improving ozone and particulate air quality in China, *Nature Geoscience*, 12, 906–910, <https://doi.org/10.1038/s41561-019-0464-x>, <https://doi.org/10.1038/s41561-019-0464-x>, 2019.
- 365 Lindeløv, J. K.: **mcp**: An R Package for Regression With Multiple Change Points, *OSF Preprints*, <https://doi.org/10.31219/osf.io/fzqxv>, 2020.
- Liu, F., Page, A., Strode, S. A., Yoshida, Y., Choi, S., Zheng, B., Lamsal, L. N., Li, C., Krotkov, N. A., Eskes, H., van der A, R., Veefkind, P., 370 Levelt, P. F., Hauser, O. P., and Joiner, J.: Abrupt decline in tropospheric nitrogen dioxide over China after the outbreak of COVID-19, *Science Advances*, <https://doi.org/10.1126/sciadv.abc2992>, <https://advances.sciencemag.org/content/early/2020/06/12/sciadv.abc2992>, 2020.
- Monks, P. S.: A review of the observations and origins of the spring ozone maximum, *Atmospheric Environment*, 34, 3545–3561, <http://www.sciencedirect.com/science/article/pii/S135223100001291>, 2000.
- NOAA: Integrated Surface Database (ISD), <https://www.ncdc.noaa.gov/isd>, 2016.
- 375 Patel, H., Talbot, N., Salmond, J., Dirks, K., Xie, S., and Davy, P.: Implications for air quality management of changes in air quality during lockdown in Auckland (New Zealand) in response to the 2020 SARS-CoV-2 epidemic, *Science of The Total Environment*, 746, 141–129, <http://www.sciencedirect.com/science/article/pii/S0048969720346581>, 2020.
- Remuzzi, A. and Remuzzi, G.: COVID-19 and Italy: what next?, *The Lancet*, [https://doi.org/10.1016/S0140-6736\(20\)30627-9](https://doi.org/10.1016/S0140-6736(20)30627-9), [https://doi.org/10.1016/S0140-6736\(20\)30627-9](https://doi.org/10.1016/S0140-6736(20)30627-9), 2020.



- 380 Ruktanonchai, N. W., Floyd, J. R., Lai, S., Ruktanonchai, C. W., Sadilek, A., Rente-Lourenco, P., Ben, X., Carioli, A., Gwinn, J., Steele, J. E., Prosper, O., Schneider, A., Oplinger, A., Eastham, P., and Tatem, A. J.: Assessing the impact of coordinated COVID-19 exit strategies across Europe, *Science*, <https://doi.org/10.1126/science.abc5096>, <https://science.sciencemag.org/content/early/2020/07/16/science.abc5096>, 2020.
- Sillman, S.: The relation between ozone, NO_x and hydrocarbons in urban and polluted rural environments, *Atmospheric Environment*, 33, 1821–1845, <http://www.sciencedirect.com/science/article/pii/S1352231098003458>, 1999.
- 385 Venter, Z. S., Aunan, K., Chowdhury, S., and Lelieveld, J.: COVID-19 lockdowns cause global air pollution declines, *Proceedings of the National Academy of Sciences*, 117, 18 984–18 990, <https://doi.org/10.1073/pnas.2006853117>, <https://www.pnas.org/content/117/32/18984>, 2020.
- Wang, T., Xue, L., Brimblecombe, P., Lam, Y. F., Li, L., and Zhang, L.: Ozone pollution in China: A review of concentrations, meteorological influences, chemical precursors, and effects, *Science of The Total Environment*, 575, 1582–1596, <https://doi.org/10.1016/j.scitotenv.2016.10.081>, <http://www.sciencedirect.com/science/article/pii/S0048969716322471>, 2017.
- 390 Wang, Y., Wen, Y., Wang, Y., Zhang, S., Zhang, K. M., Zheng, H., Xing, J., Wu, Y., and Hao, J.: Four-Month Changes in Air Quality during and after the COVID-19 Lockdown in Six Megacities in China, *Environmental Science and Technology Letters*, <https://doi.org/10.1021/acs.estlett.0c00605>, <https://doi.org/10.1021/acs.estlett.0c00605>, 2020.
- 395 Williams, M. L., Atkinson, R. W., Anderson, H. R., and Kelly, F. J.: Associations between daily mortality in London and combined oxidant capacity, ozone and nitrogen dioxide, *Air Quality, Atmosphere & Health*, 7, 407–414, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4239710/>, 2014.
- World Health Organization (WHO): Pneumonia of unknown cause—China, <https://www.who.int/csr/don/05-january-2020-pneumonia-of-unkown-cause-china/en/>, disease outbreak news. 5 January 2020, 2020a.
- 400 World Health Organization (WHO): WHO Director-General’s opening remarks at the media briefing on COVID-19—11 March 2020, <https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020>, 11 March 2020, 2020b.
- World Health Organization (WHO): Naming the coronavirus disease (COVID-19) and the virus that causes it, [https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/naming-the-coronavirus-disease-\(covid-2019\)-and-the-virus-that-causes-it](https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/naming-the-coronavirus-disease-(covid-2019)-and-the-virus-that-causes-it), disease: coronavirus disease (COVID-19). Virus: severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), 2020c.
- 405 Wu, J. T., Leung, K., and Leung, G. M.: Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study, *The Lancet*, [https://doi.org/10.1016/s0140-6736\(20\)30260-9](https://doi.org/10.1016/s0140-6736(20)30260-9), [https://doi.org/10.1016/S0140-6736\(20\)30260-9](https://doi.org/10.1016/S0140-6736(20)30260-9), 2020.
- 410 Zhou, P., Yang, X.-L., Wang, X.-G., Hu, B., Zhang, L., Zhang, W., Si, H.-R., Zhu, Y., Li, B., Huang, C.-L., Chen, H.-D., Chen, J., Luo, Y., Guo, H., Jiang, R.-D., Liu, M.-Q., Chen, Y., Shen, X.-R., Wang, X., Zheng, X.-S., Zhao, K., Chen, Q.-J., Deng, F., Liu, L.-L., Yan, B., Zhan, F.-X., Wang, Y.-Y., Xiao, G.-F., and Shi, Z.-L.: A pneumonia outbreak associated with a new coronavirus of probable bat origin, *Nature*, 579, 270–273, <https://doi.org/10.1038/s41586-020-2012-7>, 2020.