



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/172583/>

Version: Accepted Version

Article:

Jerosch-Herold, C, Bland, JDP and Horton, M (2021) Reply to: Revisiting the Functional Status Scale of the Boston Carpal Tunnel Questionnaire, a comment concerning the use of digital communication technology. *Muscle & Nerve*, 63 (6). E58-E59. ISSN: 0148-639X

<https://doi.org/10.1002/mus.27227>

© 2021 Wiley Periodicals LLC. This is the peer reviewed version of the following article: Jerosch-Herold, C, Bland, JDP and Horton, M (2021) Reply to: Revisiting the Functional Status Scale of the Boston Carpal Tunnel Questionnaire, a comment concerning the use of digital communication technology. *Muscle & Nerve*, 63 (6). E58-E59. ISSN 0148-639X, which has been published in final form at <https://doi.org/10.1002/mus.27227>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Reply to: Revisiting the Functional Severity Scale of the Boston Carpal Tunnel

Questionnaire, a comment concerning the use of digital communication technology.

Christina Jerosch-Herold

Faculty of Medicine and Health Sciences, University of East Anglia, Norwich Research Park,

Norwich, UK

Email: c.jerosch-herold@uea.ac.uk

ORCID 0000-0003-0525-1282

Jeremy DP Bland

Dept of Neurophysiology

East Kent Hospitals University NHS Foundation Trust

Canterbury, Kent, CT1 3NG

Email: Jeremy.Bland@nhs.net

ORCID 0000-0001-5591-6022

Mike Horton

Psychometric Laboratory for Health Sciences, Leeds Institute of Rheumatic and

Musculoskeletal Medicine, University of Leeds, UK.

E-mail: m.c.horton@leeds.ac.uk

ORCID 0000-0002-6675-7335

Corresponding author:

C Jerosch-Herold

School of Health Sciences

Faculty of Medicine and Health Sciences

University of East Anglia

Norwich

NR4 7TJ

Declaration of conflicting interests: The author(s) declare no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

We confirm that we have read the Journal's position on issues involved in ethical publication and affirm that this report is consistent with those guidelines."

517 words

Authors' reply:

We thank Lattré and van Holder for their letter in response to our article¹.

It is true that we live in a much more digitalised world than that in which Levine et al² first developed the BCTQ. Therefore, questions about writing and using a telephone in the Functional Status Scale (FSS) may hold a very different meaning today, along with a different operationalisation within the FSS.

Lattré and van Holder present research about possible associations between keyboard and mobile device use for writing or communicating and structural changes to the median nerve. However, the current literature on acute changes in median nerve morphology in response to activity, as assessed by ultrasound imaging, suffers from serious methodological problems with observer blinding and limitations posed by the resolving power of current ultrasound systems such that the results they describe cannot be accepted as proven so far. Also these factors are of very limited relevance to the validity of the FSS.

The crux of our argument centred around the content validity of FSS, that is, the question of whether the items that make up the FSS are relevant for assessing the functional impact of carpal tunnel syndrome (CTS), irrespective of proven or hypothesized associations. It is unclear from Levine et al's original paper² how the concepts and questionnaire items were first generated, and only by undertaking cognitive interviews³ with patients could we really begin to understand how they interpret these items in today's context.

Lattré and van Holder's proposal to revise the FSS by adding questions about digital device use and activities risks oversimplifying the process involved in patient reported outcome (PRO) instrument development, and we would urge some caution for the following reasons:

Firstly, a complete revision of the FSS should ideally follow best practice as advocated by ISPOR^{4,5} and the FDA's Guidance toward standardizing methods for developing patient-reported outcome (PRO) instruments which emphasises the importance of directly eliciting the patient perspective through qualitative research⁶. The new FSS would then have to

undergo full testing of its content, construct and criterion-validity. This is no small undertaking and a process that often takes years.

Secondly, the total score from a revised FSS would not be directly comparable to the total score of the current FSS, as the scale would be comprised of different items. Although it would be theoretically possible to co-calibrate and equate the two versions, the risk remains of these FSS scores being treated as directly equivalent.

Finally and given the above caveats, we propose as an alternative that existing and more contemporary PROs which measure the impact on hand use in daily activities are chosen based on the strongest evidence for content and construct validity, thus complementing the symptom severity scale. For example in a recent trial on sensory relearning after surgery for CTS we used 3 subscales of the Michigan Hand Questionnaire (MHQ) and which showed statistically significant improvements in unilateral and bilateral hand activities⁷. A further advantage being that the MHQ generates separate scores for the left and right hand which is particularly important in CTS where it can have a unilateral or bilateral presentation.

References:

1. Jerosch-Herold C, Bland JDP, Horton M. Is it time to revisit the Boston Carpal Tunnel Questionnaire? New insights from a Rasch model analysis. *Muscle & Nerve*; 2021; 1–6
2. Levine DW, Simmons BP, Koris MJ, Daltroy LH, Hohl GG, Fossel AH, Katz JN. A self-administered questionnaire for the assessment of severity of symptoms and

functional status in carpal tunnel syndrome. *J Bone Joint Surg Am* 1993; 75(11):1585-1592.

3. Willis, G. B. (2004). *Cognitive interviewing: A tool for improving questionnaire design*. Thousand Oaks: Sage Publications
4. Patrick DL, Burke LB, Gwaltney CJ, Leidy NK, Martin ML, Molsen E, Ring L. Content validity--establishing and reporting the evidence in newly developed patient-reported outcomes (PRO) instruments for medical product evaluation: ISPOR PRO good research practices task force report: part 1--eliciting concepts for a new PRO instrument. *Value Health*. 2011; 14(8):967-77.
5. Patrick DL, Burke LB, Gwaltney CJ, Leidy NK, Martin ML, Molsen E, Ring L. Content validity--establishing and reporting the evidence in newly developed patient-reported outcomes (PRO) instruments for medical product evaluation: ISPOR PRO Good Research Practices Task Force report: part 2--assessing respondent understanding. *Value Health*. 2011; 14(8):978-88.
6. US Food and Drug Administration Guidance for Industry Patient-Reported Outcome Measures: Use in Medical Product Development to Support Labeling Claims; Department of Health and Human Services, Food and Drug Administration, Center for Drug Evaluation and Research, Rockville, MD (2009)
7. Jerosch-Herold C, Houghton J, Miller L, Shepstone L. Does sensory relearning improve tactile function after carpal tunnel decompression? A pragmatic, assessor-blinded, randomized clinical trial. *Journal of Hand Surgery (European Volume)*. 2016; 41(9):948-956.