



This is a repository copy of *Comment on: Fraser et al. The Effects of Long-Term Oral Benfotiamine Supplementation on Peripheral Nerve Function and Inflammatory Markers in Patients With Type 1 Diabetes: A 24-Month, Double-Blind, Randomized, Placebo-Controlled Trial. Diabetes Care 2012;35:1095-1097.*

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/174189/>

Version: Published Version

Article:

Ziegler, D., Tesfaye, S. orcid.org/0000-0003-1190-1472 and Kempler, P. (2012) Comment on: Fraser et al. The Effects of Long-Term Oral Benfotiamine Supplementation on Peripheral Nerve Function and Inflammatory Markers in Patients With Type 1 Diabetes: A 24-Month, Double-Blind, Randomized, Placebo-Controlled Trial. *Diabetes Care* 2012;35:1095-1097. *Diabetes Care*, 35 (11). E79-E79. ISSN 0149-5992

<https://doi.org/10.2337/dc12-0817>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

 COMMENTS AND
 RESPONSES

Comment on: Fraser et al. The Effects of Long-Term Oral Benfotiamine Supplementation on Peripheral Nerve Function and Inflammatory Markers in Patients With Type 1 Diabetes: A 24-Month, Double-Blind, Randomized, Placebo-Controlled Trial. Diabetes Care 2012;35:1095-1097

Fraser et al. (1) reported that “high-dose benfotiamine (300 mg/day) supplementation over 24 months has no significant effects upon peripheral nerve function or soluble markers of inflammation in patients with type 1 diabetes.” Although this conclusion may be correct based on these findings, the clinical relevance remains obscure in view of the questionable study design. According to the recent consensus statement of the Toronto Diabetic Neuropathy Expert Group (2), a possible diabetic sensorimotor polyneuropathy (DSPN) can be diagnosed if symptoms or signs of DSPN are present. In the case of probable DSPN, a combination of symptoms and signs of neuropathy is present. The authors state that among the patients included, 56% had subclinical neuropathy based on abnormal nerve conduction studies (NCS), and 16% had probable DSPN. Unfortunately, how abnormal NCS was defined is unclear. In fact, this high percentage is surprising given that mean sural sensory nerve conduction velocity (NCV),

which is one of the most sensitive indicators of DSPN, was well within the normal range. Moreover, clinical neuropathy scores have been used to assume probable DSPN, but obviously the latter has not been confirmed by abnormal NCS. Because the number of subjects included with confirmed DSPN was presumably very low, this trial does not provide any information on the important question as to whether treatment with benfotiamine may improve DSPN or slow its progression. The authors note that in previous studies using benfotiamine, no effect on NCV was shown. In the study by Stracke et al. (3) cited by the authors, improvement in peroneal motor NCV in diabetic patients with DSPN was documented.

Further drawbacks of this trial are the lack of an intention-to-treat analysis to confirm the per protocol data and of an a priori-specified clinically meaningful difference in peroneal motor NCV between the two groups studied.

Recently, it has been suggested that the progression of DSPN in the placebo group within the setting of a randomized clinical trial over 4-year period was slow. Thus, for demonstration of efficacy, trials using drugs for treatment of DSPN must achieve a clinically relevant neurological improvement (4). Unfortunately, the study conducted by Fraser et al. (1) in patients with almost normal nerve function does not leave any room for a meaningful degree of improvement.

The title reads “Benfotiamine Supplementation,” but the dose applied (300 mg/day) was much higher than the recommended daily adult allowance for thiamine (1.4 mg/day). Hence, this was pharmacotherapy rather than vitamin supplementation.

Finally, as far as it can be judged from the article, the patients examined had no chronic inflammation. Therefore, it was foreseeable that no effect on inflammatory markers was shown. In conclusion, it is not surprising that 2-year treatment with benfotiamine versus placebo did not result in meaningful differences between the groups in nerve function or markers of inflammation in patients without evidence of neuropathy nor inflammation. Only trials using appropriate study

designs and end points can assess the effects of benfotiamine on nerve dysfunction resulting from DSPN.

DAN ZIEGLER, MD^{1,2}
 SOLOMON TESFAYE, MD³
 PETER KEMPLER, MD⁴

From the ¹Institute for Clinical Diabetology, German Diabetes Center at Heinrich Heine University, Leibniz Center for Diabetes Research, Düsseldorf, Germany; the ²Department of Metabolic Diseases, University Hospital, Düsseldorf, Germany; the ³Diabetes Research Unit, Sheffield Teaching Hospitals, Sheffield, U.K.; and the ⁴1st Department of Medicine, Semmelweis University, Budapest, Hungary.

Corresponding author: Dan Ziegler, dan.ziegler@ddz.uni-duesseldorf.de.

DOI: 10.2337/dc12-0817

© 2012 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Acknowledgments—D.Z. and P.K. have received honoraria from Wörwag Pharma for speaking. D.Z., P.K., and S.T. have received honoraria from Wörwag Pharma for consulting activities. No other potential conflicts of interest relevant to this article were reported.

.....

References

1. Fraser DA, Diep LM, Hovden IA, et al. The effects of long-term oral benfotiamine supplementation on peripheral nerve function and inflammatory markers in patients with type 1 diabetes: a 24-month, double-blind, randomized, placebo-controlled trial. *Diabetes Care* 2012;35:1095-1097
2. Tesfaye S, Boulton AJM, Dyck PJ, et al.; Toronto Diabetic Neuropathy Expert Group. Diabetic neuropathies: update on definitions, diagnostic criteria, estimation of severity, and treatments. *Diabetes Care* 2010;33:2285-2293
3. Stracke H, Lindemann A, Federlin K. A benfotiamine-vitamin B combination in treatment of diabetic polyneuropathy. *Exp Clin Endocrinol Diabetes* 1996;104:311-316
4. Ziegler D, Low PA, Litchy WJ, et al. Efficacy and safety of antioxidant treatment with α -lipoic acid over 4 years in diabetic polyneuropathy: the NATHAN 1 trial. *Diabetes Care* 2011;34:2054-2060