



UNIVERSITY OF LEEDS

This is a repository copy of *Parameterized Pre-Coloring Extension and List Coloring Problems*.

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/175347/>

Version: Accepted Version

Article:

Gutin, G, Majumdar, D, Ordyniak, S orcid.org/0000-0003-1935-651X et al. (1 more author) (2021) Parameterized Pre-Coloring Extension and List Coloring Problems. *SIAM Journal on Discrete Mathematics*, 35 (1). pp. 575-596. ISSN 0895-4801

<https://doi.org/10.1137/20m1323369>

© 2021, Society for Industrial and Applied Mathematics. This is an author produced version of an article published in *SIAM Journal on Discrete Mathematics*. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

1 **PARAMETERIZED PRE-COLORING EXTENSION AND LIST**
2 **COLORING PROBLEMS***

3 GREGORY GUTIN[†], DIPTAPRIYO MAJUMDAR[†], SEBASTIAN ORDYNIAK[‡], AND
4 MAGNUS WAHLSTRÖM[†]

5 **Abstract.** Golovach, Paulusma and Song (Inf. Comput. 2014) asked to determine the param-
6 eterized complexity of the following problems parameterized by k : (1) Given a graph G , a clique
7 modulator D (a *clique modulator* is a set of vertices, whose removal results in a clique) of size k for G ,
8 and a list $L(v)$ of colors for every $v \in V(G)$, decide whether G has a proper list coloring; (2) Given a
9 graph G , a clique modulator D of size k for G , and a pre-coloring $\lambda_P : X \rightarrow Q$ for $X \subseteq V(G)$, decide
10 whether λ_P can be extended to a proper coloring of G using only colors from Q . For Problem 1 we
11 design an $\mathcal{O}^*(2^k)$ -time randomized algorithm and for Problem 2 we obtain a kernel with at most
12 $3k$ vertices. Banik et al. (IWOC 2019) proved the following problem is fixed-parameter tractable
13 and asked whether it admits a polynomial kernel: Given a graph G , an integer k , and a list $L(v)$
14 of exactly $n - k$ colors for every $v \in V(G)$, decide whether there is a proper list coloring for G . We
15 obtain a kernel with $\mathcal{O}(k^2)$ vertices and colors and a compression to a variation of the problem with
16 $\mathcal{O}(k)$ vertices and $\mathcal{O}(k^2)$ colors.

17 **1. Introduction.** Graph coloring is a central topic in Computer Science and
18 Graph Theory due to its importance in theory and applications. Every text book
19 in Graph Theory has at least a chapter devoted to the topic and the monograph
20 of Jensen and Toft [25] is completely devoted to graph coloring problems focusing
21 especially on more than 200 unsolved ones. There are many survey papers on the
22 topic including recent ones such as [13, 22, 31, 33].

23 For a graph G , a *proper coloring* is a function $\lambda : V(G) \rightarrow \mathbb{N}_{\geq 1}$ such that for
24 no pair u, v of adjacent vertices of G , $\lambda(u) = \lambda(v)$. In the widely studied COLORING
25 problem, given a graph G and a positive integer p , we are to decide whether there is a
26 proper coloring $\lambda : V(G) \rightarrow [p]$, where henceforth $[p] = \{1, \dots, p\}$. In this paper, we
27 consider two extensions of COLORING: the PRE-COLORING EXTENSION problem and
28 the LIST COLORING problem. In the PRE-COLORING EXTENSION problem, given a
29 graph G , a set Q of colors, and a *pre-coloring* $\lambda_P : X \rightarrow Q$, where $X \subseteq V(G)$, we are
30 to decide whether there is a proper coloring $\lambda : V(G) \rightarrow Q$ such that $\lambda(x) = \lambda_P(x)$
31 for every $x \in X$. In the LIST COLORING problem, given a graph G and a list $L(u)$
32 of possible colors for every vertex u of G , we are to decide whether G has a proper
33 coloring λ such that $\lambda(u) \in L(u)$ for every vertex u of G . Such a coloring λ is called
34 a *proper list coloring*. Clearly, PRE-COLORING EXTENSION is a special case of LIST
35 COLORING, where all lists of vertices $x \in X$ are singletons and the lists of all other
36 vertices are equal to Q .

37 The p -COLORING problem is a special case of COLORING when p is fixed (i.e., not
38 part of input). When $Q \subseteq [p]$ ($L(u) \subseteq [p]$, respectively), PRE-COLORING EXTENSION
39 (LIST COLORING, respectively) are called p -PRE-COLORING EXTENSION (LIST p -
40 COLORING, respectively). In classical complexity, it is well-known that p -COLORING,
41 p -PRE-COLORING EXTENSION and LIST p -COLORING are polynomial-time solvable
42 for $p \leq 2$, and the three problems become NP-complete for every $p \geq 3$ [28, 31]. In this
43 paper, we solve several open problems about pre-coloring extension and list coloring
44 problems, which lie outside classical complexity, so-called parameterized problems.

*A preliminary and shortened version of this paper has been accepted at STACS 2020.

[†]Royal Holloway, University of London, UK (gutin@cs.rhul.ac.uk, dip-
tapriyo.majumdar@rhul.ac.uk, Magnus.Wahlstrom@rhul.ac.uk)

[‡]University of Leeds, UK (sordyniak@gmail.com)

45 We provide basic notions on parameterized complexity in the next section. For more
 46 information on parameterized complexity, see recent books [14, 18, 20].

47 The first two problems we study are the following ones stated by Golovach et al.
 48 [23] (see also [30]) who asked to determine their parameterized complexity. These
 49 questions were motivated by a result of Cai [10] who showed that COLORING WITH
 50 CLIQUE MODULATOR (the special case of PRE-COLORING EXTENSION WITH CLIQUE
 51 MODULATOR when $X = \emptyset$) is fixed-parameter tractable (FPT). Note that a *clique*
 52 *modulator* of a graph G is a set D of vertices such that $G - D$ is a clique. When using
 53 the size of a clique modulator as a parameter we will for convenience assume that the
 54 modulator is given as part of the input. Note that this assumption is not necessary
 55 (however it avoids having to repeat how to compute a clique modulator) as we will
 56 show in Section 2 that computing a clique modulator of size k is FPT and can be
 57 approximated to within a factor of two.

58 LIST COLORING WITH CLIQUE MODULATOR parameterized by k

Input: A graph G , a clique modulator D of size k for G , and a list $L(v)$ of
 colors for every $v \in V(G)$.

Problem: Is there a proper list coloring for G ?

59 PRE-COLORING EXTENSION WITH CLIQUE MODULATOR parameterized by k

Input: A graph G , a clique modulator D of size k for G , and a pre-coloring
 $\lambda_P : X \rightarrow Q$ for $X \subseteq V(G)$ where Q is a set of colors.

Problem: Can λ_P be extended to a proper coloring of G using only colors from
 Q ?

62
 63 In Section 3 we show that LIST COLORING WITH CLIQUE MODULATOR is FPT.
 64 We first show a randomized $\mathcal{O}^*(2^{k \log k})$ -time algorithm, then we improve the running
 65 time to $\mathcal{O}^*(2^k)$ using more refined tools and approaches. Note that all our random-
 66 ized algorithms are one-sided error algorithms having a constant probability of being
 67 wrong, when the algorithm outputs no.

68 We note that the time $\mathcal{O}^*(2^k)$ matches the best known running time of $\mathcal{O}^*(2^n)$
 69 for CHROMATIC NUMBER (where $n = |V(G)|$) [6], while applying to a more powerful
 70 parameter. It is a long-open problem whether CHROMATIC NUMBER can be solved
 71 in time $\mathcal{O}(2^{cn})$ for some $c < 1$ and Cygan et al. [15] ask whether it is possible to
 72 show that such algorithms are impossible assuming the Strong Exponential Time
 73 Hypothesis (SETH).

74 We conclude Section 3 by showing that LIST COLORING WITH CLIQUE MODU-
 75 LATOR does not admit a polynomial kernel unless $\text{NP} \subseteq \text{coNP/poly}$. The reduction
 76 used to prove this result allows us to observe that if LIST COLORING WITH CLIQUE
 77 MODULATOR could be solved in time $\mathcal{O}(2^{ck} n^{\mathcal{O}(1)})$ for some $c < 1$, then the well-
 78 known SET COVER problem could be solved in time $\mathcal{O}(2^{c|U|} |\mathcal{F}|^{\mathcal{O}(1)})$, where U and \mathcal{F}
 79 are universe and family of subsets, respectively. The existence of such an algorithm
 80 is open, and it has been conjectured that no such algorithm is possible under SETH;
 81 see Cygan et al. [15]. Thus, up to the assumption of this conjecture (called *Set Cover*
 82 *Conjecture* [27]) and SETH, our $\mathcal{O}^*(2^k)$ -time algorithm for LIST COLORING WITH
 83 CLIQUE MODULATOR is best possible w.r.t. its dependency on k .

84 In Section 4, we consider PRE-COLORING EXTENSION WITH CLIQUE MODULA-
 85 TOR, which is a subproblem of LIST COLORING WITH CLIQUE MODULATOR and prove
 86

87 that PRE-COLORING EXTENSION WITH CLIQUE MODULATOR, unlike LIST COLOR-
 88 ING WITH CLIQUE MODULATOR, admits a polynomial kernel: a linear kernel with at
 89 most $3k$ vertices. This kernel builds on a known, but counter-intuitive property of
 90 bipartite matchings (see Proposition 2.2), which was previously used in kernelization
 91 by Bodlaender et al. [8].

92 In Section 5, we study an open problem stated by Banik et al. [3]. In a classic
 93 result, Chor et al. [12] showed that COLORING has a linear vertex kernel parameterized
 94 by $k = n - p$, i.e., if the task is to “save k colors”. Arora et al. [2] consider the following
 95 as a natural extension to list coloring, and show that it is in XP. Banik et al. [3] show
 96 that the problem is FPT, but leave as an open question whether it admits a polynomial
 97 kernel.

98 $(n - k)$ -REGULAR LIST COLORING parameterized by k

Input: A graph G on n vertices, an integer k , and a list $L(v)$ of exactly $n - k$
 colors for every $v \in V(G)$.

Problem: Is there a proper list coloring for G ?

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

We answer this question in affirmative by giving a kernel with $\mathcal{O}(k^2)$ vertices and
 colors, as well as a compression to a variation of the problem with $\mathcal{O}(k)$ vertices,
 encodable in $\mathcal{O}(k^2 \log k)$ bits. We note that this compression is asymptotically almost
 tight, as even 4-COLORING does not admit a compression into $\mathcal{O}(n^{2-\varepsilon})$ bits for any
 $\varepsilon > 0$ unless the polynomial hierarchy collapses [24].

This kernel is more intricate than the above. Via known reduction rules from
 Banik et al. [3], we can compute a clique modulator of at most $2k$ vertices (hence our
 result for LIST COLORING WITH CLIQUE MODULATOR also solves $(n - k)$ -REGULAR
 LIST COLORING in $2^{\mathcal{O}(k)}$ time). However, the usual “crown rules” (as in [12] and
 in Section 4) are not easily applied here, due to complications with the color lists.
 Instead, we are able to show a set of $\mathcal{O}(k)$ vertices whose colorability make up the
 “most interesting” part of the problem, leading to the above-mentioned compression
 and kernel.

In Section 6, we consider further natural pre-coloring and list coloring variants
 of the “saving k colors” problem of Chor et al. [12]. We show that the known fixed-
 parameter tractability and linear kernelizability [12] carries over to a natural pre-
 coloring generalization but fails for a more general list coloring variant. Since $(n - k)$ -
 REGULAR LIST COLORING was originally introduced in [2] as a list coloring variant
 of the “saving k colors” problem, it is natural to consider other such variants. We
 conclude the paper in Section 7, where in particular a number of open questions are
 discussed.

2. Preliminaries.

123

124

125

126

127

128

129

130

131

2.1. Parameterized Complexity. An instance of a parameterized problem Π
 is a pair (I, k) where I is the *main part* and k is the *parameter*; the latter is usually a
 non-negative integer. A parameterized problem is *fixed-parameter tractable* (FPT) if
 there exists a computable function f such that instances (I, k) can be solved in time
 $\mathcal{O}(f(k)|I|^c)$ where $|I|$ denotes the size of I and c is an absolute constant. The class
 of all fixed-parameter tractable decision problems is called FPT and algorithms which
 run in the time specified above are called FPT algorithms. As in other literature on
 FPT algorithms, we will often omit the polynomial factor in $\mathcal{O}(f(k)|I|^c)$ and write
 $\mathcal{O}^*(f(k))$ instead. To establish that a problem under a specific parameterization is

not in FPT we prove that it is $W[1]$ -hard as it is widely believed that $FPT \neq W[1]$.

A *reduction rule* R for a parameterized problem Π is an algorithm A that given an instance (I, k) of a problem Π returns an instance (I', k') of the *same* problem. The reduction rule is said to be *safe* if it holds that $(I, k) \in \Pi$ if and only if $(I', k') \in \Pi$. If A runs in polynomial time in $|I| + k$ then R is a *polynomial-time reduction rule*. Often we omit the adjectives “safe” and “polynomial-time” in “safe polynomial-time reduction rule” as we consider only such reduction rules.

A *kernelization* (or, a *kernel*) of a parameterized problem Π is a reduction rule such that $|I'| + k' \leq f(k)$ for some computable function f . It is not hard to show that a decidable parameterized problem is FPT if and only if it admits a kernel [14, 18, 20]. The function f is called the *size* of the kernel, and we have a *polynomial kernel* if $f(k)$ is polynomially bounded in k .

A kernelization can be generalized by considering a reduction (rule) from a parameterized problem Π to another parameterized problem Π' . Then instead of a kernel we obtain a *generalized kernel* (also called a *bikernel* [1] in the literature). If the problem Π' is not parameterized, then a reduction from Π to Π' (i.e., (I, k) to I') is called a *compression*, which is *polynomial* if $|I'| \leq p(k)$, where p is a fixed polynomial in k . If there is a polynomial compression from Π to Π' and Π' is polynomial-time reducible back to Π , with a reduction $I' \mapsto (I, k)$ such that furthermore $k \leq |I'|^{O(1)}$, then combining the compression with the reduction gives a polynomial kernel for Π .

2.2. Graphs, Matchings, and Clique Modulator. We consider finite simple undirected graphs. For basic terminology on graphs, we refer to a standard textbook [16]. For an undirected graph $G = (V, E)$ we denote by $V(G)$ the vertex set of G and by $E(G)$ the edge set of G . For a vertex $v \in V(G)$, we denote by $N_G(v)$ and $N_G[v]$ the *open* respectively *closed neighborhood* of v in G , i.e., $N_G(v) := \{u \mid \{u, v\} \in E(G)\}$ and $N_G[v] := N_G(v) \cup \{v\}$. We extend this notion in the natural manner to subsets $V' \subseteq V(G)$, by setting $N_G(V') := \bigcup_{v \in V'} N_G(v)$ and $N_G[V'] := \bigcup_{v \in V'} N_G[v]$. Moreover, we omit the subscript G , if the graph G can be inferred from the context. If $V' \subseteq V(G)$, we denote by $G \setminus V'$ the graph obtained from G after deleting all vertices in V' together with their adjacent edges and we denote by $G[V']$ the graph induced by the vertices in V' , i.e., $G[V'] = G \setminus (V(G) \setminus V')$. We say that G is *bipartite with bi-partition* (A, B) , if $\{A, B\}$ partitions $V(G)$ and $G[A]$ as well as $G[B]$ have no edges.

A *matching* M is a subset of $E(G)$ such that no two edges in M share a common endpoint. We say that M is *maximal* if there is no edge $e \in E(G)$ such that $M \cup \{e\}$ is a matching and we say that M is *maximum* if it is maximal and there is no maximal matching in G containing more edges than M . We denote by $V(M)$ the set of all endpoints of the edges in M , i.e., the set $\bigcup_{e \in M} e$. We say that M *saturates* a subset $V' \subseteq V(G)$ if $V' \subseteq V(M)$. Let $H = (V, E)$ be an undirected bipartite graph with bi-partition (A, B) . We say that a set C is a *Hall set* for A or B if $C \subseteq A$ or $C \subseteq B$, respectively, and $|N_H(C)| < |C|$. We will need the following well-known properties for matchings.

PROPOSITION 2.1 (Hall’s Theorem [16]). *Let G be an undirected bipartite graph with bi-partition (A, B) . Then G has a matching saturating A if and only if there is no Hall set for A , i.e., for every $A' \subseteq A$, it holds that $|N(A')| \geq |A'|$.*

PROPOSITION 2.2 ([8, Theorem 2]). *Let G be a bipartite graph with bi-partition (X, Y) and let X_M be the set of all vertices in X that are endpoints of a maximum matching M of G . Then, for every $Y' \subseteq Y$, it holds that G contains a matching that*

180 covers Y' if and only if so does $G[X_M \cup Y]$.

181 **Clique Modulator** Let G be an undirected graph. We say that a set $D \subseteq V(G)$ is
 182 a *clique modulator* for G if $G - D$ is a clique. Since we will use the size of a smallest
 183 clique modulator as a parameter for our coloring problems, it is natural to ask whether
 184 the following problem can be solved efficiently.

185 CLIQUE MODULATOR parameterized by k
Input: A graph G and an integer k
Problem: Does G have a clique modulator of size at most k ?

186
 187 The following proposition shows that this is indeed the case. Namely, CLIQUE
 188 MODULATOR is both FPT and can be approximated within a factor of two. The former
 189 is important for our FPT algorithms and the later for our kernelization algorithms as
 190 it allows us to not depend on a clique modulator given as part of the input.

192 PROPOSITION 2.3. CLIQUE MODULATOR is fixed-parameter tractable (in time
 193 $\mathcal{O}^*(1.2738^k)$) and can be approximated within a factor of two.

194 *Proof.* It is straightforward to verify that a graph G has a clique modulator of
 195 size at most k if and only if the complement \overline{G} of G has a vertex cover of size at
 196 most k . The statement now follows from the fact that the vertex cover problem is
 197 fixed-parameter tractable [11] (in time $\mathcal{O}^*(1.2738^k)$) and can be approximated within
 198 a factor of two [21]. □

199 **2.3. Polynomial sieving.** Algorithms based on polynomial sieving and sim-
 200 ilar algebraic techniques have become an important component of the toolbox for
 201 parameterized and exact algorithms. One of the early examples within the field is
 202 the algorithm for computing CHROMATIC NUMBER in time $\mathcal{O}^*(2^n)$ by Björklund et
 203 al. [6]. Further developments include techniques such as *multilinear detection* [26]
 204 (see also [7]). We review only what we need for this paper; for more background and
 205 further techniques, see [15, 26, 7, 5].

206 For a positive integer p , $[p]$ denotes the set $\{1, 2, \dots, p\}$. For a polynomial P , we
 207 denote the coefficient of a monomial T of P by $\text{coef}_P T$.

208 The following lemma is central to the approach.

209 LEMMA 2.4. (Schwartz-Zippel [32, 36]). Let $P(x_1, \dots, x_n)$ be a multivariate poly-
 210 nomial of total degree at most d over a field \mathbb{F} , and assume that P is not identically
 211 zero. Pick r_1, \dots, r_n uniformly at random from \mathbb{F} . Then $\Pr[P(r_1, \dots, r_n) = 0] \leq$
 212 $d/|\mathbb{F}|$.

213 The general approach is to construct a polynomial whose terms enumerate po-
 214 tential solutions, and then use sieving techniques over the polynomial to ensure that
 215 undesired solutions cancel and only actual solutions remain. As long as the sieved
 216 polynomial can be evaluated in FPT time, this then gives a randomized FPT algo-
 217 rithm using the Schwartz-Zippel lemma, as above. In the case that we are working
 218 over a field of characteristic 2, we will implicitly assume that the field is large enough
 219 to allow an application of the above lemma with good success probability, e.g., by
 220 moving to an extension field or starting with a large enough field $\text{GF}(2^\ell)$.

221 We will use the following simple inclusion-exclusion based sieving technique, pre-
 222 viously used by Wahlström [34]. Let $P(x_1, \dots, x_n)$ be a polynomial and $I \subseteq [n]$ a set
 223 of indices. Define $P_{-I}(x_1, \dots, x_n) = P(y_1, \dots, y_n)$, where $y_i = 0$ for $i \in I$ and $y_i = x_i$
 224 otherwise. Then the following holds. (The variant for a field of characteristic 2 was

225 proved by Wahlström [34]. The other variant can be proved similarly.)

226 LEMMA 2.5. *Let $P(x_1, \dots, x_n)$ be a polynomial over a field of characteristic two*
 227 *(over reals, respectively), and $J \subseteq [n]$ a set of indices. Define*

$$228 \quad Q(x_1, \dots, x_n) = \sum_{I \subseteq J} P_{-I}(x_1, \dots, x_n)$$

$$229 \quad (Q(x_1, \dots, x_n) = \sum_{I \subseteq J} (-1)^{|I|} P_{-I}(x_1, \dots, x_n), \text{ respectively}).$$

230 *Then for any monomial T divisible by $\prod_{i \in J} x_i$ we have $\text{coef}_Q T = \text{coef}_P T$, and for*
 231 *every other monomial T we have $\text{coef}_Q T = 0$.*

232 We will also use the connection between permanents and bipartite matchings. Let
 233 G be a bipartite graph with balanced bi-partition (U, V) , i.e., $|U| = |V|$. The bipartite
 234 adjacency matrix of G is a matrix A , with rows are indexed by U and columns indexed
 235 by V , such that $A[u, v] = 1$ for $u \in U, v \in V$ if $uv \in E(G)$, and $A[u, v] = 0$ otherwise.
 236 It is well known that the permanent per A enumerates perfect matchings of G , but
 237 that it is hard to evaluate in general. The exception is in fields of characteristic 2,
 238 where it coincides with the determinant, but where we furthermore have to worry
 239 about cancellations due to the characteristic.

240 In order to work with determinants instead of the permanent, we define the
 241 following. The *Edmonds matrix* A of G is defined as the bipartite adjacency matrix,
 242 except every non-zero entry $A[u, v] = 1$ is replaced by a distinct variable $A[u, v] =$
 243 y_{uv} . Letting $Y = \{y_{uv} \mid uv \in E(G)\}$, we see that $\det A$ is a polynomial in Y of
 244 degree $n = |U|$. We extend this to the case when G is a bipartite multigraph. Let
 245 $Y = \{y_e \mid e \in E(G)\}$ as above, and, if G contains d edges e_1, \dots, e_d between u
 246 and v for $u \in U, v \in V$, then we let $A[u, v] = \sum_{i=1}^d y_{e_i}$. In both cases, if we view
 247 $\det A$ as a polynomial in Y , then the monomials of $\det A$ are in bijection with the
 248 perfect matchings of G . Now the Schwartz-Zippel lemma allows us to test for perfect
 249 matchings via a randomized evaluation of $\det A$. Furthermore, given a set of edge
 250 weights $w(e)$ for edges of G , we define the *weighted Edmonds matrix* in the same way
 251 as the Edmonds matrix, except every occurrence of a variable y_e for an edge $e \in E(G)$
 252 is replaced by $w(e)y_e$. In the case where the weights $w(e)$ are themselves polynomials,
 253 in a set of further variables X , this allows us to use Lemma 2.5 with $P(X, Y) = \det A$
 254 to sieve in FPT time for particular kinds of matchings in G . See Theorem 3.1 for an
 255 example.

256 **3. List Coloring with Clique Modulator.** We are ready to prove the first
 257 result of this section.

258 THEOREM 3.1. LIST COLORING WITH CLIQUE MODULATOR *can be solved by a*
 259 *randomized algorithm in time $\mathcal{O}^*(2^{k \log k})$.*

260 *Proof.* Let $L = \bigcup_{v \in V(G)} L(v)$ and $C = G - D$. We say that a proper list coloring
 261 λ for G is compatible with $(\mathcal{D}, \mathcal{D}')$ if:

- 262 • $\mathcal{D} = \{D_1, \dots, D_p\}$ is the partition of all vertices in D that do not reuse colors
 263 used by λ in C into color classes given by λ and
- 264 • $\mathcal{D}' = \{D'_1, \dots, D'_t\}$ is the partition of all vertices in D that do reuse colors
 265 used by λ in C into color classes given by λ .

266 Note that $\{D_1, \dots, D_p, D'_1, \dots, D'_t\}$ is the partition of D into color classes given by λ .

267 For a given pair $(\mathcal{D}, \mathcal{D}')$, where each set D_i and D'_i is independent in G , we will
 268 now construct a bipartite multigraph B (with weights on its edges) such that B has a

269 perfect matching satisfying certain additional properties if and only if G has a proper
 270 list coloring that is compatible with $(\mathcal{D}, \mathcal{D}')$. B has bi-partition $(C \cup \{D_1, \dots, D_p\}, L)$
 271 and edges as follows. Let $c \in C$ and $\ell \in L$ be such that $\ell \in L(c)$. Then B contains
 272 an edge $e_{c\ell}$ between c and ℓ . Furthermore, for every $j \in [t]$ there is a further edge
 273 $e_{c\ell,j}$ between c and ℓ if and only if $\ell \in (\bigcap_{d \in D'_j} L(d)) \cap L(c)$ and c is not adjacent to
 274 any vertex in D'_j . Moreover, B has an edge between a vertex D_i and a vertex $\ell \in L$
 275 if and only if $\ell \in \bigcap_{d \in D_i} L(d)$. Finally, if $|C| + p > |L|$ then λ cannot exist and we
 276 have a no-instance. Otherwise, we add $|L| - |C| - p$ dummy vertices to the partite
 277 set $C \cup \{D_1, \dots, D_p\}$ and make the dummy vertices adjacent to all vertices in L .

278 For weights, we introduce a new set of variables $X = \{x_1, \dots, x_t\}$, and for every
 279 edge $e_{c\ell,j}$ created above we set $w(e_{c\ell,j}) = x_j$. Every other edge e of B has weight
 280 $w(e) = 1$. For an illustration of B , see Figure 1.




FIG. 1. Illustration of the construction of B . $(1, (x_j)_{j \in J})$ means that there are $1 + |J|$ parallel edges between c_1 and 1 with weights $1, x_{j_1}, x_{j_2}, \dots, x_{j_{|J|}}$, where $J = \{j_1, j_2, \dots, j_{|J|}\}$.

281 Note that G has a proper list coloring that is compatible with $(\mathcal{D}, \mathcal{D}')$ if and only
 282 if B has a perfect matching F such that there is a bijection α between $[t]$ and t edges
 283 in F such that for every $i \in [t]$, the weight of the edge $\alpha(i)$ is x_i . Indeed, we have
 284 $w(\alpha(i)) = x_i$ if and only if $\alpha(i) = e_{c\ell,i}$ for some vertices c and ℓ , which in turn implies
 285 that $D'_i \cup \{c\}$ is an independent set in G and $\ell \in L(u)$ for every $u \in D'_i \cup \{c\}$. Along
 286 with the further edges of F of weight 1, this defines a proper coloring λ for G which
 287 is compatible with $(\mathcal{D}, \mathcal{D}')$.

288 Let M be the weighted Edmonds matrix of B with weights w (see Section 2.3),
 289 for simplicity constructed over a field of characteristic 2. Let $Y = \{y_e \mid e \in E(B)\}$
 290 be the set of further variables introduced in the construction of M . Then $\det M$ is
 291 a polynomial in variables $X \cup Y$, and as discussed in Section 2.3, the monomials of
 292 $\det M$ are in bijection with perfect matchings of B ; in particular, the latter holds since
 293 every weight $w(e)$ defined above is a single monomial. Furthermore, for every perfect
 294 matching F of B , the monomial of $\det M$ corresponding to F equals $\prod_{e \in F} w(e)y_e$.

295 Now it is not hard to see that $\det M$ has a monomial containing $\prod_{j=1}^t x_j$ if and
 296 only if B has a perfect matching F such that there is a bijection α between $[t]$ and t
 297 edges in F such that for every $i \in [t]$, the weight of the edge $\alpha(i)$ is x_i , which in turn
 298 is equivalent to G having a proper list coloring that is compatible with $(\mathcal{D}, \mathcal{D}')$. Note
 299 that the other $|C| - t$ edges of the form $c\ell$ contribute a factor 1 to the monomial, as
 300 do the edges of the form $D_i\ell$.

301 Hence, deciding whether G has a proper list coloring that is compatible with
 302 $(\mathcal{D}, \mathcal{D}')$ boils down to deciding whether $\det M$ has a monomial containing $\prod_{j=1}^t x_j$.
 303 For any evaluation of variables X and Y , we can compute $\det M$ in polynomial-
 304 time [9].

Now write $y = (y_1, \dots, y_m)$, and let $P(x_1, \dots, x_t, y) = \det M$. Define

$$Q(x_1, \dots, x_t) = \sum_{I \subseteq [t]} P_{-I}(x_1, \dots, x_t, y).$$

305 Note that each of P and Q is of degree at most $2n$.

306 By Lemma 2.5, $Q(x_1, \dots, x_t) \neq 0$ if and only if $\det M$ has a monomial containing
 307 $\prod_{j=1}^t x_j$. Moreover, using Lemmas 2.4 and 2.5 (with P and Q just defined), we can
 308 verify with a single evaluation of Q whether $Q(x_1, \dots, x_t) = 0$ (i.e. whether $\det M$
 309 contains a monomial containing $\prod_{j=1}^t x_j$) with probability at least $1 - \frac{2n}{|\mathbb{F}|} \geq 2/3$ for
 310 a field \mathbb{F} of characteristic 2 such that $|\mathbb{F}| \geq 6n$. Furthermore, Q can be evaluated in
 311 time $\mathcal{O}^*(2^t)$.

312 Our algorithm sets $t = k$ and for every pair $(\mathcal{D}, \mathcal{D}')$, where $\mathcal{D} \cup \mathcal{D}'$ is a partition of
 313 D into independent sets, constructs the graph B and matrix M . It then verifies in time
 314 $\mathcal{O}^*(2^t)$ whether $Q(x_1, \dots, x_t, y_1, \dots, y_m) = 0$, and if $Q(x_1, \dots, x_t, y_1, \dots, y_m) \neq 0$ it
 315 returns ‘Yes’ and terminates. If the algorithm runs to the end, it returns ‘No’.

316 Note that the time complexity of the algorithm is dominated by the number of
 317 choices for $(\mathcal{D}, \mathcal{D}')$, which is in turn dominated by $\mathcal{O}^*(\mathcal{B}_k)$, where \mathcal{B}_k is the k -th Bell
 318 number. By Berend and Tassa [4], $\mathcal{B}_k < (\frac{0.792k}{\ln(k+1)})^k$, and thus the total running time
 319 is $\mathcal{O}^*(\mathcal{B}_k 2^k) = \mathcal{O}^*(2^{k \log k})$. \square

320 **3.1. A faster FPT algorithm.** We now show a faster FPT algorithm, running
 321 in time $\mathcal{O}^*(2^k)$. It is a variation on the same algebraic sieving technique as above,
 322 but instead of guessing a partition of the modulator it works over a more complex
 323 matrix. We begin by defining the matrix, then we show how to perform the sieving
 324 step in $\mathcal{O}^*(2^k)$ time.

325 **3.1.1. Matrix definition.** As before, let $L = \bigcup_{v \in V(G)} L(v)$ be the set of all
 326 colors, and let $C = G - D$. Define an auxiliary bipartite graph $H = (U_H \cup V_H, E_H)$
 327 where initially $U_H = V(G)$ and $V_H = L$, and where $v\ell \in E_H$ for $v \in V(G)$, $\ell \in L$ if
 328 and only if $\ell \in L(v)$. Additionally, introduce a set $L' = \{\ell'_d \mid d \in D\}$ of k artificial
 329 colors, add L' to V_H , and for each $d \in D$ connect ℓ'_d to d but to no other vertex.
 330 Finally, pad U_H with $|V_H| - |U_H|$ artificial vertices connected to all of V_H ; note that
 331 this is a non-negative number, since otherwise $|L| < |V(C)|$ and we may reject the
 332 instance.

333 Next, we associate with every edge $v\ell \in E_H$ a set $S(v\ell) \subseteq 2^D$ as follows.

- 334
 - If $v \in V(C)$, then $S(v\ell)$ contains all sets $S \subseteq D$ such that the following hold:
 335
 1. S is an independent set in G
 336 2. $N(v) \cap S = \emptyset$
 337 3. $\ell \in \bigcap_{s \in S} L(s)$.
 338 - If $v \in D$ and $\ell \in L$, then $S(v\ell)$ contains all sets $S \subseteq D$ such that the following
 339 hold:
 340
 1. $v \in S$
 341 2. S is an independent set in G
 342 3. $\ell \in \bigcap_{s \in S} L(s)$.
 343 - If v or ℓ is an artificial vertex – in particular, if $\ell = \ell'_d$ for some $d \in D$ – then
 344 $S(v\ell) = \{\emptyset\}$.

345 Finally, define a matrix A of dimensions $|U_H| \times |V_H|$, with rows labelled by U_H and
 346 columns labelled by V_H , whose entries are polynomials as follows. Define a set of
 347 variables $X = \{x_d \mid d \in D\}$ corresponding to vertices of D , and additionally a set
 348 $Y = \{y_e \mid e \in E_H\}$. Then for every edge $v\ell$ in H , $v \in U_H$, $\ell \in V_H$ we define

$$349 \quad P(v\ell) = \sum_{S \in \mathcal{S}(v\ell)} \prod_{s \in S} x_s,$$

350 where as usual an empty product equals 1. Then for each edge $v\ell \in E_H$ we let $A[v, \ell] =$
 351 $y_{v\ell}P(v\ell)$, and the remaining entries of A are 0. We argue the following. (Expert
 352 readers may note although the argument can be sharpened to show the existence
 353 of a multilinear term, we do not wish to argue that there exists such a term with
 354 odd coefficient. Therefore we use the simpler sieving of Lemma 2.5 instead of full
 355 multilinear detection, cf. [14].)

356 **LEMMA 3.2.** *Let A be defined as above. Then $\det A$ (as a polynomial) contains a*
 357 *monomial divisible by $\prod_{x \in X} x$ if and only if G is properly list colorable.*

358 *Proof.* We first note that no cancellation happens in $\det A$. Note that monomials
 359 of $\det A$ correspond (many-to-one) to perfect matchings of H , and thanks to the formal
 360 variables Y , two monomials corresponding to distinct perfect matchings never interact.
 361 On the other hand, if we fix a perfect matching M in H , then the contributions of M
 362 to $\det A$ equal $\sigma_M \prod_{e \in M} y_e P(e)$, where $\sigma_M \in \{1, -1\}$ is a sign term depending only
 363 on M . Since the polynomials $P(e)$ contain only positive coefficients, no cancellation
 364 occur, and every selection of a perfect matching M of H and a factor from every
 365 polynomial $P(e)$, $e \in M$ results (many-to-one) to a monomial with non-zero coefficient
 366 in $\det A$.

367 We now proceed with the proof. On the one hand, let c be a proper list coloring of
 368 G . Define an ordering \prec on $V(G)$ such that $V(C)$ precedes D , and define a matching
 369 M as follows. For every vertex $v \in V(C)$, add $vc(v)$ to M . For every vertex $v \in D$,
 370 add $vc(v)$ to M if v is the first vertex according to \prec that uses color $c(v)$, otherwise
 371 add $v\ell'_v$ to M . Note that M is a matching in H of $|V(G)|$ edges. Pad M to a perfect
 372 matching in H by adding arbitrary edges connected to the artificial vertices in U_H ;
 373 note that this is always possible. Finally, for every edge $v\ell \in M$ with $\ell \in L$ we
 374 let $D_{v\ell} = D \cap c^{-1}(\ell)$. Observe that for every edge $v\ell$ in M , $D_{v\ell} \in \mathcal{S}(v\ell)$; indeed,
 375 this holds by construction of $\mathcal{S}(v\ell)$ and since c is a proper list coloring. Further let
 376 $p_{v\ell} = \prod_{v \in D_{v\ell}} x_v$; thus $p_{v\ell}$ is a term of $P(v\ell)$. It follows, by the discussion in the first
 377 paragraph of the proof, that

$$378 \quad \alpha \sigma_M \prod_{v\ell \in M} y_{v\ell} p_{v\ell}$$

379 is a monomial of $\det A$ for some constant $\alpha > 0$, where $\sigma_M \in \{1, -1\}$ is the sign term
 380 for M . It remains to verify that every variable $x_d \in X$ occurs in some term $p_{v\ell}$. Let
 381 $\ell = c(d)$ and let v be the earliest vertex according to \prec such that $c(v) = \ell$. Then
 382 $v\ell \in M$ and x_d occurs in $p_{v\ell}$. This finishes the first direction of the proof.

383 On the other hand, assume that $\det A$ contains a monomial T divisible by $\prod_{x \in X} x$,
 384 and let M be the corresponding perfect matching of H . Let $T = \alpha \prod_{e \in M} y_e p_e$ for
 385 some constant factor α , where p_e is a term of $P(e)$ for every $e \in M$. Clearly such
 386 a selection is possible; if it is ambiguous, make the selection arbitrarily. Now define
 387 a mapping $c: V(G) \rightarrow L$ as follows. For $v \in V(C)$, let $v\ell \in M$ be the unique edge
 388 connected to v , and set $c(v) = \ell$. For $v \in D$, let v' be the earliest vertex according
 389 to \prec such that x_v occurs in $p_{v'\ell}$, where $v'\ell \in M$. Set $c(v) = \ell$. We verify that c is

390 a proper list coloring of G . First of all, note that $c(v)$ is defined for every $v \in V(G)$
 391 and that $c(v) \in L(v)$. Indeed, if $v \in V(C)$ then $c(v) \in L(v)$ since $vc(v) \in E_H$; and if
 392 $v \in D$ then $c(v) \in L(v)$ is verified in the creation of the term $p_{vc(v)}$ in $P(vc(v))$. Next,
 393 consider two vertices $u, v \in V(G)$ with $c(u) = c(v)$. If $u, v \in D$, then u and v are
 394 represented in the same term $p_{v'c(v)}$ for some v' , hence u and v form an independent
 395 set; otherwise assume $u \in V(C)$. Note that $u, v \in V(C)$ is impossible since otherwise
 396 the matching M would contain two edges $uc(u)$ and $vc(u)$ which intersect. Thus
 397 $v \in D$, and v is represented in the term $p_{uc(u)}$. Therefore $uv \notin E(G)$, by construction
 398 of $P(uc(u))$. We conclude that c is a proper coloring respecting the lists $L(v)$, i.e., a
 399 proper list coloring. \square

400 **3.1.2. Fast evaluation.** By the above description, we can test for the existence
 401 of a list coloring of G using 2^k evaluations of $\det A$, as in Theorem 3.1; and each
 402 evaluation can be performed in $\mathcal{O}^*(2^k)$ time, including the time to evaluate the poly-
 403 nomials $P(v\ell)$, making for a running time of $\mathcal{O}^*(4^k)$ in total (or $\mathcal{O}^*(3^k)$ with more
 404 careful analysis). We show how to perform the entire sieving in time $\mathcal{O}^*(2^k)$ using
 405 fast subset convolution.

406 For $I \subseteq D$, let us define A_{-I} as A with all occurrences of variables x_i , $i \in I$
 407 replaced by 0, and for every edge $v\ell$ of H , let $P(v\ell)_{-I}$ denote the polynomial $P(v\ell)$
 408 with x_i , $i \in I$ replaced by 0. Then a generic entry (v, ℓ) of A_{-I} equals

$$409 \quad A_{-I}[v, \ell] = y_{v\ell} P_{-I}(v\ell),$$

410 and in order to construct A_{-I} it suffices to pre-compute the value of $P_{-I}(v\ell)$ for every
 411 edge $v\ell \in E_H$, $I \subseteq D$. For this, we need the *fast zeta transform* of Yates [35], which
 412 was introduced to exact algorithms by Björklund et al. [6].

413 **LEMMA 3.3** ([35, 6]). *Given a function $f : 2^N \rightarrow R$ for some ground set N and*
 414 *ring R , we may compute all values of $\hat{f} : 2^N \rightarrow R$ defined as $\hat{f}(S) = \sum_{A \subseteq S} f(A)$*
 415 *using $\mathcal{O}^*(2^{|N|})$ ring operations.*

416 We show the following lemma, which is likely to have analogues in the literature,
 417 but we provide a short proof for the sake of completeness.

418 **LEMMA 3.4.** *Given an evaluation of the variables X , the value of $P_{-I}(v\ell)$ can be*
 419 *computed for all $I \subseteq D$ and all $v\ell \in E_H$ in time and space $\mathcal{O}^*(2^k)$.*

420 *Proof.* Consider an arbitrary polynomial $P_{-I}(v\ell)$.

421 Recalling $P(v\ell) = \sum_{S \in \mathcal{S}(v\ell)} \prod_{s \in S} x_s$, we have:

$$422 \quad P_{-I}(v\ell) = \sum_{S \in \mathcal{S}(v\ell)} [S \cap I = \emptyset] \prod_{s \in S} x_s = \sum_{S \subseteq (D-I)} [S \in \mathcal{S}(v\ell)] \prod_{s \in S} x_s,$$

423 using Iverson bracket notation.¹ Using $f(S) = [S \in \mathcal{S}(v\ell)] \prod_{s \in S} x_s$, this clearly fits
 424 the form of Lemma 3.3, with $\hat{f}(D-I) = P_{-I}(v\ell)$. Hence we apply Lemma 3.3
 425 for every edge $v\ell \in E_H$, for $\mathcal{O}^*(2^k)$ time per edge, making $\mathcal{O}^*(2^k)$ time in total to
 426 compute all values. \square

427 Having access to these values, it is now easy to complete the algorithm.

428 **THEOREM 3.5.** LIST COLORING WITH CLIQUE MODULATOR *can be solved by a*
 429 *randomized algorithm in time $\mathcal{O}^*(2^k)$.*

¹Recall that for a logical proposition P , $[P] = 1$ if P is true and 0, otherwise.

430 *Proof.* Let A be the matrix defined above (but do not explicitly construct it yet).
 431 By Lemma 3.2, we need to check whether $\det A$ contains a monomial divisible by
 432 $\prod_{x \in X} x$, and by Lemma 2.5 this is equivalent to testing whether

$$433 \quad \sum_{I \subseteq D} (-1)^{|I|} \det A_{-I} \neq 0.$$

434 By the Schwartz-Zippel lemma (Lemma 2.4), it suffices to randomly evaluate the
 435 variables X and Y occurring in A and evaluate this sum once; if G has a proper list
 436 coloring and if the values of X and Y are chosen among sufficiently many values, then
 437 with high probability the result is non-zero, and if not, then the result is guaranteed
 438 to be zero. Thus the algorithm is as follows.

- 439 1. Instantiate variables of X and Y uniformly at random from $[N]$ for some
 440 sufficiently large N . Note that for an error probability of ε with $0 < \varepsilon < 1$,
 441 it suffices to use $N = \Omega(n^2(1/\varepsilon))$.
- 442 2. Use Lemma 3.4 to fill in a table with the value of $P_{-I}(v\ell)$ for all I and $v\ell$ in
 443 time $\mathcal{O}^*(2^k)$.
- 444 3. Compute

$$445 \quad \sum_{I \subseteq D} (-1)^{|I|} \det A_{-I},$$

446 constructing A_{-I} from the values $P_{-I}(v\ell)$ in polynomial time in each step.

- 447 4. Answer YES if the result is non-zero, NO otherwise.

448 Clearly this runs in total time and space $\mathcal{O}^*(2^k)$ and the correctness follows from the
 449 arguments above. \square

450 **3.2. Refuting Polynomial Kernel.** In this section, we prove that LIST COL-
 451 ORING WITH CLIQUE MODULATOR does not admit a polynomial kernel. We prove this
 452 result by a polynomial parameter transformation from HITTING SET where the param-
 453 eter is the number of sets, which is known not to have a polynomial kernel [17]. Notice
 454 that HITTING SET parameterized by number of sets is equivalent to SET COVER pa-
 455 rameterized by the universe size.

456 **THEOREM 3.6.** LIST COLORING WITH CLIQUE MODULATOR *parameterized by k*
 457 *does not admit a polynomial kernel unless $NP \subseteq coNP/poly$.*

458 *Proof.* Let us recall the formal definition of the HITTING SET problem.

459 HITTING SET parameterized by m

Input: A universe U of n elements, a family $\mathcal{F} \subseteq 2^U$ of m subsets of U , and
 an integer k .

Problem: Is there $X \subseteq U$ with at most k elements such that for every $F \in \mathcal{F}$,
 it holds that $F \cap X \neq \emptyset$?

460
 461 Let (U, \mathcal{F}, k) be an instance of HITTING SET problem where $U = [n]$, and $\mathcal{F} =$
 462 $\{F_1, \dots, F_m\}$. Now, we are ready to describe the construction.

463 **Construction:** For every $i \in [m]$, we create a vertex u_i and assign $L(u_i) =$
 464 F_i . Let $D = \{u_1, \dots, u_m\}$. In addition, we create a clique C with $n - k$ vertices
 465 $\{v_1, \dots, v_{n-k}\}$. Moreover, for every $j \in [n - k]$, we set $L(v_j) = U$ and for all $i \in [n - k]$
 466 and $j \in [m]$, let (u_i, v_j) be an edge. This completes the construction, which takes
 467 polynomial time. We denote the obtained graph by G . It remains to show that the
 468 two instances are equivalent.
 469

470 Towards showing the forward direction, let (U, \mathcal{F}, k) be an yes-instance. Then,
 471 there is a set X of at most k elements from U such that for every $F_i \in \mathcal{F}$, $X \cap F_i \neq \emptyset$.

472 Using the elements present in X , we can color D as follows. We pick an element
 473 arbitrarily from every $F_i \cap X$, and color the vertex u_i using that color. After that,
 474 we provide different colors to different vertices in C that are different from the colors
 475 used in D as well. Hence, we can color G by n colors.

476 Towards showing the backwards direction, suppose that G has a proper list col-
 477 oring. Note that all vertices of C have to get different colors. Hence, the vertices of
 478 D must be colorable using only k colors. Suppose that X is the set of k colors used
 479 to color the vertices of D . Note that the colors respect the list for every vertex in D
 480 where the list represents the sets in the family. Hence, X is a hitting set of size k . \square

481 Note that the reduction also shows that if LIST COLORING WITH CLIQUE MODULA-
 482 TOR could be solved in time $\mathcal{O}(2^{\epsilon k n^{\mathcal{O}(1)}})$ for some $\epsilon < 1$, then HITTING SET could be
 483 solved in time $\mathcal{O}(2^{\epsilon |\mathcal{F}|} |U|^{\mathcal{O}(1)})$, which in turn would imply that any instance I with
 484 universe U and set family \mathcal{F} of the well-known SET COVER problem could be solved
 485 in time $\mathcal{O}(2^{\epsilon |U|} |\mathcal{F}|^{\mathcal{O}(1)})$. The existence of such an algorithm is open, and it has been
 486 conjectured that no such algorithm is possible under SETH (the strong exponential-
 487 time hypothesis); see Cygan et al. [15]. Thus, up to the assumption of this conjecture
 488 and SETH, the algorithm for LIST COLORING WITH CLIQUE MODULATOR given in
 489 Theorem 3.5 is best possible w.r.t. its dependency on k .

490 **4. Polynomial kernel for PRE-COLORING EXTENSION WITH CLIQUE MOD-**
 491 **ULATOR.** In the following let $(G, D, k, \lambda_P, X, Q)$ be an instance of PRE-COLORING
 492 EXTENSION WITH CLIQUE MODULATOR, let $C = G - D$, let D_P be the set of all pre-
 493 colored vertices in D , and let $D' = D \setminus D_P$. W.l.o.g., we can assume that $|Q| \geq |C|$ as
 494 otherwise the instance is a trivial no-instance. In the following, we will assume that
 495 the instance will be updated with the introduction of every reduction rule, i.e., we
 496 will assume that all already introduced reduction rules have already been exhaustively
 497 applied to the current instance.

498 **Reduction Rule 1.** *Remove any vertex $v \in D'$ that has less than $|Q|$ neighbors*
 499 *in G .*

500 The proof of the following lemma is obvious and thus omitted.

501 **LEMMA 4.1.** *Reduction Rule 1 is safe and can be implemented in polynomial time.*

502 Note that if Reduction Rule 1 can no longer be applied, then every vertex in D' has
 503 at least $|Q|$ neighbors, which because of $|Q| \geq |C|$ implies that every such vertex has
 504 at most $|D| \leq k$ non-neighbors in G and hence also in C . Let C_N be the set of all
 505 vertices in C that are not adjacent to all vertices in D' and let $C' = C - C_N$. Note
 506 that $|C_N| \leq |D||D| \leq k^2$.

507 We show next how to reduce the size of C_N to k . Note that this step is optional
 508 if our aim is solely to obtain a polynomial kernel, however, it allows us to reduce
 509 the number of vertices in the resulting kernel from $\mathcal{O}(k^2)$ to $\mathcal{O}(k)$. Let J be the
 510 bipartite graph with partition (C_N, D) having an edge between $c \in C_N$ and $d \in D$ if
 511 $\{c, d\} \notin E(G)$. Our next reduction rule can be seen as a crown reduction rule that
 512 uses a crown decomposition of J with crown A and head $N_H(A)$; a similar rule has
 513 been employed previously in [3, Reduction Rule 2].

514 **Reduction Rule 2.** *If $A \subseteq C_N$ is an inclusion-wise minimal set satisfying $|A| >$
 515 $|N_J(A)|$, then remove the vertices in $D' \cap N_J(A)$ from G .*

516 Note that after the application of Reduction Rule 2, the vertices in A are implicitly

517 removed from C_N and added to C' since all their non-neighbors in D' (i.e. the vertices
518 in $D' \cap N_J(A)$) are removed from the graph.

519 LEMMA 4.2. *Reduction Rule 2 is safe and can be implemented in polynomial time.*

520 *Proof.* It is clear that the rule can be implemented in polynomial-time. Towards
521 showing the safeness of the rule, it suffices to show that G has a coloring extending λ_P
522 using only colors from Q if and only if so does $G \setminus (D' \cap N_J(A))$. Since $G \setminus (D' \cap N_J(A))$
523 is a subgraph of G , the forward direction of this statement is trivial. So assume that
524 $G \setminus (D' \cap N_J(A))$ has a coloring λ extending λ_P using only colors from Q . Because the
525 set A is inclusion-minimal, we obtain from Proposition 2.1, that there is a (maximum)
526 matching, say M , between $N_J(A)$ and A in J that saturates $N_J(A)$. Moreover, it
527 follows from the definition of J that every vertex in A is adjacent to every vertex in
528 G apart from the vertices in $N_J(A)$. Therefore, the colors in $\lambda(A)$ can only reappear
529 in $D_P \cap N_J(A)$. We can now use the matching M to reshuffle the colors in A in
530 such a way that the colors of vertices in A that are matched by M to a vertex in D'
531 appear exactly once in the graph; or in other words we reshuffle the colors in A such
532 that all colors that also appear in $D_P \cap N_J(A)$ are assigned to vertices in A that are
533 matched by M to vertices in D_P . That is, let A' be the set of all vertices a in A with
534 $\lambda(a) \in \lambda(D_P \cap N_J(A))$ such that a is matched by M to a vertex in D' . Similarly, let
535 A_P be the set of all vertices a in A with $\lambda(a) \notin \lambda(D_P \cap N_J(A))$ such that a is matched
536 by M to a vertex in D_P . Note that $|A_P| \geq |A'|$ and therefore there is a bijection
537 $\alpha : A' \rightarrow A'_P$ from A' to a subset A'_P of A_P . Now, let λ' be the coloring obtained from
538 λ by setting $\lambda'(a) = \lambda(\alpha(a))$ for every $a \in A'$, $\lambda'(a) = \lambda(\alpha^{-1}(a))$ for every $a \in A'_P$,
539 and $\lambda'(a) = \lambda(a)$ otherwise. Then, the color $\lambda'(a)$ appears exactly once for every
540 $a \in A$ that is matched by M to a vertex in D' . Therefore, we can extend λ' into a
541 coloring λ'' for G by coloring the vertices in $D' \cap N_J(A)$ according to the matching M .
542 More formally, let $\lambda_{D' \cap N_J(A)}$ be the coloring for the vertices in $D' \cap N_J(A)$ obtained
543 by setting $\lambda_{D' \cap N_J(A)}(v) = \lambda'(u)$ for every $v \in D' \cap N_J(A)$, where $\{v, u\} \in M$. Then,
544 we obtain λ'' by setting: $\lambda''(v) = \lambda'(v)$ for every $v \in V(G) \setminus (D' \cap N_J(A))$ and
545 $\lambda''(v) = \lambda_{D' \cap N_J(A)}(v)$ for every vertex $v \in D' \cap N_J(A)$. \square

546 Note that because of Proposition 2.1, we obtain that there is a set $A \subseteq C_N$ with
547 $|A| > |N_J(A)|$ as long as $|C_N| > |D|$. Moreover, since $N_J(A) \cap D' \neq \emptyset$ for every such
548 set A (due to the definition of C_N), we obtain that Reduction Rule 2 is applicable as
549 long as $|C_N| > |D|$. Hence after an exhaustive application of Reduction Rule 2, we
550 obtain that $|C_N| \leq |D'| \leq k$.

551 We now introduce our final two reduction rules, which allow us to reduce the size
552 of C' .

553 **Reduction Rule 3.** *Let $v \in V(C')$ be a pre-colored vertex with color $\lambda_P(v)$.
554 Then remove $\lambda_P^{-1}(\lambda_P(v))$, i.e., all vertices colored with the same color ($\lambda_P(v)$) as v ,
555 from G and $\lambda_P(v)$ from Q .*

556 LEMMA 4.3. *Reduction Rule 3 is safe and can be implemented in polynomial time.*

557 *Proof.* Because $v \in V(C')$, it holds that only vertices in D_P can have color $\lambda_P(v)$,
558 but these are already pre-colored. Hence in any coloring for G that extends λ_P , the
559 vertices in $\lambda_P^{-1}(\lambda_P(v))$ are the only vertices that obtain color $\lambda_P(v)$, which implies
560 the safeness of the rule. \square

561 Because of Reduction Rule 3, we can from now on assume that no vertex in C'
562 is pre-colored. Note that the only part of G , whose size is not yet bounded by a
563 polynomial in the parameter k is C' . To reduce the size of C' , we need will make use

564 of Proposition 2.2. Let $P = \lambda_P(D_P)$ and H be the bipartite graph with bi-partition
 565 (C', P) containing an edge between $c' \in C'$ and $p \in P$ if and only if c' is not adjacent
 566 to a vertex pre-colored by p in G .

567 **Reduction Rule 4.** *Let M be a maximum matching in H and let C_M be the*
 568 *endpoints of M in C' . Then remove all vertices in $C_{\overline{M}} := C' \setminus C_M$ from G and*
 569 *remove an arbitrary set of $|C_{\overline{M}}|$ colors from $Q \setminus \lambda_P(X)$. (Recall that $\lambda_P : X \rightarrow Q$.)*

570 In the following let C_M and $C_{\overline{M}}$ be as defined in the above reduction rule for an
 571 arbitrary maximum matching M of H . To show that the reduction rule is safe, we
 572 need the following auxiliary lemma, which shows that if a coloring for G reuses colors
 573 from P in C' , then those colors can be reused solely on the vertices in C_M .

574 **LEMMA 4.4.** *If there is a coloring λ for G extending λ_P using only colors in*
 575 *Q , then there is a coloring λ' for G extending λ_P using only colors in Q such that*
 576 *$\lambda'(C_{\overline{M}}) \cap P = \emptyset$.*

577 *Proof.* Let C_P be the set of all vertices v in C' with $\lambda(v) \in P$. If $C_P \cap C_{\overline{M}} =$
 578 \emptyset , then setting λ' equal to λ satisfies the claim of the lemma. Hence assume that
 579 $C_P \cap C_{\overline{M}} \neq \emptyset$. Let N be the matching in H containing the edges $\{v, \lambda(v)\}$ for every
 580 $v \in C_P$; note that N is indeed a matching in H , because C_P is a clique in G . Because
 581 of Proposition 2.2, there is a matching N' in $H[C_M \cup P]$ such that N' has exactly the
 582 same endpoints in P as N . Let $C_M[N']$ be the endpoints of N' in C_M and let λ_A be
 583 the coloring of the vertices in $C_M[N']$ corresponding to the matching N' , i.e., a vertex
 584 v in $C_M[N']$ obtains the unique color $p \in P$ such that $\{v, p\} \in N'$. Finally, let α be
 585 an arbitrary bijection between the vertices in $(V(N) \cap C') \setminus C_M[N']$ and the vertices
 586 in $C_M[N'] \setminus (V(N) \cap C')$, which exists because $|N| = |N'|$. We now obtain λ' from
 587 λ by setting $\lambda'(v) = \lambda_A(v)$ for every $v \in C_M[N']$, $\lambda'(v) = \lambda(\alpha(v))$ for every vertex
 588 $v \in (V(N) \cap C') \setminus C_M[N']$, and $\lambda'(v) = \lambda(v)$ for every other vertex. To see that λ'
 589 is a proper coloring note that $\lambda'(C') = \lambda(C')$. Moreover, all the colors in $\lambda(C') \setminus P$
 590 are “universal colors” in the sense that exactly one vertex of G obtains the color and
 591 hence those colors can be freely moved around in C' . Finally, the matching N' in H
 592 ensures that the vertices in $C_M[N']$ can be colored using the colors from P . \square

593 **LEMMA 4.5.** *Reduction Rule 4 is safe and can be implemented in polynomial time.*

594 *Proof.* Note first that the reduction can always be applied since if $Q \setminus \lambda_P(X)$
 595 contains less than $|C_{\overline{M}}|$ colors, then the instance is a no-instance. It is clear that the
 596 rule can be implemented in polynomial time using any polytime algorithm for finding
 597 a maximum matching [29]. Moreover, if the reduced graph has a coloring extending
 598 λ_P using only the colors in Q , then so does the original graph, since the vertices in
 599 $C_{\overline{M}}$ can be colored with the colors removed from the original instance.

600 Hence, it remains to show that if G has a coloring, say λ , extending λ_P using
 601 only colors in Q , then $G \setminus C_{\overline{M}}$ has a coloring extending λ_P that uses only colors in
 602 $Q' := Q \setminus Q_{\overline{M}}$, where $Q_{\overline{M}}$ is the set of $|C_{\overline{M}}|$ colors from $Q \setminus \lambda_P(X)$ that have been
 603 removed from Q .

604 Because of Lemma 4.4, we may assume that $\lambda(C_{\overline{M}}) \cap P = \emptyset$. Let B be the set of
 605 all vertices v in $G \setminus C_{\overline{M}}$ with $\lambda(v) \in Q_{\overline{M}}$. If $B = \emptyset$, then λ is a coloring extending λ_P
 606 using only colors from Q' . Hence assume that $B \neq \emptyset$. Let A be the set of all vertices
 607 v in $C_{\overline{M}}$ with $\lambda(v) \in Q'$. Then $\lambda(A) \cap \lambda_P(X) = \emptyset$, which implies that every color in
 608 $\lambda(A)$ appears only in $C_{\overline{M}}$ (and exactly once in $C_{\overline{M}}$). Moreover, $|\lambda(A)| \geq |\lambda(B)|$. Let
 609 α be an arbitrary bijection between $\lambda(B)$ and an arbitrary subset of $\lambda(A)$ (of size $|B|$)
 610 and let λ' be the coloring obtained from λ by setting $\lambda'(v) = \alpha(\lambda(v))$ for every $v \in B$,
 611 $\lambda'(v) = \alpha^{-1}(\lambda(v))$ for every $v \in A$, and $\lambda'(v) = \lambda(v)$, otherwise. Then λ' restricted

612 to $G - C_{\overline{M}}$ is a coloring for $G - C_{\overline{M}}$ extending λ_P using only colors from Q' . Note
 613 that λ' is a proper coloring because the colors in $\lambda(A)$ are not in P and hence do not
 614 appear anywhere else in G and moreover the colors in $\lambda(B)$ do not appear in $\lambda(C_{\overline{M}})$. \square

615 Note that after the application of Reduction Rule 4, it holds that $|C'| = |C_M| \leq |P| \leq$
 616 $|D_P| \leq |D| \leq k$. Together with the facts that $|D| \leq k$, $|C_N| \leq k$, we obtain that the
 617 reduced graph has at most $3k$ vertices.

618 **THEOREM 4.6.** PRE-COLORING EXTENSION WITH CLIQUE MODULATOR *admits*
 619 *a polynomial kernel with at most $3k$ vertices.*

620 **5. Polynomial kernel and Compression for $(n - k)$ -REGULAR LIST COL-**
 621 **ORING.** We now show our polynomial kernel and compression for $(n - k)$ -REGULAR
 622 LIST COLORING, which is more intricate than the one for PRE-COLORING EXTEN-
 623 SION WITH CLIQUE MODULATOR. Let (G, k, L) be an input of $(n - k)$ -REGULAR LIST
 624 COLORING. We begin by noting that we can assume that G has a clique-modulator
 625 of size at most $2k$.

626 **LEMMA 5.1** ([3]). *In polynomial-time either we can either solve (G, k, L) or*
 627 *compute a clique-modulator for G of size at most $2k$.*

628 Henceforth, we let $V(G) = C \cup D$ where $G[C]$ is a clique and D is a clique
 629 modulator, $|D| \leq 2k$. Let $T = \bigcup_{v \in V(G)} L(v)$. We note one further known reduction
 630 rule for $(n - k)$ -REGULAR LIST COLORING. Consider the bipartite graph H_G with
 631 bi-partition $(V(G), T)$ having an edge between $v \in V(G)$ and $t \in T$ if and only if
 632 $t \in L(v)$.

633 **Reduction Rule 5** ([3]). *Let T' be an inclusion-wise minimal subset of T such*
 634 *that $|N_{H_G}(T')| < |T'|$, then remove all vertices in $N_{H_G}(T')$ from G .*

635 Note that after an exhaustive application of Reduction Rule 5, it holds that $|T| \leq$
 636 $|V(G)|$ since otherwise Proposition 2.1 would ensure the applicability of the reduction
 637 rule. Hence in the following we will assume that $|T| \leq |V(G)|$.

638 With this preamble handled, let us proceed with the kernelization. We are not
 639 able to produce a direct ‘crown reduction rule’ for LIST COLORING, as for PRE-
 640 COLORING EXTENSION (e.g., we do not know of a useful generalization of Reduction
 641 Rule 2). Instead, we need to study more closely which list colorings of $G[D]$ extend
 642 to list colorings of G . For this purpose, let $H = H_G - D$ be the bipartite graph
 643 with bi-partition (C, T) having an edge $\{c, t\}$ with $c \in C$ and $t \in T$ if and only if
 644 $t \in L(c)$. Say that a partial list coloring $\lambda_0: A \rightarrow T$ is *extensible* if it can be extended
 645 to a proper list coloring λ of G . If $D \subseteq A$, then a sufficient condition for this is that
 646 $H - (A \cup \lambda_0(A))$ admits a matching saturating $C \setminus A$. (This is not a necessary condition,
 647 since some colors used in $\lambda_0(D)$ could be reused in $\lambda(C \setminus A)$, but this investigation
 648 will point in the right direction.) By Proposition 2.1, this is characterized by Hall
 649 sets in $H - (A \cup \lambda_0(A))$.

650 A Hall set $S \subseteq U$ in a bipartite graph G' with bi-partition (U, W) is *trivial* if
 651 $N(S) = W$. We start by noting that if a color occurs in sufficiently many vertex
 652 lists in H , then it behaves uniformly with respect to extensible partial colorings λ_0
 653 as above.

654 **LEMMA 5.2.** *Let $\lambda_0: A \rightarrow T$ be a partial list coloring where $|A \cap C| \leq p$ and let*
 655 *$t \in T$ be a color that occurs in at least $k + p$ lists in C . Then t is not contained in*
 656 *any non-trivial Hall set of colors in $H - (A \cup \lambda_0(A))$.*

657 *Proof.* Let $H' = H - (A \cup \lambda_0(A))$. Consider any Hall set of colors $S \subset (T \setminus \lambda_0(A))$

658 and any vertex $v \in C \setminus (A \cup N_{H'}(S))$ (which exists assuming S is non-trivial). Then
 659 $S \subseteq T \setminus L(v)$, hence $|S| \leq k$, and by assumption $|N_{H'}(S)| < |S|$. But for every $t' \in S$,
 660 we have $N_H(t') \subseteq N_{H'}(S) \cup (A \cap C)$, hence t' occurs in at most $|N_{H'}(S) \cup (A \cap C)| < k+p$
 661 vertex lists in C . Thus $t \notin S$. \square

662 In the following, we will assume that $n \geq 11k$.² This is safe, since otherwise
 663 (by Reduction Rule 5) we already have a kernel with a linear number of vertices and
 664 colors. We say that a color $t \in T$ is *rare* if it occurs in at most $6k$ lists of vertices in
 665 C .

666 **LEMMA 5.3.** *If $n \geq 11k$, then there are at most $3k$ rare colors.*

667 *Proof.* Let $S = \{t \in T \mid d_H(t) < 6k\}$. For every $t \in S$, there are $|C| - 6k$ “non-
 668 occurrences” (i.e., vertices $v \in C$ with $t \notin L(v)$), and there are $|C|k$ non-occurrences
 669 in total. Thus

$$670 \quad |S| \cdot (|C| - 6k) \leq |C|k \quad \Rightarrow \quad |S| \leq \frac{|C|}{|C| - 6k}k = \left(1 + \frac{6k}{|C| - 6k}\right)k,$$

671 where the bound is monotonically decreasing in $|C|$ and maximized (under the as-
 672 sumption that $n \geq 11k$ and hence $|C| \geq 9k$) for $|C| = 9k$ yielding $|S| \leq 3k$. \square

673 Let $T_R \subseteq T$ be the set of rare colors. Define a new auxiliary bipartite graph H^*
 674 with bi-partition $(C, D \cup T_R)$ having an edge between a vertex $c \in C$ and a vertex
 675 $d \in D$ if $\{c, d\} \notin E(G)$ and an edge between a vertex $c \in C$ and a vertex $t \in T_R$
 676 if $t \in L(c)$. Let X be a minimum vertex cover of H^* . Refer to the colors $T_R \setminus X$
 677 as *constrained* rare colors. Note that constrained rare colors only occur on lists of
 678 vertices in $D \cup (C \cap X)$. Let $T' = T \setminus (T_R \setminus X)$, $V' = (D \setminus X) \cup (C \cap X)$, and set
 679 $q = |T'| - |C \setminus X|$. Before we continue, we want to provide some useful observations
 680 about the sizes of the considered sets and numbers.

681 **Observation 1.** *It holds that:*

- 682 • $|X| \leq |D| + |T_R| \leq 5k$,
- 683 • $|V'| \leq |D| + |X| \leq 7k$,
- 684 • $q \leq |T| - |C| + |C \cap X| \leq |D| + |X| \leq 7k$; *this holds because $|T| \leq |V| =$*
 685 *$|C| + |D|$.*

686 **LEMMA 5.4.** *Assume $n \geq 11k$. Then G has a list coloring if and only if there is*
 687 *a partial list coloring $\lambda_0: V' \rightarrow T$ that uses at most $q = |T'| - |C \setminus X|$ colors from T' .*

688 *Proof.* The number of colors usable in $C \setminus X$ is $|T'| - p$ where p is the number
 689 counted above (since constrained rare colors cannot be used in $C \setminus X$ even if they
 690 are unused in λ_0). Thus it is a requirement that $|T'| - p \geq |C \setminus X|$. That is,
 691 $p \leq |T'| - |C \setminus X| = q$. Thus necessity is clear. We show sufficiency as well. That is,
 692 let λ_0 be a partial list coloring with scope $V' = (C \cap X) \cup (D \setminus X)$ which uses at most
 693 q colors of T' . We modify and extend λ_0 to a list coloring of G .

694 First let H_0 be the bipartite graph with bi-partition $(V, T_R \setminus X)$ and let M_0 be
 695 a matching saturating $T_R \setminus X$; note that this exists by reduction rule 5. We modify
 696 λ_0 to a coloring λ'_0 so that every constrained rare color is used by λ'_0 , by iterating
 697 over every color $t \in T_R \setminus X$; for every t , if t is not yet used by λ'_0 , then let $vt \in M_0$
 698 and update λ'_0 with $\lambda'_0(v) = t$. Note that the scope of λ'_0 after this modification is
 699 contained in $(C \cap X) \cup D$. Next, let M be a maximum matching in H^* . We use M

²The constants $11k$ and $6k$ in this paragraph are chosen to make the arguments work smoothly. A smaller kernel is possible with a more careful analysis and further reduction rules.

700 to further extend λ'_0 in stages to a partial list coloring λ which colors all of D and
 701 uses all rare colors. In phase 1, for every color $t \in T_R \cap X$ which is not already used,
 702 let $vt \in M$ be the edge covering t and assign $\lambda(v) = t$. Note that M matches every
 703 vertex of X in H^* with a vertex not in X , thus the edge vt exists and v has not yet
 704 been assigned in λ . Hence, at every step we maintain a partial list coloring, and at
 705 the end of the phase all rare colors have been assigned. Finally, as phase 2, for every
 706 vertex $v \in D \cap X$ not yet assigned, let $uv \in M$ where $u \in C$; necessarily $u \in C \setminus X$
 707 and u is as of yet unassigned in λ . The number of colors assigned in λ thus far is at
 708 most $|X| + |D| \leq |T_R| + 2|D| \leq 7k$, whereas $|L(u) \cap L(v)| \geq n - 2k \geq 9k$, hence there
 709 always exists an unused shared color that can be mapped to $\lambda(u) = \lambda(v)$. Let λ be
 710 the resulting partial list coloring. We claim that λ can be extended to a list coloring
 711 of G .

712 Let A be the scope of λ and let $H' = H - (A \cap \lambda(A))$. Note that $A \cap C \subseteq V(M)$,
 713 hence $|A \cap C| \leq |D| + |T_R| \leq 5k$. Thus by Lemma 5.2, no non-trivial Hall set in H' can
 714 contain a rare color. However, all rare colors are already used in λ . Thus H' contains
 715 no non-trivial Hall set of colors. Thus the only possibility that λ is not extensible is
 716 that H' has a trivial Hall set, i.e., $|T \setminus \lambda(A)| < |C \setminus A|$. However, every modification
 717 after λ'_0 added one vertex to A and one color to $\lambda(A)$, keeping the balance between
 718 the two sides. Thus already the partial coloring λ'_0 leaves behind a trivial Hall set.
 719 However, λ'_0 colors precisely $C \cap X$ in C and leaves at least $|T'| - q$ colors remaining.
 720 By design this is at least $|C \setminus X|$, yielding a contradiction. Thus we find that H'
 721 contains no Hall set, and λ is a list coloring of G . \square

722 Before we give our compression and kernelization results, we need the following aux-
 723 iliary lemma.

724 LEMMA 5.5. *T' contains at least $|T'| - |V'|k$ colors that are universal to all vertices*
 725 *in V' .*

726 *Proof.* The list of every vertex $v \in V'$ misses at most k colors from T' . Hence all
 727 but at most $|V'|k$ colors in T' are universal to all vertices in V' . \square

728 For clarity, let us define the output problem of our compression explicitly.

729

BUDGET-CONSTRAINED LIST COLORING

Input: A graph G , a set T of colors, a list $L(v) \subseteq T$ for every $v \in V(G)$, and
 a pair (T', q) where $T' \subseteq T$ and $q \in \mathbb{N}$.

Problem: Is there a proper list coloring for G that uses at most q distinct colors
 from T' ?

730

731

732 THEOREM 5.6. $(n - k)$ -REGULAR LIST COLORING admits a compression into an
 733 instance of BUDGET-CONSTRAINED LIST COLORING with at most $11k$ vertices and
 734 $\mathcal{O}(k^2)$ colors, encodable in $\mathcal{O}(k^2 \log k)$ bits.

735 *Proof.* If $|V(G)| \leq 11k$, then G itself can be used as the output (with a dummy
 736 budget constraint). Otherwise, all the bounds above apply and Lemma 5.4 shows
 737 that the existence of a list coloring in G is equivalent to the existence of a list coloring
 738 in $G[V']$ that uses at most q colors from T' . Since $|V'| \leq 7k$, it only remains to
 739 reduce the number of colors in $T_R \cup T'$. Clearly, if $|T'| < |V'|k + q$, then $|T_R \cup T'| \leq$
 740 $3k + (7k)k \in \mathcal{O}(k^2)$ and there is nothing left to show. So suppose that $|T'| \geq |V'|k + q$.
 741 Then, it follows from Lemma 5.5 that T' contains at least q colors that are universal
 742 to the vertices in V' and we obtain an equivalent instance by removing all but exactly

743 q universal colors from T' , which leaves us with an instance with at most $|T_R| + q \leq$
 744 $3k + 7k^2 \in \mathcal{O}(k^2)$ colors, as required. Finally, to describe the output concisely, note
 745 that $G[V']$ can be trivially described in $\mathcal{O}(k^2)$ bits, and the lists $L(v)$ can be described
 746 by enumerating $T \setminus L(v)$ for every vertex v , which is k colors per vertex, each color
 747 identifiable by $\mathcal{O}(\log k)$ bits. \square

748 Note that the compression is asymptotically essentially optimal, since even the
 749 basic 4-COLORING problem does not allow a compression in $\mathcal{O}(n^{2-\varepsilon})$ bits for any
 750 $\varepsilon > 0$ unless the polynomial hierarchy collapses [24]. For completeness, we also give
 751 a proper kernel, which can be obtained in a similar manner to the compression given
 752 in Theorem 5.6.

753 **THEOREM 5.7.** $(n - k)$ -REGULAR LIST COLORING admits a kernel with $\mathcal{O}(k^2)$
 754 vertices and colors.

755 *Proof.* We distinguish two cases depending on whether or not $|T'| < |V'|k + q$. If
 756 $|T'| < |V'|k + q$, then $|T| \leq |T_R| + |T'| < 3k + |V'|k + q \leq 3k + (7k)(k + 1) \in \mathcal{O}(k^2)$.
 757 Since a list coloring requires at least one distinct color for every vertex in C , it holds
 758 that $|C| \leq |T| \leq 3k + (7k)(k + 1)$ and hence $|V(G)| \leq (3 + 7k)k + 2k \in \mathcal{O}(k^2)$, implying
 759 the desired kernel.

760 If on the other hand, $|T'| \geq |V'|k + q$, then, because of Lemma 5.5 it holds that
 761 T' contains a set U of exactly q colors that are universal to the vertices in V' . Recall
 762 that Lemma 5.4 shows that the existence of a list coloring in G is equivalent to the
 763 existence of a list coloring in $G[V']$ that uses at most $q = |T'| - |C \setminus X|$ colors from T' .
 764 It follows that the graph $G[V']$ has a list coloring using only colors in $(T_R \setminus X) \cup U$
 765 if and only if G has a list coloring. Hence, it only remains to restore the regularity
 766 of the instance. We achieve this as follows. First we add a set T_N of $|(T_R \setminus X) \cup U|$
 767 novel colors. We then add these colors (arbitrarily) to the color lists of the vertices
 768 in V' such that the size of every list (for any vertex in V') is $|(T_R \setminus X) \cup U|$. This
 769 clearly already makes the instance regular, however, now we also need to ensure that
 770 no vertex in V' can be colored with any of the new colors in T_N . To achieve this
 771 we add a set C_N of $|T_N|$ novel vertices to $G[V']$, which we connect to every vertex
 772 in $(C \cap X) \cup C_N$ and whose lists all contain all the new colors in T_N . It is clear
 773 that the constructed instance is equivalent to the original instance since all the new
 774 colors in T_N are required to color the new vertices in C_N and hence no new color
 775 can be used to color a vertex in V' . Moreover, D is still a clique modulator and
 776 the number k' of missing colors (in each list of the constructed instance) is equal to
 777 $|D| + |C \cap X| \leq 2k + 5k$ because the instance is $(n - |D| - |C \cap X|)$ -regular. Finally,
 778 the instance has at most $|V' \cup C_N| \leq 7k + 3k + 7k = 17k \in \mathcal{O}(k)$ vertices and at most
 779 $2(|T_R| + |U|) \leq 2(3k + 7k) = 20k \in \mathcal{O}(k)$ colors, as required. \square

780 **6. Saving k colors: Pre-coloring and List Coloring Variants.** In this
 781 section, we consider natural pre-coloring and list coloring variants of the “saving k
 782 colors” problem, defined as:

783

$(n - k)$ -COLORING parameterized by k

Input: A graph G with n vertices and an integer k .

Problem: Does G have a proper coloring using at most $n - k$ colors?

784

785

786 This problem is known to be FPT (it even allows for a linear kernel) [12], when
 787 parameterized by k . Notably the problem provided the main motivation for the
 788 introduction of $(n - k)$ -REGULAR LIST COLORING in [3, 2].

789 We consider the following (pre-coloring and list coloring) extensions of $(n - k)$ -
 790 COLORING.

791 $(n - |Q|)$ -PRE-COLORING EXTENSION parameterized by $n - |Q|$

Input: A graph G with n vertices and a pre-coloring $\lambda_P : X \rightarrow Q$ for $X \subseteq V(G)$ where Q is a set of colors.

Problem: Can λ_P be extended to a proper coloring of G using only colors from Q ?

792

793 LIST COLORING WITH $n - k$ COLORS parameterized by k

794

Input: A graph G on n vertices with a list $L(v)$ of colors for every $v \in V(G)$ and an integer k .

Problem: Is there a proper list coloring of G using at most $n - k$ colors?

795

796 Note that the following variant seems natural, however, is trivially NP-complete
 797 even when the parameter k is equal to 0, since the problem with an empty pre-coloring
 798 then corresponds to the problem whether G can be colored by at most $|Q|$ colors.

800 $(|Q| - k)$ -PRE-COLORING EXTENSION parameterized by k

Input: A graph G with n vertices, a pre-coloring $\lambda_P : X \rightarrow Q$ for $X \subseteq V(G)$ where Q is a set of colors, and an integer k .

Problem: Can λ_P be extended to a proper coloring of G using at most $|Q| - k$ colors from Q ?

801

802 Interestingly, we show that $(n - |Q|)$ -PRE-COLORING EXTENSION is FPT and
 803 even allows a linear kernel. Thus, we generalize the above-mentioned result of Chor
 804 et al. [12] (set $Q = [n - k]$ and $X = \emptyset$). However, LIST COLORING WITH $n - k$
 805 COLORS is easily seen to be NP-hard (even for $k = 0$) using a trivial reduction from
 806 3-Coloring.

808 **THEOREM 6.1.** $(n - |Q|)$ -PRE-COLORING EXTENSION (parameterized by $n - |Q|$)
 809 has a kernel with at most $6(n - |Q|)$ vertices and is hence fixed-parameter tractable.

810 *Proof.* Let G' be the graph obtained from G after applying the following reduction
 811 rules:

812 **Reduction Rule 6.** If u and v are two distinct vertices in $G \setminus X$ such that
 813 $\lambda_P(N_G(u)) \cup \lambda_P(N_G(v)) = Q$, then we add an edge between u and v in G .

814 This rule is safe because u and v cannot be colored with the same color.

815 **Reduction Rule 7.** If u is a vertex in $G \setminus X$ that is adjacent to a vertex $v \in X$,
 816 then we can safely add all edges between u and every vertex in $\lambda_P^{-1}(\lambda_P(v))$.

817 This rule is safe because u cannot be colored by $\lambda_P(v)$.

818 **Reduction Rule 8.** If u and v are two distinct vertices in X such that $\lambda_P(u) \neq$
 819 $\lambda_P(v)$, then we can again safely add an edge between u and v .

820 This rule is safe because u and v cannot be colored with the same color.

821 Let M be a maximal matching in the complement of G' . Note that if $|M| \leq$
 822 $n - |Q|$, then $V(M)$ is a clique modulator for G' of size at most $2(n - |Q|)$ and we
 823 obtain a kernel with at most $6(n - |Q|)$ vertices using Theorem 4.6. Thus assume that

824 $|M| \geq n - |Q|$. In this case we can safe $|M| \geq n - |Q|$ colors by giving the endpoints
 825 of every edge in M the same color. Namely, let $\{u, v\} \in M$, then:

- 826 • if $u, v \notin X$, then it follows from Reduction Rule 6 that there is a color $q \in Q$
 827 that can be given to both vertices,
- 828 • if $u \notin X$ and $v \in X$, then it follows from Reduction Rule 7 that we can color
 829 u with color $\lambda_P(v)$,
- 830 • if $u, v \in X$, then by Reduction Rule 8 we have that $\lambda_P(u) = \lambda_P(v)$.

831 Note that after coloring the edges in M with the same color, removing $V(M)$ from
 832 G' , and removing the colors used for the edges in M from Q , the number of colors in
 833 the remaining instance is equal to the number of vertices in the remaining instance,
 834 implying that the remaining instance can be properly colored. \square

835 **7. Conclusions.** We have shown several results regarding the parameterized
 836 complexity of LIST COLORING and PRE-COLORING EXTENSION problems. We
 837 showed that LIST COLORING, and hence also PRE-COLORING EXTENSION, parame-
 838 terized by the size of a clique modulator admits a randomized FPT algorithm with a
 839 running time of $\mathcal{O}^*(2^k)$, matching the best known running time of the basic CHRO-
 840 MATIC NUMBER problem parameterized by the number of vertices. This answers
 841 open questions of Golovach et al. [23]. Note that also that LIST COLORING is already
 842 $W[1]$ -hard parameterized by vertex cover [23], i.e., modulator to an independent set,
 843 which excludes even quite simple generalizations of our result to, e.g., a modulator
 844 to a disjoint union of cliques. Additionally, we showed that PRE-COLORING EXTEN-
 845 SION under the same parameter admits a linear vertex kernel with at most $3k$ vertices
 846 and that $(n - k)$ -REGULAR LIST COLORING admits a compression into a problem
 847 we call BUDGET-CONSTRAINED LIST COLORING, into an instance with at most $11k$
 848 vertices, encodable in $\mathcal{O}(k^2 \log k)$ bits. The latter also admits a proper kernel with
 849 $\mathcal{O}(k^2)$ vertices and colors. This answers an open problem of Banik et al. [3].

850 One obvious open question is whether it is possible to derandomize our algorithm
 851 for LIST COLORING. This seems, however, very challenging as it would require a
 852 derandomization of Lemma 2.4, which has been an open problem for some time. It
 853 might, however, be possible (and potentially more promising) to consider a different
 854 approach than ours. Another open question is to optimize the bound $11k$ on the
 855 number of vertices in the $(n - k)$ -REGULAR LIST COLORING compression, and/or
 856 show a proper kernel with $\mathcal{O}(k)$ vertices. Finally, another set of questions is raised
 857 by Escoffier [19], who studied the MAX COLORING problem from a “saving colors”
 858 perspective. In addition to the questions explicitly raised by Escoffier, it is natural
 859 to ask whether his problems SAVING WEIGHT and SAVING COLOR WEIGHTS admit
 860 FPT algorithms with a running time of $2^{\mathcal{O}(k)}$ and/or polynomial kernels.

861

REFERENCES

- 862 [1] Noga Alon, Gregory Z. Gutin, Eun Jung Kim, Stefan Szeider, and Anders Yeo. Solving MAX-
 863 r -SAT above a tight lower bound. *Algorithmica*, 61(3):638–655, 2011.
- 864 [2] Pranav Arora, Aritra Banik, Vijay Kumar Paliwal, and Venkatesh Raman. Some (in)tractable
 865 parameterizations of coloring and list-coloring. In Jianer Chen and Pinyan Lu, editors,
 866 *Frontiers in Algorithmics - 12th International Workshop, FAW 2018, Proceedings*, volume
 867 10823 of *Lecture Notes in Computer Science*, pages 126–139. Springer, 2018.
- 868 [3] Aritra Banik, Ashwin Jacob, Vijay Kumar Paliwal, and Venkatesh Raman. Fixed-parameter
 869 tractability of $(n - k)$ List Coloring. In C.J. Colbourn, R. Grossi, and N. Pisani, edi-
 870 tors, *IWOCA 2019, Proceedings*, volume 11638 of *Lecture Notes in Computer Science*,
 871 pages 61–69, 2019. URL: https://doi.org/10.1007/978-3-030-25005-8_6, doi:10.1007/
 872 978-3-030-25005-8_6.

- 873 [4] D. Berend and T. Tassa. Improved bounds on bell numbers and on moments of sums of random
874 variables. *Probability and Mathematical Statistics*, 30:185–205, 2010.
- 875 [5] Andreas Björklund, Thore Husfeldt, Petteri Kaski, and Mikko Koivisto. Narrow sieves for
876 parameterized paths and packings. *J. Comput. Syst. Sci.*, 87:119–139, 2017.
- 877 [6] Andreas Björklund, Thore Husfeldt, and Mikko Koivisto. Set partitioning via inclusion-
878 exclusion. *SIAM J. Comput.*, 39(2):546–563, 2009.
- 879 [7] Andreas Björklund, Petteri Kaski, and Lukasz Kowalik. Constrained multilinear detection and
880 generalized graph motifs. *Algorithmica*, 74(2):947–967, 2016.
- 881 [8] Hans L. Bodlaender, Bart M. P. Jansen, and Stefan Kratsch. Kernel bounds for path and cycle
882 problems. *Theor. Comput. Sci.*, 511:117–136, 2013.
- 883 [9] James R. Bunch and John E. Hopcroft. Triangular factorization and inversion by fast matrix
884 multiplication. *Mathematics of Computation*, 28(125):231–236, 1974.
- 885 [10] Leizhen Cai. Parameterized complexity of vertex colouring. *Discrete Applied Mathematics*,
886 127(3):415–429, 2003.
- 887 [11] Jianer Chen, Iyad A. Kanj, and Ge Xia. Improved upper bounds for vertex cover. *Theor.*
888 *Comput. Sci.*, 411(40-42):3736–3756, 2010.
- 889 [12] Benny Chor, Mike Fellows, and David Juedes. Linear kernels in linear time, or how to save
890 k colors in $O(n^2)$ steps. In *Proceedings of the 30th International Conference on Graph-*
891 *Theoretic Concepts in Computer Science*, WG’04, pages 257–269. Springer-Verlag, 2004.
- 892 [13] Maria Chudnovsky. Coloring graphs with forbidden induced subgraphs. In *2014 International*
893 *Congress of Mathematicians*, volume IV, pages 291–302, 2014.
- 894 [14] M. Cygan, F.V. Fomin, L. Kowalik, D. Lokshtanov, D. Marx, M. Pilipczuk, M. Pilipczuk, and
895 S. Saurabh. *Parameterized Algorithms*. Springer, 2015.
- 896 [15] Marek Cygan, Holger Dell, Daniel Lokshtanov, Dániel Marx, Jesper Nederlof, Yoshio Okamoto,
897 Ramamohan Paturi, Saket Saurabh, and Magnus Wahlström. On problems as hard as
898 CNF-SAT. *ACM Trans. Algorithms*, 12(3):41:1–41:24, 2016.
- 899 [16] Reinhard Diestel. *Graph Theory, 4th Edition*, volume 173 of *Graduate texts in mathematics*.
900 Springer, 2012.
- 901 [17] Michael Dom, Daniel Lokshtanov, and Saket Saurabh. Kernelization Lower Bounds Through
902 Colors and IDs. *ACM Trans. Algorithms*, 11(2):13:1–13:20, 2014.
- 903 [18] R.G. Downey and M.R. Fellows. *Fundamentals of Parameterized Complexity*. Springer, 2013.
- 904 [19] Bruno Escoffier. Saving colors and max coloring: Some fixed-parameter tractability results.
905 *Theor. Comput. Sci.*, 758:30–41, 2019. URL: <https://doi.org/10.1016/j.tcs.2018.08.002>,
906 doi:10.1016/j.tcs.2018.08.002.
- 907 [20] F.V. Fomin, D. Lokshtanov, S. Saurabh, and M. Zehavi. *Kernelization*. Cambridge Univ. Press,
908 2019.
- 909 [21] M. R. Garey and David S. Johnson. *Computers and Intractability: A Guide to the Theory of*
910 *NP-Completeness*. W. H. Freeman, 1979.
- 911 [22] Petr A. Golovach, Matthew Johnson, Daniël Paulusma, and Jian Song. A survey on the
912 computational complexity of coloring graphs with forbidden subgraphs. *Journal of Graph*
913 *Theory*, 84(4):331–363, 2017.
- 914 [23] Petr A. Golovach, Daniël Paulusma, and Jian Song. Closing complexity gaps for coloring
915 problems on h -free graphs. *Inf. Comput.*, 237:204–214, 2014.
- 916 [24] Bart M. P. Jansen and Astrid Pieterse. Sparsification upper and lower bounds for graph
917 problems and not-all-equal SAT. *Algorithmica*, 79(1):3–28, 2017.
- 918 [25] Tommy R. Jensen and Bjarne Toft. *Graph Coloring Problems*. Wiley & Sons, 1994.
- 919 [26] Ioannis Koutis. *Multilinear Monomial Detection*, pages 1375–1378. Springer New York,
920 New York, NY, 2016. URL: https://doi.org/10.1007/978-1-4939-2864-4_784, doi:10.1007/
921 978-1-4939-2864-4_784.
- 922 [27] Robert Krauthgamer and Ohad Trabelsi. The set cover conjecture and subgraph isomorphism
923 with a tree pattern. In Rolf Niedermeier and Christophe Paul, editors, *36th International*
924 *Symposium on Theoretical Aspects of Computer Science, STACS 2019, March 13-16, 2019,*
925 *Berlin, Germany*, volume 126 of *LIPICs*, pages 45:1–45:15. Schloss Dagstuhl - Leibniz-
926 Zentrum fuer Informatik, 2019. URL: <https://doi.org/10.4230/LIPICs.STACS.2019.45>,
927 doi:10.4230/LIPICs.STACS.2019.45.
- 928 [28] L. Lovász. Coverings and coloring of hypergraphs. In *4th Southeastern Conference on Combi-*
929 *natorics, Graph Theory, and Computing, Utilitas Math*, pages 3–12, 1973.
- 930 [29] Silvio Micali and Vijay V. Vazirani. An $o(\sqrt{(|v|) |e|})$ algorithm for finding maximum match-
931 ing in general graphs. In *21st Annual Symposium on Foundations of Computer Science,*
932 *Syracuse, New York, USA, 13-15 October 1980*, pages 17–27. IEEE Computer Society,
933 1980.
- 934 [30] Daniël Paulusma. Open problems on graph coloring for special graph classes. In Ernst W. Mayr,

- 935 editor, *Graph-Theoretic Concepts in Computer Science - 41st International Workshop,*
936 *WG 2015, Garching, Germany, June 17-19, 2015, Revised Papers*, volume 9224 of *Lecture*
937 *Notes in Computer Science*, pages 16–30. Springer, 2015.
- [31] B. Randerath and I. Schiermeyer. Vertex coloring and forbidden subgraphs - a survey. *Graphs*
939 *Comb.*, 20:1–40, 2004.
- [32] J. T. Schwartz. Fast probabilistic algorithms for verification of polynomial identities. *J. ACM*,
940 27(4):701–717, October 1980.
- [33] Z. Tuza. Graph colorings with local restrictions - a survey. *Discussiones Mathematicae Graph*
942 *Theory*, 17:161–228, 1997.
- [34] Magnus Wahlström. Abusing the Tutte matrix: An algebraic instance compression for the
944 K -set-cycle problem. In Natacha Portier and Thomas Wilke, editors, *30th International*
945 *Symposium on Theoretical Aspects of Computer Science, STACS 2013, February 27 -*
946 *March 2, 2013, Kiel, Germany*, volume 20 of *LIPICs*, pages 341–352. Schloss Dagstuhl -
947 Leibniz-Zentrum fuer Informatik, 2013.
- [35] F. Yates. The design and analysis of factorial experiments. Technical Report Technical Com-
949 munication no. 35, Commonwealth Bureau of Soils, 1937.
- [36] Richard Zippel. Probabilistic algorithms for sparse polynomials. In *Proceedings of the Inter-*
951 *national Symposium on Symbolic and Algebraic Computation, EUROSAM '79*, pages
952 216–226, 1979.
953