

# Recent Developments of the Autoregressive Distributed Lag Modelling Framework\*

JIN SEO CHO

School of Humanities and Social Sciences, Beijing Institute of Technology, Haidan, Beijing 100081, China  
School of Economics, Yonsei University, Seodaemun-gu, Seoul 03722, Korea  
[jinseocho@yonsei.ac.kr](mailto:jinseocho@yonsei.ac.kr)

MATTHEW GREENWOOD-NIMMO

Faculty of Business and Economics, University of Melbourne, Carlton, VIC 3053, Australia  
Centre for Applied Macroeconomic Analysis, Australian National University, Canberra, ACT 2600, Australia  
[matthew.greenwood@unimelb.edu.au](mailto:matthew.greenwood@unimelb.edu.au)

YONGCHEOL SHIN

Department of Economics and Related Studies, University of York, Heslington, York, YO10 5DD, U.K.  
[yongcheol.shin@york.ac.uk](mailto:yongcheol.shin@york.ac.uk)

June 22, 2021

## Abstract

We review the literature on the Autoregressive Distributed Lag (ARDL) model, from its origins in the analysis of autocorrelated trend stationary processes to its subsequent applications in the analysis of cointegrated non-stationary time series. We then survey several recent extensions of the ARDL model, including asymmetric and nonlinear generalisations of the ARDL model, the quantile ARDL model, the pooled mean group dynamic panel data model and the spatio-temporal ARDL model.

**Key Words:** Autoregressive Distributed Lag (ARDL) Model; Asymmetry, Nonlinearity and Threshold Effects; Quantile Regression; Panel Data; Spatial Analysis.

**JEL Classifications:** C22.

---

\*Corresponding author: Matthew Greenwood-Nimmo. Cho is grateful for financial support from the Yonsei University Research Grant of 2020. The usual disclaimer applies.

# 1 Introduction

The ARDL model combines an autoregressive component (i.e. lags of a scalar dependent variable) with a distributed lag component (i.e. lags of a vector of explanatory variables). It has its origins in the analysis of autocorrelated trend stationary processes. In this context, the general practice is to model the de-trended series as a stationary distributed lag or ARDL model (Koyck, 1954; Almon, 1965). Provided that the lag structure of the ARDL model is sufficiently rich to account for the autocorrelation structure in the data, estimation can proceed by ordinary least squares (OLS) and standard inference is applicable with respect to the long-run properties of the model. The application of the ARDL model to trend stationary data has been addressed in several excellent survey articles, so we do not repeat this discussion here; rather, the reader is referred to Griliches (1967), Wallis (1969), Nerlove (1972), Sims (1974), Maddala (1977), Thomas (1977), Zellner (1979), Hendry, Pagan, and Sargan (1984) and Wickens and Breusch (1988). For a formal treatment of the probability theory underlying many of the proposed estimators, see Dhrymes (1971).

We focus our attention on the more challenging case first considered by Pesaran and Shin (1998), in which the ARDL model is applied to first-order integrated, or I(1), processes. Pesaran, Shin, and Smith (2001) provide an important generalisation that makes use of a bounds-testing framework to allow for mixed orders of integration among the variables entering the ARDL model. The methodology proposed in these two papers has proven highly influential, having given rise to thousands of empirical applications and having provided a foundation for several notable methodological extensions.

In this paper, we concisely review the specification and properties of the ARDL model of Pesaran and Shin (1998) and Pesaran et al. (2001) before surveying six important methodological extensions. First, we introduce the Nonlinear ARDL (NARDL) model developed by Shin, Yu, and Greenwood-Nimmo (2014), which makes use of partial sum decompositions of the explanatory variables to accommodate asymmetric phenomena. We review the OLS estimation procedure proposed by Shin et al. (2014) and explain how the asymptotic singularity issue mentioned by the authors arises due to the use of partial sum decompositions. This singularity problem frustrates efforts to obtain the limit distribution of the estimator proposed by Shin et al. and thus provides motivation for the development of an alternative two-step estimation framework by Cho, Greenwood-Nimmo, and Shin (2019, 2020b). In these two papers, Cho et al. employ a novel re-parameterisation of the NARDL model to eliminate the singularity problem and then develop asymptotic theory for their two-step estimator, in which the parameters of the long-run relationship are estimated in

the first step using the fully-modified OLS (FM-OLS) estimator of [Phillips and Hansen \(1990\)](#) before the short-run dynamic parameters are estimated in the second step by OLS. This procedure exploits the different convergence rates of the first and second step estimators to deliver consistent and asymptotically normal estimators of both the long- and short-run parameters. This asymptotic normality allows the authors to employ [Wald's \(1943\)](#) testing principle to develop tests for both long- and short-run asymmetry that asymptotically converge to  $\chi^2$  distributions.

The NARDL model has been widely adopted in the literature due to its ease of implementation and interpretation. Part of this is due to the simplifying assumption that the threshold parameter embedded in the NARDL model is known a priori. Typically, one uses a threshold value of zero in the construction of the partial sum processes, which gives rise to an elegant interpretation related to positive and negative changes in the vector of explanatory variables. From an inferential perspective, the use of a known threshold simplifies the analysis because it ensures that the identification problem described by [Davies \(1977, 1987\)](#) does not arise. However, in some cases, one may wish to estimate the threshold parameter. The resulting model is known as the Threshold ARDL (TARDL) model, estimation and inference on which is the focus of [Cho, Greenwood-Nimmo, and Shin \(2020c,d\)](#).

[Cho et al. \(2020c\)](#) consider a setting with a single explanatory variable that is decomposed into partial sum processes around an unknown momentum-type threshold. In this environment, the authors demonstrate that one must address a multifold identification problem in order to conduct inference on the number of regimes. The multifold identification problem arises because the null hypothesis of a single regime is composed of three sub-null hypotheses, such that the alternative hypothesis is defined by the negation of the union of these sub-null hypotheses. Testing the joint null hypothesis is complicated by the fact that each sub-null hypothesis represents an alternative of the other sub-null hypotheses. Drawing on the existing literature on the multifold identification problem, [Cho et al. \(2020c\)](#) develop a quasi likelihood ratio (QLR) test for the existence of a distinct threshold level and show that the null limit distribution of the test statistic can be represented by separable functionals of two Gaussian stochastic processes provided that the TARDL specification contains an intercept term.

[Cho et al. \(2020d\)](#) pursue a different approach with the objective of determining the number of regimes in a TARDL model with a single explanatory variable via a model selection approach. The authors consider six candidate information criteria, namely the Akaike information criterion (AIC), the Schwarz information criterion (SIC) and the Hannan-Quinn information criterion (HQIC), as well as modified versions of each

criterion defined following [Pitarakis \(2006\)](#), which we shall denote pAIC, pSIC and pHQIC, respectively. Using extensive simulation studies, [Cho et al. \(2020d\)](#) show that, in general, the standard SIC outperforms the other information criteria if the independent variable is not driven by a time drift. However, if this is not the case, then the pSIC outperforms the other information criteria in small samples, while SIC and pSIC jointly outperform the remainder in large samples.

[Cho, Kim, and Shin \(2015\)](#) were the first to apply quantile regression following [Koenker and Bassett \(1978\)](#) and [Xiao \(2009\)](#) to estimate the unknown parameters of an ARDL model. This allows researchers to evaluate the error correction relationship between a scalar dependent variable,  $y_t$ , and a vector of explanatory variables,  $\mathbf{x}_t$ , throughout the conditional distribution of  $y_t|\mathbf{x}_t$ . This contrasts with the classical estimators employed by [Pesaran and Shin \(1998\)](#) and [Pesaran et al. \(2001\)](#) that characterise the relationship at the conditional mean only. [Cho et al. \(2015\)](#) examine the large sample properties of the quantile regression estimator and show that the joint limit distribution of the long-run parameter estimator is mixed-normal, which implies that asymptotic critical values for Wald statistics testing various hypotheses on the quantile coefficients can be retrieved from  $\chi^2$  distributions. In a subsequent paper, [Cho, Greenwood-Nimmo, Kim, and Shin \(2020a\)](#) apply quantile regression in two steps to the estimation of a NARDL model.

The ARDL specification has also been applied to large  $T$  panels – that is, panels with many time series observations per group. [Pesaran, Shin, and Smith \(1999\)](#) consider a setting in which a panel ARDL specification is estimated under the assumption that the long-run parameters are homogeneous across groups, while the short-run dynamic parameters are heterogeneous. This setting has intuitive appeal, as there are often good reasons to believe that long-run equilibrium relationships should be approximately common across groups but, in general, the same cannot be said for dynamic parameters. The so-called Pooled Mean Group (PMG) estimator employs an innovative hybrid estimation approach, whereby the homogeneous long-run parameters are estimated by maximum likelihood with the data pooled over groups, while the heterogeneous short-run parameters are estimated on a group-specific basis and their group-wide distribution is summarised by taking averages across groups. In the case of stationary data, [Pesaran et al.](#) demonstrate that the PMG estimator is consistent and asymptotically normal while, under non-stationarity, it is consistent and converges to a mixed-normal distribution.

The final model that we consider is the Spatio-Temporal ARDL (STARDL) model of [Shin and Thornton \(2019\)](#). This model exploits the growing availability of spatial time series data. The STARDL model is a system of ARDL equations, each of which is augmented with variables defined using spatially-weighted

averages. [Shin and Thornton](#) develop a quasi-maximum likelihood estimator as well as an alternative estimator based on the control function framework. In both cases, the authors demonstrate that their estimators are consistent and asymptotically normal. The STARDL framework can be thought of as an omnibus model that nests several popular spatial models, including the spatial Durbin model analysed by [Lee and Yu \(2010\)](#) and [Elhorst \(2014\)](#) and the heterogeneous spatial autoregressive panel data model of [Aquaro, Bailey, and Pesaran \(2021\)](#). [Shin and Thornton](#) also note that the network structure of the STARDL model is amenable to the application of many of the popular tools of network analysis, including centrality statistics and clustering algorithms. By analogy to the dynamic multiplier effects that are widely used in the analysis of ARDL models, the authors derive diffusion multipliers from the spatial system that can be used to explore the properties of the spatial dynamic network.

This paper proceeds in eight sections. In Section 2, we briefly review the ARDL model of [Pesaran and Shin \(1998\)](#) and [Pesaran et al. \(2001\)](#). In Section 3, we introduce the classic NARDL model of [Shin et al. \(2014\)](#) based on a known threshold, before discussing the case of unknown thresholds in the context of the TARDL model. In Section 4, we outline the quantile ARDL and quantile NARDL models, while Sections 5 and 6 are devoted to the PMG estimator of [Pesaran et al. \(1999\)](#) and the STARDL model of [Shin and Thornton \(2019\)](#), respectively. In Section 7, we discuss some promising avenues for continuing development of the ARDL framework and we conclude in Section 8.

## 2 The ARDL Model with Non-Stationary Data

[Pesaran and Shin \(1998\)](#) study the following process:

$$y_t = \gamma_* + \sum_{j=1}^p \phi_{j*} y_{t-j} + \sum_{j=0}^q \theta'_{j*} \mathbf{x}_{t-j} + \varepsilon_t, \quad (1)$$

which is obtained by applying the distributed-lag form to the integrated time series  $(y_t, \mathbf{x}'_t)' \in \mathbb{R}^{1+k}$ . [Pesaran and Shin](#) refer to (1) as an *ARDL(p, q) process*. The authors go on to demonstrate that (1) can be expressed in the following alternative form:

$$y_t = \gamma_* + \mathbf{x}'_t \boldsymbol{\gamma}_* + \sum_{j=1}^p \phi_{j*} y_{t-j} + \sum_{j=0}^{q-1} \Delta \mathbf{x}'_{t-j} \boldsymbol{\delta}_{j*} + \varepsilon_t, \quad (2)$$

where  $\boldsymbol{\gamma}_* := \sum_{j=0}^q \boldsymbol{\theta}_{j*}$  and  $\boldsymbol{\delta}_{j*} := -\sum_{i=j+1}^q \boldsymbol{\theta}_{i*}$ . The long-run relationship between  $y_t$  and  $\mathbf{x}_t$  embedded

in (2) can be represented as  $y_t = \mu_* + \mathbf{x}'_t \boldsymbol{\beta}_* + u_t$ , where:

$$\mu_* := \gamma_* \left( 1 - \sum_{i=1}^p \phi_{i*} \right)^{-1}, \quad \boldsymbol{\beta}_* := \gamma_* \left( 1 - \sum_{i=1}^p \phi_{i*} \right)^{-1},$$

and  $u_t$  is a stationary process defined by  $\{\Delta \mathbf{x}_t, \varepsilon_t, \Delta \mathbf{x}_{t-1}, \varepsilon_{t-1}, \dots\}$ .

The ARDL process is an error correction process in the tradition of the London School of Economics (e.g. [Sargan, 1964](#); [Hendry and Mizon, 1978](#), among others). A typical error correction process can be written as follows:

$$\Delta y_t = \rho_* y_{t-1} + \boldsymbol{\theta}'_* \mathbf{x}_{t-1} + \gamma_* + \sum_{j=1}^{p-1} \varphi_{j*} \Delta y_{t-j} + \sum_{j=0}^{q-1} \boldsymbol{\pi}'_{j*} \Delta \mathbf{x}_{t-j} + \varepsilon_t, \quad (3)$$

where  $\{\varepsilon_t; \mathcal{F}_t^0\}$  is a martingale difference sequence and  $\mathcal{F}_t^0$  is the smallest  $\sigma$ -algebra driven by  $\{y_{t-1}, \mathbf{x}_t, y_{t-2}, \mathbf{x}_{t-1}, \dots\}$ . Note that the error correction process (3) can be derived from the ARDL process (1) by letting:

$$\rho_* := \sum_{j=1}^p \phi_{j*} - 1, \quad \boldsymbol{\theta}_* := \sum_{j=0}^q \boldsymbol{\theta}_{j*}, \quad \boldsymbol{\pi}_{0*} := \boldsymbol{\theta}_{0*}, \quad \varphi_{\ell*} := - \sum_{i=\ell+1}^p \phi_{i*}, \quad \text{and} \quad \boldsymbol{\pi}_{j*} := - \sum_{i=j+1}^q \boldsymbol{\theta}_{i*}$$

for  $\ell = 1, 2, \dots, p-1$  and  $j = 1, 2, \dots, q-1$ . Furthermore, if  $y_t$  is cointegrated with  $\mathbf{x}_t$  such that  $u_{t-1} := y_{t-1} - \boldsymbol{\beta}'_* \mathbf{x}_{t-1}$  is a stationary cointegration error, then (3) can be re-written as:

$$\Delta y_t = \rho_* u_{t-1} + \gamma_* + \sum_{j=1}^{p-1} \varphi_{j*} \Delta y_{t-j} + \sum_{j=0}^{q-1} \boldsymbol{\pi}'_{j*} \Delta \mathbf{x}_{t-j} + \varepsilon_t, \quad (4)$$

where  $\boldsymbol{\beta}_* := -(\boldsymbol{\theta}_*/\rho_*)$ , which provides motivation for the separate estimation of the long-run and short-run parameters. [Engle and Granger \(1987\)](#) estimate the long-run parameter by OLS, while [Phillips and Hansen \(1990\)](#) propose to estimate the same parameter using the fully-modified OLS (FM-OLS) estimator, which overcomes the asymptotic bias in the OLS estimator and which asymptotically follows a mixed normal distribution. Many other procedures for the consistent and efficient estimation of the long-run coefficients have been proposed in the literature (e.g. [Stock and Watson, 1993](#); [Johansen, 1988](#), among others). [Pesaran and Shin \(1998\)](#) propose a single-step procedure, in which one first estimates the coefficients of (2) by OLS and then estimates  $\boldsymbol{\beta}_*$  as  $\widehat{\boldsymbol{\beta}}_T := \widehat{\boldsymbol{\gamma}}_T \left( 1 - \sum_{i=1}^p \widehat{\phi}_{T,i} \right)^{-1}$ , where  $\widehat{\boldsymbol{\gamma}}_T$  and  $\widehat{\phi}_{T,i}$  are the OLS estimators of  $\gamma_*$  and  $\phi_{i*}$ , respectively. The authors show that  $\widehat{\boldsymbol{\beta}}_T$  converges to  $\boldsymbol{\beta}_*$  at the rate of  $T$  and asymptotically follows a mixed normal distribution, while the short-run parameter estimators converge to the unknown

parameters in (2) at the rate of  $\sqrt{T}$  and are asymptotically normal. For this reason, Pesaran and Shin (1998) are able to apply Wald's (1943) testing principle to conduct inference on the unknown long-run and short-run parameters, with simulation evidence indicating that the behaviour of the Wald test statistics is well-approximated by asymptotic results even in relatively small samples. In addition, Pesaran and Shin (1998) suggest to select the lag orders of the ARDL process using the Schwarz (1978) information criterion, although other approaches including general-to-specific lag selection are also commonly used in practice.

In an influential contribution to the ARDL literature, Pesaran et al. (2001) develop a bounds test for the null hypothesis of no cointegration using the error correction representation of the ARDL process. The authors derive the null limit distributions of the F-statistic testing the null hypothesis  $\rho_* = 0$  and  $\theta_* = \mathbf{0}$  in (3) as functionals of a Wiener process under various assumptions on the model coefficients and the order of integration of the variables in the model. Pesaran et al. (2001) tabulate critical value bounds from these asymptotic distributions for use in applied research where one is unsure of the true order of integration of the variables entering the model.

The ARDL framework developed by Pesaran and Shin (1998) and Pesaran et al. (2001) has been widely adopted in applied research. At the time of writing, Google Scholar records approximately 5,500 and 12,500 citations to these two papers, respectively. Applications of the ARDL model can be found in many areas of the social sciences, with a particular prevalence in finance, macroeconomics and energy economics. There are many reasons for the popularity of the ARDL model. Perhaps most importantly, the functional form of the ARDL( $p, q$ ) process has intuitive appeal, as it allows for partial adjustment toward an economically meaningful long-run equilibrium relationship between  $y_t$  and  $x_t$ . In addition, the ARDL model naturally accounts for the serial correlation structure that exists among the first differences of  $y_t$  and  $x_t$  and can provide consistent estimates of the long-run parameters in the presence of weak endogeneity.

In addition to the many applications of the ARDL model, a considerable body of work has been dedicated to the development of variants of the ARDL model that allow for departures from linearity, among other phenomena. The remainder of this paper is dedicated to reviewing several of these developments.

### **3 Nonlinearity and Threshold Effects**

The nonlinear autoregressive distributed lag (NARDL) and threshold autoregressive distributed lag (TARDL) models represent two notable variants of the ARDL model that employ partial sum decompositions of the

explanatory variables to accommodate asymmetry and nonlinearity, respectively. Consider the following specification:

$$y_t = \gamma_* + \sum_{j=1}^p \phi_{j*} y_{t-j} + \sum_{j=0}^q (\theta_{j*}^+ \mathbf{x}_{t-j}^+ + \theta_{j*}^- \mathbf{x}_{t-j}^-) + \varepsilon_t, \quad (5)$$

where  $\mathbf{x}_t \in \mathbb{R}^k$ ,  $\mathbf{x}_t^+ := \sum_{j=1}^t \Delta \mathbf{x}_j^+$  and  $\mathbf{x}_t^- := \sum_{j=1}^t \Delta \mathbf{x}_j^-$  with:

$$\Delta x_{tj}^+ := \begin{cases} 0, & \text{if } \tau_{j*} \geq \Delta x_{tj}; \\ \Delta x_{tj}, & \text{otherwise,} \end{cases} \quad \text{and} \quad \Delta x_{tj}^- := \begin{cases} 0, & \text{if } \tau_{j*} < \Delta x_{tj}; \\ \Delta x_{tj}, & \text{otherwise.} \end{cases}$$

In this case,  $\{\Delta \mathbf{x}_t\}$  is a strictly stationary process and we define  $\boldsymbol{\tau}_* := [\tau_{1*}, \dots, \tau_{j*}, \dots, \tau_{k*}]'$ . If the threshold level,  $\tau_*$ , is known, then although (5) is linear in the unknown parameters, it can accommodate nonlinearity in the long-run and/or the short-run relationships between  $y_t$  and  $\mathbf{x}_t$ . This is the *NARDL(p, q) model* proposed by Shin et al. (2014). If the threshold parameter,  $\tau_*$ , is unknown, then one must estimate it in order to evaluate the nonlinear relationship between  $y_t$  and  $\mathbf{x}_t$ . In this case, the estimation problem is nonlinear due to the necessity to estimate the threshold parameter. This is the *TARDL(p, q) model* advanced by Cho et al. (2020c,d).

As with the linear ARDL process discussed in the previous section, the NARDL/TARDL process in (5) can be transformed into an error correction model. Specifically, for some  $\rho_*$ ,  $\boldsymbol{\theta}_*^+$ ,  $\boldsymbol{\theta}_*^-$ ,  $\varphi_{j*}$  ( $j = 1, 2, \dots, p-1$ ),  $\boldsymbol{\pi}_{j*}^+$ , and  $\boldsymbol{\pi}_{j*}^-$  ( $j = 0, 1, \dots, q-1$ ), we can re-write (5) as follows:

$$\Delta y_t = \rho_* y_{t-1} + \boldsymbol{\theta}_*^+ \mathbf{x}_{t-1}^+ + \boldsymbol{\theta}_*^- \mathbf{x}_{t-1}^- + \gamma_* + \sum_{j=1}^{p-1} \varphi_{j*} \Delta y_{t-j} + \sum_{j=0}^{q-1} (\boldsymbol{\pi}_{j*}^+ \Delta \mathbf{x}_{t-j}^+ + \boldsymbol{\pi}_{j*}^- \Delta \mathbf{x}_{t-j}^-) + \varepsilon_t, \quad (6)$$

where  $\{\varepsilon_t, \mathcal{F}_t\}$  is a martingale difference sequence,  $\mathcal{F}_t$  is the smallest  $\sigma$ -algebra driven by  $\{y_{t-1}, \mathbf{x}_t^+, \mathbf{x}_t^-, y_{t-2}, \mathbf{x}_{t-1}^+, \mathbf{x}_{t-1}^-, \dots\}$  and the parameters of (6) are defined by the parameters of (5) as follows:

$$\rho_* := \sum_{j=1}^p \phi_{j*} - 1, \quad \boldsymbol{\theta}_*^+ := \sum_{j=0}^q \boldsymbol{\theta}_{j*}^+, \quad \boldsymbol{\theta}_*^- := \sum_{j=0}^q \boldsymbol{\theta}_{j*}^-, \quad \boldsymbol{\pi}_{0*}^+ := \boldsymbol{\theta}_{0*}^+, \quad \boldsymbol{\pi}_{0*}^- := \boldsymbol{\theta}_{0*}^-,$$

and for  $\ell = 1, 2, \dots, p-1$  and  $j = 1, 2, \dots, q-1$ :

$$\varphi_{\ell*} := - \sum_{i=\ell+1}^p \phi_{i*}, \quad \boldsymbol{\pi}_{j*}^+ := - \sum_{i=j+1}^q \boldsymbol{\theta}_{i*}^+, \quad \text{and} \quad \boldsymbol{\pi}_{j*}^- := - \sum_{i=j+1}^q \boldsymbol{\theta}_{i*}^-.$$



If  $y_t$  is cointegrated with  $(\mathbf{x}_t^+, \mathbf{x}_t^-)'$  such that  $u_{t-1} := y_{t-1} - \beta_*^+ \mathbf{x}_{t-1}^+ - \beta_*^- \mathbf{x}_{t-1}^-$  is a cointegration error, then (6) can be re-written as follows:

$$\Delta y_t = \rho_* u_{t-1} + \gamma_* + \sum_{j=1}^{p-1} \varphi_{j*} \Delta y_{t-j} + \sum_{j=0}^{q-1} \left( \pi_{j*}^+ \Delta \mathbf{x}_{t-j}^+ + \pi_{j*}^- \Delta \mathbf{x}_{t-j}^- \right) + \varepsilon_t, \quad (7)$$

where  $\{u_t\}$  is a stationary process that is possibly correlated with  $\Delta \mathbf{x}_t$ ,  $\beta_*^+ := -(\boldsymbol{\theta}_*^+ / \rho_*)$  and  $\beta_*^- := -(\boldsymbol{\theta}_*^- / \rho_*)$  are the long-run parameters of the NARDL/TARDL process and the remaining parameters capture the short-run dynamics. The error correction form in (7) has the interesting implication that, even if the population mean of  $\Delta \mathbf{x}_t$  is zero,  $y_t$  can be an integrated process with a time drift, because both  $\Delta \mathbf{x}_t^+$  and  $\Delta \mathbf{x}_t^-$  cannot have zero population mean. This is a different form of cointegrating relationship from the standard case, in which a cointegrating relationship between integrated processes with and without a time drift is driven by the presence of a non-zero intercept in (7).

The NARDL framework has been adopted widely in the social sciences, because it provides a simple method for the analysis of asymmetries of the type that may arise in pricing problems or exchange rate pass-through, for example. Furthermore, [Shin et al. \(2014\)](#) show that one may construct cumulative dynamic multipliers from the estimated parameters of the NARDL model, which provides an easily interpreted visualisation of the traverse to an equilibrium position following a shock. Unlike structural impulse response analysis, the cumulative dynamic multipliers do not rely on controversial procedures for the identification of structural shocks. Due in part to these desirable attributes, Google Scholar reports more than 1,100 citations to [Shin et al. \(2014\)](#) at the time of writing, with applications in diverse fields such as criminology (e.g. [Box, Gratzer, and Lin, 2018](#)), energy economics (e.g. [Hammoudeh, Lahiani, Nguyen, and Sousa, 2015](#)), financial economics (e.g. [He and Zhou, 2018](#)), monetary economics (e.g. [Claus and Nguyen, 2019](#)) and tourism (e.g. [Süssmuth and Woitek, 2013](#)), among others. The majority of existing NARDL applications follow [Shin et al. \(2014\)](#) and employ a single known threshold value of zero, but several studies including [Fedoseeva \(2013\)](#) and [Pal and Mitra \(2015\)](#) have used multiple thresholds. Interest in the use of non-zero thresholds was an important motivating factor behind the development of the TARDL model by [Cho et al. \(2020c,d\)](#).

### 3.1 Estimation and Inference for the NARDL Model

[Shin et al. \(2014\)](#) propose to estimate the NARDL model in a single step by OLS. However, they note that efforts to derive asymptotic theory for the single-step OLS estimator are frustrated by an asymptotic

singularity problem that arises from the presence of partial sum decompositions of the explanatory variables. Consequently, [Shin et al.](#) provide simulation evidence regarding the performance of the single-step estimator but the corresponding theory has yet to be developed. [Cho et al. \(2019\)](#) study the asymptotic singularity issue in detail and develop theory for a new two-step estimation framework. Let:

$$\mathbf{z}_t := \left[ \begin{array}{cccccccc} y_{t-1} & \mathbf{x}_{t-1}^{+'} & \mathbf{x}_{t-1}^{-'} & 1 & \Delta \mathbf{y}'_{t-1} & \Delta \mathbf{x}_t^{+'} & \dots & \Delta \mathbf{x}_{t-q+1}^{+'} & \Delta \mathbf{x}_t^{-'} & \dots & \Delta \mathbf{x}_{t-q+1}^{-'} \end{array} \right]',$$

where  $\Delta \mathbf{y}_{t-1} := [\Delta y_{t-1}, \Delta y_{t-2}, \dots, \Delta y_{t-p+1}]'$ . Under the assumption that  $\mathbb{E}[\Delta \mathbf{x}_t] = \mathbf{0}$ , [Cho et al. \(2019\)](#) show that  $\mathbf{D}_T^{-1} (\sum_{t=1}^T \mathbf{z}_t \mathbf{z}_t') \mathbf{D}_T^{-1}$  converges to a singular matrix, where  $\mathbf{D}_T := \text{diag}(T^{3/2} \mathbf{I}_{2+2k}, T^{1/2} \mathbf{I}_{p+2qk})$ . This is the singularity to which [Shin et al. \(2014\)](#) allude and which provides motivation for the development of a two-step estimation strategy by [Cho et al. \(2019\)](#). Consider the long-run relationship embedded in the NARDL model, which can be written as follows:

$$y_t = \varsigma_* + \mathbf{x}_t^{+'} \boldsymbol{\beta}_*^+ + \mathbf{x}_t^{-'} \boldsymbol{\beta}_*^- + u_t.$$

[Cho et al. \(2019\)](#) show that another asymptotic singularity issue arises here, as the matrix inverse required to obtain the OLS estimator of  $(\varsigma_*, \boldsymbol{\beta}_*^{+'}, \boldsymbol{\beta}_*^{-'})'$  is asymptotically singular, because  $\bar{\mathbf{D}}_T^{-1} \left( \sum_{t=1}^T \mathbf{v}_t \mathbf{v}_t' \right) \bar{\mathbf{D}}_T^{-1}$  converges to a singular matrix, where  $\mathbf{v}_t := (1, \mathbf{x}_t^{+'}, \mathbf{x}_t^{-'})'$  and  $\bar{\mathbf{D}}_T := \text{diag}(T^{1/2}, T^{3/2} \mathbf{I}_{2k})$ . To overcome this problem, the authors re-parameterise the long-run relationship as follows:

$$y_t = \varsigma_* + \mathbf{x}_t^{+'} \boldsymbol{\lambda}_* + \mathbf{x}_t^{-'} \boldsymbol{\eta}_* + u_t$$

where  $\mathbf{x}_t \equiv \mathbf{x}_t^+ + \mathbf{x}_t^-$ ,  $\boldsymbol{\lambda}_* = \boldsymbol{\beta}_*^-$  and  $\boldsymbol{\eta}_* + \boldsymbol{\lambda}_* = \boldsymbol{\beta}_*^+$ . Using this re-parameterisation, [Cho et al. \(2019\)](#) assume  $k$  to be unity (i.e.,  $k = 1$ ) and show that one can first estimate  $(\varsigma_*, \boldsymbol{\lambda}_*', \boldsymbol{\eta}_*')$  by OLS to obtain  $(\widehat{\varsigma}_T, \widehat{\boldsymbol{\lambda}}_T', \widehat{\boldsymbol{\eta}}_T')$  and one can then estimate  $(\boldsymbol{\beta}_*^{+'}, \boldsymbol{\beta}_*^{-'})'$  as  $\widehat{\boldsymbol{\beta}}_T^- := \widehat{\boldsymbol{\lambda}}_T$  and  $\widehat{\boldsymbol{\beta}}_T^+ := \widehat{\boldsymbol{\eta}}_T + \widehat{\boldsymbol{\lambda}}_T$ .

With the long-run parameters estimated in this way, the singularity problem is resolved. If we let  $\mathbf{q}_t := (1, \mathbf{x}_t^{+'}, \mathbf{x}_t^{-'})'$  and  $\tilde{\mathbf{D}}_T := \text{diag}(T^{1/2}, T^{3/2} \mathbf{I}_k, T \mathbf{I}_k)$ , then  $\tilde{\mathbf{D}}_T^{-1} \left( \sum_{t=1}^T \mathbf{q}_t \mathbf{q}_t' \right) \tilde{\mathbf{D}}_T^{-1}$  weakly converges to a random matrix associated with a Brownian motion. Consequently, the limit behavior of  $(\widehat{\varsigma}_T, \widehat{\boldsymbol{\lambda}}_T', \widehat{\boldsymbol{\eta}}_T')$  can be studied as usual and it is possible to show that the long-run parameter estimator has convergence rate  $T$  and has an asymptotic distribution characterised by a Brownian motion. Due to the super-consistency of the long-run parameter estimator, it is possible to estimate the short-run parameters of (7) by OLS by

defining  $\hat{u}_t := y_t - \mathbf{x}_t^+ \hat{\boldsymbol{\beta}}_T^{+'} - \mathbf{x}_t^- \hat{\boldsymbol{\beta}}_T^{-'}$  and by replacing  $u_{t-1}$  in (7) with  $\hat{u}_{t-1}$ . Because all of the variables entering (7) are stationary and  $\{\varepsilon_t\}$  is a martingale difference array, the short-run parameter estimator is asymptotically normal and converges to the unknown short-run parameters at the rate  $\sqrt{T}$ .

The procedure detailed above provides an operational framework for the estimation of the NARDL model but it suffers from the drawback that the asymptotic distribution of the long-run parameter estimator is non-normal and depends on nuisance parameters, which complicates inference on the long-run parameters. To address this issue, [Cho et al. \(2019\)](#) note that one may simply estimate  $(\boldsymbol{\beta}_*^{+'}, \boldsymbol{\beta}_*^{-'})'$  using the FM-OLS estimator of [Phillips and Hansen \(1990\)](#) rather than by OLS. If  $\tilde{\boldsymbol{\Sigma}}_T$  and  $\tilde{\boldsymbol{\Pi}}_T$  are consistent estimators for the following asymptotic covariance matrices:

$$\boldsymbol{\Sigma} := \begin{bmatrix} \Sigma^{(1,1)} & \Sigma^{(1,2)} \\ \Sigma^{(2,1)} & \sigma^{(2,2)} \end{bmatrix} := \text{acov} \left[ \frac{1}{\sqrt{T}} \sum_{t=1}^T \begin{bmatrix} \Delta \mathbf{x}_t \\ u_t \end{bmatrix}, \frac{1}{\sqrt{T}} \sum_{t=1}^T \begin{bmatrix} \Delta \mathbf{x}_t \\ u_t \end{bmatrix} \right] \quad \text{and}$$

$$\boldsymbol{\Pi}_T := \begin{bmatrix} \Pi^{(1,1)} & \Pi^{(1,2)} \\ \Pi^{(2,1)} & \pi^{(2,2)} \end{bmatrix} := \text{acov} \left[ \frac{1}{\sqrt{T}} \sum_{t=1}^T \begin{bmatrix} \Delta \mathbf{x}_t \\ u_t \end{bmatrix}, \begin{bmatrix} \Delta \mathbf{x}_0 \\ u_0 \end{bmatrix} \right],$$

then the FM-OLS estimator is defined as:

$$\tilde{\boldsymbol{\varrho}}_T := (\tilde{\zeta}_T, \tilde{\boldsymbol{\lambda}}_T', \tilde{\boldsymbol{\eta}}_T')' := \left( \sum_{t=1}^T \mathbf{q}_t \mathbf{q}_t' \right)^{-1} \left( \sum_{t=1}^T \mathbf{q}_t \tilde{y}_t - T \mathbf{S}' \tilde{\boldsymbol{\Lambda}}_T \right),$$

where  $\tilde{y}_t := y_t - \Delta \mathbf{x}_t' \left( \tilde{\boldsymbol{\Sigma}}_T^{(1,1)} \right)^{-1} \tilde{\boldsymbol{\Sigma}}_T^{(1,2)}$ ,  $\tilde{\boldsymbol{\Lambda}}_T := \tilde{\boldsymbol{\Pi}}_T^{(1,2)} - \tilde{\boldsymbol{\Pi}}_T^{(1,1)} \left( \tilde{\boldsymbol{\Sigma}}_T^{(1,1)} \right)^{-1} \tilde{\boldsymbol{\Sigma}}_T^{(1,2)}$ , and  $\mathbf{S} := [\mathbf{0}_{k \times (1+k)}, \mathbf{I}_k]$ . The long-run parameter estimator is then given by  $\tilde{\boldsymbol{\beta}}_T^- := \tilde{\boldsymbol{\lambda}}_T$  and  $\tilde{\boldsymbol{\beta}}_T^+ := \tilde{\boldsymbol{\eta}}_T + \tilde{\boldsymbol{\lambda}}_T$ . [Cho et al. \(2019\)](#) prove that the long-run parameter estimator obtained by FM-OLS has convergence rate  $T$  and asymptotically follows a mixed-normal distribution. Therefore, if one replaces  $u_{t-1}$  in (7) with  $\tilde{u}_{t-1} := y_t - \mathbf{x}_t^+ \tilde{\boldsymbol{\beta}}_T^{+'} - \mathbf{x}_t^- \tilde{\boldsymbol{\beta}}_T^{-'}$ , then the short-run parameters of (7) can be estimated by OLS, with the same asymptotic properties as the short-run parameter estimator obtained using  $(\hat{\boldsymbol{\beta}}_T^{+'}, \hat{\boldsymbol{\beta}}_T^{-'})'$ .

[Cho et al. \(2019\)](#) go on to demonstrate that the Wald test statistic asymptotically follows a  $\chi^2$  distribution under the null hypothesis, even on the long-run parameters. The authors examine two null hypotheses: first, that for some  $\mathbf{R}_\ell \in \mathbb{R}^{r \times k}$  and  $\mathbf{r} \in \mathbb{R}^r$ ,  $H_0' : \mathbf{R}_\ell (\boldsymbol{\beta}_*^+ - \boldsymbol{\beta}_*^-) = \mathbf{r}$  and, second, that for some  $\mathbf{R} \in \mathbb{R}^{r \times 2k}$ ,  $H_0'' : \mathbf{R} \boldsymbol{\beta}_* = \mathbf{r}$ , where  $\boldsymbol{\beta}_* := (\boldsymbol{\beta}_*^{+'}, \boldsymbol{\beta}_*^{-'})'$ . The first null,  $H_0'$ , is given to test the full or partial equality between  $\boldsymbol{\beta}_*^+$  and  $\boldsymbol{\beta}_*^-$ , while the second null,  $H_0''$ , is considered to test a general family of hypotheses on

the long-run parameters. [Cho et al. \(2019\)](#) derive the Wald test statistics to test both  $H'_0$  and  $H''_0$  and show that they follow chi-squared distributions under their respective hypotheses and that they exhibit consistent power against the relevant  $o(T^3)$  and  $o(T^2)$  alternatives.

In a follow-up paper, [Cho et al. \(2020b\)](#) replace the condition that  $E[\Delta \mathbf{x}_t] = \mathbf{0}$  with  $\mathbb{E}[\Delta \mathbf{x}_t] \neq \mathbf{0}$  and show that the same asymptotic singularity issues surveyed above arise in this case. [Cho et al. \(2020b\)](#) develop an extension of the two-step estimation framework developed in [Cho et al. \(2019\)](#) that can accommodate the non-zero population mean of  $\Delta \mathbf{x}_t$  and that is also effective when  $E[\Delta \mathbf{x}_t] = \mathbf{0}$  with  $k > 1$ . If we let  $\boldsymbol{\mu}_*^+ := E[\Delta \mathbf{x}_t^+]$  and  $\boldsymbol{\mu}_*^- := E[\Delta \mathbf{x}_t^-]$ , then we can write:

$$\mathbf{x}_t^+ = \boldsymbol{\gamma}_*^+ + \boldsymbol{\mu}_*^+ t + \sum_{j=1}^t \mathbf{s}_j^+ \quad \text{and} \quad \mathbf{x}_t^- = \boldsymbol{\gamma}_*^- + \boldsymbol{\mu}_*^-_t + \sum_{j=1}^t \mathbf{s}_j^-, \quad (8)$$

where  $\mathbf{s}_t^+ := \Delta \mathbf{x}_t^+ - \boldsymbol{\mu}_*^+_t$  and  $\mathbf{s}_t^- := \Delta \mathbf{x}_t^- - \boldsymbol{\mu}_*^-_t$ . The long-run relationship embedded in the NARDL model can be written as:

$$y_t = \boldsymbol{\beta}_*^{+'} (\boldsymbol{\mu}_*^+_t + \mathbf{w}_t^+) + \boldsymbol{\beta}_*^{-'} (\boldsymbol{\mu}_*^-_t + \mathbf{w}_t^-) + u_t = \boldsymbol{\beta}_*^{+'} \mathbf{w}_t^+ + \boldsymbol{\beta}_*^{-'} \mathbf{w}_t^- + \xi_* + \delta_* t + v_t, \quad (9)$$

where  $\mathbf{w}_t^+ := \sum_{j=1}^t \mathbf{s}_j^+$  and  $\mathbf{w}_t^- := \sum_{j=1}^t \mathbf{s}_j^-$ , implying that the long-run parameters can be consistently estimated by regressing  $y_t$  on  $(\mathbf{w}_t^{+'}, \mathbf{w}_t^{-'}, 1, t)'$ . However,  $\mathbf{w}_t := (\mathbf{w}_t^{+'}, \mathbf{w}_t^{-'})'$  is not observable. Therefore, [Cho et al. \(2020b\)](#) propose to first estimate (8) by regressing  $\mathbf{x}_t^+$  and  $\mathbf{x}_t^{-}$  on  $(1, t)$  to obtain regression residuals,  $\widehat{\mathbf{w}}_t^+$  and  $\widehat{\mathbf{w}}_t^-$ , that approximate  $\mathbf{w}_t^+$  and  $\mathbf{w}_t^-$ , respectively. Next, one can estimate the long-run parameters by regressing  $y_t$  on  $(\widehat{\mathbf{w}}_t^{+'}, \widehat{\mathbf{w}}_t^{-'}, 1, t)'$ . [Cho et al. \(2020b\)](#) show that the long-run parameter estimator obtained in this way converges to the unknown long-run parameters at the rate  $T$  and that the short-run parameters can be estimated by OLS in a subsequent step.

Notwithstanding the simple structure of this estimator, its asymptotic distribution is not straightforward, which complicates inference on the long-run parameters; specifically, the null limit distribution of the Wald statistic testing restrictions on the long-run parameters does not follow the standard chi-squared distribution. [Cho et al. \(2020b\)](#) show that this issue can be resolved by using FM-OLS to estimate the long-run parameters in a manner that parallels the approach of [Cho et al. \(2019\)](#). The benefit of this approach is that the long-run parameter estimator is asymptotically mixed-normal and so the associated Wald test statistic asymptotically follows a mixed chi-squared distribution under the null hypothesis.

### 3.2 Estimation and Inference for the TARDL Model

The NARDL model is relatively simple to estimate due to the fact that the threshold parameter in (5),  $\tau_*$ , is known a priori. By contrast, if the threshold is unknown, then it must be estimated. Cho et al. (2020c) and Cho et al. (2020d) both tackle this problem. Cho et al. (2020c) examine the existence of the threshold using an inferential procedure, whereas Cho et al. (2020d) examine the same problem from a model selection perspective.

Cho et al. (2020c) assume a single regressor (i.e.,  $k = 1$ ) and note that the TARDL model (5) reduces to the standard ARDL model if  $\theta_{j*}^+ = \theta_{j*}^-$  for all  $j = 0, 1, \dots, q$ . If so, the estimated TARDL model has a set of redundant parameters, which may compromise standard inferential procedures such as the  $t$ -statistic. Consequently, the authors first test for the existence of a distinct threshold level using a quasi-likelihood ratio (QLR) test statistic. Letting the original ARDL and TARDL processes in (3) and (6) be the null and alternative processes, respectively, and estimating their error variances by the least squares method, the QLR test statistic is defined as follows:

$$QLR_n := T \left( 1 - \frac{\hat{\sigma}_T^2}{\hat{\sigma}_{T,0}^2} \right),$$

where  $\hat{\sigma}_T^2$  and  $\hat{\sigma}_{T,0}^2$  are error variance estimators under the alternative and null model assumptions. Note that  $\hat{\sigma}_T^2$  is obtained by the nonlinear least squares estimator, as  $\tau_*$  is not linearly associated with the variables.

Cho et al. (2020c) examine this QLR test statistic because it has asymptotic power to distinguish the null process from the alternative process and because it may be able to handle the multifold identification problem. To understand the multifold identification problem, note that the identification problem described by Davies (1977, 1987) can arise in three different ways in the TARDL model. First, if  $\theta_*^+ = \theta_*^-$  and  $\pi_*^+ = \pi_*^-$  in (6), where  $\pi_*^+ := (\pi_{0*}^+, \pi_{1*}^+, \dots, \pi_{q-1*}^+)'$  and  $\pi_*^- := (\pi_{0*}^-, \pi_{1*}^-, \dots, \pi_{q-1*}^-)'$ , then  $\tau_*$  is not identified. Second, if  $\tau_* = \Phi^{-1}(0)$ , where  $\Phi(\cdot)$  denotes the cumulative distribution function of  $\Delta x_t$ , then it follows that  $\Delta x_t^+ \equiv \Delta x_t$  and  $\Delta x_t^- \equiv 0$ , which means that the coefficients associated with the negative regime, such as  $\theta_*^-$  and  $\pi_*^-$ , are not identified. Finally, if  $\tau_* = \Phi^{-1}(1)$ , it follows that  $\Delta x_t^- \equiv \Delta x_t$  and  $\Delta x_t^+ \equiv 0$ , such that the coefficients associated with the positive regime are not identified. Consequently, the null hypothesis of a single regime is composed of the following three sub-null hypotheses:

$$H_{01} : \theta_*^+ = \theta_*^- \quad \text{and} \quad \pi_*^+ = \pi_*^-; \quad H_{02} : \tau_* = \Phi^{-1}(0); \quad \text{and} \quad H_{03} : \tau_* = \Phi^{-1}(1),$$

and the negation of the union of the sub-null hypotheses becomes the alternative hypothesis.

Due to this multifold identification problem, testing the joint null hypothesis by standard methods such as Wald's (1943) testing principle is a challenging task, because each sub-null hypothesis represents an alternative to another sub-null hypothesis, with the result that testing the union of the sub-null hypotheses cannot proceed on the basis of testing the hypothetical locations of the null parameter space. Consequently, in keeping with the established precedent in the literature, Cho et al. (2020c) tackle the multifold identification problem using the QLR test statistic.<sup>1</sup>

Cho et al. (2020c) show that the null limit distribution of the QLR test statistic can be represented as a functional of two Gaussian stochastic processes, in contrast to results in the prior literature. As the data entering the TARDL model are cointegrated  $I(1)$  processes, an additional Gaussian process is introduced to derive the null limit distribution. Specifically, Cho et al. (2020c) first show that the null approximation of the QLR test statistic under  $H_{01}$  dominates the other null approximations obtained under  $H_{02}$  and  $H_{03}$ , such that the null limit distribution of the QLR test statistic is given by the limit of the null approximation of the QLR test statistic under  $H_{01}$ . Next, they note that the functional of the two Gaussian stochastic process is a sum of two separable functionals of the two Gaussian processes, such that the functional of the first Gaussian process is obtained while testing  $\pi_*^+ = \pi_*^-$ , whereas the functional of the second Gaussian process is obtained when testing  $\theta_*^+ = \theta_*^-$ . Furthermore, as in the prior literature on multifold identification failure, the authors discover that the first Gaussian process is indexed by the parameter  $\tau$  due to the Davies (1977, 1987) identification problem. However, the second Gaussian process is formed by a partial sum process introduced by Phillips (1991, theorem 1) when testing a hypothesis on cointegration coefficients. Based on these findings, Cho et al. (2020c) perform a simulation study to show that the limit distribution of each functional can be separately approximated by combining a bootstrap method with a chi-squared distribution. Specifically, the first functional can be approximated by the weighted bootstrap procedure of Hansen (1996), while a  $\chi^2$  distribution approximates the second functional, as demonstrated by Phillips (1991).

---

<sup>1</sup>The multifold identification problem has been shown to arise in many popular models. For example, Cho and White (2007, 2010) examine a regime-switching process and note that a multifold identification problem arises when testing for one regime versus two regimes. The multifold identification problem is also observed when testing a linear model against an artificial neural network model or a smooth transition autoregressive model and in a consistent moment specification testing context (e.g. Baek, Cho, and Phillips, 2015; Cho and Ishida, 2012; Cho, Ishida, and White, 2011, 2014; Cho and Phillips, 2018; Seong, Cho, and Teräsvirta, 2019). All of these studies show that test statistics constructed using the likelihood ratio testing principle can overcome the multifold identification problem by providing the null limit distributions of the test statistics, which can be represented as functionals of Gaussian stochastic processes.

Cho et al. (2020c) note several additional aspects of testing with TARDL models. First, as the null limit distribution of the QLR test statistic is characterised mainly by a functional of a Gaussian process indexed by  $\tau$ , the authors restrict the dimension of  $\tau$  be equal to unity (i.e.,  $k = 1$ ). This is a pragmatic decision motivated by the difficulty of obtaining the null limit distribution of  $\tau$  in higher-dimensional cases. Second, the authors show that the specification of the TARDL process also determines the form of the functional of the two Gaussian processes. For example, in the case where the intercept is excluded from the TARDL model (6) because it is known to be zero a priori, the null limit distribution is characterised by *non-separable* functionals of the two Gaussian processes. Consequently, it is recommended that one should estimate the intercept even if its value is known ex ante in order to facilitate the application of the QLR test statistic.

Cho et al. (2020d) consider a general setting of the TARDL process subject to  $S - 1$  threshold levels. The authors allow for the case in which  $k$  may differ from unity, such that there are  $\tau_{1*} < \tau_{2*} < \dots < \tau_{S-1*}$  threshold levels, which means that the long-run relationship between  $y_t$  and  $\mathbf{x}_t$  is captured by:

$$y_t = \gamma_* + \beta_*^{(1)'} \mathbf{x}_t^{(1)} + \beta_*^{(2)'} \mathbf{x}_t^{(2)} + \dots + \beta_*^{(S)'} \mathbf{x}_t^{(S)} + u_t,$$

where, for each  $j = 1, 2, \dots, S$ ,  $\mathbf{x}_t^{(j)} = \sum_{t=0}^t \Delta \mathbf{x}_j^{(j)}$  with  $\Delta \mathbf{x}_t^{(1)} := \Delta \mathbf{x}_t \circ \mathbb{1}\{\Delta \mathbf{x}_t <_{\dagger} \tau_{1*}\}$ ,  $\Delta \mathbf{x}_t^{(S)} := \Delta \mathbf{x}_t \circ \mathbb{1}\{\Delta \mathbf{x}_t \geq_{\dagger} \tau_{S-1*}\}$ , and for  $s = 2, 3, \dots, S - 2$ ,  $\Delta \mathbf{x}_t^{(s)} := \Delta \mathbf{x}_t \circ \mathbb{1}\{\tau_{s-1*} \leq_{\dagger} \Delta \mathbf{x}_t <_{\dagger} \tau_{s*}\}$ . Here,  $\circ$  denotes the Hadamard product, while the subscript ‘ $\dagger$ ’ following an inequality sign denotes the element-by-element inequality between two vectors. Cho et al. (2020d) provide the error-correction form of this process as follows:

$$\Delta y_t = \rho_* y_{t-1} + \sum_{s=1}^S \theta_*^{(s)'} \mathbf{x}_{t-1}^{(s)} + \gamma_* + \sum_{j=1}^{p-1} \varphi_{j*} \Delta y_{t-j} + \sum_{j=0}^{q-1} \sum_{s=1}^S \pi_{j*}^{(s)'} \Delta \mathbf{x}_{t-j}^{(s)} + \varepsilon_t^{(S)}, \quad (10)$$

and refer to this as the  $\text{TARDL}(\tau_*; p, q)$  process, where  $\tau_* := (\tau'_{1*}, \dots, \tau'_{S-1*})'$ . Note that this is a generalised version of the  $\text{TARDL}(p, q)$  process in (6). If  $S = 2$ , the  $\text{TARDL}(\tau_*; p, q)$  process reduces to the  $\text{TARDL}(p, q)$  process. Furthermore, if  $S = 1$ , it reduces to  $\text{ARDL}(p, q)$  process.

The objective of Cho et al. (2020d) is to estimate the number of regimes,  $S$ . To do so, the authors examine the properties of several information criteria by simulation, including the Akaike information criterion (AIC), the Schwarz information criterion (SIC) and the Hannan-Quinn information criterion (HQIC), as well as modified versions of each criterion defined following Pitarakis (2006). Specifically, for each

$S = 1, 2, \dots, \bar{S}$ , [Cho et al. \(2020d\)](#) first estimate the error variance of (10) by least squares and let:

$$\hat{\sigma}_{n(S)}^2 := \min_{\lambda} \frac{1}{T} \sum_{t=1}^T \left( \Delta y_t - \rho y_{t-1} - \sum_{s=1}^S \boldsymbol{\theta}^{(s)'} \mathbf{x}_{t-1}^{(s)}(\boldsymbol{\tau}) - \gamma - \sum_{j=1}^{p-1} \varphi_j \Delta y_{t-j} - \sum_{j=0}^{q-1} \sum_{s=1}^S \boldsymbol{\pi}_j^{(s)'} \Delta \mathbf{x}_{t-j}^{(s)}(\boldsymbol{\tau}) \right)^2,$$

where  $\boldsymbol{\lambda} := (\rho, \boldsymbol{\theta}', \gamma, \boldsymbol{\varphi}', \boldsymbol{\pi}', \boldsymbol{\tau}')$ ,  $\boldsymbol{\theta} := (\boldsymbol{\theta}^{(1)'}, \dots, \boldsymbol{\theta}^{(S)'})'$ ,  $\boldsymbol{\varphi} := (\varphi_1, \dots, \varphi_{p-1})'$ , and  $\boldsymbol{\pi} := (\boldsymbol{\pi}_0^{(1)'}, \dots, \boldsymbol{\pi}_0^{(S)'}, \dots, \boldsymbol{\pi}_{q-1}^{(1)'}, \dots, \boldsymbol{\pi}_{q-1}^{(S)'})'$ . Next, the information criteria are computed as follows:

$$IC_{TARDL(\boldsymbol{\tau}, p, q)} := \log(\hat{\sigma}_{n(S)}^2) + \frac{c_T}{T} n_{TARDL} \quad \text{and} \quad IC_{TARDL(\boldsymbol{\tau}, p, q)}^{\circ} := \log(\hat{\sigma}_{n(S)}^2) + \frac{c_T}{T} n_{TARDL}^{\circ},$$

where  $c_T$  is a deterministic penalty term that satisfies  $c_T/T \rightarrow 0$ ,  $n_{TARDL}$  is the number of parameters in the TARDL( $\boldsymbol{\tau}, p, q$ ) process and  $n_{TARDL}^{\circ}$  is the number of parameters not including  $\boldsymbol{\tau}$ . [Akaike \(1973\)](#), [Schwarz \(1978\)](#), and [Hannan and Quinn \(1979\)](#) set  $c_T$  equal to 2,  $\log(T)$ , and  $2\log(\log(T))$ , respectively, while [Pitarakis \(2006\)](#) prefers the use of  $IC_{TARDL(\boldsymbol{\tau}, p, q)}^{\circ}$  to  $IC_{TARDL(\boldsymbol{\tau}, p, q)}$ .

[Cho et al. \(2020d\)](#) conduct extensive simulation studies to determine which of the six information criteria performs best. Setting  $k = 1$ , they separately examine the cases in which  $x_t$  has no time drift and where  $x_t$  is driven by a time drift. Note that if  $\Delta x_t$  has a non-zero population mean, then  $x_t$  is driven by a time drift. The simulations reveal that the relative performance of the information criteria is influenced by a combination of factors, including the sample size, the nature of the data generating process and whether or not  $x_t$  is driven by a time drift. In general, the standard SIC outperforms the other information criteria if  $\Delta x_t$  has zero population mean. However, if  $x_t$  is driven by a time drift, the picture is more complex. In this case, the modified SIC advocated by [Pitarakis \(2006\)](#) outperforms the other information criteria when  $T$  is small. However, as the sample size increases, the standard SIC and [Pitarakis's](#) SIC converge in terms of performance, with both dominating the other four information criteria. In addition, when the regime-specific parameters of the TARDL process are similar, such that the TARDL process is locally close to the simpler ARDL process, then [Pitarakis's](#) SIC has a tendency to outperform relative to all competitors, including the standard SIC.



## 4 ARDL Estimation by Quantile Regression

In recent years, the ARDL literature has expanded to encompass the use of quantile regression. Note that (1) can be converted into the *quantile autoregressive distributed-lag (QARDL) process*. For each  $\xi \in (0, 1)$ :

$$y_t = \gamma_*(\xi) + \sum_{j=1}^p \phi_{j*}(\xi) y_{t-j} + \sum_{j=0}^q \boldsymbol{\theta}_{j*}(\xi)' \mathbf{x}_{t-j} + \varepsilon_t(\xi), \quad (11)$$

where  $(\gamma_*(\xi), \phi_{1*}(\xi), \dots, \phi_{p*}(\xi), \boldsymbol{\theta}_{0*}(\xi)', \dots, \boldsymbol{\theta}_{q*}(\xi)')$  is the quantile coefficient of the QARDL process, and  $\varepsilon_t(\xi)$  is the quantile error. Estimation of the quantile coefficients enables the researcher to conduct inference on  $y_t$  and  $\mathbf{x}_t$  at any desired conditional quantile. [Cho et al. \(2015\)](#) provide relevant theory on the estimation and inference on the parameters in (11) along with an empirical application to post-war dividend smoothing in the US.

[Cho et al. \(2020a\)](#) propose a quantile version of the NARDL process in (6). That is, for each  $\xi \in (0, 1)$ , the *quantile nonlinear autoregressive distributed lag (QNARDL) process* is defined as:

$$y_t = \gamma_*(\xi) + \sum_{j=1}^p \phi_{j*}(\xi) y_{t-j} + \sum_{j=0}^q (\boldsymbol{\theta}_{j*}^+(\xi)' \mathbf{x}_{t-j}^+ + \boldsymbol{\theta}_{j*}^-(\xi)' \mathbf{x}_{t-j}^-) + \varepsilon_t(\xi), \quad (12)$$

where  $\mathbf{x}_t^+$  and  $\mathbf{x}_t^-$  are the same as in (5) and the quantile coefficients  $(\gamma_*(\xi), \phi_{1*}(\xi), \dots, \phi_{p*}(\xi), \boldsymbol{\theta}_{0*}^+(\xi)', \dots, \boldsymbol{\theta}_{q*}^+(\xi)', \boldsymbol{\theta}_{0*}^-(\xi)', \dots, \boldsymbol{\theta}_{q*}^-(\xi)')$  can be used to explore quantile variation in the asymmetric relationship embodied by the NARDL process. In this section, we detail the main findings of [Cho et al. \(2015\)](#) and [Cho et al. \(2020a\)](#).

### 4.1 The QARDL Model

[Cho et al. \(2015\)](#) note the QARDL process (10) can be equivalently converted into two different forms. First, following [Pesaran and Shin \(1998\)](#), the authors re-write the QARDL process as follows:

$$y_t = \gamma_*(\xi) + \mathbf{x}_t' \boldsymbol{\gamma}_*(\xi) + \sum_{j=1}^p \phi_{j*}(\xi) y_{t-j} + \sum_{j=0}^{q-1} \Delta \mathbf{x}_{t-j}' \boldsymbol{\delta}_{j*}(\xi) + \varepsilon_t(\xi), \quad (13)$$

where  $\boldsymbol{\gamma}_*(\xi) := \sum_{j=0}^q \boldsymbol{\theta}_{j*}(\xi)$  and  $\boldsymbol{\delta}_{j*}(\xi) := -\sum_{i=j+1}^q \boldsymbol{\theta}_{i*}(\xi)$ . Here, they additionally note that the long-run quantile relationship is captured by a linear relationship in (13), which can be given as  $y_t =$

$\mu_*(\xi) + \mathbf{x}'_t \boldsymbol{\beta}_*(\xi) + u_t(\xi)$ , where:

$$\mu_*(\xi) := \gamma_*(\xi) \left( 1 - \sum_{i=1}^p \phi_{i*}(\xi) \right)^{-1}, \quad \boldsymbol{\beta}_*(\xi) := \gamma_*(\xi) \left( 1 - \sum_{i=1}^p \phi_{i*}(\xi) \right)^{-1},$$

and  $u_t(\xi)$  is a stationary process defined by  $\{\Delta \mathbf{x}_t, \varepsilon_t(\xi), \Delta \mathbf{x}_{t-1}, \varepsilon_{t-1}(\xi), \dots\}$ . In addition, [Cho et al. \(2015\)](#) note that the following error-correction form can be derived from (10):

$$\Delta y_t = \gamma_*(\xi) + \rho_*(\xi)(y_{t-1} - \boldsymbol{\beta}_*(\xi)' \mathbf{x}_{t-1}) + \sum_{j=1}^{p-1} \varphi_{j*}(\xi) \Delta y_{t-j} + \sum_{j=0}^{q-1} \boldsymbol{\pi}_{j*}(\xi)' \Delta \mathbf{x}_{t-j} + \varepsilon_t(\tau), \quad (14)$$

where, for  $\ell = 1, 2, \dots, p-1$  and  $j = 1, 2, \dots, q-1$ :

$$\rho_*(\xi) := \sum_{j=1}^p \phi_{j*}(\xi) - 1, \quad \boldsymbol{\pi}_{0*}(\xi) := \boldsymbol{\theta}_{0*}(\xi), \quad \varphi_{\ell*}(\xi) := - \sum_{i=\ell+1}^p \phi_{i*}(\xi), \quad \text{and} \quad \boldsymbol{\pi}_{j*} := - \sum_{i=j+1}^p \boldsymbol{\theta}_{*i}(\xi).$$

[Cho et al. \(2015\)](#) exploit these two different representations to estimate the long-run parameter,  $\boldsymbol{\beta}_*(\xi)$ , and the other short-run parameters in (14) separately. Specifically, they first estimate the parameters in (13) by quantile regression following [Koenker and Bassett \(1978\)](#). Next, they convert the estimated parameters according to the formula for  $\boldsymbol{\beta}_*(\xi)$  and show that the convergence speed for the long-run parameter estimator is  $T$  such that, if  $\widehat{\boldsymbol{\beta}}_T(\xi)$  is the long-run parameter estimator, then the other short-run parameters in (14) can be consistently estimated by replacing  $\boldsymbol{\beta}_*(\xi)$  with  $\widehat{\boldsymbol{\beta}}_T(\xi)$  and applying quantile regression. Note that this approach parallels the conditional mean estimation framework developed by [Pesaran and Shin \(1998\)](#) for the linear ARDL model.

[Cho et al. \(2015\)](#) extend the scope of the QARDL model by letting  $\xi$  be a set of multiple quantile levels and examine the large sample properties of the long-run parameter estimators for  $\xi$ . [Cho et al. \(2015\)](#) show that the long-run parameter estimator converges to the unknown long-run parameter at the speed of  $T$ , even when the long-run parameters are estimated for multiple quantile levels. Furthermore, the authors demonstrate that the joint limit distribution of the long-run parameter estimator is mixed-normal, such that a test statistic constructed using the [Wald \(1943\)](#) testing principle asymptotically follows a  $\chi^2$  distribution under the null.

## 4.2 The QNARDL Model

Cho et al. (2020a) note that, for each  $\xi \in (0, 1)$ , the QNARDL process can be written in error correction form as:

$$\begin{aligned} \Delta y_t = & \rho_*(\xi)y_{t-1} + \boldsymbol{\theta}_*^+(\xi)' \mathbf{x}_{t-1}^+ + \boldsymbol{\theta}_*^-(\xi)' \mathbf{x}_{t-1}^- + \gamma_*(\xi) \\ & + \sum_{j=1}^{p-1} \varphi_{j*}(\xi) \Delta y_{t-j} + \sum_{j=0}^{q-1} \left( \boldsymbol{\pi}_{j*}^+(\xi)' \Delta \mathbf{x}_{t-j}^+ + \boldsymbol{\pi}_{j*}^-(\xi)' \Delta \mathbf{x}_{t-j}^- \right) + \varepsilon_t(\xi), \end{aligned} \quad (15)$$

where the QNARDL quantile coefficients are defined similarly to the NARDL coefficients and so, for brevity, we do not provide them here. Letting  $u_{t-1}(\xi) := y_{t-1} - \boldsymbol{\beta}_*^+(\xi)' \mathbf{x}_{t-1}^+ - \boldsymbol{\beta}_*^-(\xi)' \mathbf{x}_{t-1}^-$  be the quantile cointegration error, we can re-write (15) as follows:

$$\Delta y_t = \rho_*(\xi)u_{t-1}(\xi) + \gamma_*(\xi) + \sum_{j=1}^{p-1} \varphi_{j*}(\xi) \Delta y_{t-j} + \sum_{j=0}^{q-1} \left( \boldsymbol{\pi}_{j*}^+(\xi)' \Delta \mathbf{x}_{t-j}^+ + \boldsymbol{\pi}_{j*}^-(\xi)' \Delta \mathbf{x}_{t-j}^- \right) + \varepsilon_t(\xi), \quad (16)$$

where  $\boldsymbol{\beta}_*^+(\xi) := -\boldsymbol{\theta}_*^+(\xi)/\rho_*(\xi)$  and  $\boldsymbol{\beta}_*^-(\xi) := -\boldsymbol{\theta}_*^-(\xi)/\rho_*(\xi)$ . This error correction form of the QNARDL process enables the researcher to estimate the unknown coefficients in a manner parallel to the NARDL model estimation outlined above. Nevertheless, as in the NARDL case, if the unknown long-run parameters are estimated by quantile regression directly, then the same singularity problem arises. Thus, Cho et al. (2020a) recommend that the QNARDL process should be re-parameterised prior to estimation. Specifically, for each  $\xi \in (0, 1)$ , let the long-run relationship be represented as follows:

$$y_t = \varsigma_*(\xi) + \boldsymbol{\lambda}_*(\xi)' \mathbf{x}_t^+ + \boldsymbol{\eta}_*(\xi)' \mathbf{x}_t + u_t(\xi),$$

where  $\mathbf{x}_t \equiv \mathbf{x}_t^+ + \mathbf{x}_t^-$ . The long-run parameters  $(\boldsymbol{\lambda}_*(\xi)', \boldsymbol{\eta}_*(\xi)')$  can be estimated by quantile regression for  $k = 1$ . With this estimator denoted by  $(\widehat{\boldsymbol{\lambda}}_T(\xi)', \widehat{\boldsymbol{\eta}}_T(\xi)')$ , the long-run parameters are obtained by  $\widehat{\boldsymbol{\beta}}_T^+(\xi) := \widehat{\boldsymbol{\lambda}}_T(\xi) + \widehat{\boldsymbol{\eta}}_T(\xi)$  and  $\widehat{\boldsymbol{\eta}}_T^-(\xi) := \widehat{\boldsymbol{\eta}}_T(\xi)$ . Under this re-parameterisation, the singularity problem does not arise and the long-run parameter estimator converges to the true parameter value at rate  $T$ . Consequently, the short-run parameters in (16) can be estimated by replacing  $u_{t-1}$  with  $\widehat{u}_{t-1}(\xi) := y_{t-1} - \widehat{\boldsymbol{\beta}}_T^+(\xi)' \mathbf{x}_{t-1}^+ - \widehat{\boldsymbol{\beta}}_T^-(\xi)' \mathbf{x}_{t-1}^-$  and then applying quantile regression. If  $k > 1$ , one can estimate the trend models first and next estimate the long and short-run equations sequentially, by analogy to Cho et al. (2020b). Due to the parallel structure of these procedures, we do not repeat the discussion here in the interest of brevity.

Despite the simplicity of the estimation procedure, [Cho et al. \(2020a\)](#) note that the asymptotic distribution of the long-run parameter estimator is not straightforward, which complicates hypothesis testing with respect to the long-run parameters. To overcome this issue, in keeping with the estimation procedure advanced by [Cho et al. \(2019\)](#), [Cho et al. \(2020a\)](#) first estimate  $(\lambda_*(\xi)', \eta_*(\xi)')$  by applying the FM-OLS estimator of [Phillips and Hansen \(1990\)](#) to the quantile regression problem and next estimate the long-run parameters as described above. The long-run parameter estimator estimated in this way is consistent, converges to the true parameter value at the rate  $T$  and asymptotically follows a mixed normal distribution. Therefore, test statistics constructed using the [Wald \(1943\)](#) principle converge to the standard  $\chi^2$  distribution. Consequently, inference on the QNARDL model proceeds in a similar manner to the NARDL model described above.

## 5 Panel Data Extensions of the ARDL Model

ARDL specifications and their derivatives have also been applied in the context of dynamic panel data analysis. Two simple estimators for panels with many time series observations for each of  $N$  groups are the traditional pooled estimators (including the fixed and random effects estimators) and the Mean Group (MG) estimator of [Pesaran and Smith \(1995\)](#). The first approach involves pooling the data together and estimating a single model under the assumption that all of the model parameters aside from the intercepts are homogeneous across groups. At the other extreme, the MG estimator of [Pesaran and Smith \(1995\)](#) involves estimating  $N$  separate models, where all of the parameters are allowed to vary over groups. To summarise the distribution of these group-specific parameters, one may simply take their mean value across groups. MG estimation of ARDL and NARDL models has been applied in the analysis of exchange rate pass-through into import prices by [Brun-Aguerre, Fuertes, and Greenwood-Nimmo \(2017\)](#).

[Pesaran et al. \(1999\)](#) pursue an intermediate approach that they refer to as the Pooled Mean Group (PMG) estimator, which mimics the structure of the ARDL model in a panel setting under the assumption of long-run homogeneity, while allowing the short-run parameters and error variances to differ across groups. The authors note that there are often good reasons to believe that equilibrium relations should be common across groups but that the same is not typically true of short-run dynamic parameters. This reasoning justifies their pursuit of a hybrid approach to estimation that makes use of pooling for the estimation of long-run parameters and averaging for the estimation of short-run parameters.

Suppose that one wishes to estimate an  $ARDL(p, q, q, \dots, q)$  model of the following form using a

panel dataset with groups indexed by  $i = 1, 2, \dots, N$  and time periods indexed by  $t = 1, \dots, T$ , where  $T$  is sufficiently large to allow the model to be estimated for each group:<sup>2</sup>

$$y_{it} = \sum_{j=1}^p \lambda_{ij*} y_{i,t-j} + \sum_{j=0}^q \delta'_{ij*} \mathbf{x}_{i,t-j} + \gamma'_{i*} \mathbf{d}_t + \varepsilon_{it}, \quad (17)$$

where  $\mathbf{x}_{it}$  and  $\mathbf{d}_t$  are  $k \times 1$  and  $s \times 1$  vectors of regressors, respectively, while the  $\lambda_{ij*}$ s are unknown scalars and the  $\delta_{ij*}$ s and  $\gamma_{i*}$ s are  $k \times 1$  and  $s \times 1$  vectors of unknown parameters to be estimated. Note that (17) can be re-parameterised as follows:

$$\Delta y_{it} = \phi_{i*} y_{i,t-1} + \beta'_{i*} \mathbf{x}_{it} + \sum_{j=1}^{p-1} \lambda_{ij}^* \Delta y_{i,t-j} + \sum_{j=0}^{q-1} \delta_{ij}^{*'} \Delta \mathbf{x}_{i,t-j} + \gamma'_{i*} \mathbf{d}_t + \varepsilon_{it}, \quad (18)$$

where  $\phi_{i*} := -(1 - \sum_{j=1}^p \lambda_{ij*})$ ,  $\beta_{i*} := \sum_{j=0}^q \delta_{ij*}$ ,  $\lambda_{ij}^* := -\sum_{m=j+1}^p \lambda_{im*}$ ,  $j = 1, \dots, p-1$  and  $\delta_{ij}^{*'} := -\sum_{m=j+1}^q \delta_{im*}$ ,  $j = 1, \dots, q-1$ ,  $i = 1, \dots, N$ . By stacking the time series observations for each group to obtain  $\mathbf{y}_i := (y_{i1}, \dots, y_{iT})'$ ,  $\mathbf{X}_i := (\mathbf{x}_{i1}, \dots, \mathbf{x}_{iT})'$ , (18) can be re-written in the following form:

$$\Delta \mathbf{y}_i = \phi_{i*} \mathbf{y}_{i,-1} + \mathbf{X}_i \beta_{i*} + \sum_{j=1}^{p-1} \lambda_{ij}^* \Delta \mathbf{y}_{i,-j} + \sum_{j=0}^{q-1} \Delta \mathbf{X}_{i,-j} \delta_{ij}^{*'} + \mathbf{D} \gamma_{i*} + \varepsilon_i, \quad (19)$$

where  $\mathbf{D} := (\mathbf{d}_1, \dots, \mathbf{d}_T)'$  is a  $T \times s$  matrix of observations on the deterministic regressors, such as intercepts and time trends, and  $\mathbf{y}_{i,-1}$  and  $\Delta \mathbf{X}_{i,-j}$  are  $T \times 1$  and  $T \times k$  matrices obtained by stacking  $y_{i,t-1}$  and  $\mathbf{x}_{i,t-j}$ , respectively.

The long-run coefficients on  $\mathbf{X}_i$  can be obtained as  $\theta_{i*} = -\beta_{i*}/\phi_{i*}$ . Pesaran et al. (1999) assume long-run homogeneity, such that  $\theta_{i*} = \theta_*$  for every  $i = 1, \dots, N$ . This allows (19) to be re-written compactly as:

$$\Delta \mathbf{y}_i = \phi_{i*} \boldsymbol{\xi}_i(\theta_*) + \mathbf{W}_i \boldsymbol{\kappa}_{i*} + \varepsilon_i, \quad i = 1, \dots, N, \quad (20)$$

where:

$$\boldsymbol{\xi}_i(\theta_*) := \mathbf{y}_{i,-1} - \mathbf{X}_i \theta_*, \quad i = 1, \dots, N,$$

is the error correction component,  $\mathbf{W}_i = (\Delta \mathbf{y}_{i,-1}, \dots, \Delta \mathbf{y}_{i,-p+1}, \Delta \mathbf{X}_i, \Delta \mathbf{X}_{i,-1}, \dots, \Delta \mathbf{X}_{i,-q+1}, \mathbf{D})$ , and  $\boldsymbol{\kappa}_{i*} := (\lambda_{i1}^*, \dots, \lambda_{i,p-1}^*, \delta_{i0}^{*'}, \delta_{i1}^{*'}, \dots, \delta_{i,q-1}^{*'}, \gamma_{i*}')'$ .

<sup>2</sup>It is straightforward to allow for different lag orders associated with each of the variables in  $\mathbf{x}_{it}$ .

Estimation of (20) is complicated by three factors: (i) the equation for each group is nonlinear in  $\phi_{i*}$  and  $\theta_*$ ; (ii) the long-run homogeneity assumption introduces cross-equation parameter restrictions; and (iii) the error variances exhibit heterogeneity across groups. Pesaran et al. (1999) propose a maximum likelihood estimation framework in which the homogeneous long-run parameters,  $\theta_*$  are estimated by pooling, while group-wide mean estimates of the heterogeneous short-run parameters,  $\kappa_{i*}$  and error-correction coefficients,  $\phi_{i*}$  are obtained by taking averages across groups. Hence the nomenclature ‘‘pooled mean group’’ estimation.

Pesaran et al. (1999) develop separate asymptotic theories for the PMG estimators in the case that the regressors,  $\mathbf{x}_{it}$ , are stationary and non-stationary. Under stationarity, the authors demonstrate the consistency and asymptotic normality of the PMG estimators and show that the long- and short-run parameter estimators share a common convergence rate of  $\sqrt{T}$ . By contrast, under the assumption that the regressors are first-order integrated processes, the asymptotic analysis is complicated by the fact that the ML estimators of the long-run and the short-run parameters converge to their true values at different rates ( $T$  and  $\sqrt{T}$ , respectively). In this case, for a fixed  $N$  and as  $T \rightarrow \infty$ , the PMG estimator asymptotically follows a mixed-normal distribution.

## 6 The Spatio-Temporal ARDL Model

In response to the increasing availability of large spatial time-series datasets, Shin and Thornton (2019) develop the spatio-temporal autoregressive distributed lag (STARDL) model. The STARDL model extends the popular spatial dynamic panel data model by allowing both spatial and temporal coefficients to differ across spatial units. The STARDL( $p, q$ ) model is specified as follows:

$$y_{it} = \sum_{\ell=1}^p \phi_{i\ell*} y_{i,t-\ell} + \sum_{\ell=0}^p \phi_{i\ell}^* y_{i,t-\ell}^* + \sum_{\ell=0}^q \pi'_{i\ell*} \mathbf{x}_{i,t-\ell} + \sum_{\ell=0}^q \pi_{i\ell}^{*'} \mathbf{x}_{i,t-\ell}^* + \alpha_{i*} + u_{it}, \quad (21)$$

for  $i = 1, \dots, N$  and  $t = 1, \dots, T$ , where  $y_{it}$  is a scalar dependent variable for the  $i$ th spatial unit at time  $t$ ,  $\mathbf{x}_{it} = (x_{it}^1, \dots, x_{it}^K)'$  is a  $K \times 1$  vector of exogenous regressors,  $\boldsymbol{\pi}_{i\ell*} = (\pi_{i\ell*}^1, \dots, \pi_{i\ell*}^K)'$  is a  $K \times 1$  vector of parameters and likewise for  $y_{i,t-\ell}$  and  $\mathbf{x}_{i,t-\ell}$  and their associated parameters. Spatial interactions between units, both contemporaneously and with lags, are captured via the spatial variables,  $y_{it}^*$  and  $\mathbf{x}_{it}^*$ , defined as:

$$y_{it}^* \equiv \sum_{j=1}^N w_{ij} y_{jt} = \mathbf{w}'_i \mathbf{y}_t \quad \text{with} \quad \mathbf{y}_t = (y_{1t}, \dots, y_{Nt})', \quad (22)$$

$$\mathbf{x}_{K \times 1}^* = (x_{it}^{1*}, \dots, x_{it}^{K*})' \equiv \left( \sum_{j=1}^N w_{ij} x_{jt}^1, \dots, \sum_{j=1}^N w_{ij} x_{jt}^K \right)' = (\mathbf{w}'_i \otimes \mathbf{I}_K) \mathbf{x}_t; \quad \mathbf{x}_t = \begin{pmatrix} \mathbf{x}_{1t} \\ \vdots \\ \mathbf{x}_{Nt} \end{pmatrix} \quad (23)$$

where  $\mathbf{w}'_i = (w_{i1}, \dots, w_{iN})$  denotes a  $1 \times N$  vector of non-stochastic spatial weights that are assumed to be known a priori, with  $w_{ii} = 0$ .

Stacking the  $N$  individual STARDL  $(p, q)$  equations in (21), Shin and Thornton (2019) obtain the following spatial system:

$$\mathbf{y}_t = \sum_{\ell=1}^p \Phi_{\ell*} \mathbf{y}_{t-\ell} + \sum_{\ell=0}^p \Phi_{\ell}^* \mathbf{W} \mathbf{y}_{t-\ell} + \sum_{\ell=0}^q \Pi_{\ell*} \mathbf{x}_{t-\ell} + \sum_{\ell=0}^q \Pi_{\ell}^* (\mathbf{W} \otimes \mathbf{I}_K) \mathbf{x}_{t-\ell} + \boldsymbol{\alpha}_* + \mathbf{u}_t, \quad (24)$$

where  $\mathbf{W}$  is the following  $N \times N$  spatial weights matrix:

$$\mathbf{W} = \begin{bmatrix} w_{11} & \cdots & w_{1N} \\ \vdots & \ddots & \vdots \\ w_{N1} & \cdots & w_{NN} \end{bmatrix} = \begin{bmatrix} \mathbf{w}'_1 \\ \vdots \\ \mathbf{w}'_N \end{bmatrix}, \quad (25)$$

$\boldsymbol{\alpha}_* = (\alpha_{1*}, \dots, \alpha_{N*})'$  and  $\Phi_{\ell*}$ ,  $\Phi_{\ell}^*$ ,  $\Pi_{\ell*}$ ,  $\Pi_{\ell}^*$  are the following diagonal matrices:

$$\Phi_{\ell*} = \begin{bmatrix} \phi_{1\ell*} & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \phi_{N\ell*} \end{bmatrix}, \ell = 1, \dots, p; \quad \Phi_{\ell}^* = \begin{bmatrix} \phi_{1\ell}^* & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \phi_{N\ell}^* \end{bmatrix}, \ell = 0, 1, \dots, p$$

$$\Pi_{\ell*} = \begin{bmatrix} \pi'_{1\ell*} & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \pi'_{N\ell*} \end{bmatrix}, \quad \Pi_{\ell}^* = \begin{bmatrix} \pi^*_{1\ell} & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \pi^*_{N\ell} \end{bmatrix} \quad \text{for } \ell = 0, 1, \dots, q.$$

Shin and Thornton (2019) propose the use of both quasi-maximum likelihood and control function-based IV estimators<sup>3</sup> to obtain consistent and asymptotically normal estimates of the parameters of the STARDL model (24). Note that the STARDL model is both relatively simple to estimate and also highly adaptable, as it nests several popular spatial dynamic panel data models, including the dynamic spatial Durbin model

<sup>3</sup>Here we obtain the instrumental variables internally from the spatially lagged exogenous and predetermined variables, say  $(\mathbf{W}^2 \mathbf{y}_{t-1}, \mathbf{W}^2 \mathbf{x}_t, \dots)$

analysed by [Lee and Yu \(2010\)](#) and [Elhorst \(2014\)](#) and the heterogeneous spatial autoregressive panel data model of [Aquaro et al. \(2021\)](#).

Spatial models, including the STARDL model, have an implicit network structure and so many of the popular tools of network analysis, including centrality statistics and clustering algorithms, can be used to facilitate their interpretation. Their use is particularly advantageous in the context of the STARDL model, where spatial dynamic interactions are governed by an array of parameters, unlike simple homogeneous parameter models in which interest primarily centres on a single spatial parameter. [Shin and Thornton \(2019\)](#) make use of two quantities for this purpose: (i) individual spatio-temporal dynamic multipliers based on work by [Shin et al. \(2014\)](#); and (ii) system diffusion multipliers. The diffusion multipliers represent a novel technique to measure the joint impacts of  $\mathbf{x}_t$  on  $\mathbf{y}_{t+h}$  in space and time for  $h = 0, 1, 2, \dots$ . To obtain the diffusion multipliers, it is first necessary to re-write (24) as follows:

$$\tilde{\Phi}(L) \mathbf{y}_t = \tilde{\Pi}(L) \mathbf{x}_t + \tilde{\mathbf{u}}_t, \quad (26)$$

where  $\tilde{\Phi}(L) = \mathbf{I}_N - \sum_{\ell=1}^p \tilde{\Phi}_{\ell*} L^\ell$ ,  $\tilde{\Pi}(L) = \sum_{\ell=0}^q \tilde{\Pi}_{\ell*} L^\ell$ ,  $\tilde{\mathbf{u}}_t = (\mathbf{I}_N - \Phi_0^* \mathbf{W}(L))^{-1} \mathbf{u}_t$  and  $\tilde{\Phi}_{\ell*}$  and  $\tilde{\Pi}_{\ell*}$  are the coefficients of the endogenous and exogenous variables obtained from (24), respectively. Premultiplying (26) by  $[\tilde{\Phi}(L)]^{-1}$ , we have:

$$\mathbf{y}_t = \mathbf{B}(L) \mathbf{x}_t + [\tilde{\Phi}(L)]^{-1} \tilde{\mathbf{u}}_t,$$

where  $\mathbf{B}(L) := [\tilde{\Phi}(L)]^{-1} \tilde{\Pi}(L) = \sum_{j=0}^{\infty} \mathbf{B}_{j*} L^j$  and where the diffusion multipliers,  $\mathbf{B}_{j*}$  for  $j = 0, 1, \dots$ , can be evaluated as follows:

$$\mathbf{B}_{j*} = \tilde{\Phi}_{1*} \mathbf{B}_{j-1*} + \tilde{\Phi}_{2*} \mathbf{B}_{j-2*} + \dots + \tilde{\Phi}_{j-1*} \mathbf{B}_{1*} + \tilde{\Phi}_{j*} \mathbf{B}_{0*} + \tilde{\Pi}_{j*}, \quad j = 1, 2, \dots,$$

where  $\mathbf{B}_{0*} := \tilde{\Pi}_{0*}$  and  $\mathbf{B}_{j*} = \mathbf{0}$  for  $j < 0$  by construction. The  $N \times NK$  matrix of cumulative diffusion multiplier effects is given by:

$$\mathbf{d}_{x*}^H = \sum_{h=0}^H \frac{\partial \mathbf{y}_{t+h}}{\partial \mathbf{x}'_t} = \sum_{h=0}^H \mathbf{B}_{h*}, \quad H = 0, 1, 2, \dots$$

[Shin and Thornton \(2019\)](#) leverage recent developments in the econometric analysis of networks and



connectedness due to [Diebold and Yilmaz \(2014\)](#) and [Greenwood-Nimmo, Nguyen, and Shin \(in press\)](#) to provide a simple and intuitive summary of the impacts of  $\{x_{jt}\}_{j=1}^N$  on  $\{y_{it}\}_{i=1}^N$ . In this way, [Shin and Thornton](#) show that the diffusion multipliers can be used to obtain two simple and easily interpreted measures of the role of each node within the network: (i) its *external motivation*, which captures the extent and direction to which each node is steered by the network; and (ii) its *systemic influence*, which captures the relative importance of each node within the network. The authors apply their framework to analyse the effect of enemy casualties on civilian deaths across the 18 Governorates of Iraq in the aftermath of the 2003 invasion. Their results indicate that violence is used both as a means to maintain influence in emergent political institutions and also in reprisal for past acts.

## 7 Potential Avenues for Further Development

The large body of research that we survey above highlights the adaptability of the ARDL specification but it should not be taken as evidence that all of the worthwhile avenues for development have already been explored. There are three areas in particular that have yet to be developed and that hold considerable promise. The first is to develop theoretical and applied methods that combine the ARDL and NARDL specifications with other popular regime-switching mechanisms, such as Markov-switching and smooth transition models. The second objective is the development of a system extension of the NARDL/TARDL model, which would provide a valuable framework for the analysis of asymmetric and nonlinear phenomena in multivariate systems. The last is the development of new panel ARDL models that can jointly accommodate both spatial and factor dependence. Developments in this area promise substantial contributions to the fast-growing literature on the unified modelling of cross-section dependence in panels (e.g. [Bai and Li, 2013](#); [Shi and Lee, 2017](#); [Kuersteiner and Prucha, 2020](#); [Chen, Shin, and Zheng, 2020](#)).

## 8 Concluding Remarks

In this paper, we survey the literature on the ARDL model. Given the existence of several excellent surveys focusing on the case of stationary distributed lags (e.g. [Griliches, 1967](#); [Nerlove, 1972](#); [Hendry et al., 1984](#); [Wickens and Breusch, 1988](#)), our starting point is the more challenging setting in which the ARDL specification is applied to cointegrated non-stationary time series ([Pesaran and Shin, 1998](#)) or time series with mixed orders of integration ([Pesaran et al., 2001](#)). We present several recent extensions of this

model, including the NARDL and TARDL models associated with [Shin et al. \(2014\)](#) and [Cho et al. \(2019, 2020b,c,d\)](#), the QARDL and QNARDL models developed by [Cho et al. \(2015\)](#) and [Cho et al. \(2020a\)](#), the pooled mean group panel data estimator of [Pesaran et al. \(1999\)](#) and the spatio-temporal ARDL model proposed by [Shin and Thornton \(2019\)](#).

## References

- AKAIKE, H. (1973): “Information Theory and an Extension of the Maximum Likelihood Principle,” in *Proceedings of the Second International Symposium on Information Theory*, ed. by B. N. Petrov and F. Csáki, Budapest: Akademiai Kiado, 267–281.
- ALMON, S. (1965): “The Distributed Lag Between Capital Appropriations and Expenditures,” *Econometrica*, 33, 178–196.
- AQUARO, M., N. BAILEY, AND M. H. PESARAN (2021): “Estimation and Inference for Spatial Models with Heterogeneous Coefficients: An Application to US House Prices,” *Journal of Applied Econometrics*, 36, 18–44.
- BAEK, Y. I., J. S. CHO, AND P. C. PHILLIPS (2015): “Testing Linearity Using Power Transforms of Regressors,” *Journal of Econometrics*, 187, 376–384.
- BAI, J. AND K. LI (2013): “Spatial Panel Data Models with Common Shocks,” MPRA Paper 52786, University Library of Munich, Germany.
- BOX, M., K. GRATZER, AND X. LIN (2018): “The Asymmetric Effect of Bankruptcy Fraud in Sweden: A Long-Term Perspective,” *Journal of Quantitative Criminology*, in press.
- BRUN-AGUERRE, R. X., A.-M. FUERTES, AND M. J. GREENWOOD-NIMMO (2017): “Heads I Win; Tails You Lose: Asymmetry In Exchange Rate Pass-through into Import Prices,” *Journal of the Royal Statistical Society Series A*, 180, 587–612.
- CHEN, J., Y. SHIN, AND C. ZHENG (2020): “Estimation and Inference in Heterogeneous Spatial Panel Data Models with a Multifactor Error Structure,” University of York Discussion Papers in Economics 20/03, York.

- CHO, J. S., M. J. GREENWOOD-NIMMO, T.-H. KIM, AND Y. SHIN (2020a): “Hawks, Doves and Asymmetry in US Monetary Policy: Evidence from a Dynamic Quantile Regression Model,” Mimeo: University of York.
- CHO, J. S., M. J. GREENWOOD-NIMMO, AND Y. SHIN (2019): “Two-Step Estimation of the Nonlinear Autoregressive Distributed Lag Model,” Working Paper 2019rwp-154, Yonsei University, Seoul.
- (2020b): “Estimating the Nonlinear Autoregressive Distributed Lag Model for Time Series Data with Drifts,” Mimeo: University of York.
- (2020c): “Testing for the Threshold Autoregressive Distributed Lag Model,” Mimeo: University of York.
- (2020d): “The Threshold Autoregressive Distributed Lag Model,” Mimeo: University of York.
- CHO, J. S. AND I. ISHIDA (2012): “Testing for the Effects of Omitted Power Transformations,” *Economics Letters*, 117, 287–290.
- CHO, J. S., I. ISHIDA, AND H. WHITE (2011): “Revisiting Tests for Neglected Nonlinearity Using Artificial Neural Networks,” *Neural Computation*, 23, 1133–1186.
- (2014): “Testing for Neglected Nonlinearity Using Twofold Unidentified Models under the Null and Hexic Expansions,” in *Essays on Nonlinear Time Series Econometrics: A Festschrift in Honor of Timo Teräsvirta*, ed. by N. Haldrup, M. Meitz, and P. Saikkonen, Oxford: Oxford University Press, chap. 1, 3–27.
- CHO, J. S., T.-H. KIM, AND Y. SHIN (2015): “Quantile Cointegration in the Autoregressive Distributed Lag Modeling Framework,” *Journal of Econometrics*, 188, 281–300.
- CHO, J. S. AND P. C. PHILLIPS (2018): “Sequentially Testing Polynomial Model Hypothesis Using the Power Transform of Regressors,” *Journal of Applied Econometrics*, 33, 141–159.
- CHO, J. S. AND H. WHITE (2007): “Testing for Regime Switching,” *Econometrica*, 75, 1671–1720.
- (2010): “Testing for Unobserved Heterogeneity in Exponential and Weibull Duration Models,” *Journal of Econometrics*, 157, 458–480.

- CLAUS, E. AND V. H. NGUYEN (2019): “Monetary Policy Shocks from the Consumer Perspective,” *Journal of Monetary Economics*, in press.
- DAVIES, R. B. (1977): “Hypothesis Testing when a Nuisance Parameter is Present only under the Alternative,” *Biometrika*, 64, 247–254.
- (1987): “Hypothesis Testing when a Nuisance Parameter is Present only under the Alternative,” *Biometrika*, 74, 33–43.
- DHRYMES, P. J. (1971): *Distributed Lags: Problems of Estimation and Formulation*, Edinburgh: Oliver and Boyd.
- DIEBOLD, F. X. AND K. YILMAZ (2014): “On the Network Topology of Variance Decompositions: Measuring the Connectedness of Financial Firms,” *Journal of Econometrics*, 182, 119–134.
- ELHORST, J. P. (2014): *Spatial Econometrics: From Cross-Sectional Data to Spatial Panels*, Berlin: Springer.
- ENGLE, R. F. AND C. W. GRANGER (1987): “Co-integration and Error Correction: Representation, Estimation and Testing,” *Econometrica*, 55, 251–276.
- FEDOSEEVA, S. (2013): “(A)symmetry, (Non)linearity and Hysteresis of Pricing-To-Market: Evidence from German Sugar Confectionery Exports,” *Journal of Agricultural and Food Industrial Organization*, 11, 69–85.
- GREENWOOD-NIMMO, M. J., V. H. NGUYEN, AND Y. SHIN (in press): “Measuring the Connectedness of the Global Economy,” *International Journal of Forecasting*.
- GRILICHES, Z. (1967): “Distributed Lags: A Survey,” *Econometrica*, 35, 16–49.
- HAMMOUDEH, S., A. LAHIANI, D. K. NGUYEN, AND R. M. SOUSA (2015): “An Empirical Analysis of Energy Cost Pass-Through to CO2 Emission Prices,” *Energy Economics*, 49, 149–156.
- HANNAN, E. J. AND B. G. QUINN (1979): “The Determination of the Order of an Autoregression,” *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 41, 190–195.
- HANSEN, B. E. (1996): “Inference when a Nuisance Parameter is not Identified under the Null Hypothesis,” *Econometrica*, 64, 413–430.

- HE, Z. AND F. ZHOU (2018): “Time-Varying and Asymmetric Effects of the Oil-Specific Demand Shock on Investor Sentiment,” *PLOS ONE*, 31.
- HENDRY, D. F. AND G. E. MIZON (1978): “Serial Correlation as a Convenient Simplification, Not a Nuisance: A Comment on a Study of the Demand for Money by the Bank of England,” *Economic Journal*, 88, 549–563.
- HENDRY, D. F., A. R. PAGAN, AND J. D. SARGAN (1984): “Dynamic Specifications,” in *Handbook of Econometrics*, ed. by Z. Griliches and M. D. Intriligator, Amsterdam: North Holland, vol. 2, chap. 18, 1023–1100.
- JOHANSEN, S. (1988): “Statistical Analysis of Cointegration Vectors,” *Journal of Economic Dynamics and Control*, 12, 231–255.
- KOENKER, R. AND G. BASSETT (1978): “Regression Quantiles,” *Econometrica*, 46, 33–50.
- KOYCK, L. M. (1954): *Distributed Lags and Investment Analysis*, Amsterdam: North-Holland.
- KUERSTEINER, G. M. AND I. R. PRUCHA (2020): “Dynamic Spatial Panel Models: Networks, Common Shocks, and Sequential Exogeneity,” *Econometrica*, 88, 2109–2146.
- LEE, L. AND J. YU (2010): “Some Recent Developments in Spatial Panel Data Models,” *Regional Science and Urban Economics*, 40.
- MADDALA, G. S. (1977): *Econometrics*, New York: McGraw-Hill.
- NERLOVE, M. L. (1972): “Lags in Economic Behavior,” *Econometrica*, 40, 221–251.
- PAL, D. AND S. MITRA (2015): “Asymmetric Impact of Crude Price on Oil Product Pricing in the United States: An Application of Multiple Threshold Nonlinear Autoregressive Distributed Lag Model,” *Economics Modelling*, 51, 436–443.
- PESARAN, M. H. AND Y. SHIN (1998): “An Autoregressive Distributed Lag Modelling Approach to Cointegration Analysis,” in *Econometrics and Economic Theory: The Ragnar Frisch Centennial Symposium*, ed. by S. Strom, Cambridge: Cambridge University Press, Econometric Society Monographs, 371–413.

- PESARAN, M. H., Y. SHIN, AND R. J. SMITH (2001): “Bounds Testing Approaches to the Analysis of Level Relationships,” *Journal of Applied Econometrics*, 16, 289–326.
- PESARAN, M. H., Y. SHIN, AND R. P. SMITH (1999): “Pooled Mean Group Estimation of Dynamic Heterogeneous Panels,” *Journal of the American Statistical Association*, 94, 621–634.
- PESARAN, M. H. AND R. P. SMITH (1995): “Estimating Long-run Relationships from Dynamic Heterogeneous Panels,” *Journal of Econometrics*, 68, 79–113.
- PHILLIPS, P. C. AND B. E. HANSEN (1990): “Statistical Inference in Instrumental Variable Regression with I(1) Processes,” *Review of Economic Studies*, 57, 99–125.
- PHILLIPS, P. C. B. (1991): “Optimal Inference in Cointegrated Systems,” *Econometrica*, 59, 283–306.
- PITARAKIS, J.-Y. (2006): “Model Selection Uncertainty and Detection of Threshold Effects,” *Studies in Nonlinear Dynamics and Econometrics*, 10, 1–30.
- SARGAN, J. D. (1964): “Wages and Prices in the United Kingdom: A Study in Econometric Methodology (with Discussion),” in *Econometric Analysis for National Economic Planning*, ed. by P. Hart, G. Mills, and J. Whitaker, London: Butterworth and Co., 25–63.
- SCHWARZ, G. (1978): “Estimating the Dimension of a Model,” *Annals of Statistics*, 6, 461–464.
- SEONG, D., J. S. CHO, AND T. TERÄSVIRTA (2019): “Comprehensive Testing of Linearity against the Smooth Transition Autoregressive Model,” CREATES Research Papers 2019-17, Aarhus University, Aarhus.
- SHI, W. AND L. LEE (2017): “Spatial Dynamic Panel Data Models with Interactive Fixed Effects,” *Journal of Econometrics*, 197, 323–347.
- SHIN, Y. AND M. THORNTON (2019): “The Spatio-Temporal Autoregressive Distributed Lag Modelling Approach to an Analysis of the Spatial Heterogeneity and Diffusion Dependence,” Mimeo: University of York.
- SHIN, Y., B. YU, AND M. J. GREENWOOD-NIMMO (2014): “Modelling Asymmetric Cointegration and Dynamic Multipliers in a Nonlinear ARDL Framework,” in *Festschrift in Honor of Peter Schmidt*:

- Econometric Methods and Applications*, ed. by W. Horrace and R. Sickles, New York (NY): Springer Science & Business Media, 281–314.
- SIMS, C. A. (1974): “Distributed Lags,” in *Frontiers of Quantitative Economics*, ed. by D. A. Kendrick and M. D. Intriligator, Amsterdam: North Holland, vol. 2.
- STOCK, J. H. AND M. W. WATSON (1993): “A Simple Estimator of Cointegrating Vectors in Higher Order Integrated Systems,” *Econometrica*, 61, 783–820.
- SÜSSMUTH, B. AND U. WOITEK (2013): “Estimating Dynamic Asymmetries in Demand at the Munich Oktoberfest,” *Tourism Economics*, 19, 653–674.
- THOMAS, J. (1977): “Some Problems in the use of Almon’s Technique in the Estimation of Distributed Lags,” *Empirical Economics*, 2, 175–193.
- WALD, A. (1943): “Tests of Statistical Hypotheses Concerning Several Parameters When the Number of Observations is Large,” *Transactions of the American Mathematical Society*, 54, 426–482.
- WALLIS, K. F. (1969): “Some Recent Developments in Applied Econometrics: Dynamic Models and Simultaneous Equation Systems,” *Journal of Economic Literature*, 7, 771–796.
- WICKENS, M. R. AND T. S. BREUSCH (1988): “Dynamic Specification, the Long Run Estimation of the Transformed Regression Models,” *Economic Journal*, 98, 189–205.
- XIAO, Z. (2009): “Quantile Cointegrating Regression,” *Journal of Econometrics*, 150, 248–260.
- ZELLNER, A. (1979): “Statistical Analysis of Econometric Models,” *Journal of the American Statistical Association*, 74, 628–643.