White Rose Research Online

eprints@whiterose.ac.uk
https://eprints.whiterose.ac.uk
Universities of Leeds, Sheffield and York

Deposited via The University of Leeds.

White Rose Research Online URL for this paper:
https://eprints.whiterose.ac.uk/id/eprint/176906/

Version: Supplemental Material

Article:
VilÀ, M, Dunn, AM, Essl, F et al. (2021) Viewing Emerging Human Infectious Epidemics through the Lens of Invasion Biology. BioScience, 71 (7). pp. 722-740. ISSN: 0006-3568

https://doi.org/10.1093/biosci/biab047

© The Author(s) 2021. Published by Oxford University Press on behalf of the American Institute of Biological Sciences. This is an author produced version of an article published in Bioscience. Uploaded in accordance with the publisher's self-archiving policy.

Reuse
Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown
If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

UNIVERSITY OF LEEDS
University of Sheffield
UNIVERSITY of York

Vilà M, AM Dunn, F Essl, E Gómez-Díaz, PE Hulme, JM Jeschke, MA Núñez, RS Ostfeld, A Pauchard, A Ricciardi & B Gallardo. Viewing emerging human infectious epidemics through the lens of invasion biology. BioScience

Supplementary material

Table S1. Hypotheses in biological invasions and their parallels in human epidemics. The list of hypotheses follows the classification provided in Enders et al. (2020).

Hypothesis	**Description for biological invasions**	**Parallelism with epidemics**	**References**
Disturbance	The invasion of non-native species is higher in disturbed than in relatively undisturbed ecosystems	There is a positive relation between disease outbreaks and ecosystem disturbance; host susceptibility also increases with altered body conditions	Elton (1958), Hobbs and Huenneke (1992), Wolfe et al. (2005), Keesing et al. (2010), Myers et al. (2013)
Reckless invader *aka* 'boom-bust'	A population of a non-native species that is highly successful shortly after its introduction can decline or disappear over time due to different reasons such as competition with other introduced species or adaptation of native species	The incidence of a pathogen can decline because of different reasons such as the extinction of the wildlife reservoir or acquired immunity of the host	Simberloff and Gibbons (2004), Yates et al. (2006)
Colonization pressure	As the number of introduced non-native species increases, the number of invasive species in that location is predicted to increase	Coinfections create heterogeneity in the host population, which can lead to evolutionary branching in the parasite population and the emergence of a hyper-virulent parasite strategy	Lockwood et al. (2009), Alizon et al. (2008), Alizon et al. (2013)
Ecological naivety *aka* eco-evolutionary naivety	The impact of a non-native species on biodiversity is influenced by experience of the invaded community. Thus, the evolutionary largest impacts are caused by species invading systems where no phylogenetically or functionally similar species exist	Pathogens are more virulent in naïve hosts than in previous infected hosts who have developed resistance to infection	Diamond and Case (1986), Ricciardi and Atkinson (2004), Pascual et al. (2008), Alizon et al. (2009), Domínguez-Andrés and Netea (2019)
Enemy release	The absence of enemies in the exotic range facilitates invasion	The absence of virophages and bacteriophagues in novel infested populations facilitates infections	Keane and Crawley (2002), Dalmasso et al. (2014)
Enemy reduction	The partial release of enemies in the exotic range facilitates invasion	The partial absence of virophages and bacteriophagues in novel infested populations increase infections	Colautti et al. (2004), Dalmasso et al. (2014)

Human commensalism	Species that live in close proximity to humans are more successful in invading new areas than other species	There is higher probability of zoonosis and pathogen spill over in human populations that are in close proximity to animal reservoirs and/or vectors populations	Jeschke and Strayer (2006), Gallardo et al. (2015)
Ideal weed	The invasion success of a non-native species depends on particular life-history traits (e.g. seed bank, clonality)	Human pathogens have life-history traits that increase their establishment and persistence (e.g. transmission modes, standing genetic variation, plasticity, production of exoenzimes)	Baker (1965), Rejmánek and Richardson (1996), Alcamí and Koszinowski (2000), Woolhouse et al. (2005), Easterday (2020)
Propagule pressure	The number and frequency of introduced individuals of a non-native species increases the probability of invasion	The amount of the pathogen exposed to the human host increase the likelihood of infection	Lockwood et al. (2005), Horrocks et al. (2011), Hartfield and Allizon (2013)
Darwin's naturalization	The invasion of non-native species is higher in areas that are poor in closely related species than in areas that are rich in closely related species	Pathogens that invoke an immune response immediately after infection can also provide partial cross-protection against other strains of the same or closely related pathogens	Daehler (2001), Horimoto and Kawaoka (2005), Bhattacharyya et al. (2015)
Adaptation	The likelihood of invasion depends on non-native species preadaptation to the conditions in the exotic range. Invasive species that are related to native species are more successful in this adaptation	Introduced pathogens that are more related to native pathogens will be more successful as they will be preadapted to the native hosts due to long-term co-evolutionary interactions	Duncan and Williams (2002), Gagneux et al. (2006), Prohaska et al. (2019)
Ecological imbalance	Invasion patterns are a function of the evolutionary characteristics of both the recipient region and potential donor regions. Species from regions with highly diverse evolutionary lineages are more likely to become successful invaders in less diverse regions	Pathogen evolution depend on human modification of the habitat	Fridley and Sax (2014), Myers et al. (2013)
Evolution of increased competitive ability	There is a trade-off between defense and competitive ability. After having been released from natural enemies, non-native species allocate more energy to growth and/or reproduction, which makes them more competitive	There is a trade-off between high transmission rates and long infectious periods. Selection for pathogen virulence and horizontal transmission is highest at the onset of an epidemic but decreases thereafter, as the epidemic depletes the pool of susceptible hosts	Blossey and Nötzold (1995), Bolker et al. (2010), Berngruber et al. (2013)

Empty niche	The invasion of non-native species increases with the availability of empty niches in the exotic range	Infections increase in human populations not previously exposed to the pathogens	MacArthur (1970), Conn (2009)
Invasional meltdown	The presence of non- native species in an ecosystem facilitates invasion by additional species, increasing their likelihood of survival or ecological impact	Coinfections migh lead to increased suscepibilityto other infections though inmunosuppresion, but not always	Simberloff and Holle (1999), Graham (2008), Alizon and van Baalen (2008)
Tens rule	Approximately 10% of species successfully take consecutive steps of the invasion process	To be tested	Williamson and Brown (1986)
Biotic acceptance *aka* 'the rich get richer'	Ecosystems tend to accommodate the establishment and coexistence of non-native species despite the presence and abundance of native species	Several pathogens can accumulate and coexist in the body after competitive interactions. Hyper-virulent strains cannot coexist	Stohlgren (2003), Alizon and van Baalen (2008), Mideo (2009)
Biotic indirect effects	Non-native species benefit from different indirect effects triggered by native species	New pathogens can profit from endemic pathogens (e.g. that have induced immunodepression, immune evasion)	Callaway (2004), Alizon and van Baalen (2008), Mideo (2009)
Biotic resistance *aka* diversity invasibility hypothesis	An ecosystem with high biodiversity is more resistant against non-native species than an ecosystem with lower biodiversity	Microbial diversity reduce the establishment of pathogens	Elton (1958), Levine and D'Antonio (1999), Mallon et al. (2015), Libertucci and Young (2019)
Dynamic equilibrium model	The establishment of a non-native species depends on natural fluctuations of the ecosystem, which influence the level of competition from local species	Altered physiological, immunological or physiological conditions affect host susceptibility to infection and the severity of the disease	Huston (1979), Plowright et al. (2017)
Enemy of my enemy *aka* accumulation-of-local pathogens hypothesis	Introduced enemies of a non-native species are less harmful to the non-native than to the native species	Pathogens are less virulent in the hosts with whom they have coevolved than in new hosts	Eppinga et al. (2006), Bolker et al. (2010)
Enemy inversion	Introduced enemies of non-native species are less harmful in the exotic than the native range, due to altered biotic and abiotic conditions	This contradicts the hypothesis that pathogens are less virulent in the hosts with whom they have coevolved than in new hosts	Colautti et al. (2004), Bolker et al. (2010)
Environmental heterogeneity	The invasion of non-native species is high if the introduced range has a highly heterogeneous environment	An epidemic spreads more rapidly in contact with heterogeneous human populations (genetic variation, age, sex, nutritional status, drug	Melbourne et al. (2007), Stecher et al. (2007), Cousineau and Alizon (2014)

		exposure, etc.) and also within host heterogeneities	
Global competition	A large number of different non-native species is more successful than a small number	This is controversial. An infected host might be immune-compromised and be more susceptible to new hosts. However, an infected host might also be prevented to be further infested because not providing the required resources or environmental conditions.	Colautti et al. (2006), Kamiya et al. (2018)
Habitat filtering	The invasion of non-native species in the new area is high if they are pre-adapted to the habitat conditions of the area	The host immune system acts as an important environmental filter to limit the extent of available niches to pathogens.	Weiher and Keddy (1995), Costello et al. (2012)
Increased resource availability	The invasion of non-native species increases with the availability of resources	Disease outbreaks increase with availability of resources	Sher and Hyatt (1999), Watson et al. (2007)
Increased susceptibility	If a non-native species has a lower genetic diversity than the native species, there will be a low probability that the non-native species establishes itself	Slow evolving pathogens and with low genetic variation, would infest less efectively. To be tested	Colautti et al. (2004)
Island susceptibility	Non-native species are more likely to become established and have major ecological impacts on islands than on continents	Emerging pathogens are more likely to become established and have major impacts on islands than on continents. To be tested	Jeschke (2008)
Limiting similarity	The invasion of non-native species is high if they strongly differ from native species, and low if they are similar to native species	The lack of evolutionary adaptation to a novel pathogen cause higher infections	MacArthur and Levins (1967), Brown et al. (2012), Harimoto and Kawaoka (2005)
Missed mutualisms	In their exotic range, non-native species suffer from missing mutualists	In a new range, missing vectors or reservoirs can deter the infection process	Mitchell et al. (2006), Plowright et al. (2017)
New associations	New relationships between non-native and native species can positively or negatively influence the establishment of the non-native species	In zoonotic diseases, the existence of encountering new reservoirs can definitely influence the spread of the pathogen	Colautti et al. (2006), Plowright et al. (2017)

Novel weapons	In the exotic range, non-native species can have a competitive advantage against native species because they possess a novel weapon, that is, a trait that is new to the resident community of native species and, therefore, affects them negatively	Emerging pathogens have new weapons to which the host has not been exposed before	Callaway and Ridenour (2004), Alizon et al. (2009)
Opportunity windows	The invasion of non-native species increases with the availability of empty niches in the exotic range, and the availability of these niches fluctuates spatio-temporally	The environment created by the host determines the number of potential niche opportunities	Johnstone (1986), Costello et al. (2012), Buffie and Pamer (2013)
Plasticity hypothesis	Invasive species are more phenotypically plastic than non-invasive or native ones	Antigenic variation is a similar strategy as the phenotypic variation of invasive species to cope with different environmental conditions. Phenotypic plasticity in pathogens facilitate hosts shifts and rapid adaptation to new and unpredictable host environments	Davidson et al. (2011), Palmer et al. (2016), De Fine Licht (2018)
Polyploidy	Polyploid organisms, particularly plants, are predicted to have an increased invasion likelihood, since polyploidy can lead to higher fitness during the establishment phase and/or increased potential for subsequent adaptation	To be tested	Beest et al. (2012)
Resource-enemy release	The non-native species is released from its natural enemies and can spend more energy in its reproduction, and invasion increases with the availability of resources	Selection for pathogen virulence and horizontal transmission is highest at the onset of an epidemic but decreases thereafter, as the epidemic depletes the pool of susceptible hosts	Blumenthal (2006), Bolker et al. (2010), Berngruber et al. (2013)
Shifting defense hypothesis	After having been released from natural specialist enemies, non-native species will allocate more energy to cheap (energy inexpensive) defenses against generalist enemies and less energy to expensive defenses against specialist enemies; the energy gained in this way will be invested in growth and/or reproduction, which makes the non-native species more competitive	Specialist pathogens would become generalist after host switch	Doorduin and Vrieling (2011), Woodcock et al. (2017)
Specialist-generalist	Non-native species are more invasive in a new region if the local predators are specialists and local mutualists are generalists	Generalist parasites use generalist cell receptors	Callaway et al. (2004), Baranowski et al. (2001)

Sampling	A large number of different non-native species is more likely to become invasive than a small number due to interspecific competition. Also, the species identity of the locals is more important than the richness in terms of the invasion of an area	Exposure to many pathogens, increase the chances to get infected by someone	Crawley et al. (1999), Alizon et al. (2013)

References

Alcami A, Koszinowski UH. 2000. Viral mechanisms of immune evasion. Trends in Microbiology 8: 410–418.

Alizon S, van Baalen M. 2008. Multiple infections, immune dynamics, and the evolution of virulence. The American Naturalist 172: E150–E168.

Alizon S, Mideo N, van Baalen M. 2009. Virulence evolution and the trade-off hypothesis: history, current state of affairs and the future. Journal of Evolutionary Biology 22: 245–259.

Alizon S, Roode JC de, Michalakis Y. 2013. Multiple infections and the evolution of virulence. Ecology Letters 16: 556–567.

Baker HG. 1965. Characteristics and modes of origin of weeds. Pages 147–172 in Baker HG and Stebbins GL, eds. The genetics of colonizing species. Academic Press Inc., N.Y.

Baranowski E, Ruiz-Jarabo CM, Domingo E. 2001. Evolution of cell recognition by viruses. Science 292: 1102–1105.

Berngruber TW, Froissart R, Choisy M, Gandon S. 2013. Evolution of virulence in emerging epidemics. PLOS Pathogens 9: e1003209.

Bhattacharyya S, Gesteland PH, Korgenski K, Bjørnstad ON, Adler FR. 2015. Cross-immunity between strains explains the dynamical pattern of paramyxoviruses. Proceedings of the National Academy of Sciences 112: 13396–13400.

Blossey B, Notzold R. 1995. Evolution of increased competitive ability in invasive nonindigenous plants: a hypothesis. Journal of Ecology 83: 887–889.

Blumenthal DM. 2006. Interactions between resource availability and enemy release in plant invasion. Ecology Letters 9: 887–895.

Bolker BM, Nanda A, Shah D. 2010. Transient virulence of emerging pathogens. Journal of the Royal Society Interface 7: 811–822.

Brown SP, Cornforth DM, Mideo N. 2012. Evolution of virulence in opportunistic pathogens: generalism, plasticity, and control. Trends in Microbiology 20: 336–342.

Buffie CG, Pamer EG. 2013. Microbiota-mediated colonization resistance against intestinal pathogens. Nature Reviews Immunology 13: 790–801.

Callaway RM, Ridenour WM. 2004. Novel weapons: invasive success and the evolution of increased competitive ability. Frontiers in Ecology and the Environment 2: 436–443.

Callaway RM, Thelen GC, Rodriguez A, Holben WE. 2004. Soil biota and exotic plant invasion. Nature 427: 731–733.

Colautti RI, Grigorovich IA, MacIsaac HJ. 2006. Propagule pressure: a null model for biological invasions. Biological Invasions 8: 1023–1037.

Colautti RI, MacIsaac HJ. 2004. A neutral terminology to define 'invasive'species. Diversity and Distributions 10: 135–141.

Colautti RI, Ricciardi A, Grigorovich IA, MacIsaac HJ. 2004. Is invasion success explained by the enemy release hypothesis? Ecology Letters 7: 721–733.

Conn DB. 2009. Presidential address: parasites on a shrinking planet. Journal of Parasitology 95: 1253–1263.

Costello EK, Stagaman K, Dethlefsen L, Bohannan BJM, Relman DA. 2012. The application of ecological theory toward an understanding of the human microbiome. Science 336: 1255–1262.

Cousineau SV, Alizon S. 2014. Parasite evolution in response to sex-based host heterogeneity in resistance and tolerance. Journal of Evolutionary Biology 27: 2753–2766.

Crawley MJ, Brown SL, Heard MS, Edwards GR. 1999. Invasion-resistance in experimental grassland communities: species richness or species identity? Ecology Letters 2: 140–148.

Daehler CC. 2001. Darwin's naturalization hypothesis revisited. The American Naturalist 158: 324–330.

Dalmasso M, Hill C, Ross RP. 2014. Exploiting gut bacteriophages for human health. Trends in Microbiology 22: 399–405.

Davidson AM, Jennions M, Nicotra AB. 2011. Do invasive species show higher phenotypic plasticity than native species and, if so, is it adaptive? A meta-analysis. Ecology letters 14: 419–431.

De Fine Licht HH. 2018. Does pathogen plasticity facilitate host shifts? PLOS Pathogens 14: e1006961.

Diamond J. 1986. Overview: introductions, extinctions, exterminations, and invasions. Pages 65–79 in Diamond J and Case TJ, eds. Community Ecology. New York: Harper & Row.

Domínguez-Andrés J, Netea MG. 2019. Impact of historic migrations and evolutionary processes on human immunity. Trends in Immunology 40: 1105–1119.

Doorduin LJ, Vrieling K. 2011. A review of the phytochemical support for the shifting defence hypothesis. Phytochemistry Reviews 10: 99–106.

Duncan RP, Williams PA. 2002. Ecology: Darwin's naturalization hypothesis challenged. Nature 417: 608–609.

Easterday WR. 2020. The first step in the success or failure of emerging pathogens. Proceedings of the National Academy of Sciences 117: 29271–29273.

Elton CS. 1958. The ecology of invasions by animals and plants. Springer, Cham.

Enders M, Havemann F, Ruland F, Bernard-Verdier M, Catford JA, Gómez-Aparicio L, Haider S, Heger T, Kueffer C, Kühn I, Meyerson L, Musseau C, Novoa A, Ricciardi A, Sagouis A, Schittko C, Strayer DL, Vilà M, Essl F, Hulme PE, van Kleunen M, Kumschick S, Lockwood JL, Mabey AL, McGeoch MA, Palma E, Pysek P, Saul W-C, Yannelli FA, Jeschke JM. 2020. A conceptual map of invasion biology: Integrating hypotheses into a consensus network. Global Ecology and Biogeography 29: 978–991.

Eppinga MB, Rietkerk M, Dekker SC, De Ruiter PC, Van der Putten WH, Van der Putten WH. 2006. Accumulation of local pathogens: a new hypothesis to explain exotic plant invasions. Oikos 114: 168–176.

Fridley JD, Sax DF. 2014. The imbalance of nature: revisiting a Darwinian framework for invasion biology. Global Ecology and Biogeography 23: 1157–1166.

Gagneux S, DeRiemer K, Van T, Kato-Maeda M, Jong BC de, Narayanan S, Nicol M, Niemann S, Kremer K, Gutierrez MC, Hilty M, Hopewell PC, Small PM. 2006. Variable host–pathogen compatibility in Mycobacterium tuberculosis. Proceedings of the National Academy of Sciences 103: 2869–2873.

Gallardo B, Zieritz A, Aldridge DC. 2015. The importance of the human footprint in shaping the global distribution of terrestrial, freshwater and marine invaders. Plos One 10: e0125801.

Graham AL. 2008. Ecological rules governing helminth–microparasite coinfection. Proceedings of the National Academy of Sciences 105: 566–570.

Hartfield M, Alizon S. 2013. Introducing the outbreak threshold in epidemiology. PLOS Pathogens 9: e1003277.

Hobbs RJ, Huenneke LF. 1992. Disturbance, diversity, and invasion: implications for conservation. Conservation Biology 6: 324–337.

Horimoto T, Kawaoka Y. 2005. Influenza: lessons from past pandemics, warnings from current incidents. Nature Reviews Microbiology 3: 591–600.

Horrocks NP, Matson KD, Tieleman BI. 2011. Pathogen pressure puts immune defense into perspective. Oxford University Press.

Huston M. 1979. A general hypothesis of species diversity. The American Naturalist 113: 81–101.

Jeschke JM. 2008. Across islands and continents, mammals are more successful invaders than birds. Diversity and Distributions 14: 913–916.

Jeschke JM, Strayer DL. 2006. Determinants of vertebrate invasion success in Europe and North America. Global Change Biology 12: 1608–1619.

Johnstone I. 1986. Plant invasion windows: a time-based classification of invasion potential. Biological Reviews 61: 369–394.

Kamiya T, Mideo N, Alizon S. 2018. Coevolution of virulence and immunosuppression in multiple infections. Journal of Evolutionary Biology 31: 995–1005.

Keane RM, Crawley MJ. 2002. Exotic plant invasions and the enemy release hypothesis. Trends in Ecology & Evolution 17: 164–170.

Keesing F, Belden LK, Daszak P, Dobson A, Harvell CD, Holt RD, Hudson P, Jolles A, Jones KE, Mitchell CE, Myers SS, Bogich T, Ostfeld RS. 2010. Impacts of biodiversity on the emergence and transmission of infectious diseases. Nature 468: 647–652.

Levine JM, D'Antonio CM. 1999. Elton revisited: A review of evidence linking diversity and invasibility. Oikos 87: 15–26.

Libertucci J, Young VB. 2019. The role of the microbiota in infectious diseases. Nature Microbiology 4: 35–45.

Lockwood JL, Cassey P, Blackburn T. 2005. The role of propagule pressure in explaining species invasions. Trends in Ecology & Evolution 20: 223–228.

Lockwood JL, Cassey P, Blackburn TM. 2009. The more you introduce the more you get: the role of colonization pressure and propagule pressure in invasion ecology. Diversity and Distributions 15: 904–910.

MacArthur R. 1970. Species packing and competitive equilibrium for many species. Theoretical Population Biology 1: 1–11.

MacArthur R, Levins R. 1967. The limiting similarity, convergence, and divergence of coexisting species. The American Naturalist 101: 377–385.

Mallon CA, Van Elsas JD, Salles JF. 2015. Microbial invasions: the process, patterns, and mechanisms. Trends in Microbiology 23: 719–729.

Melbourne BA, Cornell HV, Davies KF, Dugaw CJ, Elmendorf S, Freestone AL, Hall RJ, Harrison S, Hastings A, Holland M, Holyoak M, Lambrinos J, Moore K, Yokomizo H. 2007. Invasion in a heterogeneous world: resistance, coexistence or hostile takeover? Ecology Letters 10: 77–94.

Mideo N. 2009. Parasite adaptations to within-host competition. Trends in Parasitology 25: 261–268.

Mitchell CE, Agrawal AA, Bever JD, Gilbert GS, Hufbauer RA, Klironomos JN, Maron JL, Morris WF, Parker IM, Power AG, Seabloom EW, Torchin ME, Vazquez DP. 2006. Biotic interactions and plant invasions. Ecology Letters 9: 726–740.

Myers SS, Gaffikin L, Golden CD, Ostfeld RS, H. Redford K, H. Ricketts T, Turner WR, Osofsky SA. 2013. Human health impacts of ecosystem alteration. Proceedings of the National Academy of Sciences of the United States of America 110: 18753–18760.

Palmer GH, Bankhead T, Seifert HS. 2016. Antigenic variation in bacterial pathogens. Pages 445–480 in Kudva IT, Cornick NA, Plummer PJ, Zhang Q, Nicholson TL, Bannantine JP, and Bellaire BH, eds. Virulence mechanisms of bacterial pathogens. American Society for Microbiology.

Pascual M, Cazelles B, Bouma MJ, Chaves LJ, Koelle K. 2008. Shifting patterns: malaria dynamics and rainfall variability in an African highland. Proceedings of the Royal Society B: Biological Sciences 275: 123–132.

Plowright RK, Parrish CR, McCallum H, Hudson PJ, Ko AI, Graham AL, Lloyd-Smith JO. 2017. Pathways to zoonotic spillover. Nature Reviews Microbiology 15: 502–510.

Prohaska A, Racimo F, Schork AJ, Sikora M, Stern AJ, Ilardo M, Allentoft ME, Folkersen L, Buil A, Moreno-Mayar JV, Korneliussen T, Geschwind D, Ingason A, Werge T, Nielsen R, Willerslev E. 2019. Human disease variation in the light of population genomics. Cell 177: 115–131.

Rejmanek M, Richardson DM. 1996. What attributes make some plant species more invasive? Ecology 77: 1655–1661.

Ricciardi A, Atkinson SK. 2004. Distinctiveness magnifies the impact of biological invaders in aquatic ecosystems. Ecology Letters 7: 781–784.

Sher AA, Hyatt LA. 1999. The disturbed resource-flux invasion matrix: a new framework for patterns of plant invasion. Biological Invasions 1: 107–114.

Simberloff D, Gibbons L. 2004. Now you see them, now you don't! – Population Crashes of Established Introduced Species. Biological Invasions 6: 161–172.

Simberloff D, von Holle B. 1999. Positive interactions of nonindigenous species: invasional meltdown? Biological Invasions 1: 21–32.

Stecher B, Robbiani R, Walker AW, Westendorf AM, Barthel M, Kremer M, Chaffron S, Macpherson AJ, Buer J, Parkhill J, Dougan G, von Mering C, Hardt W-D. 2007. Salmonella enterica serovar typhimurium exploits inflammation to compete with the intestinal microbiota. PLoS Biology 5: e244.

Stohlgren TJ, Barnett DT, Kartesz JT. 2003. The rich get richer: patterns of plant invasions in the United States. Frontiers in Ecology and the Environment 1: 11–14.

Te Beest M, Le Roux JJ, Richardson DM, Brysting AK, Suda J, Kubešová M, Pyšek P. 2012. The more the better? The role of polyploidy in facilitating plant invasions. Annals of Botany 109: 19–45.

Watson JT, Gayer M, Connolly MA. 2007. Epidemics after natural disasters. Emerging Infectious Diseases 13: 1–5.

Weiher E, Keddy PA. 1995. Assembly rules, null models, and trait dispersion: new questions from old patterns. Oikos 159–164.

Williamson MH, Brown KC. 1986. The analysis and modelling of British invasions. Philosophical Transactions of the Royal Society of London. B, Biological Sciences 314: 505–522.

Wolfe ND, Daszak P, Kilpatrick AM, Burke DS. 2005. Bushmeat hunting, deforestation, and prediction of zoonotic disease. Emerging Infectious Diseases 11: 1822–1827.

Woodcock DJ, Krusche P, Strachan NJC, Forbes KJ, Cohan FM, Méric G, Sheppard SK. 2017. Genomic plasticity and rapid host switching can promote the evolution of generalism: a case study in the zoonotic pathogen Campylobacter. Scientific Reports 7: 9650.

Woolhouse MEJ, Haydon DT, Antia R. 2005. Emerging pathogens: the epidemiology and evolution of species jumps. Trends in Ecology & Evolution 20: 238–244.

Yates A, Antia R, Regoes RR. 2006. How do pathogen evolution and host heterogeneity interact in disease emergence? Proceedings of the Royal Society B: Biological Sciences 273: 3075–3083.