



This is a repository copy of *Was the implementation of standardised tobacco packaging legislation in England associated with changes in smoking prevalence? A segmented regression analysis between 2006 and 2019.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/178269/>

Version: Accepted Version

---

**Article:**

Opazo Breton, M. [orcid.org/0000-0002-1226-7541](https://orcid.org/0000-0002-1226-7541), Britton, J., Brown, J. et al. (2 more authors) (2023) Was the implementation of standardised tobacco packaging legislation in England associated with changes in smoking prevalence? A segmented regression analysis between 2006 and 2019. *Tobacco Control*, 32 (2). pp. 195-204. ISSN 0964-4563

<https://doi.org/10.1136/tobaccocontrol-2021-056694>

---

© 2021 Author(s) (or their employer(s)). Reuse of this manuscript version (excluding any databases, tables, diagrams, photographs and other images or illustrative material included where another copyright owner is identified) is permitted strictly pursuant to the terms of the Creative Commons Attribution-NonCommercial 4.0 International (CC-BY-NC 4.0) (<http://creativecommons.org/licenses/by-nc/4.0/>).

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial (CC BY-NC) licence. This licence allows you to remix, tweak, and build upon this work non-commercially, and any new works must also acknowledge the authors and be non-commercial. You don't have to license any derivative works on the same terms. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

**Was the implementation of standardised tobacco packaging legislation in England associated with changes in smoking prevalence? A segmented regression analysis between 2006 and 2019**

**Magdalena Opazo Breton<sup>1</sup>, John Britton<sup>2</sup>, Jamie Brown<sup>3,4</sup>, Emma Beard<sup>3,4</sup>, Ilze Bogdanovica<sup>2</sup>**

**Affiliations:**

1 School of Health and Related Research (ScHARR), University of Sheffield, Sheffield, UK

2 School of Medicine, University of Nottingham/ UK Centre for Tobacco and Alcohol Studies

3 Department of Clinical, Educational and Health Psychology, University College London, London, UK

4 Cancer Research UK Health Behaviour Research Centre, University College London, London, UK

**Running head:** Standardised packaging and smoking prevalence

**Word count:** 3,420 words

**Declaration of competing interests:** Jamie Brown and Emma Beard have received unrestricted research grants from Pfizer related to smoking cessation. John Britton, Magdalena Opazo Breton & Ilze Bogdanovica have no competing interests to declare. All authors declare no financial links with tobacco companies or e-cigarette manufacturers or their representatives.

**Keywords:** smoking, prevalence, standardized packaging

## **ABSTRACT**

**Background and aim:** In 2016 England initiated the implementation of standardised tobacco packaging, introduced in conjunction with minimum pack sizes and other measures included in the 2014 European Tobacco Products Directive, over the course of a one-year sell-off period ending in May 2017. These measures have been shown to have been associated with increases in tobacco prices and product diversity. We now investigate the association between implementation of the new legislation and smoking status in England.

**Design:** Segmented regression analysis of repeated cross-sectional surveys using a GLM model with individual-level data to test for a change in trend and immediate step change.

**Setting:** England.

**Participants:** Participants in the Smoking Toolkit Study, which involves repeated, cross-sectional household surveys of individuals aged 16 years and older in England. The sample included 278,219 individual observations collected between November 2006 and December 2019.

**Intervention:** Implementation of standardised packaging legislation (May 2016 and May 2017).

**Measurements:** Individual level current smoking status adjusted for implementation of tobacco control policies, cigarette price, seasonality and autocorrelation.

**Findings:** The implementation of standardised packaging was associated with a significant step reduction in the odds of being a smoker after May 2017 (Odds Ratio (OR) 0.93; 95% Confidence Interval (CI) 0.87 to 0.99). The magnitude of the association was similar when modelling the step change in May 2016 at the start of the one-year policy implementation period (OR 0.90; 95% CI: 0.83 to 0.97).

**Conclusions:** This is the first independent study demonstrating that implementation of standardised packaging was associated with a reduction in smoking in England which occurred in anticipation of, rather than after, full policy implementation. It appears that the odds of being a smoker was affected by the prospect of the move to standardised packs and accompanying legislation.

## INTRODUCTION

Over the past decade the prevalence of current tobacco smoking in England has been in sustained decline, falling from 19.8% in 2011 to 13.9%, or approximately 5.7 million smokers, in 2019 [1]. This reduction in prevalence has been particularly marked among children and young adults [1, 2] and those in higher socio-economic groups [1], and has been achieved by a range of tobacco control policies implemented by the UK government over the past two decades, the most recent of which was the introduction of standardised packaging legislation in May 2016 [3]. This legislation determined that after a one-year transition period to May 2017, manufactured cigarettes and roll-your-own tobacco products could be sold in England only if packaged in generic drab dark brown packs with brand names and a single descriptor presented in a standard font. These requirements were implemented alongside the 2014 European Tobacco Products Directive, which among other measures mandated minimum pack sizes and larger pictorial health warnings [4].

The primary aim of the standardised packaging legislation was to make smoking less appealing to and discourage smoking uptake among young people [5], but there is evidence that standardised packaging legislation might also reduce the prevalence of smoking among adults [6, 7]. Evidence from Australia, which in 2012 became the first country to introduce standardised packaging, suggests that implementation led to an increase in quitline calls [8] and increased the rate of decline in smoking prevalence [9]. Research on the introduction of standardised packaging in England has demonstrated that implementation has been associated with considerable increases in the price of tobacco products, switching to less expensive tobacco products and increased use of e-cigarettes among smokers [10-13]. However, the effect of standardised packaging on smoking prevalence in England has not yet been explored by researchers independent of the tobacco industry's funding. In 2018 Tobacco Manufacturers' Association published analysis using Smoking Toolkit Study (STS) data suggesting that implementation of standardised packaging was associated with an increase in smoking prevalence [14]. However, the data analysis was based on a very basic comparison of three months rolling average with the data from the same time period in the previous year. The analysis was considerably underpowered and did not consider any potential confounders. The aim of this study was therefore 1) to investigate the effect of the introduction of standardised packaging on smoking prevalence by estimating step and trend changes in individual current smoking status after the policy was implemented in England, and aggregating these results to estimate changes in smoking prevalence; 2) to explore whether differences in step and trend

changes were observed in different population subgroups defined by age, sex and socio-economic status.

## **METHODS**

### ***Data sources and research design***

We used data from Smoking Toolkit Study (STS), a monthly interview-based household survey of smoking status in representative samples of ~1,700 adults aged 16 and over in England that has collected data since 2006 [15]. The survey used a random location sampling design to select grouped output areas (~300 households) stratified by socio-demographic characteristics, while interviewers selected households within areas based on quotas targeted to the characteristics of the output area. Face-to-face computer-assisted interviews were then carried out with one household member. More details about the methods and the data can be found elsewhere [16-19]. Comparisons to other national surveys suggests the design produces a sample of the population in England representative of key demographic variables, smoking prevalence, and cigarette consumption [15].

We used individual-level data from November 2006 to December 2019 to carry out a segmented regression analysis to study level and post-slope changes [20, 21] in current smoking status after the implementation of the standardised packaging policy in England, using a before and after approach [21, 22]. Our analysis first explored effects before and after May 2017, the end of the one-year implementation period, as the main implementation point, and then before and after May 2016 and explored effects after the start of that implementation period. To check the robustness of our results, we performed a sensitivity analysis using implementation dates based on the proportion of tobacco sales in standardised packs, derived from Nielsen sales data, as a marker of the extent to which standardised packaging had been implemented during the implantation period [11]. These indicated that standardised packs began to appear on the UK market in July 2016 [11], and that sales of branded packs probably did not end until July 2017 [11, 13].

### ***Measures***

#### ***Outcome variable***

The outcome variable was current smoking status. All participants from STS were classified as current smokers (with the value '1') if they responded affirmatively to any of the statements

‘I smoke cigarettes (including hand-rolled) every day’, ‘I smoke cigarettes (including hand-rolled), but not every day’, or ‘I do not smoke cigarettes at all, but I do smoke tobacco of some kind (e.g. Pipe, cigar or shisha)’; and as a non-smoker (with the value ‘0’) if they responded affirmatively to any of the statements ‘I stopped smoking completely in the last year’, ‘I stopped smoking completely more than a year ago’, or ‘I have never been a smoker’. This question differs from the question used in other nationally representative surveys. Hence, two actions were taken as a precautionary measure. First, we compared quarterly trends in smoking prevalence from STS data to Annual Population Survey (APS), and second, we compared general monthly smoking prevalence in STS to that of different tobacco product users, such as manufactured cigarettes only, hand-rolled tobacco only, combined manufactured cigarettes and hand-rolled tobacco users, and other tobacco product users, which would include those using pipe cigar or cigarillos among others. These two comparisons can be found in Figure S1 and Figure S2 in the Online Supplementary Material and show that trends in STS were fairly similar for most quarters when compared to APS data and that most of the smoking prevalence figures refer to only manufactured cigarette and only hand-rolled tobacco users, which were the most affected by standardised packaging legislation.

#### *Segmented regression analysis variables*

For our analysis after full implementation of the policy we created a level variable that took the value ‘0’ for all observations from months up to and including May 2017 (before), and the value ‘1’ for all observations after May 2017 (after). For our analysis after the policy start date we added a before and after variable for May 2016, while our sensitivity analysis studied level changes using July 2016 and July 2017 as the start and full implementation dates. We also created a slope variable (post intervention) with values between zero and one that increases in equal amounts each month after full implementation up to eighteen months to study changes in smoking status trend after full policy implementation (post-slope after May 2017 and post-slope after July 2017 in our sensitivity analysis). An equivalent slope variable was created for the implementation period (between May 2016 and May 2017) as well as for the period between July 2016 and July 2017. In addition to the level and post intervention slope variables, the model included a linear time-trend variable with equally increasing values starting in the first month of the data until the last month of our study period [23].

#### *Subgroup analysis by age and socio-economic status*

We studied changes in the likelihood of being a current smoker among six population subgroups defined by sex (males and females), age (individuals aged 16-25 and those above 25 years of age) and socioeconomic status (routine and manual occupations vs non-routine occupations).

#### *Control for the effects of other tobacco control policies*

During our study data period the following tobacco control policies were implemented: smoke-free public places legislation in July 2007, an increase in the minimum age of sale from 16 to 18 years in October 2007, and a ban on point of sale displays which applied to large shops from April 2012 and small shops from April 2015. We created a dummy variable for each of these policies which assigned the value '0' for all months up to and including the implementation month, and the value '1' for all subsequent months. To adjust for the effects of tobacco tax and other price rises [24-28] we used the average monthly price for a 20 cigarette pack [29], adjusted for inflation using the Consumer Price Index (CPI) [30].

#### *Seasonality and autocorrelation*

Two additional variables were included to adjust our models: seasonality and autocorrelation. Regarding the first, evidence suggests that smoking has a seasonal pattern [31, 32]. Hence, we used a categorical 'month of the year variable' to account for possible differences in smoking status specific to the month of the year in which the survey took place. For autocorrelation, as we were using individual level data to estimate grouped policy effects we used robust standard errors and created a variable with lagged values (one lag) of smoking prevalence (general population smoking prevalence and subgroup smoking prevalence to use accordingly) [33].

#### *Statistical analysis*

We initially plotted aggregated monthly trends in current smoking prevalence in the general population and in the subgroups defined by sex, age and socioeconomic status using weighted STS data from November 2006 to December 2019 to illustrate overall prevalence trends for the population in England [15]. We compared smoking prevalence during the year the policy was implemented to the year before using a t-test to have a simple estimate of difference in prevalence before and after policy implementation.

We then used a generalized linear model (GLM) to estimate changes in level and slope of the likelihood of being a smoker after implementation of the standardised packaging policy using individual level data on a binary smoking status variable. We did not use survey weights for this analysis, but we performed the same analysis using quarterly data from Annual Population Survey (2010-2019) and run the general population analysis for different types of tobacco users in order to check the robustness of our results. The results from these two analyses can be found in Table S1 and Table S2 in the Supplementary Material. Our GLM models were defined using binomial family and logit link to estimate odds ratio (OR) and 95% confidence intervals for each of the variables included in the regression. Only results for level (before/after) and slope (implementation period and 18 months after implementation) were reported here, while full list of results for the main analyses can be found in the Online Supplementary Material (Table S4 to Table S1). We used robust standard errors, and adjustments by seasonality and autocorrelation since we estimated aggregate before/after and slope effects.

We estimated unadjusted models, which only included seasonality and autocorrelation, and adjusted models, which added other tobacco control policies implemented during the period 2006-2019 (smoking ban, change in minimum age of sale, and tobacco display ban in small and large shops), as well as our monthly average retail price variable. We first estimated the model exploring changes after May 2017, the full implementation date (level and post intervention slope). We then added May 2016, the start of policy implementation period, and explored level changes for before/after May 2016, the slope for the implementation period (slope May 2016-May 2017), level changes before/after May 2017, and post intervention slope 18 months after May 2017.

We estimated changes in level and slope among population subgroups by running the same models described for each population subgroup (the four model specifications six times), in order to study each group's smoking status separately. Interaction effects by subgroup were also investigated for the main analysis. These results can be found in the Online Supplementary Material (Table S3).

We performed a sensitivity analysis using the same models described above but exploring level and slope changes before/after July 2017, and before/after July 2016. Using July 2017 instead of May 2017 also allowed us to disentangle policy effects from any tax effect that were not



captured in our price variable, since in 2017 there were changes in tobacco taxes in March, May and November.

Finally, we plotted the linear predictions of our model for the whole sample of England against a counterfactual prediction reflecting the hypothetical situation ‘if the policy was not in place’ to visually compare smoking prevalence trends with and without (counterfactual) standardised packaging policy. To obtain the standardised packaging policy predicted trends we ran our unadjusted models, removing adjustments by seasonality to compute linear trends. To obtain the counterfactual predicted trend we estimated our GLM model only including a time-trend variable and limited the regression to the period before full implementation (May 2017), and to the period before implementation start date (May 2016). Then, we aggregated individual level predicted values from the unadjusted models, and from the counterfactual model to generate scatter plots of smoking prevalence combined with line graphs for the linear predictions. We performed all analyses in Stata 16.0, and the confidence level was set to 95%.

## RESULTS

Our sample included 278,219 individual observations collected between November 2006 and December 2019, of which 48.6% were from males and 51.5% from females, 15.8% from persons aged up to 25 years and 84.2% above 25 years old, while 40.3% were classified as manual workers and 59.7% as non-manual workers.

### Figure 1 here

There was a secular downward trend in smoking prevalence throughout the study period for the general population of England (Figure 1a). During the standardised packaging policy implementation period (May 2016 to May 2017) the prevalence of smoking was on average 17.9% (95% CI: 17.2 to 18.6%), while it was 19.2% (95% CI: 18.6 to 19.9%) in the year before the policy was implemented. Trends within age, sex and socioeconomic subgroups were similar to those in the total population, though among those aged 16-25, the prevalence of smoking declined rapidly from November 2006 until May 2012, then remained at around 24.9% (95% CI: 24.2% to 25.7%) until May 2016, declined to an average of 22.5% (95% CI: 21.1% to 23.8%) between May 2016 and May 2017, and then after a brief increase again assumed a decreasing trend (Figure 1c).

### *Regression results*

Our model before and after full implementation demonstrated a statistically significant level decrease in the odds of being a smoker after May 2017 (adjusted OR 0.93; 95% CI: 0.87 to 0.99) with no statistically significant change in post intervention slope. However, when May 2016 was included in the model (before/after policy start date), the observed level decrease in the odds of being a smoker was similar to the analysis after full implementation in both the unadjusted (OR 0.91; 95% CI: 0.85 to 0.98) and adjusted models (OR 0.90; 95% CI: 0.83 to 0.97), again with no significant change in post intervention slope. Hence, our results indicate that the level decrease in the odds of being a smoker was associated with the onset of standardized packaging in May 2016, and not the full implementation of, standardised packaging and other TPD measures.

### Table 1 here

Our subgroup analyses explored each population subgroup's smoking status and showed statistically significant step changes for females, males, those aged over 25 years old and manual occupations. For females there was a significant level decrease after May 2016 (adjusted OR: 0.89; 95% CI: 0.79 to 1.00), which was also observed among males (unadjusted OR 0.89; 95% CI: 0.80 to 0.98). Males also showed a statistically significant level decrease in our model exploring effects before/after May 2017 (unadjusted OR: 0.92; 95% CI: 0.85 to 1.00; adjusted OR: 0.90; 95% CI: 0.83 to 0.98). For population aged over 25 years old there was a significant level decrease after May 2017 in our model exploring effects before/after full implementation (unadjusted OR 0.92; 95% CI: 0.87 to 0.98; and adjusted OR 0.90; 95% CI: 0.84 to 0.96) and after May 2016 (unadjusted OR 0.90; 95% CI: 0.83 to 0.98; and adjusted OR 0.90; 95% CI: 0.82 to 0.98) in our model exploring effects before/after the policy start date. Finally, there was a significant decrease after May 2016 for manual occupations in our model exploring effects after the policy start date (adjusted OR: OR 0.85; 95% CI: 0.75 to 0.96). The only increase in the odds of being a smoker was observed among population aged 25 years old or younger after May 2017 in our model exploring effects before/after the policy start date (adjusted OR: 1.29; 95% CI: 1.03 to 1.62).

### ***Sensitivity analysis***

Our sensitivity analysis of step and trend changes between July 2016 and July 2017 (Table 2) was consistent with our two models for the general population of England with similar step changes observed at both the beginning and end of the policy implementation period, though only statistically significant after May 2016 (unadjusted OR: 0.93; 95% CI: 0.86 to 1.00; adjusted OR: 0.92; 95% CI: 0.85 to 0.99).

Subgroup results in our sensitivity analyses differed in that the only step decreases observed were among males after May 2016 in our model exploring effects after the policy start date (unadjusted OR: 0.89; 95% CI: 0.80 to 0.98), among population aged over 25 years old after May 2016 in our model exploring effects after full implementation (adjusted OR: 0.93; 95% CI: 0.87 to 0.99) and among manual occupations after May 2016 in our model exploring effects after the policy start date (adjusted OR: 0.85; 95% CI: 0.75 to 0.96), and in that there was a decreasing trend observed between July 2016 and July 2017 among non-manual occupations (adjusted OR 0.79; 95% CI: 0.64 to 0.97).

**Table 2 here**

***Model predictions***

Figure 2 shows predictions of smoking prevalence obtained from the unadjusted model, excluding autocorrelation and seasonality, allowing us to explore linear changes before and after the implementation of standardised packaging. The two models show that there was a step decrease in prevalence and no significant change in trends eighteen months after full implementation –in line with our regression results. Moreover, both the predictions from our models after full implementation and after the policy start date show no complete return to the pre-policy level of smoking prevalence, and a larger difference between the counterfactual trend and the predicted trend when the model accounts for the start of the implementation date (May 2016) in Figure 2b.

## DISCUSSION

To our knowledge, this is the first study evaluating the possible impact of standardised packaging on smoking prevalence in England. Our individual level findings found that the implementation of standardised packaging legislation was associated with a step decrease in the odds of being a smoker which was associated with the onset of standardised packaging after May 2016, when the transition to the new policy officially began, rather than May 2017 when the policy was fully implemented. Insofar that the association reflected a causal impact, the suggestion is that smokers were influenced more by the prospect of standardised packs, and possibly also of minimum pack sizes and other TPD measures, or of changes in the tobacco market introduced by the tobacco industry in advance of standardised packaging, than the actual adoption of standardised packaging. Within this study we were unable to investigate what the underlying reasons were though we hypothesize that smokers might have been aware of the policy due to media coverage and had consider what effect this specific policy might have on their smoking behaviour (costs, no brand loyalty, lack of appealing packaging). We found no strong evidence in our subgroup analyses that this effect differed markedly between the sexes, between older and younger smokers, or those of high or low occupational socioeconomic status.

The main limitation of our study was that we were only able to include policy implementation as May 2016 and May 2017 though the policy was gradually implemented over nine months of the one-year transition period [11], and our model did not account for that. The standardised packaging policy was actually implemented across the UK but the results of this study were based on individual level data from a large sample representative to population in England only. Therefore, generalisability of the findings to other UK countries and elsewhere in the world is limited and further research exploring differences in the UK countries and globally is warranted. To our knowledge studies similar to this have not been carried out elsewhere. Also, time series analysis using ARIMA models at the aggregated level would be more suitable for assessing the effect of policy on smoking prevalence though at the time of analysis the power was too low to produce such analysis. Therefore, we will use aggregated level data to evaluate longer term effects.

Although UK law did not require tobacco manufacturers to adopt standardised packaging simultaneously with minimum pack sizes, updated health warnings and other measures, in practice the changes were introduced simultaneously within individual cigarette brands. Consequently, we were unable to determine the effect of each of these policy elements

separately. However the findings of our study, which used individual level data and estimated odds of being a smokers in various population groups instead of measuring the effect of policy on aggregated prevalence data, are consistent with previous research [7-9] and provide further evidence that standardised packaging, or at least the prospect of standardised packaging, influences smoking status. In this study we were unable to determine whether the reduction in odds of smoking occurred due to changes in quitting or smoking uptake though detailed analysis of each of these aspects is necessary in further research.

A possible explanation for this is that prospect of standardised packs proved to be a stimulus for smokers intending to quit smoking to act on that intention, rather than a direct visual effect of the pack itself. We have previously described substantive changes in the diversity of products available on the market both in advance of and after standard packs appeared in the UK [11] and described substantial price increases with the adoption of standardised packaging [11, 13]. These included the introduction of, and widespread consumption of, low price cigarettes in packs of less than 20 in advance of the change to standardised packaging [11]. Whether any of these changes, introduced by the tobacco industry in preparation to standardised packs and the loss of distinctive branding and brand descriptors contributed to the step change in smoking prevalence when the law mandating the change to standardised packs came into force is not known and will be difficult to determine. The fact remains however that standardised packaging occurred in the UK in conjunction with a wide range of other legislative and market changes, so the precise contribution of standardised packs *per se* to the reduction in smoking prevalence we observed is impossible to determine.

In the long run, more pronounced effects on uptake of smoking and smoking cessation might be observed related to reduced appeal of packaging and more prominent health warnings. However, our analysis focusing on short to medium term effects provides clear evidence that these marked changes in packaging policy have had an appreciable beneficial effect on smoking prevalence in England.

**Acknowledgments:** Department of Behavioural Science and Health, University College London, London, UK.

**Funding:**

The work was supported by Cancer Research UK grant number C45256/A20606.

Cancer Research UK (C1417/A22962) supported the data collection and EB's salary. All authors are members of SPECTRUM, a UK Prevention Research Partnership Consortium (MR/S037519/1). UKPRP is an initiative funded by the UK Research and Innovation Councils, the Department of Health and Social Care (England) and the UK devolved administrations, and leading health research charities.

## References

1. Office for National Statistics, *Adult smoking habits in the UK: 2019*. 2020: Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/healthandlifeexpectancies/bulletins/adultsmokinghabitsingreatbritain/2019>.
2. NHS Digital, *Smoking, Drinking and Drug Use among Young People in England 2018 [NS]*. 2019: <https://digital.nhs.uk/data-and-information/publications/statistical/smoking-drinking-and-drug-use-among-young-people-in-england/2018>.
3. *The Standardised Packaging of Tobacco Products Regulations 2015*. Available from: <https://www.legislation.gov.uk/ukdsi/2015/9780111129876> (accessed 31 July 2018).
4. *Directive 2014/40/EU of the European Parliament and of the Council of 3 April 2014 on the approximation of the laws, regulations and administrative provisions of the Member States concerning the manufacture, presentation and sale of tobacco and related products and repealing Directive 2001/37/EC Text with EEA relevance*. Available from: <http://eur-lex.europa.eu/legal-content/en/TXT/?uri=CELEX%3A32014L0040> (accessed 31 March 2017). Archived at: <http://www.webcitation.org/6vOW4O45R>.
5. *Explanatory Memorandum to The Standardised Packaging of Tobacco Products Regulations 2015*. Available from: [https://www.legislation.gov.uk/ukdsi/2015/829/pdfs/uksem\\_20150829\\_en.pdf](https://www.legislation.gov.uk/ukdsi/2015/829/pdfs/uksem_20150829_en.pdf) (accessed 01 June 2021).
6. *Standardised packaging of tobacco. Report of the independent review undertaken by Sir Cyril Chantler*. April 2014. Available from: <https://www.kcl.ac.uk/health/10035-TSO-2901853-Chantler-Review-ACCESSIBLE.PDF> (accessed 30 Sept 2017). Archived at: <http://www.webcitation.org/6vOXktOpa>.
7. McNeill, A., et al., *Tobacco packaging design for reducing tobacco use*. Cochrane Database Syst Rev, 2017. **4**: p. CD011244.
8. Young, J.M., et al., *Association between tobacco plain packaging and Quitline calls: a population-based, interrupted time-series analysis*. Med J Aust, 2014. **200**(1): p. 29-32.
9. Chipty, T., *Study of the Impact of the Tobacco Plain Packaging Measure on Smoking Prevalence in Australia*. 2016: <https://www.health.gov.au/sites/default/files/study-of-the-impact-of-the-tobacco-plain-packaging-measure-on-smoking-prevalence-in-australia.pdf> [Accessed 28 May 2020].
10. Hiscock, R., et al., *Standardised packaging, minimum excise tax, and RYO focussed tax rise implications for UK tobacco pricing*. PLOS ONE, 2020. **15**(2): p. e0228069.
11. Opazo Breton, M., et al., *Cigarette brand diversity and price changes during the implementation of plain packaging in the United Kingdom*. Addiction, 2018. **113**(10): p. 1883-1894.
12. Opazo Breton, M., J. Britton, and I. Bogdanovica, *Effect of UK plain tobacco packaging and minimum pack size legislation on tobacco and nicotine product switching behaviour*. Addiction, 2020. **115**(10): p. 1913-1923.
13. Opazo Breton, M., J. Britton, and I. Bogdanovica, *Changes in roll-your-own tobacco and cigarette sales volume and prices before, during and after plain packaging legislation in the UK*. Tobacco Control, 2019: p. tobaccocontrol-2018-054734.
14. Tobacco Manufacturers' Association. *Plain packaging failing one year after full introduction*. 2018 [10 November 2020]; Available from: <http://the-tma.org.uk/2018/05/14/plain-packaging-failing-one-year-after-full-introduction/>.
15. Fidler, J.A., et al., *'The smoking toolkit study': a national study of smoking and smoking cessation in England*. BMC Public Health, 2011. **11**(1): p. 479.
16. Beard, E., et al., *Understanding and using time series analyses in addiction research*. Addiction, 2019. **114**(10): p. 1866-1884.



17. Beard, E., et al., *Trends in Attempts to Quit Smoking in England Since 2007: A Time Series Analysis of a Range of Population-Level Influences*. Nicotine & Tobacco Research, 2019.
18. Beard, E., et al., *Association of prevalence of electronic cigarette use with smoking cessation and cigarette consumption in England: a time-series analysis between 2006 and 2017*. Addiction, 2019.
19. Beard, E., et al., *Association between electronic cigarette use and changes in quit attempts, success of quit attempts, use of smoking cessation pharmacotherapy, and use of stop smoking services in England: time series analysis of population trends*. BMJ, 2016. **354**: p. i4645.
20. Kuipers, M.A.G., et al., *Impact on smoking of England's 2012 partial tobacco point of sale display ban: a repeated cross-sectional national study*. Tobacco Control, 2017. **26**(2): p. 141.
21. Hallingberg, B., et al., *Have e-cigarettes renormalised or displaced youth smoking? Results of a segmented regression analysis of repeated cross sectional survey data in England, Scotland and Wales*. Tobacco Control, 2020. **29**(2): p. 207.
22. Moore, G., et al., *Young people's use of e-cigarettes in Wales, England and Scotland before and after introduction of EU Tobacco Products Directive regulations: a mixed-method natural experimental evaluation*. International Journal of Drug Policy, 2020. **85**: p. 102795.
23. Bernal, J.L., S. Cummins, and A. Gasparrini, *Interrupted time series regression for the evaluation of public health interventions: a tutorial*. International Journal of Epidemiology, 2016. **46**(1): p. 348-355.
24. He, Y., C. Shang, and F.J. Chaloupka, *The association between cigarette affordability and consumption: An update*. PLOS ONE, 2018. **13**(12): p. e0200665.
25. Kostova, D., F. Chaloupka, and C. Shang, *A duration analysis of the role of cigarette prices on smoking initiation and cessation in developing countries*. The European Journal of Health Economics, 2014: p. 1-10.
26. Ross, H. and F.J. Chaloupka, *The Effect of Public Policies and Prices on Youth Smoking*. Southern Economic Journal, 2004. **70**(4): p. 796-815.
27. Chaloupka, F.J. and K.E. Warner, *The economics of smoking*. Handbook of health economics, 2000. **1**: p. 1539-1627.
28. Chaloupka, F.J. and H. Wechsler, *Price, tobacco control policies and smoking among young adults*. Journal of health economics, 1997. **16**(3): p. 359-373.
29. Office for National Statistics, *RPI: Ave price - Cigarettes 20 king size filter*, in *Monthly*. 2020: <https://www.ons.gov.uk/economy/inflationandpriceindices/timeseries/czmp> [Accessed on April 8, 2020].
30. Office for National Statistics, *Consumer price inflation time series*. 2020: <https://www.ons.gov.uk/economy/inflationandpriceindices/datasets/consumerpriceindices> [Accessed April 8, 2020].
31. Chandra, S. and F.J. Chaloupka, *Seasonality in cigarette sales: patterns and implications for tobacco control*. Tobacco Control, 2003. **12**(1): p. 105.
32. Wellman, R.J. and J.R. DiFranza, *Seasonality in onset of youth smoking parallels seasonality in cigarette sales*. Tobacco Control, 2003. **12**(3): p. 339.
33. Beck, N. and J.N. Katz, *Nuisance vs. Substance: Specifying and Estimating Time-Series-Cross-Section Models*. Political Analysis, 1996. **6**: p. 1-36.

Table 1: Regression results for level and slope changes in the odds of being a smoker after full implementation of standardised packaging in **May 2017** (before/after full implementation) and accounting for standardised packaging start date in **May 2016** (before/after policy start date) in England (Smoking Toolkit Study data January 2007 to December 2019)

	Before/after full implementation		Before/after policy start date	
	Unadjusted	Adjusted	Unadjusted	Adjusted
	OR (p-value) [95% CI]	OR (p-value) [95% CI]	OR (p-value) [95% CI]	OR (p-value) [95% CI]
a. General population				
Level after May 2016			<b>0.91</b>	<b>0.90</b>
			(0.018)	(0.009)
			0.85 - 0.98	0.83 - 0.97
Slope May 2016-May 2017			1.02	0.96
			(0.728)	(0.578)
			0.90 - 1.17	0.84 - 1.10
Level after May 2017	0.95	<b>0.93</b>	0.99	1.00
	(0.073)	(0.015)	(0.813)	(0.975)
	0.90 - 1.00	0.87 - 0.99	0.90 - 1.09	0.91 - 1.10
Post-slope 18 months after May 2017	1.02	1.00	1.02	1.00
	(0.474)	(0.892)	(0.591)	(0.940)
	0.96 - 1.10	0.94 - 1.08	0.95 - 1.09	0.93 - 1.07
Observations	276,416	276,416	276,416	276,416
b. Females only subgroup				
Level after May 2016			0.95	<b>0.89</b>
			(0.342)	(0.046)
			0.85 - 1.06	0.79 - 1.00
Slope May 2016-May 2017			1.08	1.08
			(0.438)	(0.430)
			0.89 - 1.30	0.89 - 1.32
Level after May 2017	0.99	0.97	0.96	0.97
	(0.802)	(0.480)	(0.572)	(0.617)
	0.91 - 1.07	0.89 - 1.06	0.84 - 1.10	0.84 - 1.11
Post-slope 18 months after May 2017	0.98	0.97	0.98	0.96
	(0.640)	(0.522)	(0.636)	(0.459)
	0.89 - 1.08	0.87 - 1.07	0.89 - 1.08	0.87 - 1.07
Observations	142,107	142,107	142,107	142,107
c. Males only subgroup				
Level after May 2016			<b>0.89</b>	0.91
			(0.022)	(0.083)
			0.80 - 0.98	0.81 - 1.01
Slope May 2016-May 2017			1.00	0.88
			(0.969)	(0.204)
			0.83 - 1.20	0.73 - 1.07
Level after May 2017	<b>0.92</b>	<b>0.90</b>	1.01	1.03
	(0.046)	(0.013)	(0.932)	(0.706)
	0.85 - 1.00	0.83 - 0.98	0.88 - 1.15	0.90 - 1.18
Post-slope 18 months after May 2017	1.07	1.04	1.05	1.03
	(0.176)	(0.444)	(0.262)	(0.556)
	0.97 - 1.17	0.94 - 1.15	0.96 - 1.16	0.93 - 1.14
Observations	134,254	134,254	134,254	134,254
d. Population aged 18 to 25 years old only subgroup				
Level after May 2016			1.02	0.95
			(0.842)	(0.541)
			0.86 - 1.21	0.79 - 1.13

Slope May 2016-May 2017			0.87	0.84
			(0.377)	(0.277)
			0.64 - 1.18	0.61 - 1.15
Level after May 2017	1.12	1.10	1.25	<b>1.29</b>
	(0.078)	(0.183)	(0.052)	(0.028)
	0.99 - 1.28	0.96 - 1.27	1.00 - 1.57	1.03 - 1.62
Post-slope 18 months after May 2017	0.87	0.87	0.87	0.86
	(0.088)	(0.088)	(0.074)	(0.068)
	0.75 - 1.02	0.73 - 1.02	0.74 - 1.01	0.72 - 1.01
Observations	43,729	43,729	43,729	43,729
e. Population over 25 years old only subgroup				
Level after May 2016			<b>0.90</b>	<b>0.90</b>
			(0.014)	(0.016)
			0.83 - 0.98	0.82 - 0.98
Slope May 2016-May 2017			1.07	1.00
			(0.386)	(0.976)
			0.92 - 1.23	0.86 - 1.17
Level after May 2017	<b>0.92</b>	<b>0.90</b>	0.94	0.94
	(0.013)	(0.002)	(0.228)	(0.283)
	0.87 - 0.98	0.84 - 0.96	0.84 - 1.04	0.85 - 1.05
Post-slope 18 months after May 2017	1.06	1.04	1.06	1.04
	(0.109)	(0.304)	(0.147)	(0.382)
	0.99 - 1.15	0.96 - 1.13	0.98 - 1.14	0.96 - 1.12
Observations	232,687	232,687	232,687	232,687
f. Routine and manual occupations only subgroup				
Level after May 2016			0.89	<b>0.83</b>
			(0.065)	(0.004)
			0.79 - 1.01	0.73 - 0.94
Slope May 2016-May 2017			1.08	1.11
			(0.499)	(0.361)
			0.87 - 1.33	0.89 - 1.38
Level after May 2017	1.07	1.07	1.08	1.09
	(0.149)	(0.171)	(0.315)	(0.289)
	0.98 - 1.17	0.97 - 1.18	0.93 - 1.27	0.93 - 1.27
Post-slope 18 months after May 2017	1.02	1.00	1.01	1.00
	(0.721)	(0.939)	(0.798)	(0.948)
	0.91 - 1.14	0.89 - 1.13	0.91 - 1.13	0.89 - 1.12
Observations	95,770	95,770	95,770	95,770
g. Non-routine occupations only subgroup				
Level after May 2016			1.03	1.00
			(0.578)	(0.961)
			0.92 - 1.16	0.89 - 1.13
Slope May 2016-May 2017			0.99	0.93
			(0.942)	(0.457)
			0.82 - 1.21	0.75 - 1.13
Level after May 2017	0.98	0.94	0.97	0.99
	(0.669)	(0.149)	(0.666)	(0.846)
	0.90 - 1.07	0.85 - 1.02	0.84 - 1.12	0.85 - 1.14
Post-slope 18 months after May 2017	1.00	0.96	1.00	0.96
	(0.999)	(0.453)	(0.973)	(0.435)
	0.91 - 1.10	0.86 - 1.07	0.91 - 1.11	0.86 - 1.07
Observations	141,844	141,844	141,844	141,844

Note: Values in **bold** refer to statistically significant OR (p-value < 0.05). Unadjusted model includes adjustment by seasonality and serial correlation, while the adjusted model includes other tobacco control policies implemented during the period studied and monthly average real retail price, in addition to seasonality and serial correlation.

Table 2: Regression results for level and slope changes in the odds of being a smoker after full implementation of standardised packaging using **July 2017** (before/after full implementation) and accounting for standardised packaging start date using **July 2016** (before/after policy start date) in England (Smoking Toolkit Study data January 2007 to December 2019)

	Before/after full implementation		Before/after policy start date	
	Unadjusted	Adjusted	Unadjusted	Adjusted
	OR (p-value) [95% CI]	OR (p-value) [95% CI]	OR (p-value) [95% CI]	OR (p-value) [95% CI]
a. General Population				
Level after July 2016			<b>0.93</b>	<b>0.92</b>
			(0.042)	(0.027)
			0.86 - 1.00	0.85 - 0.99
Slope July 2016-July 2017			1.01	0.95
			(0.878)	(0.445)
			0.89 - 1.15	0.82 - 1.09
Level after July 2017	0.96	0.94	1.00	1.02
	(0.165)	(0.054)	(0.956)	(0.664)
	0.91 - 1.02	0.89 - 1.00	0.91 - 1.10	0.93 - 1.13
Post-slope 18 months after July 2017	1.01	0.99	1.01	0.99
	(0.679)	(0.836)	(0.793)	(0.738)
	0.95 - 1.09	0.92 - 1.07	0.94 - 1.08	0.92 - 1.06
Observations	276,416	276,416	276,416	276,416
b. Females only subgroup				
Level after July 2016			0.98	0.93
			(0.768)	(0.239)
			0.89 - 1.09	0.83 - 1.05
Slope July 2016-July 2017			1.05	1.06
			(0.645)	(0.598)
			0.87 - 1.26	0.86 - 1.29
Level after July 2017	0.97	0.95	0.95	0.94
	(0.511)	(0.242)	(0.443)	(0.408)
	0.90 - 1.06	0.87 - 1.04	0.82 - 1.09	0.82 - 1.08
Post-slope 18 months after July 2017	0.99	0.98	0.99	0.98
	(0.854)	(0.748)	(0.866)	(0.719)
	0.90 - 1.09	0.88 - 1.09	0.90 - 1.10	0.88 - 1.09
Observations	142,107	142,107	142,107	142,107
c. Males only subgroup				
Level after July 2016			<b>0.89</b>	0.90
			(0.020)	(0.070)
			0.80 - 0.98	0.81 - 1.01
Slope July 2016-July 2017			0.99	0.87
			(0.917)	(0.152)
			0.82 - 1.19	0.71 - 1.05
Level after July 2017	0.96	0.94	1.05	1.10
	(0.264)	(0.155)	(0.490)	(0.197)
	0.88 - 1.03	0.87 - 1.02	0.92 - 1.20	0.95 - 1.26
Post-slope 18 months after July 2017	1.04	1.00	1.02	0.99
	(0.479)	(0.957)	(0.628)	(0.903)
	0.94 - 1.14	0.91 - 1.11	0.93 - 1.13	0.90 - 1.10
Observations	134,254	134,254	134,254	134,254
d. Population aged 18 to 25 years old only subgroup				
Level after July 2016			0.93	0.85
			(0.393)	(0.084)
			0.78 - 1.10	0.71 - 1.02

Slope July 2016-July 2017			1.10	1.11
			(0.553)	(0.533)
			0.81 - 1.50	0.80 - 1.54
Level after July 2017	1.09	1.06	1.06	1.06
	(0.211)	(0.441)	(0.639)	(0.614)
	0.95 - 1.24	0.92 - 1.21	0.84 - 1.32	0.85 - 1.33
Post-slope 18 months after July 2017	0.89	0.89	0.89	0.88
	(0.169)	(0.171)	(0.162)	(0.140)
	0.76 - 1.05	0.75 - 1.05	0.76 - 1.05	0.74 - 1.04
Observations	43,729	43,729	43,729	43,729
e. Population over 25 years old only subgroup				
Level after July 2016			0.94	0.94
			(0.124)	(0.174)
			0.86 - 1.02	0.86 - 1.03
Slope July 2016-July 2017			0.99	0.92
			(0.919)	(0.277)
			0.86 - 1.15	0.79 - 1.07
Level after July 2017	0.94	<b>0.93</b>	0.99	1.01
	(0.071)	(0.025)	(0.886)	(0.809)
	0.89 - 1.00	0.87 - 0.99	0.89 - 1.10	0.91 - 1.13
Post-slope 18 months after July 2017	1.04	1.02	1.04	1.02
	(0.269)	(0.630)	(0.338)	(0.687)
	0.97 - 1.13	0.94 - 1.11	0.96 - 1.12	0.94 - 1.10
Observations	232,687	232,687	232,687	232,687
f. Routine and manual occupations only subgroup				
Level after July 2016			0.91	<b>0.85</b>
			(0.106)	(0.011)
			0.80 - 1.02	0.75 - 0.96
Slope July 2016-July 2017			1.14	1.19
			(0.243)	(0.129)
			0.92 - 1.41	0.95 - 1.49
Level after July 2017	1.07	1.07	1.03	1.02
	(0.146)	(0.185)	(0.747)	(0.818)
	0.98 - 1.17	0.97 - 1.18	0.88 - 1.20	0.87 - 1.19
Post-slope 18 months after July 2017	1.01	1.00	1.01	0.99
	(0.832)	(0.953)	(0.854)	(0.887)
	0.90 - 1.13	0.89 - 1.12	0.90 - 1.13	0.88 - 1.12
Observations	95,770	95,770	95,770	95,770
g. Non-routine occupations only subgroup				
Level after July 2016			1.09	1.07
			(0.147)	(0.265)
			0.97 - 1.22	0.95 - 1.21
Slope July 2016-July 2017			0.87	<b>0.79</b>
			(0.159)	(0.026)
			0.71 - 1.06	0.64 - 0.97
Level after July 2017	1.01	0.97	1.08	1.12
	(0.783)	(0.507)	(0.281)	(0.140)
	0.93 - 1.10	0.89 - 1.06	0.94 - 1.25	0.96 - 1.29
Post-slope 18 months after July 2017	0.97	0.93	0.97	0.93
	(0.556)	(0.187)	(0.534)	(0.178)
	0.88 - 1.07	0.83 - 1.04	0.87 - 1.07	0.83 - 1.03
Observations	141,844	141,844	141,844	141,844

Note: Values in **bold** refer to statistically significant OR (p-value < 0.05). Unadjusted model includes adjustment by seasonality and serial correlation, while the adjusted model includes other tobacco control policies implemented during the period studied and monthly average real retail price, in addition to seasonality and serial correlation.

Figure 1: Monthly smoking prevalence in England, general population, and by sex (males/females), age (population age 18-25 years old/ population age over 25 years old) and socioeconomic status (routine and manual occupations/ non-routine occupations), STS data (January 2007-December 2019)





Figure 2: Model predictions on smoking prevalence for general population without adjustments, STS data (January 2007-December 2019)



### Supplementary material

Figure S1: Percentage of current regular cigarette smokers in England (2007-2019) using Smoking Toolkit Study (STS) and Annual Population Survey (APS) quarterly data (2007q1-2019q4).





Figure S2: Percentage of smokers by type of tobacco product used: manufactured cigarettes (MF), hand-rolled tobacco (HRT), other (including pipe, cigar, cigarillos and other) and all tobacco products (Smoking Toolkit Study data November 2007\* to December 2019)



\* The variable 'type of cigarette smoked' was only available from November 2007. This implies that the smoke-free in public spaces legislation or the increase in minimum age of sale.



Table S1: Regression results for level and slope changes in the odds of being a smoker after full implementation of plain packaging in **May 2017** (before/after full implementation) and accounting for plain packaging start date in May 2016 (before/after policy start date) in England (Annual Population Survey, 2010q1 to 2019q4)

	Primary analysis		Secondary analysis	
	Unadjusted	Adjusted	Unadjusted	Adjusted
	OR (p-value) [95% CI]	OR (p-value) [95% CI]	OR (p-value) [95% CI]	OR (p-value) [95% CI]
a. General population				
Level after May 2016			<b>0.96</b>	<b>0.97</b>
			(0.007)	(0.036)
			0.93 - 0.99	0.94 - 1.00
Slope May 2016-May 2017			1.02	1.00
			(0.563)	(0.935)
			0.96 - 1.07	0.94 - 1.06
Level after May 2017	0.98	0.98	0.98	0.97
	(0.051)	(0.125)	(0.231)	(0.213)
	0.96 - 1.00	0.95 - 1.01	0.94 - 1.02	0.93 - 1.02
Post-slope 18 months after May 2017	1.02	1.01	1.02	1.01
	(0.125)	(0.500)	(0.090)	(0.586)
	0.99 - 1.05	0.98 - 1.04	1.00 - 1.05	0.98 - 1.04
Observations	1,555,452	1,555,452	1,555,452	1,555,452
b. Females only subgroup				
Level after May 2016			<b>0.95</b>	0.96
			(0.015)	(0.076)
			0.92 - 0.99	0.92 - 1.00
Slope May 2016-May 2017			1.01	1.01
			(0.831)	(0.841)
			0.94 - 1.09	0.93 - 1.09
Level after May 2017	<b>0.97</b>	0.99	0.98	0.98
	(0.043)	(0.617)	(0.410)	(0.539)
	0.94 - 1.00	0.95 - 1.03	0.92 - 1.03	0.92 - 1.04
Post-slope 18 months after May 2017	1.03	1.02	1.03	1.01
	(0.187)	(0.477)	(0.186)	(0.549)
	0.99 - 1.06	0.97 - 1.06	0.99 - 1.06	0.97 - 1.06
Observations	818,331	818,331	818,331	818,331
c. Males only subgroup				
Level after May 2016			<b>0.95</b>	0.96
			(0.030)	(0.062)
			0.91 - 1.00	0.92 - 1.00
Slope May 2016-May 2017			1.02	0.98
			(0.622)	(0.638)
			0.95 - 1.10	0.91 - 1.06
Level after May 2017	0.98	<b>0.96</b>	0.97	0.96
	(0.111)	(0.047)	(0.329)	(0.203)
	0.95 - 1.01	0.92 - 1.00	0.92 - 1.03	0.91 - 1.02
Post-slope 18 months after May 2017	1.03	1.02	1.03	1.01
	(0.128)	(0.405)	(0.108)	(0.586)
	0.99 - 1.06	0.98 - 1.06	0.99 - 1.07	0.97 - 1.05
Observations	737,121	737,121	737,121	737,121
d. Population aged 18 to 25 years old only subgroup				
Level after May 2016			0.93	0.97
			(0.100)	(0.427)
			0.86 - 1.01	0.89 - 1.05

Slope May 2016-May 2017			0.95	0.92
			(0.546)	(0.325)
			0.82 - 1.11	0.78 - 1.09
Level after May 2017	0.98	0.99	1.03	1.03
	(0.529)	(0.818)	(0.602)	(0.634)
	0.92 - 1.05	0.90 - 1.08	0.92 - 1.16	0.91 - 1.16
Post-slope 18 months after May 2017	1.01	0.99	1.00	0.98
	(0.868)	(0.885)	(0.963)	(0.587)
	0.93 - 1.09	0.91 - 1.08	0.93 - 1.08	0.89 - 1.07
Observations	154,924	154,924	154,924	154,924
<b>e. Population over 25 years old only subgroup</b>				
Level after May 2016			<b>0.96</b>	<b>0.97</b>
			(0.012)	(0.041)
			0.93 - 0.99	0.94 - 1.00
Slope May 2016-May 2017			1.03	1.01
			(0.366)	(0.654)
			0.97 - 1.08	0.95 - 1.08
Level after May 2017	<b>0.97</b>	0.98	0.97	0.97
	(0.036)	(0.152)	(0.122)	(0.126)
	0.95 - 1.00	0.94 - 1.01	0.93 - 1.01	0.92 - 1.01
Post-slope 18 months after May 2017	1.02	1.01	1.03	1.01
	(0.088)	(0.358)	(0.070)	(0.377)
	1.00 - 1.05	0.98 - 1.05	1.00 - 1.05	0.98 - 1.05
Observations	1,400,528	1,400,528	1,400,528	1,400,528
<b>f. Manual and routine occupations only subgroup</b>				
Level after May 2016			0.97	0.98
			(0.153)	(0.470)
			0.92 - 1.01	0.93 - 1.03
Slope May 2016-May 2017			1.04	1.02
			(0.382)	(0.730)
			0.95 - 1.14	0.92 - 1.12
Level after May 2017	0.97	0.97	0.96	0.96
	(0.179)	(0.217)	(0.221)	(0.249)
	0.94 - 1.01	0.92 - 1.02	0.89 - 1.03	0.89 - 1.03
Post-slope 18 months after May 2017	1.02	0.99	1.03	0.99
	(0.287)	(0.819)	(0.273)	(0.834)
	0.98 - 1.07	0.94 - 1.05	0.98 - 1.07	0.94 - 1.05
Observations	368,993	368,993	368,993	368,993
<b>g. Non-routine occupations only subgroup</b>				
Level after May 2016			0.96	0.96
			(0.112)	(0.079)
			0.92 - 1.01	0.91 - 1.01
Slope May 2016-May 2017			1.02	0.99
			(0.589)	(0.880)
			0.94 - 1.11	0.91 - 1.09
Level after May 2017	0.99	0.98	0.99	0.98
	(0.786)	(0.442)	(0.652)	(0.491)
	0.96 - 1.03	0.93 - 1.03	0.92 - 1.05	0.91 - 1.04
Post-slope 18 months after May 2017	<b>1.06</b>	1.04	<b>1.05</b>	1.03
	(0.007)	(0.075)	(0.008)	(0.146)
	1.02 - 1.10	1.00 - 1.09	1.01 - 1.10	0.99 - 1.08
Observations	735,612	735,612	735,612	735,612

Note: Values in **bold** refer to statistically significant OR (p-value < 0.05). Unadjusted model includes adjustment by seasonality and serial correlation, while the adjusted model includes other tobacco control policies implemented during the period studied and monthly average real retail price, in addition to seasonality and serial correlation.

Table S2: Regression results for level and slope changes in the odds of being a smoker after full implementation of plain packaging in **May 2017** (before/after full implementation) and accounting for plain packaging start date in May 2016 (before/after policy start date) in England by type of smoking product (Smoking Toolkit Study data November 2007\* to December 2019)

	Primary analysis		Secondary analysis	
	Unadjusted	Adjusted	Unadjusted	Adjusted
	OR (p-value) [95% CI]	OR (p-value) [95% CI]	OR (p-value) [95% CI]	OR (p-value) [95% CI]
<b>h. Smokers of manufactured cigarettes</b>				
Level after May 2016			<b>0.73</b>	<b>0.83</b>
			(0.000)	(0.001)
			0.65 - 0.81	0.74 - 0.93
Slope May 2016-May 2017			1.04	1.18
			(0.683)	(0.088)
			0.86 - 1.26	0.98 - 1.43
Level after May 2017	<b>0.81</b>	0.92	0.97	0.89
	(0.000)	(0.067)	(0.652)	(0.104)
	0.75 - 0.88	0.85 - 1.01	0.84 - 1.11	0.78 - 1.02
Post-slope 18 months after May 2017	0.94	<b>1.13</b>	0.92	<b>1.12</b>
	(0.209)	(0.019)	(0.074)	(0.028)
	0.86 - 1.03	1.02 - 1.25	0.84 - 1.01	1.01 - 1.24
Observations	240,251	225,270	240,251	225,270
<b>i. Smokers of hand-rolled tobacco</b>				
Level after May 2016			<b>0.68</b>	0.82
			(0.009)	(0.193)
			0.51 - 0.91	0.60 - 1.11
Slope May 2016-May 2017			0.98	1.12
			(0.929)	(0.681)
			0.57 - 1.67	0.66 - 1.89
Level after May 2017	<b>0.73</b>	0.87	0.97	0.89
	(0.005)	(0.268)	(0.889)	(0.546)
	0.59 - 0.91	0.68 - 1.11	0.66 - 1.44	0.61 - 1.30
Post-slope 18 months after May 2017	0.96	1.22	0.93	1.21
	(0.747)	(0.185)	(0.581)	(0.205)
	0.73 - 1.25	0.91 - 1.63	0.71 - 1.21	0.90 - 1.61
Observations	221,890	206,909	221,890	206,909
<b>a. Smokers of manufactured cigarettes and hand-rolled tobacco</b>				
Level after May 2016			<b>0.68</b>	0.82
			(0.009)	(0.193)
			0.51 - 0.91	0.60 - 1.11
Slope May 2016-May 2017			0.98	1.12
			(0.929)	(0.681)
			0.57 - 1.67	0.66 - 1.89
Level after May 2017	<b>0.73</b>	0.87	0.97	0.89
	(0.005)	(0.268)	(0.889)	(0.546)
	0.59 - 0.91	0.68 - 1.11	0.66 - 1.44	0.61 - 1.30
Post-slope 18 months after May 2017	0.96	1.22	0.93	1.21
	(0.747)	(0.185)	(0.581)	(0.205)
	0.73 - 1.25	0.91 - 1.63	0.71 - 1.21	0.90 - 1.61
Observations	221,890	206,909	221,890	206,909
<b>h. Smokers of other tobacco products (including pipe, cigar, cigarillos, etc.)</b>				
Level after May 2016			<b>6.47</b>	<b>1.36</b>
			(0.000)	(0.020)

			5.08 - 8.24	1.05 - 1.77
Slope May 2016-May 2017			1.28	<b>0.59</b>
			(0.212)	(0.031)
			0.87 - 1.87	0.37 - 0.95
Level after May 2017	<b>3.29</b>	1.00	0.72	1.27
	(0.000)	(0.978)	(0.050)	(0.175)
	2.56 - 4.23	0.80 - 1.25	0.51 - 1.00	0.90 - 1.79
Post-slope 18 months after May 2017	<b>2.65</b>	0.82	<b>2.95</b>	0.83
	(0.000)	(0.122)	(0.000)	(0.143)
	1.98 - 3.54	0.63 - 1.06	2.21 - 3.94	0.64 - 1.07
Observations	226,983	226,983	226,983	226,983

\* The variable 'type of cigarette smoked' was only available from November 2007. This implies that the smoke-free in public spaces legislation or the increase in minimum age of sale.

Table S3: Regression results step and trend changes in the odds of being a smoker after full implementation of plain packaging in May 2017 (before/after full implementation) and accounting for plain packaging start date in May 2016 (before/after policy start date) in England using interaction effect (Smoking Toolkit Study data January 2007 to December 2019)

	After full implementation		After policy start date	
	Unadjusted	Adjusted	Unadjusted	Adjusted
	OR (p-value) [95% CI]	OR (p-value) [95% CI]	OR (p-value) [95% CI]	OR (p-value) [95% CI]
a. Interaction with female gender (reference: male)				
Step change after May 2016			0.91	<b>0.89</b>
			(0.056)	(0.031)
			0.82 - 1.00	0.80 - 0.99
Female's step change after May 2016			1.02	1.02
			(0.758)	(0.762)
			0.89 - 1.17	0.89 - 1.17
Trend May 2016-May 2017			0.98	0.92
			(0.806)	(0.358)
			0.81 - 1.17	0.76 - 1.10
Female's trend May 2016-May 2017			1.00 - 1.00	1.00 - 1.00
			1.11	1.11
			(0.435)	(0.431)
Step change after May 2017	0.93	<b>0.91</b>	1.02	1.04
	(0.068)	(0.017)	(0.740)	(0.594)
	0.86 - 1.01	0.84 - 0.98	0.90 - 1.17	0.91 - 1.19
Female's step change after May 2017	1.05	1.05	0.93	0.93
	(0.358)	(0.358)	(0.468)	(0.463)
	0.95 - 1.17	0.95 - 1.17	0.77 - 1.12	0.77 - 1.12
Trend 18 months after May 2017	1.07	1.05	1.06	1.04
	(0.175)	(0.356)	(0.218)	(0.443)
	0.97 - 1.17	0.95 - 1.15	0.97 - 1.16	0.94 - 1.14
Female's trend 18 months after May 2017	0.92	0.92	0.92	0.92
	(0.224)	(0.226)	(0.224)	(0.227)
	0.81 - 1.05	0.81 - 1.05	0.81 - 1.05	0.81 - 1.05
Observations	276,361	276,361	276,361	276,361
b. Interaction with population aged 18 to 25 years old (reference: population > 25 years old)				
Step change after May 2016			0.93	<b>0.91</b>
			(0.075)	(0.033)
			0.86 - 1.01	0.84 - 0.99
18-25's step change after May 2016			0.97	0.97
			(0.714)	(0.708)

			0.81 - 1.15	0.81 - 1.15
Trend			1.04	0.98
May 2016-May 2017			(0.557)	(0.804)
			0.90 - 1.21	0.84 - 1.14
18-25's trend			0.93	0.93
May 2016-May 2017			(0.641)	(0.649)
			0.67 - 1.28	0.67 - 1.28
Step change after	0.94	<b>0.92</b>	0.95	0.96
May 2017	(0.066)	(0.010)	(0.366)	(0.510)
	0.89 - 1.00	0.86 - 0.98	0.86 - 1.06	0.87 - 1.07
18-25's step change	1.08	1.08	1.20	1.20
after May 2017	(0.238)	(0.238)	(0.128)	(0.131)
	0.95 - 1.24	0.95 - 1.24	0.95 - 1.53	0.95 - 1.53
Trend 18 months	1.07*	1.05	1.06	1.04
after May 2017	(0.099)	(0.266)	(0.129)	(0.350)
	0.99 - 1.15	0.97 - 1.13	0.98 - 1.14	0.96 - 1.12
18-25's trend 18	<b>0.82</b>	<b>0.82</b>	<b>0.82</b>	<b>0.82</b>
months after May	(0.024)	(0.024)	(0.024)	(0.024)
2017	0.69 - 0.97	0.70 - 0.97	0.69 - 0.97	0.70 - 0.97
Observations	276,416	276,416	276,416	276,416
c. Interaction with manual and routine occupations (reference: non-routine occupations)				
Step change after			1.04	0.98
May 2016			(0.503)	(0.743)
			0.93 - 1.15	0.88 - 1.10
Manual's step change			0.86	0.86
after May 2016			(0.051)	(0.052)
			0.74 - 1.00	0.74 - 1.00
Trend			0.99	0.96
May 2016-May 2017			(0.881)	(0.716)
			0.81 - 1.19	0.79 - 1.17
Manual's trend			1.10	1.10
May 2016-May 2017			(0.492)	(0.496)
			0.83 - 1.46	0.83 - 1.46
Step change after	1.00	0.97	0.97	0.99
May 2017	(0.969)	(0.543)	(0.713)	(0.855)
	0.92 - 1.08	0.90 - 1.06	0.85 - 1.12	0.86 - 1.14
Manual's step change	1.06	1.06	1.11	1.10
after May 2017	(0.344)	(0.349)	(0.336)	(0.339)
	0.94 - 1.18	0.94 - 1.18	0.90 - 1.36	0.90 - 1.36
Trend 18 months	1.00	0.97	1.00	0.97
after May 2017	(0.955)	(0.573)	(0.937)	(0.495)
	0.90 - 1.10	0.88 - 1.08	0.90 - 1.10	0.87 - 1.07
Manual's trend 18	1.02	1.02	1.02	1.02
months after May	(0.771)	(0.772)	(0.771)	(0.771)
2017	0.88 - 1.18	0.88 - 1.18	0.88 - 1.18	0.88 - 1.18
Observations	237,614	237,614	237,614	237,614

Note: Values in **bold** refer to statistically significant OR (p-value < 0.05). Unadjusted model includes adjustment by seasonality and serial correlation, while the adjusted model includes other tobacco control policies implemented during the period studied and monthly average real retail price, in addition to seasonality and serial correlation.

Table S4: General population regression results for step and trend changes in the odds of being a smoker after full implementation of plain packaging in May 2017 (before/after full implementation) and accounting for plain packaging start date in May 2016 (before/after policy start date) in England (Smoking Toolkit Study data January 2007 to December 2019)

	After full implementation		After policy start date	
	Unadjusted	Adjusted	Unadjusted	Adjusted
	OR (p-value) [95% CI]	OR (p-value) [95% CI]	OR (p-value) [95% CI]	OR (p-value) [95% CI]
General population				
Level after May 2016			<b>0.91</b>	<b>0.90</b>
			(0.018)	(0.009)
			0.85 - 0.98	0.83 - 0.97
Slope May 2016-May 2017			1.02	0.96
			(0.728)	(0.578)
			0.90 - 1.17	0.84 - 1.10
Level after May 2017	0.95	<b>0.93</b>	0.99	1.00
	(0.073)	(0.015)	(0.813)	(0.975)
	0.90 - 1.00	0.87 - 0.99	0.90 - 1.09	0.91 - 1.10
Post-slope 18 months after May 2017	1.02	1.00	1.02	1.00
	(0.474)	(0.892)	(0.591)	(0.940)
	0.96 - 1.10	0.94 - 1.08	0.95 - 1.09	0.93 - 1.07
Smoke free ban		1.08		1.07
		(0.056)		(0.062)
		1.00 - 1.16		1.00 - 1.16
Minimum age of sale		<b>0.84</b>		<b>0.82</b>
		(0.000)		(0.000)
		0.78 - 0.90		0.76 - 0.88
Display ban in small shops		0.98		0.97
		(0.407)		(0.205)
		0.93 - 1.03		0.92 - 1.02
Display ban in large shops		0.98		1.03
		(0.372)		(0.242)
		0.93 - 1.03		0.98 - 1.09
Price per cigarette		1.00		<b>1.00</b>
		(0.090)		(0.031)
		1.00 - 1.00		1.00 - 1.00
Lagged smoking prevalence	Yes	Yes	Yes	Yes
Seasonality	Yes	Yes	Yes	Yes
Observations	276,416	276,416	276,416	276,416

Note: Values in **bold** refer to statistically significant OR (p-value < 0.05)

Table S5: Females only subgroup regression results for step and trend changes in the odds of being a smoker after full implementation of plain packaging in May 2017 (before/after full implementation) and accounting for plain packaging start date in May 2016 (before/after policy start date) in England (Smoking Toolkit Study data January 2007 to December 2019)

	After full implementation		After policy start date	
	Unadjusted	Adjusted	Unadjusted	Adjusted
	OR (p-value) [95% CI]	OR (p-value) [95% CI]	OR (p-value) [95% CI]	OR (p-value) [95% CI]
Females only subgroup				
Level after May 2016			0.95	<b>0.89</b>
			(0.342)	(0.046)

			0.85 - 1.06	0.79 - 1.00
Slope May 2016-May 2017			1.08	1.08
			(0.438)	(0.430)
			0.89 - 1.30	0.89 - 1.32
Level after May 2017	0.99	0.97	0.96	0.97
	(0.802)	(0.480)	(0.572)	(0.617)
	0.91 - 1.07	0.89 - 1.06	0.84 - 1.10	0.84 - 1.11
Post-slope 18 months after May 2017	0.98	0.97	0.98	0.96
	(0.640)	(0.522)	(0.636)	(0.459)
	0.89 - 1.08	0.87 - 1.07	0.89 - 1.08	0.87 - 1.07
Smoke free ban		1.10		1.09
		(0.076)		(0.088)
		0.99 - 1.22		0.99 - 1.21
Minimum age of sale		<b>0.87</b>		<b>0.86</b>
		(0.005)		(0.005)
		0.78 - 0.96		0.78 - 0.95
Display ban in small shops		0.99		0.98
		(0.856)		(0.640)
		0.92 - 1.07		0.91 - 1.06
Display ban in large shops		1.04		1.08
		(0.215)		(0.053)
		0.97 - 1.12		1.00 - 1.16
Price per cigarette		1.00		1.00
		(0.765)		(0.744)
		1.00 - 1.00		1.00 - 1.00
Lagged smoking prevalence	Yes	Yes	Yes	Yes
Seasonality	Yes	Yes	Yes	Yes
Observations	142,107	142,107	142,107	142,107

Note: Values in **bold** refer to statistically significant OR (p-value < 0.05)

Table S6: Males only subgroup regression results for level and slope changes in the odds of being a smoker after full implementation of plain packaging in May 2017 (before/after full implementation) and accounting for plain packaging start date in May 2016 (before/after policy start date) in England (Smoking Toolkit Study data January 2007 to December 2019)

	After full implementation		After policy start date	
	Unadjusted OR (p-value) [95% CI]	Adjusted OR (p-value) [95% CI]	Unadjusted OR (p-value) [95% CI]	Adjusted OR (p-value) [95% CI]
Males only subgroup				
Level after May 2016			<b>0.89</b>	0.91
			(0.022)	(0.083)
			0.80 - 0.98	0.81 - 1.01
Slope May 2016-May 2017			1.00	0.88
			(0.969)	(0.204)
			0.83 - 1.20	0.73 - 1.07
Level after May 2017	<b>0.92</b>	<b>0.90</b>	1.01	1.03
	(0.046)	(0.013)	(0.932)	(0.706)
	0.85 - 1.00	0.83 - 0.98	0.88 - 1.15	0.90 - 1.18
Post-slope 18 months after May 2017	1.07	1.04	1.05	1.03
	(0.176)	(0.444)	(0.262)	(0.556)

	0.97 - 1.17	0.94 - 1.15	0.96 - 1.16	0.93 - 1.14
Smoke free ban		1.04		1.04
		(0.541)		(0.515)
		0.93 - 1.16		0.93 - 1.16
Minimum age of sale		<b>0.84</b>		<b>0.80</b>
		(0.003)		(0.000)
		0.75 - 0.94		0.72 - 0.90
Display ban in small shops		0.97		0.95
		(0.362)		(0.211)
		0.90 - 1.04		0.88 - 1.03
Display ban in large shops		<b>0.93</b>		1.00
		(0.036)		(0.902)
		0.88 - 1.00		0.93 - 1.07
Price per cigarette		<b>1.00</b>		<b>1.00</b>
		(0.010)		(0.001)
		1.00 - 1.00		1.00 - 1.00
Lagged smoking prevalence	Yes	Yes	Yes	Yes
Seasonality	Yes	Yes	Yes	Yes
Observations	134,254	134,254	134,254	134,254

Note: Values in **bold** refer to statistically significant OR (p-value < 0.05)

Table S7: Population aged 18 to 25 years old only regression results for step and trend changes in the odds of being a smoker after full implementation of plain packaging in May 2017 (before/after full implementation) and accounting for plain packaging start date in May 2016 (before/after policy start date) in England (Smoking Toolkit Study data January 2007 to December 2019)

	After full implementation		After policy start date	
	Unadjusted OR (p-value) [95% CI]	Adjusted OR (p-value) [95% CI]	Unadjusted OR (p-value) [95% CI]	Adjusted OR (p-value) [95% CI]
Population aged 18-25 years old only				
Level after May 2016			1.02	0.95
			(0.842)	(0.541)
			0.86 - 1.21	0.79 - 1.13
Slope May 2016-May 2017			0.87	0.84
			(0.377)	(0.277)
			0.64 - 1.18	0.61 - 1.15
Level after May 2017	1.12	1.10	1.25	<b>1.29</b>
	(0.078)	(0.183)	(0.052)	(0.028)
	0.99 - 1.28	0.96 - 1.27	1.00 - 1.57	1.03 - 1.62
Post-slope 18 months after May 2017	0.87	0.87	0.87	0.86
	(0.088)	(0.088)	(0.074)	(0.068)
	0.75 - 1.02	0.73 - 1.02	0.74 - 1.01	0.72 - 1.01
Smoke free ban		1.09		1.09
		(0.347)		(0.366)
		0.91 - 1.31		0.91 - 1.30
Minimum age of sale		0.85		<b>0.82</b>
		(0.070)		(0.031)
		0.71 - 1.01		0.69 - 0.98
Display ban in small shops		1.02		1.01
		(0.725)		(0.818)



		0.90 - 1.16		0.90 - 1.15
Display ban in large shops		1.07		<b>1.13</b>
		(0.211)		(0.033)
		0.96 - 1.19		1.01 - 1.28
Price per cigarette		1.00		1.00
		(0.936)		(0.705)
		1.00 - 1.00		1.00 - 1.00
Lagged smoking prevalence	Yes	Yes	Yes	Yes
Seasonality	Yes	Yes	Yes	Yes
Observations	43,729	43,729	43,729	43,729

Note: Values in **bold** refer to statistically significant OR (p-value < 0.05)

Table S9: Population over 25 years old only regression results for step and trend changes in the odds of being a smoker after full implementation of plain packaging in May 2017 (before/after full implementation) and accounting for plain packaging start date in May 2016 (before/after policy start date) in England (Smoking Toolkit Study data January 2007 to December 2019)

	After full implementation		After policy start date	
	Unadjusted OR (p-value) [95% CI]	Adjusted OR (p-value) [95% CI]	Unadjusted OR (p-value) [95% CI]	Adjusted OR (p-value) [95% CI]
Population over 25 years old only				
Level after May 2016			<b>0.90</b>	<b>0.90</b>
			(0.014)	(0.016)
			0.83 - 0.98	0.82 - 0.98
Slope May 2016-May 2017			1.07	1.00
			(0.386)	(0.976)
			0.92 - 1.23	0.86 - 1.17
Level after May 2017	<b>0.92</b>	<b>0.90</b>	0.94	0.94
	(0.013)	(0.002)	(0.228)	(0.283)
	0.87 - 0.98	0.84 - 0.96	0.84 - 1.04	0.85 - 1.05
Post-slope 18 months after May 2017	1.06	1.04	1.06	1.04
	(0.109)	(0.304)	(0.147)	(0.382)
	0.99 - 1.15	0.96 - 1.13	0.98 - 1.14	0.96 - 1.12
Smoke free ban		1.07		1.06
		(0.139)		(0.151)
		0.98 - 1.16		0.98 - 1.16
Minimum age of sale		<b>0.85</b>		<b>0.83</b>
		(0.000)		(0.000)
		0.78 - 0.92		0.77 - 0.91
Display ban in small shops		0.97		0.96
		(0.296)		(0.153)
		0.91 - 1.03		0.90 - 1.02
Display ban in large shops		0.97		1.01
		(0.242)		(0.659)
		0.92 - 1.02		0.96 - 1.07
Price per cigarette		1.00		<b>1.00</b>
		(0.060)		(0.035)
		1.00 - 1.00		1.00 - 1.00

Lagged smoking prevalence	Yes	Yes	Yes	Yes
Seasonality	Yes	Yes	Yes	Yes
Observations	232,687	232,687	232,687	232,687

Note: Values in **bold** refer to statistically significant OR (p-value < 0.05)

Table S10: Manual and routine occupations only regression results for step and trend changes in the odds of being a smoker after full implementation of plain packaging in May 2017 (before/after full implementation) and accounting for plain packaging start date in May 2016 (before/after policy start date) in England (Smoking Toolkit Study data January 2007 to December 2019)

	After full implementation		After policy start date	
	Unadjusted	Adjusted	Unadjusted	Adjusted
	OR	OR	OR	OR
	(p-value)	(p-value)	(p-value)	(p-value)
	[95% CI]	[95% CI]	[95% CI]	[95% CI]
Manual and routine occupations only				
Level after May 2016			0.89	<b>0.83</b>
			(0.065)	(0.004)
			0.79 - 1.01	0.73 - 0.94
Slope			1.08	1.11
May 2016-May 2017			(0.499)	(0.361)
			0.87 - 1.33	0.89 - 1.38
Level after May 2017	1.07	1.07	1.08	1.09
	(0.149)	(0.171)	(0.315)	(0.289)
	0.98 - 1.17	0.97 - 1.18	0.93 - 1.27	0.93 - 1.27
Post-slope 18 months	1.02	1.00	1.01	1.00
after May 2017	(0.721)	(0.939)	(0.798)	(0.948)
	0.91 - 1.14	0.89 - 1.13	0.91 - 1.13	0.89 - 1.12
Smoke free ban		1.09		1.09
		(0.149)		(0.173)
		0.97 - 1.23		0.96 - 1.22
Minimum age of sale		<b>0.89</b>		<b>0.88</b>
		(0.045)		(0.027)
		0.79 - 1.00		0.78 - 0.99
Display ban in small		0.99		0.97
shops		(0.729)		(0.423)
		0.91 - 1.07		0.89 - 1.05
Display ban in large		1.00		1.06
shops		(0.945)		(0.179)
		0.93 - 1.08		0.98 - 1.14
Price per cigarette		1.00		1.00
		(0.215)		(0.234)
		1.00 - 1.00		1.00 - 1.00
Lagged smoking prevalence	Yes	Yes	Yes	Yes
Seasonality	Yes	Yes	Yes	Yes
Observations	95,770	95,770	95,770	95,770

Note: Values in **bold** refer to statistically significant OR (p-value < 0.05)

Table S11: Non-routine occupations only regression results for step and trend changes in the odds of being a smoker after full implementation of plain packaging in May 2017 (before/after full implementation) and accounting for plain packaging start date in May 2016 (before/after policy start date) in England (Smoking Toolkit Study data January 2007 to December 2019)

	After full implementation		After policy start date	
	Unadjusted	Adjusted	Unadjusted	Adjusted
	OR (p-value) [95% CI]	OR (p-value) [95% CI]	OR (p-value) [95% CI]	OR (p-value) [95% CI]
Non-routine occupations only				
Level after May 2016			1.03 (0.578) 0.92 - 1.16	1.00 (0.961) 0.89 - 1.13
Slope May 2016-May 2017			0.99 (0.942) 0.82 - 1.21	0.93 (0.457) 0.75 - 1.13
Level after May 2017	0.98 (0.669) 0.90 - 1.07	0.94 (0.149) 0.85 - 1.02	0.97 (0.666) 0.84 - 1.12	0.99 (0.846) 0.85 - 1.14
Post-slope 18 months after May 2017	1.00 (0.999) 0.91 - 1.10	0.96 (0.453) 0.86 - 1.07	1.00 (0.973) 0.91 - 1.11	0.96 (0.435) 0.86 - 1.07
Smoke free ban		1.04 (0.573) 0.91 - 1.17		1.04 (0.567) 0.92 - 1.18
Minimum age of sale		<b>0.79</b> (0.000) 0.70 - 0.90		<b>0.78</b> (0.000) 0.69 - 0.89
Display ban in small shops		0.94 (0.206) 0.86 - 1.03		0.94 (0.203) 0.86 - 1.03
Display ban in large shops		1.02 (0.544) 0.95 - 1.11		1.04 (0.344) 0.96 - 1.13
Price per cigarette		1.00 (0.311) 1.00 - 1.00		1.00 (0.211) 1.00 - 1.00
Lagged smoking prevalence	Yes	Yes	Yes	Yes
Seasonality	Yes	Yes	Yes	Yes
Observations	141,844	141,844	141,844	141,844

Note: Values in **bold** refer to statistically significant OR (p-value < 0.05)