



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/179893/>

Version: Accepted Version

Conference or Workshop Item:

Tang, G, Tan, P, Zhang, Q et al. (2021) Physical activity, transportation mode and sedentary time on the development of overweight and obesity in school-age children in China: China Health and Nutrition Survey (CHNS) 2010–2012. In: Nutrition Society Irish Section Conference 2021, 22-24 Jun 2021.

<https://doi.org/10.1017/s0029665121002688>

© The Authors 2021. This is an author produced version of a conference abstract published in Proceedings of the Nutrition Society. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Physical activity, transportation mode and sedentary time on the development of overweight and obesity in school-age children in China: China Health and Nutrition Survey (CHNS) 2010–2012

Published online by Cambridge University Press: 17 August 2021

GuYuan Tang¹, PuiYee Tan¹, Qian Zhang², JunSheng Huo² and YunYun Gong¹

¹ School of Food Science and Nutrition, Faculty of Environment, University of Leeds, Leeds LS2 9JT, United Kingdom and

² National Institute for Nutrition and Health (NINH), Chinese Center for Disease Control and Prevention (China CDC), Beijing

Worldwide prevalence of childhood obesity has increased substantially in the past decades and was associated with significant impact on child's physical and social development. High energy-dense foods and physically inactive are the major lifestyle risk factors for childhood obesity(1). Therefore, this study aimed to determine the prevalence of overweight and obesity among the Chinese schoolaged children, and the relationships between childhood obesity and lifestyle factors, particularly on physical activity and dietary intake. Data for this study was obtained from Chinese Health and Nutrition Survey (CHNS) 2010–2012(2). Social-demographic and physical activity data were collected using structured questionnaires. Dietary intake was assessed using a 100-item FFQ. Overweight and obesity were defined as BMI at 85th–95th and >95th percentiles, respectively, specified for age and gender(3). 24,722 children aged 6–17 years old from 31 provinces were recruited using a multistage random-cluster sampling process². Of those, 2,617 participants who had complete data on the aforementioned parameters were included in this study. Statistics: Multivariate logistic regression analysis was performed to determine the associations between lifestyle factors and the odds of overweight and obesity. Mann-Whitney test was performed to assess the difference in average daily food consumption between the low and high physical activity groups. 16.3% of our study participants were found to be overweight, and of those, 7.3% were obese. Children using motorized transportation method (including car, bus, taxi or other public transport) to school and who spent ≥ 3 h/day of sedentary activity were associated with increased odds of overweight and obesity (OR = 1.296, CI = 1.030–1.631; $p = 0.027$ and OR = 1.485, CI = 1.020–2.164; $p = 0.039$, respectively) compared to non-motorized transportation and < 0.001 compared to < 0.05). Moreover, ≥ 41.5 g/day of egg, and ≥ 113.3 g/day of dairy product consumption were associated with increased odds of overweight and obesity (OR = 1.645, CI = 1.119–2.417; $p = 0.011$ and OR = 1.432, CI = 1.000–2.053; $p = 0.050$, respectively), compared to lower consumption. However, such associations did not reach statistical significance after adjusting for covariates ($p > 0.05$). No other dietary factor alone was found significantly associated with overweight and obesity. Motorised transportation to school and sedentary time were significantly associated with increased risk of childhood overweight and obesity. Similar finding was found in physical activity; however, this may be due to the increase in total energy intake associated with higher physical activity level. Dietary factor alone was not found to be significantly associated with overweight and obesity. Acknowledgements The authors are grateful to Professor Junsheng Huo and Professor Jing Sun from Chinese Centre for Disease Control and Prevention's (CDC) National institute of Health and Nutrition (NINH) for providing the CHNS 2010–2012 data. The authors acknowledge the funding support from the Biotechnology and Biological Sciences Research Council (BBSRC) (Grant number: BB/T008989/1) through Global Challenge Research Fund (GCRF) 2020 granted to the University of Leeds project titled "Addressing micronutrient deficiencies associated with the double burden of childhood malnutrition in China, a combined food system framework".

References

1. Mistry SK & Puthussery S (2015) *Public Health* 129, 200–209
2. Popkin BM, Du S, Zhai F et al. (2010) *Int J Epidemiol* 39, 1435–1440
3. Group of China Obesity Task Force (2004) *Zhonghua Liu Xing Bing Xue Za Zhi* 25, 97–02